

code of federal regulations

Food and Drugs

21

PARTS 500 TO 599

Revised as of April 1, 1999

CONTAINING
A CODIFICATION OF DOCUMENTS
OF GENERAL APPLICABILITY
AND FUTURE EFFECT

AS OF APRIL 1, 1999

With Ancillaries

Published by
the Office of the Federal Register
National Archives and Records
Administration

as a Special Edition of
the Federal Register



U.S. GOVERNMENT PRINTING OFFICE
WASHINGTON : 1999

For sale by U.S. Government Printing Office
Superintendent of Documents, Mail Stop: SSOP, Washington, DC 20402-9328

Table of Contents

Explanation	<i>Page</i> v
Title 21:	
Chapter I—Food and Drug Administration, Department of Health and Human Services (Continued)	3
Finding Aids:	
Material Approved for Incorporation by Reference	519
Table of CFR Titles and Chapters	521
Alphabetical List of Agencies Appearing in the CFR	539
Redesignation Table	549
List of CFR Sections Affected	551

Cite this Code: CFR

*To cite the regulations in
this volume use title,
part and section num-
ber. Thus, 21 CFR
500.23 refers to title 21,
part 500, section 23.*

Explanation

The Code of Federal Regulations is a codification of the general and permanent rules published in the Federal Register by the Executive departments and agencies of the Federal Government. The Code is divided into 50 titles which represent broad areas subject to Federal regulation. Each title is divided into chapters which usually bear the name of the issuing agency. Each chapter is further subdivided into parts covering specific regulatory areas.

Each volume of the Code is revised at least once each calendar year and issued on a quarterly basis approximately as follows:

Title 1 through Title 16.....	as of January 1
Title 17 through Title 27.....	as of April 1
Title 28 through Title 41.....	as of July 1
Title 42 through Title 50.....	as of October 1

The appropriate revision date is printed on the cover of each volume.

LEGAL STATUS

The contents of the Federal Register are required to be judicially noticed (44 U.S.C. 1507). The Code of Federal Regulations is prima facie evidence of the text of the original documents (44 U.S.C. 1510).

HOW TO USE THE CODE OF FEDERAL REGULATIONS

The Code of Federal Regulations is kept up to date by the individual issues of the Federal Register. These two publications must be used together to determine the latest version of any given rule.

To determine whether a Code volume has been amended since its revision date (in this case, April 1, 1999), consult the “List of CFR Sections Affected (LSA),” which is issued monthly, and the “Cumulative List of Parts Affected,” which appears in the Reader Aids section of the daily Federal Register. These two lists will identify the Federal Register page number of the latest amendment of any given rule.

EFFECTIVE AND EXPIRATION DATES

Each volume of the Code contains amendments published in the Federal Register since the last revision of that volume of the Code. Source citations for the regulations are referred to by volume number and page number of the Federal Register and date of publication. Publication dates and effective dates are usually not the same and care must be exercised by the user in determining the actual effective date. In instances where the effective date is beyond the cut-off date for the Code a note has been inserted to reflect the future effective date. In those instances where a regulation published in the Federal Register states a date certain for expiration, an appropriate note will be inserted following the text.

OMB CONTROL NUMBERS

The Paperwork Reduction Act of 1980 (Pub. L. 96-511) requires Federal agencies to display an OMB control number with their information collection request.

Many agencies have begun publishing numerous OMB control numbers as amendments to existing regulations in the CFR. These OMB numbers are placed as close as possible to the applicable recordkeeping or reporting requirements.

OBSOLETE PROVISIONS

Provisions that become obsolete before the revision date stated on the cover of each volume are not carried. Code users may find the text of provisions in effect on a given date in the past by using the appropriate numerical list of sections affected. For the period before January 1, 1986, consult either the List of CFR Sections Affected, 1949–1963, 1964–1972, or 1973–1985, published in seven separate volumes. For the period beginning January 1, 1986, a “List of CFR Sections Affected” is published at the end of each CFR volume.

INCORPORATION BY REFERENCE

What is incorporation by reference? Incorporation by reference was established by statute and allows Federal agencies to meet the requirement to publish regulations in the Federal Register by referring to materials already published elsewhere. For an incorporation to be valid, the Director of the Federal Register must approve it. The legal effect of incorporation by reference is that the material is treated as if it were published in full in the Federal Register (5 U.S.C. 552(a)). This material, like any other properly issued regulation, has the force of law.

What is a proper incorporation by reference? The Director of the Federal Register will approve an incorporation by reference only when the requirements of 1 CFR part 51 are met. Some of the elements on which approval is based are:

- (a) The incorporation will substantially reduce the volume of material published in the Federal Register.
- (b) The matter incorporated is in fact available to the extent necessary to afford fairness and uniformity in the administrative process.
- (c) The incorporating document is drafted and submitted for publication in accordance with 1 CFR part 51.

Properly approved incorporations by reference in this volume are listed in the Finding Aids at the end of this volume.

What if the material incorporated by reference cannot be found? If you have any problem locating or obtaining a copy of material listed in the Finding Aids of this volume as an approved incorporation by reference, please contact the agency that issued the regulation containing that incorporation. If, after contacting the agency, you find the material is not available, please notify the Director of the Federal Register, National Archives and Records Administration, Washington DC 20408, or call (202) 523–4534.

CFR INDEXES AND TABULAR GUIDES

A subject index to the Code of Federal Regulations is contained in a separate volume, revised annually as of January 1, entitled CFR INDEX AND FINDING AIDS. This volume contains the Parallel Table of Statutory Authorities and Agency Rules (Table I), and Acts Requiring Publication in the Federal Register (Table II). A list of CFR titles, chapters, and parts and an alphabetical list of agencies publishing in the CFR are also included in this volume.

An index to the text of “Title 3—The President” is carried within that volume.

The Federal Register Index is issued monthly in cumulative form. This index is based on a consolidation of the “Contents” entries in the daily Federal Register.

A List of CFR Sections Affected (LSA) is published monthly, keyed to the revision dates of the 50 CFR titles.

REPUBLICATION OF MATERIAL

There are no restrictions on the republication of material appearing in the Code of Federal Regulations.

INQUIRIES

For a legal interpretation or explanation of any regulation in this volume, contact the issuing agency. The issuing agency's name appears at the top of odd-numbered pages.

For inquiries concerning CFR reference assistance, call 202-523-5227 or write to the Director, Office of the Federal Register, National Archives and Records Administration, Washington, DC 20408.

SALES

The Government Printing Office (GPO) processes all sales and distribution of the CFR. For payment by credit card, call 202-512-1800, M-F, 8 a.m. to 4 p.m. e.s.t. or fax your order to 202-512-2233, 24 hours a day. For payment by check, write to the Superintendent of Documents, Attn: New Orders, P.O. Box 371954, Pittsburgh, PA 15250-7954. For GPO Customer Service call 202-512-1803.

ELECTRONIC SERVICES

The full text of the Code of Federal Regulations, the LSA (List of CFR Sections Affected), The United States Government Manual, the Federal Register, Public Laws, Weekly Compilation of Presidential Documents and the Privacy Act Compilation are available in electronic format at www.access.gpo.gov/nara ("GPO Access"). For more information, contact Electronic Information Dissemination Services, U.S. Government Printing Office. Phone 202-512-1530, or 888-293-6498 (toll-free). E-mail, gpoaccess@gpo.gov.

The Office of the Federal Register also offers a free service on the National Archives and Records Administration's (NARA) World Wide Web site for public law numbers, Federal Register finding aids, and related information. Connect to NARA's web site at www.nara.gov/fedreg. The NARA site also contains links to GPO Access.

RAYMOND A. MOSLEY,
Director,
Office of the Federal Register.

April 1, 1999.

THIS TITLE

Title 21—FOOD AND DRUGS is composed of nine volumes. The parts in these volumes are arranged in the following order: Parts 1–99, 100–169, 170–199, 200–299, 300–499, 500–599, 600–799, 800–1299 and 1300–end. The first eight volumes, containing parts 1–1299, comprise Chapter I—Food and Drug Administration, Department of Health and Human Services. The ninth volume, containing part 1300 to end, includes Chapter II—Drug Enforcement Administration, Department of Justice, and Chapter III—Office of National Drug Control Policy. The contents of these volumes represent all current regulations codified under this title of the CFR as of April 1, 1999.

Redesignation tables for Chapter I—Food and Drug Administration appear in the Finding Aids section for the volumes containing parts 170–199 and 500–599.

For this volume, Melanie L. Marcec was Chief Editor. The Code of Federal Regulations publication program is under the direction of Frances D. McDonald, assisted by Alomha S. Morris.

if any changes have been made to the *Code of Federal Regulations* or what documents have been published in the *Federal Register* without reading the *Federal Register* every day? If so, you may wish to subscribe to the *LSA* (List of CFR Sections Affected), the *Federal Register Index*, or both.

The LSA (List of CFR Sections Affected) is designed to lead users of the *Code of Federal Regulations* to amendatory actions published in the *Federal Register*. The LSA is issued monthly in cumulative form. Entries indicate the nature of the changes—such as revised, removed, or corrected. **\$27 per year.**

The index, covering the contents of the daily *Federal Register*, is issued monthly in cumulative form. Entries are carried primarily under the names of the issuing agencies. Significant subjects are carried as cross-references.

\$25 per year.

A finding aid is included in each publication which lists *Federal Register* page numbers with the date of publication in the *Federal Register*.



Order Processing Code: ***5421** ☐ **YES**, send me the following indicated subscriptions for one year:
 _____ **LSA (List of CFR Sections Affected)**, (LCS) for \$27 per year.
 _____ **Federal Register Index (FRSU)** \$25 per year.

Purchase order No. (optional)

Thank you for your order!

Mail To: Superintendent of Documents
PO Box 371954
Pittsburgh PA 15250-7954



Fax your orders (202) 512-2250
Phone your orders (202) 512-1800

Title 21—Food and Drugs

(This book contains parts 500 to 599)

Part

CHAPTER 1—Food and Drug Administration, Department of Health and Human Services (Continued)	500

CROSS REFERENCES: Animal and Plant Health Inspection Service, Department of Agriculture:
See Animals and Animal Products, 9 CFR chapter I.
Food Safety and Inspection Service, Meat and Poultry Inspection, Department of Agriculture: See Animals and Animal Products, 9 CFR chapter III.
Federal Trade Commission: See Commercial Practices, 16 CFR chapter I.
U.S. Customs Service, Department of the Treasury: See Customs Duties, 19 CFR chapter I.
Internal Revenue Service, Department of the Treasury: See Internal Revenue, 26 CFR chapter I.
Bureau of Alcohol, Tobacco, and Firearms, Department of the Treasury: See Alcohol, Tobacco Products and Firearms, 27 CFR chapter I.

CHAPTER I—FOOD AND DRUG ADMINISTRATION, DEPARTMENT OF HEALTH AND HUMAN SERVICES—(Continued)

SUBCHAPTER E—ANIMAL DRUGS, FEEDS, AND RELATED PRODUCTS

<i>Part</i>		<i>Page</i>
500	General	5
501	Animal food labeling	14
502	Common or usual names for nonstandardized ani- mal foods	31
509	Unavoidable contaminants in animal food and food-packaging material	33
510	New animal drugs	37
511	New animal drugs for investigational use	54
514	New animal drug applications	58
520	Oral dosage form new animal drugs	84
522	Implantation or injectable dosage form new ani- mal drugs	205
524	Ophthalmic and topical dosage form new animal drugs	284
526	Intramammary dosage forms	312
529	Certain other dosage form new animal drugs	318
530	Extralabel drug use in animals	323
556	Tolerances for residues of new animal drugs in food	329
558	New animal drugs for use in animal feeds	348
564	[Reserved]	
570	Food additives	457
571	Food additive petitions	464
573	Food additives permitted in feed and drinking water of animals	469
579	Irradiation in the production, processing, and han- dling of animal feed and pet food	486
582	Substances generally recognized as safe	487
584	Food substances affirmed as generally recognized as safe in feed and drinking water of animals	511

21 CFR Ch. I (4-1-99 Edition)

<i>Part</i>		<i>Page</i>
589	Substances prohibited from use in animal food or feed	512
590-599	[Reserved]	

EDITORIAL NOTE: For nomenclature changes to Chapter I, see 59 FR 14366, Mar. 28, 1994.

SUBCHAPTER E—ANIMAL DRUGS, FEEDS, AND RELATED PRODUCTS

PART 500—GENERAL

Subpart A [Reserved]

Subpart B—Specific Administrative Rulings and Decisions

Sec.

- 500.23 Thermally processed low-acid foods packaged in hermetically sealed containers.
- 500.24 Emergency permit control.
- 500.25 Anthelmintic drugs for use in animals.
- 500.26 Timed-release dosage form drugs.
- 500.27 Methylene blue-containing drugs for use in animals.
- 500.29 Gentian violet for use in animal feed.
- 500.30 Gentian violet for animal drug use.
- 500.35 Animal feeds contaminated with *Salmonella* microorganisms.
- 500.45 Use of polychlorinated biphenyls (PCB's) in the production, handling, and storage of animal feed.
- 500.46 Hexachlorophene in animal drugs.
- 500.50 Propylene glycol in or on cat food.

Subpart C—Animal Drug Labeling Requirements

- 500.51 Labeling of animal drugs; misbranding.
- 500.52 Use of terms such as “tonic”, “tone”, “toner”, or “conditioner” in the labeling of preparations intended for use in or on animals.
- 500.55 Exemption from certain drug-labeling requirements.

Subpart D—Requirements for Specific Animal Drugs

- 500.65 Epinephrine injection 1:1,000 in 10-milliliter containers for emergency treatment of anaphylactoid shock in cattle, horses, sheep, and swine.

Subpart E—Regulation of Carcinogenic Compounds Used in Food-Producing Animals

- 500.80 Scope of this subpart.
- 500.82 Definitions.
- 500.84 Operational definition of “no residue”.
- 500.86 Marker residue and target tissue.
- 500.88 Regulatory method.
- 500.90 Waiver of requirements.
- 500.92 Implementation.

AUTHORITY: 21 U.S.C. 321, 331, 342, 343, 348, 351, 352, 353, 360b, 371.

SOURCE: 40 FR 13802, Mar. 27, 1975, unless otherwise noted.

Subpart A [Reserved]

Subpart B—Specific Administrative Rulings and Decisions

§ 500.23 Thermally processed low-acid foods packaged in hermetically sealed containers.

The provisions of part 113 of this chapter shall apply to the manufacture, processing or packing of low-acid foods in hermetically sealed containers, and intended for use as food for animals.

[61 FR 37681, July 19, 1996]

§ 500.24 Emergency permit control.

The provisions of part 108 of this chapter shall apply to the issuance of emergency control permits for the manufacturer or packer of thermally processed low-acid foods packaged in hermetically sealed containers, and intended for use as food for animals.

[61 FR 37681, July 19, 1996]

§ 500.25 Anthelmintic drugs for use in animals.

(a) The Commissioner of Food and Drugs has determined that, in order to assure that anthelmintic drugs, including animal feeds bearing or containing such drugs, which do not carry the prescription statement are labeled to provide adequate directions for their effective use, labeling of these anthelmintic drugs shall bear, in addition to other required information, a statement that a veterinarian should be consulted for assistance in the diagnosis, treatment, and control of parasitism.

(b) The label and any labeling furnishing or purporting to furnish directions for use, shall bear conspicuously the following statement: “Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.”

§ 500.26

(c) For drugs covered by approved new animal drug applications, the labeling revisions required for compliance with this section may be placed into effect without prior approval as provided for in § 514.8 (d) and (e) of this chapter. For animal feeds bearing or containing anthelmintic drugs covered by approved applications, the labeling revisions required for compliance with this section may be placed into effect without the submission of supplemental applications as provided for in § 514.9 of this chapter.

(d) Labeling revisions required for compliance with this section shall be placed into effect by February 25, 1975, following which, any such drugs that are introduced into interstate commerce and not in compliance with this section will be subject to regulatory proceedings.

§ 500.26 Timed-release dosage form drugs.

(a) Drugs are being offered in dosage forms that are designed to release the active ingredients over a prolonged period of time. There is a possibility of unsafe overdosage or ineffective dosage if such products are improperly made and the active ingredients are released at one time, over too short or too long a period of time, or not released at all. Drugs marketed in this form, which are referred to by such terms as timed-release, controlled-release, prolonged-release, sustained-release, or delayed-release drugs, are regarded as new animal drugs within the meaning of section 201(v) of the Federal Food, Drug, and Cosmetic Act.

(b) Timed-release dosage form animal drugs that are introduced into interstate commerce are deemed to be adulterated within the meaning of section 501(a)(5) of the act and subject to regulatory action unless such animal drug is the subject of an approved new animal drug application as required by paragraph (a) of this section.

(c) The fact that the labeling of this kind of drug may claim delayed, prolonged, controlled, or sustained-release of all or only some of the active ingredients does not affect the new animal drug status of such articles. A new animal drug application is required in any such case.

21 CFR Ch. I (4–1–99 Edition)

(d) New animal drug applications for timed-release dosage form animal drugs must contain, among other things, data to demonstrate safety and effectiveness by establishing that the article is manufactured using procedures and controls to ensure release of the total dosage at a safe and effective rate. Data submitted in the new animal drug application must demonstrate that the formulation of the drug and the procedures used in its manufacture will ensure release of the active ingredient(s) of the drug at a safe and effective rate and that these release characteristics will be maintained until the expiration date of the drug. When the drug is intended for use in food-producing animals, data submitted must also demonstrate that, with respect to possible residues of the drug, food derived from treated animals is safe for consumption.

[42 FR 8635, Feb. 11, 1977, as amended at 60 FR 38480, July 27, 1995]

§ 500.27 Methylene blue-containing drugs for use in animals.

(a) New information requires a re-evaluation of the status of drugs containing methylene blue (tetramethylthionine chloride) for oral use in cats or dogs.

(1)(i) It has been demonstrated that two orally administered urinary antiseptic-antispasmodic preparations that contained methylene blue cause Heinz body hemolytic anemia in cats when used according to label directions. The specific cause of the reaction was determined to be the methylene blue contained in the preparations. The reaction can be severe enough to cause death of treated animals.

(ii) The Heinz body hemolytic anemia reaction to methylene blue has also been demonstrated in dogs under laboratory conditions. The precise mechanism by which methylene blue produces the characteristic erythrocytic inclusion bodies (Heinz bodies) and associated hemolytic anemia is unclear.

(2) The effectiveness of orally administered methylene blue as a urinary antiseptic is open to question. It appears that following oral administration, methylene blue is poorly and erratically absorbed and also slowly and erratically excreted in the urine. Studies

Food and Drug Administration, HHS

§ 500.35

in the dog indicate it is excreted in the urine essentially as leukomethylene blue stabilized in some manner. Methylene blue itself is stepwise demethylated in alkaline solutions (alkaline urine being a frequent consequence of urinary infection) to Azure B, Azure A, and Azure C. The antiseptic efficacy of all of these excretion products is unsubstantiated.

(3) In view of the foregoing, the Commissioner has concluded that animal drugs containing methylene blue for oral use in cats or dogs are neither safe nor generally recognized as effective within the meaning of section 201(v) of the act and are therefore considered new animal drugs. Accordingly, all prior formal and informal opinions expressed by the Food and Drug Administration that such drugs are “not new drugs” or “no longer new drugs” are hereby revoked.

(b) Animal drugs that contain methylene blue for oral use in cats or dogs and not the subject of an approved new animal drug application (NADA) are deemed to be adulterated under the provisions of section 501(a) (5) and/or (6) and/or misbranded under section 502(a) of the act and subject to regulatory action as of April 10, 1978.

(c) Sponsors of animal drugs that contain methylene blue for oral use in cats or dogs and not the subject of an approved new animal drug application (NADA) may submit an application in conformity with §514.1 of this chapter. Such applications will be processed in accordance with section 512 of the act. Submission of an NADA will not constitute grounds for continued marketing of this drug substance until such application is approved.

(d) New animal drug applications required by this regulation pursuant to section 512 of the act shall be submitted to the Food and Drug Administration, Center for Veterinary Medicine, Office of New Animal Drug Evaluation (HFV-100), 7500 Standish Pl., Rockville, MD 20855.

[43 FR 9803, Mar. 10, 1978; 43 FR 12310, Mar. 24, 1978, as amended at 54 FR 18279, Apr. 28, 1989; 57 FR 6475, Feb. 25, 1992; 60 FR 38480, July 27, 1995]

§ 500.29 Gentian violet for use in animal feed.

The Food and Drug Administration has determined that gentian violet is not generally recognized as safe for use in animal feed and is a food additive subject to section 409 of the Federal Food, Drug, and Cosmetic Act (the act), unless it is intended for use as a new animal drug, in which case it is subject to section 512 of the act. The Food and Drug Administration has determined that gentian violet is not prior sanctioned for any use in animal feed.

[56 FR 40506, Aug. 15, 1991]

§ 500.30 Gentian violet for animal drug use.

The Food and Drug Administration (FDA) has determined that gentian violet is not generally recognized as safe and effective for any veterinary drug use in food animals and is a new animal drug subject to section 512 of the Federal Food, Drug, and Cosmetic Act. FDA has determined that gentian violet is not exempted from new animal drug status under the “grandfather” provisions of the Drug Amendments of 1962 (21 U.S.C. 342).

[56 FR 40507, Aug. 15, 1991]

§ 500.35 Animal feeds contaminated with *Salmonella* microorganisms.

(a) Investigations by the Food and Drug Administration, the Centers for Disease Control of the U.S. Public Health Service, the Animal Health Division of the Agricultural Research Service, U.S. Department of Agriculture, and by various State public health agencies have revealed that processed fish meal, poultry meal, meat meal, tankage, and other animal byproducts intended for use in animal feed may be contaminated with *Salmonella* bacteria, an organism pathogenic to man and animals. Contamination of these products may occur through inadequate heat treatment of the product during its processing or through recontamination of the heat-treated product during a time of improper storage or handling subsequent to processing.

(b) Articles used in food for animals are included within the definition of

§ 500.45

21 CFR Ch. I (4–1–99 Edition)

food in section 201(f) of the Federal Food, Drug, and Cosmetic Act. Further, *Salmonella* contamination of such animal feeds having the potentiality for producing infection and disease in animals must be regarded as an adulterant within the meaning of section 402(a) of the act. Therefore, the Food and Drug Administration will regard as adulterated within the meaning of section 402(a) of the act shipments of the following when intended for animal feed and encountered in interstate commerce and found upon examination to be contaminated with *Salmonella* microorganisms: Bone meal, blood meal, crab meal, feather meal, fish meal, fish solubles, meat scraps, poultry meat meal, tankage, or other similar animal byproducts, or blended mixtures of these.

[40 FR 13802, Mar. 27, 1975, as amended at 54 FR 18279, Apr. 28, 1989]

§ 500.45 Use of polychlorinated biphenyls (PCB's) in the production, handling, and storage of animal feed.

(a) Polychlorinated biphenyls (PCB's) represent a class of toxic industrial chemicals manufactured and sold under a variety of trade names, including: Aroclor (United States); Phenoclor (France); Colphen (Germany); and Kanaclor (Japan). PCB's are highly stable, heat resistant, and nonflammable chemicals. Industrial uses of PCB's include, or did include in the past, their use as electrical transformer and capacitor fluids, heat transfer fluids, hydraulic fluids, plasticizers, and in formulations of lubricants, coatings, and inks. Their unique physical and chemical properties and widespread, uncontrolled industrial applications have caused PCB's to be a persistent and ubiquitous contaminant in the environment, causing the contamination of certain foods. In addition, incidents have occurred in which PCB's have directly contaminated animal feeds as a result of industrial accidents (leakage or spillage of PCB fluids from plant equipment). These accidents in turn cause the contamination of food intended for human consumption (meat, milk, and eggs). Investigations by the Food and Drug Administration have revealed that heat exchange fluids for

certain pasteurization equipment used in processing animal feed contain PCB's. Although heat exchange fluids in such equipment are considered to be in *closed systems*, leakage has occurred that resulted in direct contamination of animal feed with PCB's and subsequently resulted in the transfer of PCB's to human food produced by animals consuming the contaminated feed. The use of PCB-containing coatings on the inner walls of silos has resulted in the contamination of silage which has in turn caused PCB residues in the milk of dairy cows consuming the contaminated silage. Since PCB's are toxic chemicals, the PCB contamination of food as a result of these and other incidents represent a hazard to public health. It is therefore necessary to place certain restrictions on the industrial uses of PCB's in the production, handling, and storage of animal feed.

(b) The following special provisions are necessary to preclude accidental PCB contamination of animal feed:

(1) Coatings or paints for use on the contact surfaces of feed storage areas may not contain PCB's or any other harmful or deleterious substances likely to contaminate feed.

(2) New equipment or machinery for handling or processing feed in or around an establishment producing animal feed shall not contain PCB's.

(3) On or before Sept. 4, 1973, the management of establishments producing animal feed shall:

(i) Have the heat exchange fluid used in existing equipment or machinery for handling and processing feed sampled and tested to determine whether it contains PCB's, or verify the absence of PCB's in such formulations by other appropriate means. On or before Sept. 4, 1973, any such fluid formulated with PCB's must to the fullest extent possible commensurate with current good manufacturing practices, be replaced with a heat exchange fluid that does not contain PCB's.

(ii) Eliminate to the fullest extent possible commensurate with current good manufacturing practices from the animal feed producing establishment any PCB-containing lubricants for equipment or machinery used for handling or processing animal feed.

(iii) Eliminate to the fullest extent possible commensurate with current good manufacturing practices from the animal feed producing establishment any other PCB-containing materials, whenever there is a reasonable expectation that such materials could cause animal feed to become contaminated with PCB's either as a result of normal use or as a result of accident, breakage, or other mishap.

(iv) The toxicity and other characteristics of fluids selected as PCB replacements must be adequately determined so that the least potentially hazardous replacement should be used. In making this determination with respect to a given fluid, consideration should be given to (a) its toxicity; (b) the maximum quantity that could be spilled onto a given quantity of food before it would be noticed, taking into account its color and odor; (c) possible signaling devices in the equipment to indicate a loss of fluid, etc.; (d) and its environmental stability and tendency to survive and be concentrated through the food chain. The judgment as to whether a replacement fluid is sufficiently non-hazardous is to be made on an individual installation and operation basis.

(c) For the purpose of this section, the provisions do not apply to electrical transformers and condensers containing PCB's in sealed containers.

(d) For the purpose of this section, the term *animal feed* includes all articles used for food or drink for animals other than man.

§ 500.46 Hexachlorophene in animal drugs.

(a) The Commissioner of Food and Drugs has determined that there are no adequate data to establish that animal drugs containing hexachlorophene are safe and effective for any animal use other than in topical products for use on non-food-producing animals as part of a product preservative system at a level not to exceed 0.1 percent; that there is no information on the potential risk to humans from exposure to hexachlorophene by persons who apply animal products containing the drug at levels higher than 0.1 percent; and that there is likewise no information on human exposure to animals on which these animal drugs have been used and

no information on possible residues of hexachlorophene in edible products of food-producing animals treated with new animal drugs that contain any quantity of hexachlorophene.

(b) Animal drugs containing hexachlorophene for other than preservative use on non-food-producing animals at levels not exceeding 0.1 percent are considered new animal drugs and shall be the subject of new animal drug applications (NADA's).

(c) Any person currently marketing animal drugs that contain hexachlorophene other than as part of a product preservative system for products used on non-food-producing animals at a level not exceeding 0.1 percent shall submit a new animal drug application, supplement an existing application, or reformulate the product by September 29, 1977. Each application or supplemental application shall include adequate data to establish that the animal drug is safe and effective. If the animal drug is currently subject to an approved new animal drug application, each reformulation shall require an approved supplemental application. The interim marketing of these animal drugs may continue until the application has been approved, until it has been determined that the application is not approvable under the provisions of § 514.111 of this chapter, or until an existing approved application has been withdrawn.

(d) After September 29, 1977, animal drugs that contain hexachlorophene other than for preservative use on non-food-producing animals at a level not exceeding 0.1 percent that are introduced into interstate commerce shall be deemed to be adulterated within the meaning of section 501(a)(5) of the act (21 U.S.C. 351(a)(5)) unless such animal drug is the subject of a new animal drug application submitted pursuant to paragraph (c) of this section. Action to withdraw approval of new animal drug applications will be initiated if supplemental new animal drug applications have not been submitted in accordance with this section.

(e) New animal drug applications submitted for animal drugs containing hexachlorophene for use in or on food-

§ 500.50

producing animals shall include adequate data to assure that edible products from treated animals are safe for human consumption under the labeled conditions of use.

[42 FR 33725, July 1, 1977; 42 FR 37975, July 26, 1977]

§ 500.50 Propylene glycol in or on cat food.

The Food and Drug Administration has determined that propylene glycol in or on cat food is not generally recognized as safe and is a food additive subject to section 409 of the Federal Food, Drug, and Cosmetic Act (the act). The Food and Drug Administration also has determined that this use of propylene glycol is not prior sanctioned.

[61 FR 19544, May 2, 1996]

Subpart C—Animal Drug Labeling Requirements

§ 500.51 Labeling of animal drugs; misbranding.

(a) Among the representations on the label or labeling of an animal drug which will render the drug misbranded are any broad statements suggesting or implying that the drug is not safe and effective for use when used in accordance with labeling direction, or suggesting or implying that the labeling does not contain adequate warnings or adequate directions for use. Such statements include, but are not limited to:

(1) Any statement that disclaims liability when the drug is used in accordance with directions for use contained on the label or labeling.

(2) Any statement that disclaims liability when the drug is used under “abnormal” or “unforeseeable” conditions.

(3) Any statement limiting the warranty for the products to a warranty that the drug in the package contains the ingredients listed on the label.

(b) This regulation is not intended to prohibit any liability disclaimer that purports to limit the amount of damages or that sets forth the legal theory under which damages are to be recovered.

(c) Any person wishing to obtain an evaluation of an animal drug liability disclaimer under this regulation may

21 CFR Ch. I (4–1–99 Edition)

submit it to Division of Compliance, (HFV–230), Center for Veterinary Medicine, Food and Drug Administration, 7500 Standish Pl., Rockville, MD 20855. A supplemental NADA providing appropriately revised labeling shall be submitted for any approved new animal drug the labeling of which is not in compliance with this regulation.

[41 FR 8473, Feb. 27, 1976, as amended at 54 FR 18279, Apr. 28, 1989; 57 FR 6475, Feb. 25, 1992]

§ 500.52 Use of terms such as “tonic”, “tone”, “toner”, or “conditioner” in the labeling of preparations intended for use in or on animals.

(a) The use of terms such as *tonic*, *tone*, *toner*, and similar terms in the labeling of a product intended for use in or on animals implies that such product is capable of a therapeutic effect(s) and causes such a product to be a drug within the meaning of section 201(g) of the Federal Food, Drug, and Cosmetic Act. The unqualified use of such terms in a product’s labeling fails to provide adequate directions and indications for use of such product and causes it to be misbranded within the meaning of section 502(a) and (f)(1) of the act. The terms *tonic*, *tone*, *toner*, and similar terms may be used in labeling only when appropriately qualified so as to fully inform the user regarding the intended use(s) of the product.

(b) The unqualified use of the term *conditioner* and similar terms in the labeling of a product intended for use in or on animals implies that such product is capable of a therapeutic effect(s) and causes such a product to be a drug within the meaning of section 201(g) of the act. The unqualified use of such terms in a product’s labeling fails to provide adequate directions and indications for use of such product and causes it to be misbranded within the meaning of section 502(a) and (f)(1) of the act. The term *conditioner* and similar terms may be used in labeling only when appropriately qualified so as to fully inform the user regarding the intended use(s) of the product. A product labeled as a “conditioner” or with a similar term can be either a food or drug depending upon the manner in

Food and Drug Administration, HHS

§ 500.65

which the term is qualified in the labeling to reflect the product's intended use.

(c) An article so qualified as to be represented as a drug must be the subject of an approved new animal drug application unless the use of the article under the conditions set forth in its labeling is generally recognized as safe and effective among experts qualified by scientific training and experience to evaluate the safety and effectiveness of animal drugs.

§ 500.55 Exemption from certain drug-labeling requirements.

(a) Section 201.105(c) of this chapter provides that in the case of certain drugs for which directions, hazards, warnings, and use information are commonly known to practitioners licensed by law, such information may be omitted from the dispensing package. Under this proviso, the Commissioner of Food and Drugs will offer an opinion, upon written request, stating reasonable grounds therefore on a proposal to omit such information from the dispensing package.

(b) The Commissioner of Food and Drugs has considered submitted material covering a number of drug products and has offered the opinion that the following drugs when intended for those veterinary uses for which they are now generally employed by the veterinary medical profession, should be exempt from the requirements of § 201.105(c) of this chapter, provided that they meet the conditions prescribed in this paragraph. Preparations that are not in dosage unit form (for example, solutions) will be regarded as meeting the conditions with respect to the maximum quantity of drug per dosage unit if they are prepared in a manner that enables accurate and ready administration of a quantity of drug not in excess of the stated maximum per dosage unit:

Atropine sulfate. As an injectable for cattle, goats, horses, pigs, and sheep, not in excess of 15 milligrams per dosage unit; as an injectable for cats and dogs, not in excess of 0.6 milligram per dosage unit.

Barbital sodium. For oral use in cats and dogs, not in excess of 300 milligrams per dosage unit.

Epinephrine injection. 1:1,000. For cats, dogs, cattle, goats, horses, pigs, and sheep (except as provided in § 500.65).

Morphine sulfate. As an injectable for dogs, not in excess of 15 milligrams per dosage unit.

Pentobarbital sodium. For oral use in cats and dogs, not in excess of 100 milligrams per dosage unit.

Phenobarbital sodium. For oral use in cats and dogs, not in excess of 100 milligrams per dosage unit.

Procaine hydrochloride injection. Containing not in excess of 2 percent procaine hydrochloride, with or without epinephrine up to a concentration of 1:50,000. For use in cats, dogs, cattle, goats, horses, pigs, and sheep.

Thyroid. For oral use in dogs, not in excess of 60 milligrams per dosage unit.

Subpart D—Requirements for Specific Animal Drugs

§ 500.65 Epinephrine injection 1:1,000 in 10-milliliter containers for emergency treatment of anaphylactoid shock in cattle, horses, sheep, and swine.

(a) Anaphylactoid reactions in cattle, horses, sheep, and swine occur occasionally from the injection of antibiotics, bacterins, and vaccines. Adequate directions for use of these antibiotics, bacterins, and vaccines can generally be written for use by the laity and thus are available to livestock producers. Epinephrine injection is effective for the treatment of anaphylactoid reactions in animals and would be of value in saving lives of animals if it were readily available at the time of administration of the causative agents. In connection with this problem the Food and Drug Administration has obtained the views of the Advisory Committee on Veterinary Medicine, and other experts, and has concluded that adequate directions for over-the-counter sale of epinephrine injection 1:1,000 can be prepared.

(b) In view of the above, the Commissioner of Food and Drugs has concluded that it is in the public interest to make epinephrine injection 1:1,000 available for sale without a prescription provided that it is packaged in vials not exceeding 10 milliliters and its label bears, in addition to other required information, the following statements in a prominent and conspicuous manner: "For emergency use only in treating

§ 500.80

anaphylactoid shock. Usual Dosage: Cattle, horses, sheep, and swine—1 cubic centimeter per 100 pounds of body weight. Inject subcutaneously”.

(c) The labeling must also bear a description of the symptoms of anaphylactoid shock including glassy eyes, increased salivation, grinding of the teeth, rapid breathing, muscular tremors, staggering gait, and collapse with death following. These symptoms may appear shortly after injection of a bacterin, vaccine, or antibiotic.

Subpart E—Regulation of Carcinogenic Compounds Used in Food-Producing Animals

SOURCE: 52 FR 49586, Dec. 31, 1987, unless otherwise noted.

§ 500.80 Scope of this subpart.

(a) The Federal Food, Drug, and Cosmetic Act requires that sponsored compounds intended for use in food-producing animals be shown to be safe and that food produced from animals exposed to these compounds be shown to be safe for consumption by people. The statute prohibits the use in food-producing animals of any compound found to induce cancer when ingested by people or animals unless it can be determined by methods of examination prescribed or approved by the Secretary (a function delegated to the Commissioner of Food and Drugs under § 5.10 of this chapter) that no residue of that compound will be found in the food produced from those animals under conditions of use reasonably certain to be followed in practice. This subpart provides an operational definition of no residue and identifies the steps a sponsor of a compound shall follow to secure the approval of the compound. FDA guidelines contain the procedures and protocols FDA recommends for the implementation of this subpart. These guidelines are available from the Dockets Management Branch (HFA-305), Food and Drug Administration, rm. 1-23, 12420 Parklawn Dr., Rockville, MD 20857. Requests for these guidelines should be identified with Docket No. 83D-0288.

(b) If FDA concludes on the basis of the threshold assessment that a sponsor shall conduct carcinogenicity test-

21 CFR Ch. I (4-1-99 Edition)

ing on the sponsored compound, FDA will also determine whether and to what extent the sponsor shall conduct carcinogenicity testing on metabolites of the sponsored compound. The bioassays that a sponsor conducts must be designed to assess carcinogenicity and to determine the quantitative aspects of any carcinogenic response.

(c) If FDA concludes on the basis of the threshold assessment or at a later time during the approval process that the data show that the sponsored compound and its metabolites should not be subject to this subpart, FDA will continue to consider the compound for approval under the general safety provisions of the act for risks other than cancer.

(d) This subpart does not apply to essential nutrients.

[52 FR 49586, Dec. 31, 1987, as amended at 59 FR 14365, Mar. 28, 1994; 62 FR 66983, Dec. 23, 1997]

§ 500.82 Definitions.

(a) The definitions and interpretations contained in section 201 of the act apply to those terms when used in this subpart.

(b) The following definitions apply to this subpart:

Act means the Federal Food, Drug, and Cosmetic Act (sections 201-901, 52 Stat. 1040 *et seq.* as amended (21 U.S.C. 301-392)).

Essential nutrients means compounds that are found in the tissues of untreated, healthy target animals and not produced in sufficient quantity to support the animal's growth, development, function, or reproduction, e.g., vitamins, *essential* minerals, *essential* amino acids, and *essential* fatty acids. These compounds must be supplied from external sources.

FDA means the Food and Drug Administration.

Marker residue means the residue selected for assay whose concentration is in a known relationship to the concentration of the residue of carcinogenic concern in the last tissue to deplete to its permitted concentration.

Preslaughter withdrawal period or *milk discard time* means the time after cessation of administration of the sponsored compound for the residue of carcinogenic concern in the edible product

to deplete to the concentration that will satisfy the operational definition of no residue.

Regulatory method means the aggregate of all experimental procedures for measuring and confirming the presence of the marker residue of the sponsored compound in the target tissue of the target animal.

R_m means the concentration of the marker residue in the target tissue when the residue of carcinogenic concern is equal to S_m in the last tissue to deplete to its permitted concentration.

Residue means any compound present in edible tissues of the target animal which results from the use of the sponsored compound, including the sponsored compound, its metabolites, and any other substances formed in or on food because of the sponsored compound's use.

Residue of carcinogenic concern means all compounds in the total residue of a demonstrated carcinogen excluding any compounds judged by FDA not to present a carcinogenic risk.

S_m means the permitted concentration of residue of carcinogenic concern for a specific edible tissue.

S_o means the concentration of the test compound in the total diet of test animals that corresponds to a maximum lifetime risk of cancer in the test animals of 1 in 1 million. For the purpose of this subpart, FDA will also assume that this S_o will correspond to the concentration of residue of carcinogenic concern in the total human diet that represents no significant increase in the risk of cancer to people.

Sponsor means the person or organization proposing or holding an approval by FDA for the use of a sponsored compound.

Sponsored compound means any drug or food additive or color additive proposed for use, or used, in food-producing animals or in their feed.

Target animals means the production class of animals in which a sponsored compound is proposed or intended for use.

Target tissue means the edible tissue selected to monitor for residues in the target animals, including, where appropriate, milk or eggs.

Test animals means the species selected for use in the toxicity tests.

Threshold assessment means FDA's review of data and information about a sponsored compound to determine whether chronic bioassays in test animals are necessary to resolve questions concerning the carcinogenicity of the compound.

§ 500.84 Operational definition of "no residue".

(a) On the basis of the results of the chronic bioassays and other information, FDA will determine whether any of the substances tested are carcinogenic.

(b) If FDA concludes that the results of the bioassays do not establish carcinogenicity, then FDA will not subject the sponsored compound to the remainder of the requirements of this subpart.

(c) For each sponsored compound that FDA decides should be regulated as a carcinogen, FDA will analyze the data from the bioassays using a statistical extrapolation procedure.

(1) For each substance tested in separate bioassays, FDA will calculate the concentration of the residue of carcinogenic concern that corresponds to a maximum lifetime risk to the test animal of 1 in 1 million. FDA will designate the lowest value obtained as S_o .

(2) FDA will consider that "no residue" of the compound remains in the edible tissue when conditions of use of the sponsored compound, including any required preslaughter withdrawal period or milk discard time, ensure that the concentration of the residue of carcinogenic concern in the total diet of people will not exceed S_o . Because the total diet is not derived from food-producing animals, FDA will make corrections for food intake. FDA will designate as S_m the concentration of residue of carcinogenic concern that is permitted in a specific edible product.

§ 500.86 Marker residue and target tissue.

(a) For each edible tissue, the sponsor shall measure the depletion of the residue of carcinogenic concern until its concentration is at or below S_m .

(b) In one or more edible tissues, the sponsor shall also measure the depletion of one or more potential marker residues until the concentration of the

§ 500.88

residue of carcinogenic concern is at or below S_m .

(c) From these data, FDA will select a target tissue and a marker residue and designate the concentration of marker residue (R_m) that the regulatory method must be capable of measuring in the target tissue. FDA will select R_m such that the absence of the marker residue in the target tissue above R_m can be taken as confirmation that the residue of carcinogenic concern does not exceed S_m in each of the edible tissues and, therefore, that the residue of carcinogenic concern in the diet of people does not exceed S_o .

(d) When a compound is to be used in milk- or egg-producing animals, milk or eggs must be the target tissue in addition to the tissue selected to monitor for residues in the edible carcass.

(Approved by the Office of Management and Budget under control number 0910-0228)

§ 500.88 Regulatory method.

(a) The sponsor shall submit for evaluation and validation a regulatory method developed to monitor compliance with FDA's operational definition of no residue.

(b) The regulatory method must reliably measure and confirm the identity of the marker residue in the target tissue at concentrations equal to and above R_m .

(c) FDA will publish in the FEDERAL REGISTER the complete regulatory method for measuring the marker residue in the target tissue in accordance with the provisions of sections 409(c)(3)(A), 512(d)(1)(H) and (i), and 721(b)(5)(B) of the act.

(Approved by the Office of Management and Budget under control number 0910-0228)

§ 500.90 Waiver of requirements.

In response to a petition or on the Commissioner's own initiative, the Commissioner may waive, in whole or in part, the requirements of this subpart except those provided under § 500.88. A petition for this waiver may be filed by any person who would be adversely affected by the application of the requirements to a particular compound. The petition shall explain and document why the requirements from which a waiver is requested are not reasonably applicable to the com-

21 CFR Ch. I (4-1-99 Edition)

pound, and set forth clearly the reasons why the alternative procedures will provide the basis for concluding that approval of the compound satisfies the requirements of the anticancer provisions of the act. If the Commissioner determines that waiver of any of the requirements of this subpart is appropriate, the Commissioner will state the basis for that determination in the regulation approving marketing of the sponsored compound.

(Approved by the Office of Management and Budget under control number 0910-0228)

§ 500.92 Implementation.

(a) This subpart E applies to all new animal drug applications, food additive petitions, and color additive petitions concerning any compound intended for use in food-producing animals (including supplemental applications and amendments to petitions).

(b) This subpart E also applies in the following manner to compounds already approved:

(1) For those compounds that FDA determines may induce cancer when ingested by man or animals, i.e., suspect carcinogens, §§ 500.80(b), 500.82, and 500.90 apply.

(2) For those compounds that FDA determines have been shown to induce cancer when ingested by man or animals, §§ 500.82 through 500.90 apply.

PART 501—ANIMAL FOOD LABELING

Subpart A—General Provisions

Sec.

501.1 Principal display panel of package form animal food.

501.2 Information panel of package for animal food.

501.3 Identity labeling of animal food in package form.

501.4 Animal food; designation of ingredients.

501.5 Animal food; name and place of business of manufacturer, packer, or distributor.

501.8 Labeling of animal food with number of servings.

501.15 Animal food; prominence of required statements.

501.17 Animal food labeling warning statements.

501.18 Misbranding of animal food.

Food and Drug Administration, HHS

§ 501.2

Subpart B—Specific Animal Food Labeling Requirements

501.22 Animal foods; labeling of spices, flavorings, colorings, and chemical preservatives.

Subparts C–E [Reserved]

Subpart F—Exemptions From Animal Food Labeling Requirements

501.100 Animal food; exemptions from labeling.

501.103 Petitions requesting exemptions from or special requirements for label declaration of ingredients.

501.105 Declaration of net quantity of contents when exempt.

501.110 Animal feed labeling; collective names for feed ingredients.

AUTHORITY: 15 U.S.C. 1453, 1454, 1455; 21 U.S.C. 321, 331, 342, 343, 348, 371.

SOURCE: 41 FR 38619, Sept. 10, 1976, unless otherwise noted.

Subpart A—General Provisions

§ 501.1 Principal display panel of package form animal food.

The term *principal display panel* as it applies to food in package form and as used in this part, means the part of a label that is most likely to be displayed, presented, shown, or examined under customary conditions of display for retail sale. The principal display panel shall be large enough to accommodate all the mandatory label information required to be placed thereon by this part with clarity and conspicuousness and without obscuring design, vignettes, or crowding. Where packages bear alternate principal display panels, information required to be placed on the principal display panel shall be duplicated on each principal display panel. For the purpose of obtaining uniform type size in declaring the quantity of contents for all packages of substantially the same size, the term *area of the principal display panel* means the area of the side or surface that bears the principal display panel, which area shall be:

(a) In the case of a rectangular package where one entire side properly can be considered to be the principal display panel side, the product of the height times the width of that side;

(b) In the case of a cylindrical or nearly cylindrical container, 40 percent of the product of the height of the container times the circumference;

(c) In the case of any otherwise shaped container, 40 percent of the total surface of the container: *Provided, however,* That where such container presents an obvious *principal display panel* such as the top of a triangular or circular package, the area shall consist of the entire top surface. In determining the area of the principal display panel, exclude tops, bottoms, flanges at tops and bottoms of cans, and shoulders and necks of bottles or jars. In the case of cylindrical or nearly cylindrical containers, information required by this part to appear on the principal display panel shall appear within that 40 percent of the circumference which is most likely to be displayed, presented, shown, or examined under customary conditions of display for retail sale.

§ 501.2 Information panel of package for animal food.

(a) The term *information panel* as it applies to packaged food means that part of the label immediately contiguous and to the right of the principal display panel as observed by an individual facing the principal display panel with the following exceptions:

(1) If the part of the label immediately contiguous and to the right of the principal display panel is too small to accommodate the necessary information or is otherwise unusable label space, e.g., folded flaps or can ends, the panel immediately contiguous and to the right of this part of the label may be used.

(2) If the package has one or more alternate principal display panels, the information panel is immediately contiguous and to the right of any principal display panel.

(3) If the top of the container is the principal display panel and the package has no alternate principal display panel, the information panel is any panel adjacent to the principal display panel.

(b) All information required to appear on the label of any package of food pursuant to §§ 501.4, 501.5, 501.8 and

§ 501.2

21 CFR Ch. I (4–1–99 Edition)

501.17 shall appear either on the principal display panel or on the information panel, unless otherwise specified by regulations in this chapter.

(c) All information appearing on the principal display panel or the information panel pursuant to this section shall appear prominently and conspicuously, but in no case may the letters and/or numbers be less than $\frac{1}{16}$ inch in height unless an exemption pursuant to paragraph (f) of this section is established. The requirements for conspicuousness and legibility shall include the specifications of §§ 501.15 and 501.105(h) (1) and (2).

(1) Packaged foods are exempt from the type size requirements of this paragraph: *Provided, That:*

(i) The package is designed such that it has a surface area that can bear an information panel and/or an alternate principal display panel.

(ii) The area of surface available for labeling on the principal display panel of the package as this term is defined in § 501.1 is less than 10 square inches.

(iii) The label information includes a full list of ingredients in accordance with regulations in this part.

(iv) The information required by paragraph (b) of this section appears on the principal display panel or information panel label in accordance with the provisions of this paragraph (c) except that the type size is not less than $\frac{3}{64}$ inch in height.

(2) Packaged foods are exempt from the type size requirements of this paragraph: *Provided, That:*

(i) The package is designed such that it has a single *obvious principal display panel* as this term is defined in § 501.1 and has no other available surface area for an information panel or alternate principal display panel.

(ii) The area of surface available for labeling on the principal display panel of the package as this term is defined in § 501.1 is less than 12 square inches and bears all labeling appearing on the package.

(iii) The label information includes a full list of ingredients in accordance with regulations in this part.

(iv) The information required by paragraph (b) of this section appears on the single, obvious principal display panel in accordance with the provisions

of this paragraph (c) except that the type size is not less than $\frac{1}{32}$ inch in height.

(3) Packaged foods are exempt from the type size requirements of this paragraph: *Provided, That:*

(i) The package is designed such that it has a total surface area available to bear labeling of less than 12 square inches.

(ii) The label information includes a full list of ingredients in accordance with regulations in this part.

(iii) The information required by paragraph (b) of this section appears on the principal display panel or information panel label in accordance with the provisions of this paragraph (c) except that the type size is not less than $\frac{1}{32}$ inch in height.

(d) All information required to appear on the principal display panel or on the information panel pursuant to this section shall appear on the same panel unless there is insufficient space. In determining the sufficiency of the available space, any vignettes, design, and other nonmandatory label information shall not be considered. If there is insufficient space for all of this information to appear on a single panel, it may be divided between these two panels except that the information required pursuant to any given section or part shall all appear on the same panel. A food whose label is required to bear the ingredient statement on the principal display panel may bear all other information specified in paragraph (b) of this section on the information panel.

(e) All information appearing on the information panel pursuant to this section shall appear in one place without other intervening material.

(f) If the label of any package of food is too small to accommodate all of the information required by §§ 501.4, 501.5, 501.8, and 501.17, the Commissioner may establish by regulation an acceptable alternative method of disseminating such information to the public, e.g., a type size smaller than one-sixteenth inch in height, or labeling attached to or inserted in the package or available at the point of purchase. A petition requesting such a regulation, as an amendment to this paragraph shall be

Food and Drug Administration, HHS

§ 501.3

submitted pursuant to part 10 of this chapter.

[41 FR 38619, Sept. 10, 1976, as amended at 42 FR 4716, Jan. 25, 1977; 42 FR 15675, Mar. 22, 1977]

§ 501.3 Identity labeling of animal food in package form.

(a) The principal display panel of a food in package form shall bear as one of its principal features a statement of the identity of the commodity.

(b) Such statement of identity shall be in terms of:

(1) The name now or hereafter specified in or required by any applicable Federal law or regulation; or, in the absence thereof,

(2) The common or usual name of the food; or, in the absence thereof,

(3) An appropriately descriptive term, or when the nature of the food is obvious, a fanciful name commonly used by the public for such food.

(c) Where a food is marketed in various optional forms (whole, slices, diced, etc.), the particular form shall be considered to be a necessary part of the statement of identity and shall be declared in letters of a type size bearing a reasonable relation to the size of the letters forming the other components of the statement of identity; except that if the optional form is visible through the container or is depicted by an appropriate vignette, the particular form need not be included in the statement. This specification does not affect the required declarations of identity under definitions and standards for foods promulgated pursuant to section 401 of the act.

(d) This statement of identity shall be presented in bold type on the principal display panel, shall be in a size reasonably related to the most prominent printed matter on such panel, and shall be in lines generally parallel to the base on which the package rests as it is designed to be displayed.

(e) Under the provisions of section 403(c) of the Federal Food, Drug, and Cosmetic Act, a food shall be deemed to be misbranded if it is an imitation of another food unless its label bears, in type of uniform size and prominence, the word *imitation* and, immediately thereafter, the name of the food imitated.

(1) A food shall be deemed to be an imitation and thus subject to the requirements of section 403(c) of the act if it is a substitute for and resembles another food but is nutritionally inferior to that food.

(2) A food that is a substitute for and resembles another food shall not be deemed to be an imitation provided it meets each of the following requirements:

(i) It is not nutritionally inferior to the food for which it substitutes and which it resembles.

(ii) Its label bears a common or usual name that complies with the provisions of § 502.5 of this chapter and that is not false or misleading, or in the absence of an existing common or usual name, an appropriately descriptive term that is not false or misleading. The label may, in addition, bear a fanciful name which is not false or misleading.

(3) A food for which a common or usual name is established by regulation (e.g., in a standard of identity pursuant to section 401 of the act, in a common or usual name regulation and may, in addition, bear a fanciful name which is not false or misleading, and established pursuant to part 502 of this chapter), and which complies with all of the applicable requirements of such regulation(s), shall not be deemed to be an imitation.

(4) Nutritional inferiority includes:

(i) Any reduction in the content of an essential nutrient that is present in a measurable amount.

(ii) If the Commissioner concludes that a food is a substitute for and resembles another food but is inferior to the food imitated for reasons other than those set forth in this paragraph, he may propose appropriate revisions to this regulation or he may propose a separate regulation governing the particular food.

(f) A label may be required to bear the percentage(s) of a characterizing ingredient(s) or information concerning the presence or absence of an ingredient(s) or the need to add an ingredient(s) as part of the common or usual name of the food pursuant to part 502 of this chapter.

[41 FR 38619, Sept. 10, 1976, as amended at 42 FR 14091, Mar. 15, 1977; 54 FR 18279, Apr. 28, 1989]

§ 501.4

21 CFR Ch. I (4–1–99 Edition)

§ 501.4 Animal food; designation of ingredients.

(a) Ingredients required to be declared on the label of a food, including foods that comply with standards of identity that require labeling in compliance with this part 501, except those exempted by § 501.100, shall be listed by common or usual name in descending order of predominance by weight on either the principal display panel or the information panel in accordance with the provisions of § 501.2.

(b) The name of an ingredient shall be a specific name and not a collective (generic) name, except that:

(1) Spices, flavorings, colorings and chemical preservatives shall be declared according to the provisions of § 501.22.

(2) An ingredient which itself contains two or more ingredients and which has an established common or usual name, conforms to a standard established pursuant to the Meat Inspection or Poultry Products Inspection Acts by the U.S. Department of Agriculture, or conforms to a definition and standard of identity established pursuant to section 401 of the Federal Food, Drug, and Cosmetic Act, shall be designated in the statement of ingredients on the label of such food by either of the following alternatives:

(i) By declaring the established common or usual name of the ingredient followed by a parenthetical listing of all ingredients contained therein in descending order of predominance except that, if the ingredient is a food subject to a definition and standard of identity established in this subchapter E, only the ingredients required to be declared by the definition and standard of identity need be listed; or

(ii) By incorporating into the statement of ingredients in descending order of predominance in the finished food, the common or usual name of every component of the ingredient without listing the ingredient itself.

(3) Skim milk, concentrated skim milk, reconstituted skim milk, and nonfat dry milk may be declared as *skim milk* or *nonfat milk*.

(4) Milk, concentrated milk, reconstituted milk, and dry whole milk may be declared as *milk*.

(5) Bacterial cultures may be declared by the word *cultured* followed by the name of the substrate, e.g., *made from cultured skim milk* or *cultured buttermilk*.

(6) Sweetcream buttermilk, concentrated sweetcream buttermilk, reconstituted sweetcream buttermilk, and dried sweetcream buttermilk may be declared as *buttermilk*.

(7) Whey, concentrated whey, reconstituted whey, and dried whey may be declared as *whey*.

(8) Cream, reconstituted cream, dried cream, and plastic cream (sometimes known as concentrated milkfat) may be declared as *cream*.

(9) Butteroil and anhydrous butterfat may be declared as *butterfat*.

(10) Dried whole eggs, frozen whole eggs, and liquid whole eggs may be declared as *eggs*.

(11) Dried egg whites, frozen egg whites, and liquid egg whites may be declared as *egg whites*.

(12) Dried egg yolks, frozen egg yolks, and liquid egg yolks may be declared as *egg yolks*.

(13) A livestock or poultry feed may be declared by a collective name listed in § 501.110 if it is an animal feed within the meaning of section 201(w) of the act and meets the requirements for the use of a collective name as prescribed in § 501.110 for certain feed ingredients.

(14) [Reserved]

(15) When all the ingredients of a wheat flour are declared in an ingredient statement, the principal ingredient of the flour shall be declared by the name(s) specified in §§ 137.105, 137.200, 137.220, 137.225 of this chapter, i.e., the first ingredient designated in the ingredient list of flour, or bromated flour, or enriched flour, or self-rising flour is *flour*, *white flour*, *wheat flour*, or *plain flour*; the first ingredient designated in the ingredient list of durum flour is *durum flour*; the first ingredient designated in the ingredient list of whole wheat flour, or bromated whole wheat flour is *whole wheat flour*, *graham flour*, or *entire wheat flour*; and the first ingredient designated in the ingredient list of whole durum wheat flour is *whole durum wheat flour*.

Food and Drug Administration, HHS

§ 501.15

(c) When water is added to reconstitute, completely or partially, an ingredient permitted by paragraph (b) of this section to be declared by a class name, the position of the ingredient class name in the ingredient statement shall be determined by the weight of the unreconstituted ingredient plus the weight of the quantity of water added to reconstitute that ingredient, up to the amount of water needed to reconstitute the ingredient to single strength. Any water added in excess of the amount of water needed to reconstitute the ingredient to single strength shall be declared as *water* in the ingredient statement.

[41 FR 38619, Sept. 10, 1976, as amended at 42 FR 14091, Mar. 15, 1977; 60 FR 38480, July 27, 1995]

§ 501.5 Animal food; name and place of business of manufacturer, packer, or distributor.

(a) The label of a food in packaged form shall specify conspicuously the name and place of business of the manufacturer, packer, or distributor.

(b) The requirement for declaration of the name of the manufacturer, packer, or distributor shall be deemed to be satisfied, in the case of a corporation, only by the actual corporate name, which may be preceded or followed by the name of the particular division of the corporation. In the case of an individual, partnership, or association, the name under which the business is conducted shall be used.

(c) Where the food is not manufactured by the person whose name appears on the label, the name shall be qualified by a phrase that reveals the connection such person has with such food; such as "Manufactured for _____," "Distributed by _____," or any other wording that expresses the facts.

(d) The statement of the place of business shall include the street address, city, state, and ZIP Code; however, the street address may be omitted if it is shown in a current city directory or telephone directory. The requirement for inclusion of the ZIP Code shall apply only to consumer commodity labels developed or revised after the effective date of this section. In the case of nonconsumer packages,

the ZIP Code shall appear either on the label or the labeling (including invoice).

(e) If a person manufactures, packs, or distributes a food at a place other than his principal place of business, the label may state the principal place of business in lieu of the actual place where such food was manufactured or packed or is to be distributed, unless such statement would be misleading.

§ 501.8 Labeling of animal food with number of servings.

(a) The label of any package of a food which bears a representation as to the number of servings contained in such package shall bear in immediate conjunction with such statement, and in the same size type as is used for such statement, a statement of the net quantity (in terms of weight, measure, or numerical count) of each such serving; however, such statement may be expressed in terms that differ from the terms used in the required statement of net quantity of contents (for example, cupfuls, tablespoonfuls, etc.) when such differing term is common to cookery and describes a constant quantity. Such statement may not be misleading in any particular. A statement of the number of units in a package is not in itself a statement of the number of servings.

(b) If there exists a voluntary product standard promulgated pursuant to the procedures found in 15 CFR part 10 by the Department of Commerce, quantitatively defining the meaning of the term *serving* with respect to a particular food, then any label representation as to the number of servings in such packaged food shall correspond with such quantitative definition. (Copies of published standards are available upon request from the National Bureau of Standards, Department of Commerce, Washington, DC 20234.)

§ 501.15 Animal food; prominence of required statements.

(a) A word, statement, or other information required by or under authority of the act to appear on the label may lack that prominence and conspicuousness required by section 403(f) of the act by reason (among other reasons) of:

§ 501.17

21 CFR Ch. I (4–1–99 Edition)

(1) The failure of such word, statement, or information to appear on the part or panel of the label which is presented or displayed under customary conditions of purchase;

(2) The failure of such word, statement, or information to appear on two or more parts or panels of the label, each of which has sufficient space therefor, and each of which is so designed as to render it likely to be, under customary conditions of purchase, the part or panel displayed;

(3) The failure of the label to extend over the area of the container or package available for such extension, so as to provide sufficient label space for the prominent placing of such word, statement, or information;

(4) Insufficiency of label space (for the prominent placing of such word, statement, or information) resulting from the use of label space for any word, statement, design, or device which is not required by or under authority of the act to appear on the label;

(5) Insufficiency of label space (for the prominent placing of such word, statement, or information) resulting from the use of label space to give materially greater conspicuousness to any other word, statement, or information, or to any design or device; or

(6) Smallness or style of type in which such word, statement, or information appears, insufficient background contrast, obscuring designs or vignettes, or crowding with other written, printed, or graphic matter.

(b) No exemption depending on insufficiency of label space, as prescribed in regulations promulgated under section 403(e) or (i) of the act, shall apply if such insufficiency is caused by:

(1) The use of label space for any word, statement, design, or device which is not required by or under authority of the act to appear on the label;

(2) The use of label space to give greater conspicuousness to any word, statement, or other information that is required by section 403(f) of the act; or

(3) The use of label space for any representation in a foreign language.

(c)(1) All words, statements, and other information required by or under authority of the act to appear on the

label or labeling shall appear thereon in the English language: *Provided, however,* That in the case of articles distributed solely in the Commonwealth of Puerto Rico or in a territory where the predominant language is one other than English, the predominant language may be substituted for English.

(2) If the label contains any representation in a foreign language, all words, statements, and other information required by or under authority of the act to appear on the label shall appear thereon in the foreign language.

(3) If any article of labeling (other than a label) contains any representation in a foreign language, all words, statements, and other information required by or under authority of the act to appear on the label or labeling shall appear on such article of labeling.

§ 501.17 Animal food labeling warning statements.

(a) *Self-pressurized containers.* (1) The label of a food packaged in a self-pressurized container and intended to be expelled from the package under pressure shall bear the following warning:

Warning Avoid spraying in eyes. Contents under pressure. Do not puncture or incinerate. Do not store at temperature above 120° F. Keep out of reach of children.

(2) In the case of products intended for use by children, the phrase “except under adult supervision” may be added at the end of the last sentence in the warning required by paragraph (a)(1) of this section.

(3) In the case of products packaged in glass containers, the word “break” may be substituted for the word “puncture” in the warning required by paragraph (a)(1) of this section.

(4) The words “Avoid spraying in eyes” may be deleted from the warning required by paragraph (a)(1) of this section in the case of a product not expelled as a spray.

(b) *Self-pressurized containers with halocarbon or hydrocarbon propellants.* (1) In addition to the warning required by paragraph (a) of this section, the label of a food packaged in a self-pressurized container in which the propellant consists in whole or in part of a halocarbon or a hydrocarbon shall bear the following warning:

Food and Drug Administration, HHS

§ 501.22

Warning Use only as directed. Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

(2) The warning required by paragraph (b)(1) of this section is not required for the following products:

(i) Products expelled in the form of a foam or cream, which contain less than 10 percent propellant in the container.

(ii) Products in a container with a physical barrier that prevents escape of the propellant at the time of use.

(iii) Products of a net quantity of contents of less than 2 ozs that are designed to release a measured amount of product with each valve actuation.

(iv) Products of a net quantity of contents of less than ½ oz.

(c) *Animal food containing or manufactured with a chlorofluorocarbon or other ozone-depleting substance.* Labeling requirements for animal foods that contain or are manufactured with a chlorofluorocarbon or other ozone-depleting substance designated by the Environmental Protection Agency (EPA) are set forth in 40 CFR part 82.

[41 FR 38619, Sept. 10, 1976, as amended at 42 FR 22033, Apr. 29, 1977; 61 FR 20101, May 3, 1996]

§ 501.18 Misbranding of animal food.

(a) Among representations in the labeling of a food which render such food misbranded is a false or misleading representation with respect to another food or a drug, device, or cosmetic.

(b) The labeling of a food which contains two or more ingredients may be misleading by reason (among other reasons) of the designation of such food in such labeling by a name which includes or suggests the name of one or more but not all such ingredients, even though the names of all such ingredients are stated elsewhere in the labeling.

(c) Among representations in the labeling of a food which render such food misbranded is any representation that expresses or implies a geographical origin of the food or any ingredient of the food except when such representation is either:

(1) A truthful representation of geographical origin.

(2) A trademark or trade name provided that as applied to the article in

question its use is not deceptively misdescriptive. A trademark or trade name comprised in whole or in part of geographical words shall not be considered deceptively misdescriptive if it:

(i) Has been so long and exclusively used by a manufacturer or distributor that it is generally understood by the consumer to mean the product of a particular manufacturer or distributor; or

(ii) Is so arbitrary or fanciful that it is not generally understood by the consumer to suggest geographic origin.

(3) A part of the name required by applicable Federal law or regulation.

(4) A name whose market significance is generally understood by the consumer to connote a particular class, kind, type, or style of food rather than to indicate geographical origin.

Subpart B—Specific Animal Food Labeling Requirements

§ 501.22 Animal foods; labeling of spices, flavorings, colorings, and chemical preservatives.

(a)(1) The term *artificial flavor* or *artificial flavoring* means any substance, the function of which is to impart flavor, which is not derived from a spice, fruit or fruit juice, vegetable or vegetable juice, edible yeast, herb, bark, bud, root, leaf or similar plant material, meat, fish, poultry, eggs, dairy products, or fermentation products thereof. Artificial flavor includes the substances listed in §§ 172.515(b) and 582.60 of this chapter except where these are derived from natural sources.

(2) The term *spice* means any aromatic vegetable substance in the whole, broken, or ground form, except for those substances which have been traditionally regarded as foods, such as onions, garlic and celery; whose significant function in food is seasoning rather than nutritional; that is true to name; and from which no portion of any volatile oil or other flavoring principle has been removed. Spices include the spices listed in subpart A of part 582 of this chapter, such as the following:

Allspice, Anise, Basil, Bay leaves, Caraway seed, Cardamon, Celery seed, Chervil, Cinamon, Cloves, Coriander, Cumin seed, Dill seed, Fennel seed, Fenugreek, Ginger, Horseradish, Mace, Marjoram, Mustard

§ 501.22

flour, Nutmeg, Oregano, Paprika, Parsley, Pepper, black; Pepper, white; Pepper, red; Rosemary, Saffron, Sage, Savory, Star aniseed, Tarragon, Thyme, Turmeric.

Paprika, turmeric, and saffron or other spices which are also colors, shall be declared as *spice and coloring* unless declared by their common or usual name.

(3) The term *natural flavor* or *natural flavoring* means the essential oil, oleoresin, essence or extractive, protein hydrolysate, distillate, or any product of roasting, heating or enzymolysis, which contains the flavoring constituents derived from a spice, fruit or fruit juice, vegetable or vegetable juice, edible yeast, herb, bark, bud, root, leaf or similar plant material, meat, seafood, poultry, eggs, dairy products, or fermentation products thereof, whose significant function in food is flavoring rather than nutritional. Natural flavors, include the natural essence or extractives obtained from plants listed in subpart A of part 582 of this chapter, and the substances listed in § 172.510 of this chapter.

(4) The term *artificial color* or *artificial coloring* means any *color additive* as defined in § 70.3(f) of this chapter.

(5) The term *chemical preservative* means any chemical that, when added to food, tends to prevent or retard deterioration thereof, but does not include common salt, sugars, vinegars, spices, or oils extracted from spices, substances added to food by direct exposure thereof to wood smoke, or chemicals applied for their insecticidal or herbicidal properties.

(b) A food which is subject to the requirements of section 403(k) of the act shall bear labeling, even though such food is not in package form.

(c) A statement of artificial flavoring, artificial coloring, or chemical preservative shall be placed on the food, or on its container or wrapper, or on any two or all of these, as may be necessary to render such statement likely to be read by the ordinary individual under customary conditions of purchase and use of such food.

(d) A food shall be exempt from compliance with the requirements of section 403(k) of the act if it is not in package form and the units thereof are so small that a statement of artificial flavoring, artificial coloring, or chem-

21 CFR Ch. I (4–1–99 Edition)

ical preservative, as the case may be, cannot be placed on such units with such conspicuousness as to render it likely to be read by the ordinary individual under customary conditions of purchase and use.

(e) A food shall be exempt while held for sale from the requirements of section 403(k) of the act (requiring label statement of any artificial flavoring, artificial coloring, or chemical preservatives) if said food, having been received in bulk containers at a retail establishment, is displayed to the purchaser with either (1) the labeling of the bulk container plainly in view or (2) a counter card, sign, or other appropriate device bearing prominently and conspicuously the information required to be stated on the label pursuant to section 403(k) of the act.

(f) A fruit or vegetable shall be exempt from compliance with the requirements of section 403(k) of the act with respect to a chemical preservative applied to the fruit or vegetable as a pesticide chemical prior to harvest.

(g) A flavor shall be labeled in the following way when shipped to a food manufacturer or processor (but not a consumer) for use in the manufacture of a fabricated food, unless it is a flavor for which a standard of identity has been promulgated, in which case it shall be labeled as provided in the standard:

(1) If the flavor consists of one ingredient, it shall be declared by its common or usual name.

(2) If the flavor consists of two or more ingredients, the label either may declare each ingredient by its common or usual name or may state "All flavor ingredients contained in this product are approved for use in a regulation of the Food and Drug Administration." Any flavor ingredient not contained in one of these regulations, and any non-flavor ingredient, shall be separately listed on the label.

(3) In cases where the flavor contains a solely natural flavor(s), the flavor shall be so labeled, e.g., *strawberry flavor*, *banana flavor*, or *natural strawberry flavor*. In cases where the flavor contains both a natural flavor and an artificial flavor, the flavor shall be so labeled, e.g., *natural and artificial strawberry flavor*. In cases where the flavor

contains a solely artificial flavor(s), the flavor shall be so labeled, e.g., *artificial strawberry flavor*.

(h) The label of a food to which flavor is added shall declare the flavor in the statement of ingredients in the following way:

(1) Spice, natural flavor, and artificial flavor may be declared as *spice*, *natural flavor*, or *artificial flavor*, or any combination thereof, as the case may be.

(2) An incidental additive in a food, originating in a spice or flavor used in the manufacture of the food, need not be declared in the statement of ingredients if it meets the requirements of § 501.100(a)(3).

(3) Substances obtained by cutting, grinding, drying, pulping, or similar processing of tissues derived from fruit, vegetable, meat, fish, or poultry, e.g., powdered or granulated onions, garlic powder, and celery powder, are commonly understood by consumers to be food rather than flavor and shall be declared by their common or usual name.

(4) Any salt (sodium chloride) used as an ingredient in food shall be declared by its common or usual name *salt*.

(5) Any monosodium glutamate used as an ingredient in food shall be declared by its common or usual name *monosodium glutamate*.

(6) Any pyroligneous acid or other artificial smoke flavors used as an ingredient in a food may be declared as *artificial flavor* or *artificial smoke flavor*. No representation may be made, either directly or implied, that a food flavored with pyroligneous acid or other artificial smoke flavor has been smoked or has a true smoked flavor, or that a seasoning sauce or similar product containing pyroligneous acid or other artificial smoke flavor and used to season or flavor other foods will result in a smoked product or one having a true smoked flavor.

(i) If the label, labeling, or advertising of a food makes any direct or indirect representations with respect to the primary recognizable flavor(s), by word, *vignette*, e.g., depiction of a fruit, or other means, or if for any other reason the manufacturer or distributor of a food wishes to designate the type of flavor in the food other than through the statement of ingredients, such fla-

vor shall be considered the characterizing flavor and shall be declared in the following way:

(1) If the food contains no artificial flavor which simulates, resembles or reinforces the characterizing flavor, the name of the food on the principal display panel or panels of the label shall be accompanied by the common or usual name of the characterizing flavor in letters not less than one-half the height of the letters used in the name of the food, except that:

(i) If the food is one that is commonly expected to contain a characterizing food ingredient, and the food contains natural flavor derived from such ingredient and an amount of characterizing ingredient insufficient to independently characterize the food, or the food contains no such ingredient, the name of the characterizing flavor may be immediately preceded by the word *natural* and shall be immediately followed by the word *flavored* in letters not less than one-half the height of the letters in the name of the characterizing flavor.

(ii) If none of the natural flavor used in the food is derived from the product whose flavor is simulated, the food in which the flavor is used shall be labeled either with the flavor of the product from which the flavor is derived or as *artificially flavored*.

(iii) If the food contains both a characterizing flavor from the product whose flavor is simulated and other natural flavor which simulates, resembles or reinforces the characterizing flavor, the food shall be labeled in accordance with the introductory text and paragraph (i)(1)(i) of this section and the name of the food shall be immediately followed by the words *with other natural flavor* in letters not less than one-half the height of the letters used in the name of the characterizing flavor.

(2) If the food contains any artificial flavor which simulates, resembles or reinforces the characterizing flavor, the name of the food on the principal display panel or panels of the label shall be accompanied by the common or usual name(s) of the characterizing flavor, in letters not less than one-half the height of the letters used in the name of the food and the name of the

characterizing flavor shall be accompanied by the word(s) *artificial* or *artificially flavored*, in letters not less than one-half the height of the letters in the name of the characterizing flavor.

(3) Wherever the name of the characterizing flavor appears on the label (other than in the statement of ingredients) so conspicuously as to be easily seen under customary conditions of purchase, the words prescribed by this paragraph shall immediately and conspicuously precede or follow such name, without any intervening written, printed, or graphic matter, except:

(i) Where the characterizing flavor and a trademark or brand are presented together, other written, printed, or graphic matter that is a part of or is associated with the trademark or brand may intervene if the required words are in such relationship with the trademark or brand as to be clearly related to the characterizing flavor; and

(ii) If the finished product contains more than one flavor subject to the requirements of this paragraph, the statements required by this paragraph need appear only once in each statement of characterizing flavors present in such food.

(iii) If the finished product contains three or more distinguishable characterizing flavors, or a blend of flavors with no primary recognizable flavor, the flavor may be declared by an appropriately descriptive generic term in lieu of naming each flavor.

(4) A flavor supplier shall certify, in writing, that any flavor he supplies which is designated as containing no artificial flavor does not, to the best of his knowledge and belief, contain any artificial flavor, and that he has added no artificial flavor to it. The requirement for such certification may be satisfied by a guarantee under section 303(c)(2) of the act which contains such a specific statement. A flavor used shall be required to make such a written certification only where he adds to or combines another flavor with a flavor which has been certified by a flavor supplier as containing no artificial flavor, but otherwise such user may rely upon the supplier's certification and need make no separate certification. All such certifications shall be retained by the certifying party through-

out the period in which the flavor is supplied and for a minimum of 3 years thereafter, and shall be subject to the following conditions:

(i) The certifying party shall make such certifications available upon request at all reasonable hours to any duly authorized officer, or employee of the Food and Drug Administration or any other employee acting on behalf of the Secretary of Health and Human Services. Such certifications are regarded by the Food and Drug Administration as reports to the government and as guarantees or other undertakings within the meaning of section 301(h) of the act and subject the certifying party to the penalties for making any false report to the government under 18 U.S.C. 1001 and any false guarantee or undertaking under section 303(a) of the act. The defenses provided under section 303(c)(2) of the act shall be applicable to the certifications provided for in this section.

(ii) Wherever possible, the Food and Drug Administration shall verify the accuracy of a reasonable number of certifications made pursuant to this section, constituting a representative sample of such certifications, and shall not request all such certifications.

(iii) Where no person authorized to provide such information is reasonably available at the time of inspection, the certifying party shall arrange to have such person and the relevant materials and records ready for verification as soon as practicable; provided that, whenever the Food and Drug Administration has reason to believe that the supplier or user may utilize this period to alter inventories or records, such additional time shall not be permitted. Where such additional time is provided, the Food and Drug Administration may require the certifying party to certify that relevant inventories have not been materially disturbed and relevant records have not been altered or concealed during such period.

(iv) The certifying party shall provide, to an officer or representative duly designated by the Secretary, such qualitative statement of the composition of the flavor or product covered by the certification as may be reasonably

expected to enable the Secretary's representatives to determine which relevant raw and finished materials and flavor ingredient records are reasonably necessary to verify the certifications. The examination conducted by the Secretary's representative shall be limited to inspection and review of inventories and ingredient records for those certifications which are to be verified.

(v) Review of flavor ingredient records shall be limited to the qualitative formula and shall not include the quantitative formula. The person verifying the certifications may make only such notes as are necessary to enable him to verify such certification. Only such notes or such flavor ingredient records as are necessary to verify such certification or to show a potential or actual violation may be removed or transmitted from the certifying party's place of business: *Provided*, That, where such removal or transmittal is necessary for such purposes the relevant records and notes shall be retained as separate documents in Food and Drug Administration files, shall not be copied in other reports, and shall not be disclosed publicly other than in a judicial proceeding brought pursuant to the act or 18 U.S.C. 1001.

(j) A food to which a chemical preservative(s) is added shall, except when exempt pursuant to § 501.100, bear a label declaration stating both the common or usual name of the ingredient(s) and a separate description of its function, e.g., *preservative, to retard spoilage, a mold inhibitor, to help protect flavor or to promote color retention*.

[41 FR 38619, Sept. 10, 1976, as amended at 42 FR 14091, Mar. 15, 1977; 42 FR 15675, Mar. 22, 1977]

Subparts C-E [Reserved]

Subpart F—Exemptions From Animal Food Labeling Requirements

§ 501.100 Animal food; exemptions from labeling.

(a) The following foods are exempt from compliance with the requirements of section 403(i)(2) of the act (re-

quiring a declaration on the label of the common or usual name of each ingredient when the food is fabricated from two or more ingredients).

(1) An assortment of different items of food, when variations in the items that make up different packages packed from such assortment normally occur in good packing practice and when such variations result in variations in the ingredients in different packages, with respect to any ingredient that is not common to all packages. Such exemption, however, shall be on the condition that the label shall bear, in conjunction with the names of such ingredients as are common to all packages, a statement (in terms that are as informative as practicable and that are not misleading) indicating by name other ingredients which may be present.

(2) A food having been received in bulk containers at a retail establishment, if displayed to the purchaser with either (i) the labeling of the bulk container plainly in view or (ii) a counter card, sign, or other appropriate device bearing prominently and conspicuously the information required to be stated on the label pursuant to section 403(i)(2) of the act.

(3) Incidental additives that are present in a food at insignificant levels and do not have any technical or functional effect in that food. For the purposes of this paragraph (a)(3), incidental additives are:

(i) Substances that have no technical or functional effect but are present in a food by reason of having been incorporated into the food as an ingredient of another food, in which the substance did have a functional or technical effect.

(ii) Processing aids, which are as follows:

(a) Substances that are added to a food during the processing of such food but are removed in some manner from the food before it is packaged in its finished form.

(b) Substances that are added to a food during processing, are converted into constituents normally present in the food, and do not significantly increase the amount of the constituents naturally found in the food.

(c) Substances that are added to a food for their technical or functional effect in the processing but are present in the finished food at insignificant levels and do not have any technical or functional effect in that food.

(iii) Substances migrating to food from equipment or packaging or otherwise affecting food that are not food additives as defined in section 201(s) of the act; or if they are food additives as so defined, they are used in conformity with regulations established pursuant to section 409 of the act.

(b) A food repackaged in a retail establishment is exempt from the following provisions of the act if the conditions specified are met.

(1) Section 403(e)(1) of the act (requiring a statement on the label of the name and place of business of the manufacturer, packer, or distributor).

(2) Section 403(g)(2) of the act (requiring the label of a food which purports to be or is represented as one for which a definition and standard of identity has been prescribed to bear the name of the food specified in the definition and standard and, insofar as may be required by the regulation establishing the standard the common names of the optional ingredients present in the food), if the food is displayed to the purchaser with its interstate labeling clearly in view, or with a counter card, sign, or other appropriate device bearing prominently and conspicuously the information required by these provisions.

(3) Section 403(i)(1) of the act (requiring the label to bear the common or usual name of the food), if the food is displayed to the purchaser with its interstate labeling clearly in view, or with a counter card, sign, or other appropriate device bearing prominently and conspicuously the common or usual name of the food, or if the common or usual name of the food is clearly revealed by its appearance.

(c) [Reserved]

(d) Except as provided by paragraphs (e) and (f) of this section, a shipment or other delivery of a food which is, in accordance with the practice of the trade, to be processed, labeled, or repacked in substantial quantity at an establishment other than that where originally processed or packed, shall be exempt,

during the time of introduction into and movement in interstate commerce and the time of holding in such establishment, from compliance with the labeling requirements of section 403 (c), (e), (g), (h), (i), (j) and (k) of the act if:

(1) The person who introduced such shipment or delivery into interstate commerce is the operator of the establishment where such food is to be processed, labeled, or repacked; or

(2) In case such person is not such operator, such shipment or delivery is made to such establishment under a written agreement, signed by and containing the post office addresses of such person and such operator, and containing such specifications for the processing, labeling, or repacking, as the case may be, of such food in such establishment as will ensure, if such specifications are followed, that such food will not be adulterated or misbranded within the meaning of the act upon completion of such processing, labeling, or repacking. Such person and such operator shall each keep a copy of such agreement until 2 years after the final shipment or delivery of such food from such establishment, and shall make such copies available for inspection at any reasonable hour to any officer or employee of the Department who requests them.

(e) Conditions affecting expiration of exemptions.

(1) An exemption of a shipment or other delivery of a food under paragraph (d)(1) of this section shall, at the beginning of the act of removing such shipment or delivery, or any part thereof, from such establishment become void ab initio if the food comprising such shipment, delivery, or part is adulterated or misbranded within the meaning of the act when so removed.

(2) An exemption of a shipment or other delivery of a food under paragraph (d)(2) of this section shall become void ab initio with respect to the person who introduced such shipment or delivery into interstate commerce upon refusal by such person to make available for inspection a copy of the agreement, as required by paragraph (d)(2) of this section.

(3) An exemption of a shipment or other delivery of a food under paragraph (d)(2) of this section shall expire:

(i) At the beginning of the act of removing such shipment or delivery, or any part thereof, from such establishment if the food comprising such shipment, delivery, or part is adulterated or misbranded within the meaning of the act when so removed; or

(ii) Upon refusal by the operator of the establishment where such food is to be processed, labeled, or repacked, to make available for inspection a copy of the agreement as required by such paragraph.

(f) [Reserved]

(g) The label declaration of a harmless marker used to identify a particular manufacturer's product may result in unfair competition through revealing a trade secret. Exemption from the label declaration of such a marker is granted, therefore, provided that the following conditions are met:

(1) The person desiring to use the marker without label declaration of its presence has submitted to the Commissioner of Food and Drugs full information concerning the proposed usage and the reasons why he believes label declaration of the marker should be subject to this exemption; and

(2) The person requesting the exemption has received from the Commissioner of Food and Drugs a finding that the marker is harmless and that the exemption has been granted.

§ 501.103 Petitions requesting exemptions from or special requirements for label declaration of ingredients.

The Commissioner of Food and Drugs, either on his own initiative or on behalf of any interested person who has submitted a petition pursuant to part 10 of this chapter may issue a proposal to amend § 501.4 to specify the manner in which an ingredient(s) shall be declared, i.e., by specific or class name, or § 501.100 to exempt an ingredient(s) from the requirements for label declaration.

[41 FR 38619, Sept. 10, 1976, as amended at 42 FR 15675, Mar. 22, 1977]

§ 501.105 Declaration of net quantity of contents when exempt.

(a) The principal display panel of a food in package form shall bear a declaration of the net quantity of contents. This shall be expressed in the terms of weight, measure, numerical count, or a combination of numerical count and weight or measure. The statement shall be in terms of fluid measure if the food is liquid, or in terms of weight if the food is solid, semisolid, or viscous, or a mixture of solid and liquid; except that such statement may be in terms of dry measure if the food is a fresh fruit, fresh vegetable, or other dry commodity that is customarily sold by dry measure. If there is a firmly established general consumer usage and trade custom of declaring the contents of a liquid by weight, or a solid, semisolid, or viscous product by fluid measure, it may be used. Whenever the Commissioner determines that an existing practice of declaring net quantity of contents by weight, measure, numerical count, or a combination in the case of a specific packaged food does not facilitate value comparisons by consumers and offers opportunity for consumer confusion, he will by regulation designate the appropriate term or terms to be used for such commodity.

(b)(1) Statements of weight shall be in terms of avoirdupois pound and ounce.

(2) Statements of fluid measure shall be in terms of the U.S. gallon of 231 cubic inches and quart, pint, and fluid ounce subdivisions thereof, and shall:

(i) In the case of frozen food that is sold and consumed in a frozen state, express the volume at the frozen temperature.

(ii) In the case of refrigerated food that is sold in the refrigerated state, express the volume at 40 °F (4 °C).

(iii) In the case of other foods, express the volume at 68° F (20° C).

(3) Statements of dry measure shall be in terms of the U.S. bushel of 2,150.42 cubic inches and peck, dry quart, and dry pint subdivisions thereof.

(c) When the declaration of quantity of contents by numerical count does not give adequate information as to the quantity of food in the package, it

§ 501.105

21 CFR Ch. I (4–1–99 Edition)

shall be combined with such statement of weight, measure, or size of the individual units of the foods as will provide such information.

(d) The declaration may contain common or decimal fractions. A common fraction shall be in terms of halves, quarters, eighths, sixteenths, or thirty-seconds; except that if there exists a firmly established general consumer usage and trade custom of employing different common fractions in the net quantity declaration of a particular commodity, they may be employed. A common fraction shall be reduced to its lowest terms; a decimal fraction shall not be carried out to more than two places. A statement that includes small fractions of an ounce shall be deemed to permit smaller variations than one which does not include such fractions.

(e) The declaration shall be located on the principal display panel of the label, and with respect to packages bearing alternate principal panels it shall be duplicated on each principal display panel.

(f) The declaration shall appear as a distinct item on the principal display panel, shall be separated (by at least a space equal to the height of the lettering used in the declaration) from other printed label information appearing above or below the declaration and (by at least a space equal to twice the width of the letter “N” of the style of type used in the quantity of contents statement) from other printed label information appearing to the left or right of the declaration. It shall not include any term qualifying a unit of weight, measure, or count (such as *jumbo quart* and *full gallon*) that tends to exaggerate the amount of the food in the container. It shall be placed on the principal display panel within the bottom 30 percent of the area of the label panel in lines generally parallel to the base on which the package rests as it is designed to be displayed: *Provided*, That on packages having a principal display panel of 5 square inches or less, the requirement for placement within the bottom 30 percent of the area of the label panel shall not apply when the declaration of net quantity of contents meets the other requirements of this part.

(g) The declaration shall accurately reveal the quantity of food in the package exclusive of wrappers and other material packed therewith; provided that in the case of foods packed in containers designed to deliver the food under pressure, the declaration shall state the net quantity of the contents that will be expelled when the instructions for use as shown on the container are followed. The propellant is included in the net quantity declaration.

(h) The declaration shall appear in conspicuous and easily legible boldface print or type in distinct contrast (by typography, layout, color, embossing, or molding) to other matter on the package; except that a declaration of net quantity blown, embossed, or molded on a glass or plastic surface is permissible when all label information is so formed on the surface. Requirements of conspicuousness and legibility shall include the specifications that:

(1) The ratio of height to width (of the letter) shall not exceed a differential of 3 units to 1 unit (no more than 3 times as high as it is wide).

(2) Letter heights pertain to upper case or capital letters. When upper and lower case or all lower case letters are used, it is the lower case letter “o” or its equivalent that shall meet the minimum standards.

(3) When fractions are used, each component numeral shall meet one-half the minimum height standards.

(i) The declaration shall be in letters and numerals in a type size established in relationship to the area of the principal display panel of the package and shall be uniform for all packages of substantially the same size by complying with the following type specifications:

(1) Not less than $\frac{1}{16}$ inch in height on packages the principal display panel of which has an area of 5 square inches or less.

(2) Not less than $\frac{1}{8}$ inch in height on packages the principal display panel of which has an area of more than 5 but not more than 25 square inches.

(3) Not less than $\frac{3}{16}$ inch in height on packages the principal display panel of which has an area of more than 25 but not more than 100 square inches.

(4) Not less than $\frac{1}{4}$ inch in height on packages the principal display panel of

which has an area of more than 100 square inches, except not less than $\frac{1}{2}$ inch in height if the area is more than 400 square inches.

Where the declaration is blown, embossed, or molded on a glass or plastic surface rather than by printing, typing, or coloring, the lettering sizes specified in paragraphs (i) (1) through (4) of this section shall be increased by $\frac{1}{16}$ of an inch.

(j) On packages containing less than 4 pounds or 1 gallon and labeled in terms of weight or fluid measure:

(1) The declaration shall be expressed both in ounces, with identification by weight or by liquid measure and, if applicable (1 pound or 1 pint or more) followed in parentheses by a declaration in pounds for weight units, with any remainder in terms of ounces or common or decimal fractions of the pound (see examples set forth in paragraphs (m) (1) and (2) of this section), or in the case of liquid measure, in the largest whole units (quarts, quarts and pints, or pints, as appropriate) with any remainder in terms of fluid ounces or common or decimal fractions of the pint or quart (see examples in paragraphs (m) (3) and (4) of this section).

(2) If the net quantity of contents declaration appears on a random package, that is a package which is one of a lot, shipment, or delivery of packages of the same consumer commodity with varying weights and with no fixed weight pattern, it may, when the net weight exceeds 1 pound, be expressed in terms of pounds and decimal fractions of the pound carried out to not more than two decimal places. When the net weight does not exceed 1 pound, the declaration on the random package may be in decimal fractions of the pound in lieu of ounces (see example in paragraph (m)(5) of this section).

(3) The declaration may appear in more than one line. The term *net weight* shall be used when stating the net quantity of contents in terms of weight. Use of the terms *net* or *net contents* in terms of fluid measure or numerical count is optional. It is sufficient to distinguish avoirdupois ounce from fluid ounce through association of terms; for example, *Net wt. 6 oz.* or *6 oz. net wt.* and *6 fl. oz.* or *net contents 6 fl. oz.*

(k) On packages containing 4 pounds or 1 gallon or more and labeled in terms of weight or fluid measure, the declaration shall be expressed in pounds for weight units with any remainder in terms of ounces or common or decimal fraction of the pound, or in the case of fluid measure, it shall be expressed in the largest whole unit (gallons followed by common or decimal fraction of a gallon or by the next smaller whole unit or units (quarts, or quarts and pints)) with any remainder in terms of fluid ounces or common or decimal fractions of the pint or quart (see paragraph (m)(6) of this section).

(l) [Reserved]

(m) *Examples:* (1) A declaration of $1\frac{1}{2}$ pounds weight shall be expressed as *Net Wt. 24 oz. (1 lb. 8 oz.)*, *Net Wt. 24 oz. (1½ lb.)*, or *Net Wt. 24 oz. (1.5 lb.)*.

(2) A declaration of $\frac{3}{4}$ pound avoirdupois weight shall be expressed as *Net Wt. 12 oz.*

(3) A declaration of 1 quart liquid measure shall be expressed as *Net 32 fl. oz. (1 qt.)*.

(4) A declaration of $1\frac{3}{4}$ quarts liquid measure shall be expressed as *Net contents 56 fluid ounces (1 quart 1½ pints)* or as *Net 56 fluid oz. (1 qt. 1 pt. 8 oz.)*, but not in terms of quart and ounce such as *Net 56 fluid oz. (1 quart 24 ounces)*.

(5) On a random package, declaration of $\frac{3}{4}$ pound avoirdupois may be expressed as *Net Wt. .75 lb.*

(6) A declaration of $2\frac{1}{2}$ gallons liquid measure shall be expressed as *Net contents 2½ gallons*, *Net contents 2.5 gallons*, or *Net contents 2 gallons 2 quarts* and not as *2 gallons 4 pints*.

(n) For quantities, the following abbreviations and none other may be employed (periods and plural forms are optional):

weight wt.	pint pt.
ounce oz.	quart qt.
pound lb.	fluid fl.
gallon gal.	

(o) Nothing in this section shall prohibit supplemental statements at locations other than the principal display panel(s) describing in nondeceptive terms the net quantity of contents; provided, that such supplemental statements of net quantity of contents shall not include any term qualifying a unit of weight, measure, or count that tends to exaggerate the amount of the

food contained in the package; for example, *jumbo quart* and *full gallon*. Dual or combination declarations of net quantity of contents as provided for in paragraphs (a), (c), and (j) of this section (for example, a combination of net weight plus numerical count, net contents plus dilution directions of a concentrate, etc.) are not regarded as supplemental net quantity statements and may be located on the principal display panel.

(p) A separate statement of the net quantity of contents in terms of the metric system is not regarded as a supplemental statement and an accurate statement of the net quantity of contents in terms of the metric system of weight or measure may also appear on the principal display panel or on other panels.

(q) The declaration of net quantity of contents shall express an accurate statement of the quantity of contents of the package. Reasonable variations caused by loss or gain of moisture during the course of good distribution practice or by unavoidable deviations in good manufacturing practice will be recognized. Variations from stated quantity of contents shall not be unreasonably large.

(r) [Reserved]

(s) On a multiunit retail package, a statement of the quantity of contents shall appear on the outside of the package and shall include the number of individual units, the quantity of each individual unit, and, in parentheses, the total quantity of contents of the multiunit package in terms of avoirdupois or fluid ounces, except that such declaration of total quantity need not be followed by an additional parenthetical declaration in terms of the largest whole units and subdivisions thereof, as required by paragraph (j)(1) of this section. A multiunit retail package may thus be properly labeled: *6–16 oz. bottles—(96 fl. oz.)* or *3–16 oz. cans—(net wt. 48 oz.)*. For the purposes of this section, *multiunit retail package* means a package containing two or more individually packaged units of the identical commodity and in the same quantity, intended to be sold as part of the multiunit retail package but capable of being individually sold in full compliance with all requirements of the regu-

lations in this part. Open multiunit retail packages that do not obscure the number of units nor prevent examination of the labeling on each of the individual units are not subject to this paragraph if the labeling of each individual unit complies with the requirements of paragraphs (f) and (i) of this section.

(t) Where the declaration of net quantity of contents is in terms of net weight and/or drained weight or volume and does not accurately reflect the actual quantity of the contents or the product falls below the applicable standard of fill of container because of equipment malfunction or otherwise unintentional product variation, and the label conforms in all other respects to the requirements of this chapter (except the requirement that food falling below the applicable standard of fill of container shall bear the general statement of substandard fill specified in § 564.14(b) of this chapter), the mislabeled food product, including any food product that fails to bear the general statement of substandard fill specified in § 564.14(b) of this chapter, may be sold by the manufacturer or processor directly to institutions operated by Federal, State or local governments: *Provided, That:*

(1) The purchaser shall sign a statement at the time of sale stating that he is aware that the product is mislabeled to include acknowledgement of the nature and extent of the mislabeling, e.g., “Actual net weight may be as low as —% below labeled quantity” and that any subsequent distribution by him of said product except for his own institutional use is unlawful. This statement shall be kept on file at the principal place of business of the manufacturer or processor for 2 years subsequent to the date of shipment of the product and shall be available to the Food and Drug Administration upon request.

(2) The product shall be labeled on the outside of its shipping container with the statement(s):

(i) When the variation concerns net weight and/or drained weight or volume—“Product Mislabeled. Actual net weight (drained weight or volume where appropriate) may be as low as —% below labeled quantity. This

Product Not for Retail Distribution,” the blank to be filled in with the maximum percentage variance between the labeled and actual weight or volume of contents of the individual packages in the shipping container, and

(ii) When the variation is in regard to a fill of container standard—“Product Mislabeled. Actual fill may be as low as —% below standard of fill. This Product Not for Retail Distribution.”

(3) The statements required by paragraphs (t)(2)(i) and (ii) of this section, which may be consolidated where appropriate, shall appear prominently and conspicuously as compared to other printed matter on the shipping container and in boldface print or type on a clear, contrasting background in order to render them likely to be read and understood by the purchaser under ordinary conditions of purchase.

[41 FR 38619, Sept. 10, 1976, as amended at 54 FR 18279, Apr. 28, 1989]

§501.110 Animal feed labeling; collective names for feed ingredients.

(a) An animal feed shall be exempt from the requirements of section 403(i)(2) of the act with respect to its label bearing the common or usual names of the animal feed ingredients listed in paragraph (b) of this section under the following prescribed conditions:

(1) The animal feed is intended solely for livestock and poultry.

(2) The label of the animal feed bears the collective name(s) prescribed in paragraph (b) of this section in lieu of the corresponding common or usual names of the individual feed ingredients contained therein.

(3) The label of the animal feed otherwise conforms to the requirements of section 403(i)(2) of the act.

(4) The ingredients of any feed listed in paragraph (b) of this section neither contain nor are food additives as defined in section 201(s) of the act unless provided for by and in conformity with applicable regulations established pursuant to section 409 of the act.

(b) Each collective name referred to in this paragraph may be used for the purpose of labeling where one or more of the ingredients listed for that collective name are present. The animal feed ingredients listed under each of the

collective names are the products defined by the Association of American Feed Control Officials. The collective names are as follows:

(1) *Animal protein products* include one or more of the following: Animal products, marine products, and milk products.

(2) *Forage products* include one or more of the following: Alfalfa meals, entire plant meals, hays, and stem meals.

(3) *Grain products* include one or more of the following: Barley, grain sorghums, maize (corn), oats, rice, rye, and wheat.

(4) *Plant protein products* include one or more of the following: Algae meals, coconut meals (copra), cottonseed meals, guar meal, linseed meals, peanut meals, safflower meals, soybean meals, sunflower meals, and yeasts.

(5) *Processed grain byproducts* include one or more of the following: Brans, brewers dried grains, distillers grains, distillers solubles, flours, germ meals, gluten feeds, gluten meals, grits, groats, hominy feeds, malt sprouts, middlings, pearled, polishings, shorts, and wheat mill run.

(6) *Roughage products* include one or more of the following: Cobs, hulls, husks, pulps, and straws.

PART 502—COMMON OR USUAL NAMES FOR NONSTANDARDIZED ANIMAL FOODS

Sec.

502.5 General principles.

502.19 Petitions.

AUTHORITY: 21 U.S.C. 321, 343, 371.

§502.5 General principles.

(a) The common or usual name of a food, which may be a coined term, shall accurately identify or describe, in as simple and direct terms as possible, the basic nature of the food or its characterizing properties or ingredients. The name shall be uniform among all identical or similar products and may not be confusingly similar to the name of any other food that is not reasonably encompassed within the same name. Each class or subclass of food shall be given its own common or usual name that states, in clear terms, what it is in

§ 502.5

21 CFR Ch. I (4–1–99 Edition)

a way that distinguishes it from different foods.

(b) The common or usual name of a food shall include the percentage(s) of any characterizing ingredient(s) or component(s) when the proportion of such ingredient(s) or component(s) in the food has a material bearing on price or consumer acceptance or when the labeling or the appearance of the food may otherwise create an erroneous impression that such ingredient(s) or component(s) is present in an amount greater than is actually the case. The following requirements shall apply unless modified by a specific regulation in this part.

(1) The percentage of a characterizing ingredient or component shall be declared on the basis of its quantity in the finished product (i.e., weight/weight in the case of solids, or volume/volume in the case of liquids).

(2) The percentage of a characterizing ingredient or component shall be declared by the words “containing (or contains) — percent (or %) —” or “— percent (or %) —” with the first blank filled in with the percentage expressed as a whole number not greater than the actual percentage of the ingredient or component named and the second blank filled in with the common or usual name of the ingredient or component. The word “containing” (or “contains”), when used, shall appear on a line immediately below the part of the common or usual name of the food required by paragraph (a) of this section. For each characterizing ingredient or component, the words “— percent (or %) —” shall appear following or directly below the word “containing” (or “contains”), or directly below the part of the common or usual name of the food required by paragraph (a) of this section when the word “containing” (or “contains”) is not used, in easily legible boldface print or type in distinct contrast to other printed or graphic matter, and in a height not less than the larger of the following alternatives:

(i) Not less than one-sixteenth inch in height on packages having a principal display panel with an area of 5 square inches or less and not less than one-eighth inch in height if the area of

the principal display panel is greater than 5 square inches; or

(ii) Not less than one-half the height of the largest type appearing in the part of the common or usual name of the food required by paragraph (a) of this section.

(c) The common or usual name of a food shall include a statement of the presence or absence of any characterizing ingredient(s) or component(s) and/or the need for the user to add any characterizing ingredient(s) or component(s) when the presence or absence of such ingredient(s) or component(s) in the food has a material bearing on price or consumer acceptance or when the labeling or the appearance of the food may otherwise create an erroneous impression that such ingredient(s) or component(s) is present when it is not, and consumers may otherwise be misled about the presence or absence of the ingredient(s) or component(s) in the food. The following requirements shall apply unless modified by a specific regulation in this part.

(1) The presence or absence of a characterizing ingredient or component shall be declared by the words “containing (or contains) —” or “containing (or contains) no —” or “no —” or “does not contain —,” with the blank being filled in with the common or usual name of the ingredient or component.

(2) The need for the user of a food to add any characterizing ingredient(s) or component(s) shall be declared by an appropriate informative statement.

(3) The statement(s) required under paragraph (c) (1) and/or (2) of this section shall appear following or directly below the part of the common or usual name of the food required by paragraphs (a) and (b) of this section, in easily legible boldface print or type in distinct contrast to other printed or graphic matter, and in a height not less than the larger of the alternatives established under paragraph (b)(2) (i) and (ii) of this section.

(d) A common or usual name of a food may be established by common

Food and Drug Administration, HHS

usage or by establishment of a regulation in this part, in a standard of identity, or in other regulations in this chapter.

[41 FR 38627, Sept. 10, 1976. Redesignated at 42 FR 14091, Mar. 15, 1977]

§ 502.19 Petitions.

(a) The Commissioner of Food and Drugs, either on his own initiative or on behalf of any interested person who has submitted a petition, may publish a proposal to issue, amend, or revoke, under this part, a regulation prescribing a common or usual name for a food, pursuant to part 10 of this chapter.

(b) If the principal display panel of a food for which a common or usual name regulation is established is too small to accommodate all mandatory requirements, the Commissioner may establish by regulation an acceptable alternative, e.g., a smaller type size. A petition requesting such a regulation, which would amend the applicable regulation, shall be submitted pursuant to part 10 of this chapter.

[42 FR 4716, Jan. 25, 1977; 42 FR 10980, Feb. 25, 1977. Redesignated at 42 FR 14091, Mar. 15, 1977, and amended at 42 FR 15675, Mar. 22, 1977; 42 FR 24254, May 13, 1977]

PART 509—UNAVOIDABLE CONTAMINANTS IN ANIMAL FOOD AND FOOD-PACKAGING MATERIAL

Subpart A—General Provisions

Sec.

509.3 Definitions and interpretations.

509.4 Establishment of tolerances, regulatory limits, and action levels.

509.5 Petitions.

509.6 Added poisonous or deleterious substances.

509.7 Unavoidability.

509.15 Use of polychlorinated biphenyls (PCB's) in establishments manufacturing food-packaging materials.

Subpart B—Tolerances for Unavoidable Poisonous or Deleterious Substances

509.30 Temporary tolerances for polychlorinated biphenyls (PCB's).

§ 509.4

Subpart C—Regulatory Limits for Added Poisonous or Deleterious Substances [Reserved]

Subpart D—Naturally Occurring Poisonous or Deleterious Substances [Reserved]

AUTHORITY: 21 U.S.C. 336, 342, 346, 346a, 348, 371.

SOURCE: 42 FR 52821, Sept. 30, 1977, unless otherwise noted.

Subpart A—General Provisions

§ 509.3 Definitions and interpretations.

(a) *Act* means the Federal Food, Drug, and Cosmetic Act.

(b) The definitions of terms contained in section 201 of the act are applicable to such terms when used in this part unless modified in this section.

(c) A *naturally occurring poisonous or deleterious substance* is a poisonous or deleterious substance that is an inherent natural constituent of a food and is not the result of environmental, agricultural, industrial, or other contamination.

(d) An *added poisonous or deleterious substance* is a poisonous or deleterious substance that is not a naturally occurring poisonous or deleterious substance. When a naturally occurring poisonous or deleterious substance is increased to abnormal levels through mishandling or other intervening acts, it is an added poisonous or deleterious substance to the extent of such increase.

(e) *Food* includes pet food, animal feed, and substances migrating to food from food-contact articles.

§ 509.4 Establishment of tolerances, regulatory limits, and action levels.

(a) When appropriate under the criteria of § 509.6, a tolerance for an added poisonous or deleterious substance, which may be a food additive, may be established by regulation in subpart B of this part under the provisions of section 406 of the act. A tolerance may prohibit any detectable amount of the substance in food.

(b) When appropriate under the criteria of § 509.6, and under section 402(a)(1) of the act, a regulatory limit for an added poisonous or deleterious

§ 509.5

substance, which may be a food additive, may be established by regulation in subpart C of this part under the provisions of sections 402(a)(1) and 701(a) of the act. A regulatory limit may prohibit any detectable amount of the substance in food. The regulatory limit established represents the level at which food is adulterated within the meaning of section 402(a)(1) of the act.

(c)(1) When appropriate under the criteria of § 509.6, an action level for an added poisonous or deleterious substance, which may be a food additive, may be established to define a level of contamination at which a food may be regarded as adulterated.

(2) Whenever an action level is established or changed, a notice shall be published in the FEDERAL REGISTER as soon as practicable thereafter. The notice shall call attention to the material supporting the action level which shall be on file with the Dockets Management Branch before the notice is published. The notice shall invite public comment on the action level.

(d) A regulation may be established in subpart D of this part to identify a food containing a naturally occurring poisonous or deleterious substance which will be deemed to be adulterated under section 402(a)(1) of the act. These regulations do not constitute a complete list of such foods.

[42 FR 52821, Sept. 30, 1977, as amended at 55 FR 20786, May 21, 1990]

§ 509.5 Petitions.

The Commissioner of Food and Drugs, either on his own initiative or on behalf of any interested person who has submitted a petition, may issue a proposal to establish, revoke, or amend a regulation under this part. Any such petition shall include an adequate factual basis to support the petition, shall be in the form set forth in § 10.30 of this chapter, and will be published in the FEDERAL REGISTER for comment if it contains reasonable grounds for the proposed regulation.

[42 FR 52821, Sept. 30, 1977, as amended at 54 FR 18280, Apr. 28, 1989]

21 CFR Ch. I (4–1–99 Edition)

§ 509.6 Added poisonous or deleterious substances.

(a) Use of an added poisonous or deleterious substance, other than a pesticide chemical, that is also a food additive will be controlled by a regulation issued under section 409 of the act when possible. When such a use cannot be approved under the criteria of section 409 of the act, or when the added poisonous or deleterious substance is not a food additive, a tolerance, regulatory limit, or action level may be established pursuant to the criteria in paragraphs (b), (c), or (d) of this section. Residues resulting from the use of an added poisonous or deleterious substance that is also a pesticide chemical will ordinarily be controlled by a tolerance established in a regulation issued under sections 406, 408, or 409 of the act by the U.S. Environmental Protection Agency (EPA). When such a regulation has not been issued, an action level for an added poisonous or deleterious substance that is also a pesticide chemical may be established by the Food and Drug Administration. The Food and Drug Administration will request EPA to recommend such an action level pursuant to the criteria established in paragraph (d) of this section.

(b) A tolerance for an added poisonous or deleterious substance in any food may be established when the following criteria are met:

(1) The substance cannot be avoided by good manufacturing practice.

(2) The tolerance established is sufficient for the protection of the public health, taking into account the extent of which the presence of the substance cannot be avoided and the other ways in which the consumer may be affected by the same or related poisonous or deleterious substances.

(3) No technological or other changes are foreseeable in the near future that might affect the appropriateness of the tolerance established. Examples of changes that might affect the appropriateness of the tolerance include anticipated improvements in good manufacturing practice that would change the extent to which use of the substance is unavoidable and anticipated

studies expected to provide significant new toxicological or use data.

(c) A regulatory limit for an added poisonous or deleterious substance in any food may be established when each of the following criteria is met:

(1) The substance cannot be avoided by current good manufacturing practices.

(2) There is no tolerance established for the substance in the particular food under sections 406, 408, or 409 of the act.

(3) There is insufficient information by which a tolerance may be established for the substance under section 406 of the act or technological changes appear reasonably possible that may affect the appropriateness of a tolerance. The regulatory limit established represents the level at which food is adulterated within the meaning of section 402(a)(1) of the act.

(d) An action level for an added poisonous or deleterious substance in any food may be established when the criteria in paragraph (b) of this section are met, except that technological or other changes that might affect the appropriateness of the tolerance are foreseeable in the near future. An action level for an added poisonous or deleterious substance in any food may be established at a level at which the Food and Drug Administration may regard the food as adulterated within the meaning of section 402(a)(1) of the act, without regard to the criteria in paragraph (b) of this section or in section 406 of the act. An action level will be withdrawn when a tolerance or regulatory limit for the same substance and use has been established.

(e) Tolerances will be established under authority appropriate for action levels (sections 306, 402(a), and 701(a) of the act, together with section 408 or 409 of the act, if appropriate) as well as under authority appropriate for tolerances (sections 406 and 701 of the act). In the event the effectiveness of a tolerance is stayed pursuant to section 701(e)(2) of the act by the filing of an objection, the order establishing the tolerance shall be deemed to be an order establishing an action level until final action is taken upon such objection.

[42 FR 52821, Sept. 30, 1977, as amended at 55 FR 20786, May 21, 1990]

§ 509.7 Unavoidability.

(a) Tolerances and action levels in this part are established at levels based on the unavoidability of the poisonous or deleterious substance concerned and do not establish a permissible level of contamination where it is avoidable.

(b) Compliance with tolerances, regulatory limits, and action levels does not excuse failure to observe either the requirement in section 402(a)(4) of the act that food may not be prepared, packed, or held under insanitary conditions or the other requirements in this chapter that food manufacturers must observe current good manufacturing practices. Evidence obtained through factory inspection or otherwise indicating such a violation renders the food unlawful, even though the amounts of poisonous or deleterious substances are lower than the currently established tolerances, regulatory limits, or action levels. The manufacturer of food must at all times utilize quality control procedures which will reduce contamination to the lowest level currently feasible.

[42 FR 52821, Sept. 30, 1977, as amended at 55 FR 20786, May 21, 1990]

§ 509.15 Use of polychlorinated biphenyls (PCB's) in establishments manufacturing food-packaging materials.

(a) Polychlorinated biphenyls (PCB's) represent a class of toxic industrial chemicals manufactured and sold under a variety of trade names, including: Aroclor (United States); Phenoclor (France); Colphen (Germany); and Kanaclor (Japan). PCB's are highly stable, heat resistant, and nonflammable chemicals. Industrial uses of PCB's include, or did include in the past, their use as electrical transformer and capacitor fluids, heat transfer fluids, hydraulic fluids, and plasticizers, and in formulations of lubricants, coatings, and inks. Their unique physical and chemical properties and widespread, uncontrolled industrial applications have caused PCB's to be a persistent and ubiquitous contaminant in the environment, causing the contamination of certain foods. In addition, incidents have occurred in which PCB's have directly contaminated animal feeds as a result of industrial accidents (leakage

§ 509.30

or spillage of PCB fluids from plant equipment). These accidents in turn caused the contamination of food products intended for human consumption (meat, milk and eggs). Investigations by the Food and Drug Administration have revealed that a significant percentage of paper food-packaging material contains PCB's which can migrate to the packaged food. The origin of PCB's in such material is not fully understood. Reclaimed fibers containing carbonless copy paper (contains 3 to 5 percent PCB's) have been identified as a primary source of PCB's in paper products. Some virgin paper products have also been found to contain PCB's, the source of which is generally attributed to direct contamination from industrial accidents from the use of PCB-containing equipment and machinery in food-packaging manufacturing establishments. Since PCB's are toxic chemicals, the PCB contamination of food-packaging materials as a result of industrial accidents, which can cause the PCB contamination of food, represents a hazard to public health. It is therefore necessary to place certain restrictions on the industrial uses of PCB's in establishments manufacturing food-packaging materials.

(b) The following special provisions are necessary to preclude the accidental PCB contamination of food-packaging materials:

(1) New equipment or machinery for manufacturing food-packaging materials shall not contain or use PCB's.

(2) On or before September 4, 1973, the management of establishments manufacturing food-packaging materials shall:

(i) Have the heat exchange fluid used in existing equipment for manufacturing food-packaging materials sampled and tested to determine whether it contains PCB's or verify the absence of PCB's in such formulations by other appropriate means. On or before Sept. 4, 1973, any such fluid formulated with PCB's must to the fullest extent possible commensurate with current good manufacturing practices be replaced with a heat exchange fluid that does not contain PCB's.

(ii) Eliminate to the fullest extent possible commensurate with current good manufacturing practices from the

21 CFR Ch. I (4-1-99 Edition)

establishment any other PCB-containing equipment, machinery and materials wherever there is a reasonable expectation that such articles could cause food-packaging materials to become contaminated with PCB's either as a result of normal use or as a result of accident, breakage, or other mishap.

(iii) The toxicity and other characteristics of fluids selected as PCB replacements must be adequately determined so that the least potentially hazardous replacement is used. In making this determination with respect to a given fluid, consideration should be given to (a) its toxicity; (b) the maximum quantity that could be spilled onto a given quantity of food before it would be noticed, taking into account its color and odor; (c) possible signaling devices in the equipment to indicate a loss of fluid, etc.; and (d) its environmental stability and tendency to survive and be concentrated through the food chain. The judgment as to whether a replacement fluid is sufficiently non-hazardous is to be made on an individual installation and operation basis.

(c) The provisions of this section do not apply to electrical transformers and condensers containing PCB's in sealed containers.

Subpart B—Tolerances for Unavoidable Poisonous or Deleterious Substances

§ 509.30 Temporary tolerances for polychlorinated biphenyls (PCB's).

(a) Polychlorinated biphenyls (PCB's) are toxic, industrial chemicals. Because of their widespread, uncontrolled industrial applications, PCB's have become a persistent and ubiquitous contaminant in the environment. As a result, certain foods and animal feeds, principally those of animal and marine origin, contain PCB's as unavoidable, environmental contaminants. PCB's are transmitted to the food portion (meat, milk, and eggs) of food producing animals ingesting PCB contaminated animal feed. In addition, a significant percentage of paper food-packaging materials contain PCB's which may migrate to the packaged food. The source of PCB's in paper food-packaging materials is primarily of certain types of carbonless copy

paper (containing 3 to 5 percent PCB's) in waste paper stocks used for manufacturing recycled paper. Therefore, temporary tolerances for residues of PCB's as unavoidable environmental or industrial contaminants are established for a sufficient period of time following the effective date of this paragraph to permit the elimination of such contaminants at the earliest practicable time. For the purposes of this paragraph, the term *polychlorinated biphenyls* (PCB's) is applicable to mixtures of chlorinated biphenyl compounds, irrespective of which mixture of PCB's is present as the residue. The temporary tolerances for residues of PCB's are as follows:

(1) 0.2 part per million in finished animal feed for food-producing animals (except the following finished animal feeds: feed concentrates, feed supplements, and feed premixes).

(2) 2 parts per million in animal feed components of animal origin, including fishmeal and other by-products of marine origin and in finished animal feed concentrates, supplements, and premixes intended for food-producing animals.

(3) 10 parts per million in paper food-packaging material intended for or used with finished animal feed and any components intended for animal feeds. The tolerance shall not apply to paper food-packaging material separated from the food therein by a functional barrier which is impermeable to migration of PCB's.

(b) A compilation entitled "Analytical Methodology for Polychlorinated Biphenyls, February 1973" for determining compliance with the tolerances established in this section is available from the Dockets Management Branch, Food and Drug Administration, rm. 1-23, 12420 Parklawn Dr., Rockville, MD 20857.

[42 FR 52821, Sept. 30, 1977, as amended at 46 FR 8460, Jan. 27, 1981; 59 FR 14365, Mar. 28, 1994]

Subpart C—Regulatory Limits for Added Poisonous or Deleterious Substances [Reserved]

Subpart D—Naturally Occurring Poisonous or Deleterious Substances [Reserved]

PART 510—NEW ANIMAL DRUGS

Subpart A—General Provisions

Sec.

510.3 Definitions and interpretations.

510.4 Biologics; products subject to license control.

510.7 Consignees of new animal drugs for use in the manufacture of animal feed.

510.45 Packaging requirements for drugs for animal use.

510.95 Designated journals.

Subpart B—Specific Administrative Rulings and Decisions

510.105 Labeling of drugs for use in milk-producing animals.

510.106 Labeling of antibiotic and antibiotic-containing drugs intended for use in milk-producing animals.

510.110 Antibiotics used in food-producing animals.

510.112 Antibiotics used in veterinary medicine and for nonmedical purposes; required data.

Subpart C [Reserved]

Subpart D—Records and Reports

510.300 Records and reports concerning experience with new animal drugs for which an approved application is in effect.

510.301 Records and reports concerning experience with animal feeds bearing or containing new animal drugs for which an approved application is in effect.

510.302 Reporting forms.

510.305 Maintenance of copies of approved applications for animal feed bearing or containing new animal drugs.

Subpart E—Requirements for Specific New Animal Drugs

510.410 Corticosteroids for oral, injectable, and ophthalmic use in animals; warnings and labeling requirements.

§ 510.3

21 CFR Ch. I (4–1–99 Edition)

510.440 Injectable iron preparations.

510.455 New animal drug requirements regarding free-choice administration in feeds.

Subpart F—Animal Use Exemptions From Certification and Labeling Requirements

510.515 Animal feeds bearing or containing new animal drugs subject to the provisions of section 512(n) of the act.

Subpart G—Sponsors of Approved Applications

510.600 Names, addresses, and drug labeler codes of sponsors of approved applications.

AUTHORITY: 21 U.S.C. 321, 331, 351, 352, 353, 360b, 371, 379e.

SOURCE: 40 FR 13807, Mar. 27, 1975, unless otherwise noted.

Subpart A—General Provisions

§ 510.3 Definitions and interpretations.

As used in this part:

(a) The term *act* means the Federal Food, Drug, and Cosmetic Act, as amended (secs. 201–902, 52 Stat. 1040 *et seq.*, as amended; 21 U.S.C. 321–392).

(b) *Department* means the Department of Health and Human Services.

(c) *Secretary* means the Secretary of Health and Human Services.

(d) *Commissioner* means the Commissioner of Food and Drugs.

(e) *Person* means individuals, partnerships, corporations, and associations.

(f) The definitions and interpretations of terms contained in section 201 of the act shall be applicable to such terms when used in the regulations in this part.

(g) The term *new animal drug* means any drug intended for use for animals other than man, including any drug intended for use in animal feed but not including such animal feed:

(1) The composition of which is such that such drug is not generally recognized, among experts qualified by scientific training and experience to evaluate the safety and effectiveness of animal drugs, as safe and effective for use under the conditions prescribed, recommended, or suggested in the labeling thereof; except that such a drug not so recognized shall not be deemed to be a *new animal drug* if at any time prior to June 25, 1938, it was subject to

the Food and Drug Act of June 30, 1906, as amended, and if at such time its labeling contained the same representations concerning the conditions of its use; or

(2) The composition of which is such that such drug, as a result of investigations to determine its safety and effectiveness for use under such conditions, has become so recognized but which has not, otherwise than in such investigations, been used to a material extent or for a material time under such conditions.

(h) The term *animal feed* means an article which is intended for use for food for animals other than man and which is intended for use as a substantial source of nutrients in the diet of the animal, and is not limited to a mixture intended to be the sole ration of the animal.

(i) The newness of an animal drug, including a new animal drug intended for use in or on animal feed, may arise by reason of: (1) The newness for its intended drug use of any substance of which the drug is comprised, in whole or in part, whether it be an active substance or a menstruum, excipient, carrier, coating, or other component; (2) the newness for its intended drug use of a combination of two or more substances, none of which is itself a new animal drug; (3) the newness for its intended drug use of the proportion of a substance in a combination, even though such combination containing such substance in other proportion is not a new animal drug; (4) the newness for its intended drug use in a different species of animal; (5) the newness of its intended drug use in diagnosing, curing, mitigating, treating, or preventing a disease, or to affect a structure or function of the animal body, even though such drug is not a new animal drug when used in another disease or to affect another structure or function of the body; or (6) the newness of a dosage, or method or duration of administration or application, or any other condition of use prescribed, recommended, or suggested in the labeling of such drug, even though such drug or animal feed containing such drug when used in another dosage, or another method or duration of administration

Food and Drug Administration, HHS

§ 510.95

or application, or different condition, is not a new animal drug.

(j) *Animals used only for laboratory research and laboratory research animals* mean individual animals or groups of animals intended for use and used solely for laboratory research purposes, regardless of species, and does not include animals intended to be used for any food purposes or animals intended to be kept as livestock.

(k) The term *sponsor* means the person responsible for an investigation of a new animal drug, including responsibility for compliance with applicable provisions of the act and regulations. The *sponsor* may be an individual, partnership, corporation, or Government agency or may be a manufacturer, scientific institution, or an investigator regularly and lawfully engaged in the investigation of new animal drugs.

(l) *Designated journal(s)* means journals listed in § 510.95.

[40 FR 13807, Mar. 27, 1975, as amended at 50 FR 7517, Feb. 22, 1985; 54 FR 22741, May 26, 1989]

§ 510.4 Biologics; products subject to license control.

An animal drug produced and distributed in full conformance with the animal virus, serum, and toxin law of March 4, 1913 (37 Stat. 832; 21 U.S.C. 151 *et seq.*) and any regulations issued thereunder shall not be deemed to be subject to section 512 of the Federal Food, Drug, and Cosmetic Act.

§ 510.7 Consignees of new animal drugs for use in the manufacture of animal feed.

(a) A new animal drug intended for use in the manufacture of animal feed shall be deemed to be unsafe unless at the time of its removal from the establishment of a manufacturer, packer, or distributor of such drug, such manufacturer, packer, or distributor has an unrevoked written statement from the consignee of such drug, or a notice from the Secretary, to the effect that with respect to the use of such drug in animal feed the consignee:

(1) Is the holder of an approved application under § 514.2 of this chapter; or

(2) Will, if the consignee is not a user of the drug, ship such drug only to a

holder of an approved application under § 514.2 of this chapter.

(b) The requirements of paragraph (a) of this section do not apply:

(1) Where such drugs are intended for export and/or

(2) When the use of such drug in the manufacture of a finished feed has been exempted from the requirements of section 512(m) of the act under the conditions specified by regulations published in part 558 of this chapter.

§ 510.45 Packaging requirements for drugs for animal use.

The packaging requirements for antibiotic drugs for veterinary use are described under § 432.1 of this chapter, except that antibiotic drugs for veterinary use need not be packaged for dispensing in containers of colorless, transparent glass.

EFFECTIVE DATE NOTE: At 64 FR 403, Jan. 5, 1999, § 510.45 was removed, effective May 20, 1999.

§ 510.95 Designated journals.

The following journals are available to the Food and Drug Administration and thus permit waiving of the submission of reprints and summaries covering reports contained in these journals to the extent that such requirements are waived in the regulations in this part:

All Pet's Magazine (Jersey City).
American Journal of Veterinary Research (Chicago).
Animal Health (Journal of the Animal Health Trust) (London).
Animal Nutrition & Health (Sausalito, CA).
Animal Production (Edinburgh).
Avian Diseases (Amherst).
British Poultry Science (Edinburgh).
Canadian Journal of Comparative Medicine and Veterinary Science (Gardenvale, Quebec).
Canadian Veterinary Journal (Guelph, Ontario).
Cornell Veterinarian (Ithaca).
Experimental Parasitology (New York).
The Feed Bag (Milwaukee).
Feedstuffs (Minneapolis).
Hoard's Dairyman (Fort Atkinson).
Journal of the American Veterinary Medical Association (Chicago).
Journal of Animal Science (Albany).
Journal of Dairy Science (Champaign).
Journal of Economic Entomology (Baltimore).
Journal of Small Animal Practice (London).

§ 510.105

Modern Veterinary Practice (formerly North American Veterinarian) (Wheaton, IL).
National Hog Farmer (Grundy Center, IA).
New Zealand Veterinary Journal (Wellington).
Poultry Science (Guelph, Ontario).
Praktische Tierarzt (Postfach, Germany).
Research in Veterinary Science (Chicago).
Small Animal Clinician (Kansas City, MO).
Veterinaermedizin (Konstanz, Germany).
Veterinarian (London).
Veterinarian (International) (New York).
The Veterinary Bulletin (Farnham Royal, England).
Veterinary Medicine (Kansas City, MO).
Veterinary Record (Croydon, England).
Zentralblatt Fuer Veterinaermedizin Zentr. Veterinaermed (Berlin).

[40 FR 13807, Mar. 27, 1975, as amended at 50 FR 7517, Feb. 22, 1985]

Subpart B—Specific Administrative Rulings and Decisions

§ 510.105 Labeling of drugs for use in milk-producing animals.

(a) Part 540 of this chapter provides for new animal drugs intended for intramammary use in animals and includes conditions of use intended to prevent the contamination of milk from the use of such drugs.

(b) Preparations containing antibiotics and other potent drugs labeled with directions for use in milk-producing animals will be misbranded under section 502(f)(2) of the act unless their labeling bears appropriate warnings and directions for use to avoid adulteration of milk under section 402(a)(2)(D) of the act.

(c) It is the position of the Food and Drug Administration that the labeling for such preparations should bear a clear warning that either:

(1) The article should not be administered to animals producing milk, since to do so would result in contamination of the milk; or

(2) The label should bear the following statement: “Warning: Milk that has been taken from animals during treatment and for _____ hours after the latest treatment must not be used for food”, the blank being filled in with the figure that the manufacturer has determined by appropriate investigation is needed to insure that the milk will not carry violative residues resulting from use of the preparation. If the use of the preparation as recommended

21 CFR Ch. I (4–1–99 Edition)

does not result in contamination of the milk, neither of the above warning statements is required.

[40 FR 13807, Mar. 27, 1975, as amended at 63 FR 32980, June 17, 1998]

§ 510.106 Labeling of antibiotic and antibiotic-containing drugs intended for use in milk-producing animals.

Whenever the labeling of an antibiotic drug included in the regulations in this chapter suggests or recommends its use in milk-producing animals, the label of such drugs shall bear either the statement “Warning: Not for use in animals producing milk, since this use will result in contamination of the milk” or the statement “Warning: Milk that has been taken from animals during treatment and for _____ hours after the latest treatment must not be used for food”, the blank being filled in with the figure that the Commissioner has authorized the manufacturer of the drug to use. The Commissioner shall determine what such figures shall be from information submitted by the manufacturer and which the Commissioner considers is adequate to prove that period of time after the latest treatment that the milk from treated animals will contain no violative residues from use of the preparation. If the Commissioner determines from the information submitted that the use of the antibiotic drug as recommended does not result in its appearance in the milk, the Commissioner may exempt the drug from bearing either of the above warning statements.

[63 FR 32980, June 17, 1998]

§ 510.110 Antibiotics used in food-producing animals.

(a) The Food and Drug Administration in the interest of fulfilling its responsibilities with regard to protection of the public health has requested an evaluation of the public health aspects of the use of antibiotics in veterinary medical and nonmedical uses. There is particular concern with regard to the potential hazards associated with the extensive use of antibiotics administered to food-producing animals. Accordingly, an ad hoc committee on the Veterinary Medical and Nonmedical Uses of Antibiotics was established by the Food and Drug Administration to

Food and Drug Administration, HHS

§ 510.112

study and advise the Commissioner of Food and Drugs on the uses of antibiotics in veterinary medicine and for various nonmedical purposes as such uses may affect the enforcement of the Federal Food, Drug, and Cosmetic Act with respect to their safety and effectiveness.

(b) Based upon an evaluation of the conclusions of said Committee and other relevant material, § 510.112 was published in the FEDERAL REGISTER of August 23, 1966 (31 FR 11141), asking sponsors of drugs containing any antibiotic intended for use in food-producing animals to submit data to establish whether such antibiotic and its metabolites are present as residues in edible tissues, milk, and eggs from treated animals. The data on the residues of antibiotics in milk from intramammary infusion preparations were requested within 60 days and the data on all other products were requested within 180 days following the date of publication of § 510.112 in the FEDERAL REGISTER.

(c) An evaluation of the data now available shows that use of many antibiotic preparations cause residues in edible products of treated animals for varying and, in some cases, for long periods of time following the last administration. Because of the accumulation of new information with regard to the development of resistance of bacteria to antibiotics, the ability of bacteria to transfer this resistance, and the development of sensitivity to antibiotics in humans, unauthorized and unsafe residues of antibiotics cannot be permitted in food obtained from treated animals.

(d) Based on evaluation of information available, including the conclusions of the aforementioned ad hoc Committee, the Commissioner concludes that antibiotic preparations intended for use in food-producing animals, other than topical and ophthalmic preparations, are not generally recognized among qualified experts as having been shown to be safe for their intended use(s) within the meaning of section 201(s) of the Federal Food, Drug, and Cosmetic Act.

(e) Therefore, all exemptions from the provisions of section 409 of the act for use of antibiotics in food-producing animals based on sanctions or approv-

als granted prior to enactment of the Food Additives Amendment of 1958 (Pub. L. 85-929; 72 Stat. 1784) will be revoked and the uses which are concluded to be safe will be covered by food additive regulations. On those products for which there are inadequate residue data, actions will be initiated to withdraw approval of new-drug applications under the provisions of section 505 of the act. Antibiotic preparations, other than those for topical and ophthalmic application in food-producing animals, which are not covered by food additive regulations will be subject to regulatory action within 180 days after publication of the forthcoming revocation order.

(f) Because of the variation in the period of time that antibiotic residues may remain in edible products from treated animals, all injectable, intramammary infusion, intrauterine, and oral preparations, including medicated premixes intended for use in food-producing animals, are deemed to be new drugs as well as food additives.

[40 FR 13807, Mar. 27, 1975, as amended at 54 FR 18280, Apr. 28, 1989; 64 FR 403, Jan. 5, 1999]

EFFECTIVE DATE NOTE: At 64 FR 403, Jan. 5, 1999, § 510.110 was amended by removing the phrase "to amend or revoke antibiotic regulations under the provisions of section 507 of the act, or" from the second sentence of paragraph (e), by removing the phrase "(except certifiable antibiotics)" from the first sentence of paragraph (f) and by removing the last sentence of paragraph (f), effective May 20, 1999. For the convenience of the user, the superseded text is set forth as follows:

§ 510.110 Antibiotics used in food-producing animals.

* * * * *

(f) * * * An antibiotic application (see § 431.50 of this chapter) will be required for all medicated premixes containing certifiable antibiotics.

§ 510.112 Antibiotics used in veterinary medicine and for nonmedical purposes; required data.

(a) An ad hoc committee, Committee on the Veterinary Medical and Non-medical Uses of Antibiotics, was formed by the Food and Drug Administration to study, and advise the Commissioner on, the use of antibiotics in veterinary medicine and for various

§ 510.300

nonmedical purposes as such uses may affect the enforcement of the Federal Food, Drug, and Cosmetic Act with respect to the safety and effectiveness of such substances. A copy of the report may be obtained from the Food and Drug Administration, Office of Public Affairs, Room 15-05, Parklawn Building, 5600 Fishers Lane, Rockville, MD 20857.

(b) On the basis of the report of the Committee and other information, sponsors of drugs containing any antibiotic intended for use in food-producing animals shall submit data for determining whether or not such antibiotics and their metabolites are present as residues in edible tissues, milk, and eggs from treated animals; however, in the case of a drug for which such data have already been submitted and for which a regulation has been promulgated under section 409 of the act, only such data as has been accumulated since the issuance of the regulation need be submitted.

(c) The required data shall be submitted within 180 days of the date of publication of this section in the FEDERAL REGISTER; except that in the case of data on intramammary infusion preparations the data shall be submitted within 60 days of such publication. Data demonstrating the absence in milk of residues of intramammary infusion preparations when used as directed in their labeling are needed within the 60-day period because of the importance of milk in the human diet.

(d) Regulatory proceedings including revocation of prior sanctions, or actions to suspend or amend new drug or antibiotic approvals granted prior to passage of the Food Additives Amendment of 1958 (72 Stat. 1784), may be initiated with regard to the continued marketing of any antibiotic preparation on which the required information is not submitted within the period of time prescribed by paragraph (c) of this section.

(e) Questions relating to the acceptability of proposed research protocols and assay methods for determining the amount of antibiotic residues in food should be directed to the Director, Center for Veterinary Medicine, Food and

21 CFR Ch. I (4-1-99 Edition)

Drug Administration, 7500 Standish Pl., Rockville, MD 20855.

[40 FR 13807, Mar. 27, 1975, as amended at 46 FR 8460, Jan. 27, 1981; 54 FR 18280, Apr. 28, 1989; 57 FR 6475, Feb. 25, 1992]

Subpart C [Reserved]

Subpart D—Records and Reports

§ 510.300 Records and reports concerning experience with new animal drugs for which an approved application is in effect.

(a) On receiving notification that an application submitted pursuant to § 514.1 of this chapter for a new animal drug is approved, the applicant shall establish and maintain such records and make such reports as are specified in this section to facilitate a determination as to whether there may be grounds for suspending or withdrawing approval of the application or whether any applicable regulation should be amended or repealed. The applicant shall maintain adequately organized and indexed files containing full reports of information pertinent to the safety or effectiveness of the new animal drug that have not previously been submitted as part of his application for the drug and which are received or otherwise obtained by him from any source, as follows:

(1) Unpublished reports of clinical or other animal experience, studies, investigations, and tests conducted by the applicant or reported to him by any person involving the new animal drug that is the subject of the application or any related drugs. An adequate summary and bibliography of reports in the scientific literature would ordinarily suffice. (The application must identify at the time of each report submission, each drug he considers related to the subject drug.)

(2) Experience, investigations, studies, or tests involving the chemical or physical properties or any other properties of the new animal drug, such as its behavior or properties in relation to microorganisms, including both the effects of the drug on microorganisms and the effect of microorganisms on the drug.

(3) For information required by this section, adequate identification of its

source, when known, including the name and post office address of the person who furnishes such information.

(4) Copies of all mailing pieces and other labeling, and, if it is a prescription new animal drug, all advertising other than that contained in the application used in promoting the drug, and copies of the currently used package labeling that gives full information for use of the drug whether or not such labeling is contained in the application.

(5) Information concerning the quantity of the new animal drug distributed in a manner and form that facilitates estimates of the incidence of any adverse effects reported to be associated with the use of the drug. This does not require disclosure of financial, pricing, or sales data.

(6) Information concerning any previously unreported changes from the conditions described in an application conforming to the conditions of § 514.8(a)(5) of this chapter.

(b) The applicant shall submit to the Food and Drug Administration copies of the records and reports described in paragraph (a) of this section, except routine assay and control records, appropriately identified with the new animal drug application(s) to which they relate, as follows:

(1) Immediately upon receipt by the applicant, complete records or reports covering information of the following kinds:

(i) Information concerning a mixup in the new animal drug or its labeling with another article.

(ii) Information concerning any bacteriological or significant physical or other change or deterioration in the new animal drug, or any failure of one or more distributed batches of the drug to meet the specifications established for it in the new animal drug application.

(2) As soon as possible, and in any event within 15 working days of its receipt by the applicant, complete records of reports concerning any information of the following kinds:

(i) Information concerning any unexpected side effects, injury, toxicity, or sensitivity reaction or any unexpected incidence or severity thereof associated with clinical use, studies, investigations, or tests, whether or not de-

termined to be attributable to the new animal drug, except that this requirement shall not apply to the submission of information described in a written communication to the applicant from the Food and Drug Administration as types of information that may be submitted at other designated intervals. *Unexpected* as used in this subdivision refers to conditions or developments not previously submitted as part of the new animal drug application, or conditions and developments occurring at a rate higher than that shown by information previously submitted as part of the application.

(ii) Information concerning any unusual failure of the new animal drug to exhibit its expected pharmacological activities.

(3) When mailing pieces, any other labeling, and advertising are devised for promotion of the new animal drug, specimens shall be submitted at the time of initial dissemination of such labeling and at the time of initial publication of any advertisement for a prescription drug. Mailing pieces and labeling designed to contain samples of a drug shall be complete except for the omission of the drug.

(4) All the kinds of information described in paragraph (a) of this section, other than that submitted under the provisions of paragraphs (b) (1), (2), and (3) of this section, shall be submitted as follows unless otherwise ordered in a written communication from the Commissioner:

(i) At intervals within 6 months beginning with the date of approval of the new animal drug application during the first year following such date, and at yearly intervals thereafter.

(ii) Whenever an applicant is required to submit reports under the provisions of paragraph (b)(4)(i) of this section with respect to more than one approved application for preparations containing the same new animal drug so that the same item(s) of information is (are) required to be reported for more than one application, he may elect to submit as a part of the report for one such application all the information common to such applications in lieu of reporting separately and repetitively on each. The applicant shall state when this is done and identify all

§ 510.301

21 CFR Ch. I (4–1–99 Edition)

the new animal drug applications for which the reports are submitted.

(iii) The submitted copies of records and reports shall include all the required information that was received or otherwise obtained by the applicant during the designated intervals.

(5) On written order of the Commissioner, within the time stated in such order or agreed to by the applicant and the Commissioner, any designated records or reports containing the kinds of information described in this section shall be submitted.

(c) The applicant shall, upon request of any properly authorized officer or employee of the Department at reasonable times, permit such officers to have access to and copy and verify any records and reports established and maintained under the provisions of this section.

(d) If the Food and Drug Administration finds that the applicant has failed to establish a system for maintaining required records or has repeatedly or deliberately failed to maintain such records or to make required reports in accordance with the provisions of this section, or that the applicant has refused to permit access to or copying of, or verification of such records or reports, the Commissioner shall give the applicant notice and opportunity for a hearing on the question of whether to withdraw the approval of the application, as provided in § 514.200 of this chapter.

(e) Upon written request of the applicant stating reasonable grounds therefor, the Commissioner will make available any information in possession of the Food and Drug Administration of the kinds the applicant is required to maintain under the provisions of this section, except information readily available to the applicant from other sources or information which the Commissioner concludes is confidential.

(f) The *applicant* required to establish and maintain records and make reports required by this section includes any person whose name appears on the labeling of the drug as its manufacturer, packer, or distributor under an approval or who is engaged in the manufacturing, processing, packing, or labeling of the drug under an approval of the new animal drug application or any

supplement to it; however, to avoid unnecessary duplication in the submission of reports, any such applicant's obligation to submit a report may be met by its submission on his behalf, designated as such, by another person responsible for reporting.

§ 510.301 Records and reports concerning experience with animal feeds bearing or containing new animal drugs for which an approved application is in effect.

Records and reports of clinical and other experience with the new animal drug will be maintained and reported, appropriately identified with the new animal drug application(s) to which they relate, to the Center for Veterinary Medicine in duplicate in accordance with the following:

(a) Immediately upon receipt by the applicant, complete records or reports covering information of the following kinds:

(1) Information concerning any mixup in the new animal drug or its labeling with another article.

(2) Information concerning any bacteriological, or any significant chemical, physical, or other change or deterioration in the drug, or any failure of one or more distributed batches of the drug to meet the specifications established for it in the new animal drug application.

(b) As soon as possible, and in any event within 15 working days of its receipt by the applicant, complete records or reports concerning any information of the following kinds:

(1) Information concerning any unexpected side effect, injury, toxicity, or sensitivity reaction or any unexpected incidence or severity thereof associated with clinical uses, studies, investigations, or tests, whether or not determined to be attributable to the new animal drug, except that this requirement shall not apply to the submission of information described in a written communication to the applicant from the Food and Drug Administration as types of information that may be submitted at other designated intervals. *Unexpected* as used in this paragraph refers to conditions or developments not previously submitted as part of the new animal drug application or not encountered during clinical trials of the

Food and Drug Administration, HHS

§ 510.410

drug, or conditions or developments occurring at a rate higher than shown by information previously submitted as part of the new animal drug application or at a rate higher than encountered during such clinical trials.

(2) Information concerning any unusual failure of the new animal drug to exhibit its expected pharmacological activity.

[40 FR 13807, Mar. 27, 1975, as amended at 54 FR 18280, Apr. 28, 1989]

§ 510.302 Reporting forms.

(a) The information described in § 510.300, except that described in paragraphs (b) (1) and (2) of that section, shall be submitted appropriately identified with the new animal drug application(s) to which they relate in duplicate on Form FD-2301 "Transmittal of Periodic Reports and Promotional Material for New Animal Drugs."

(b) All adverse experiences with new animal drugs as described in § 510.300(b)(2) or § 510.301(b) whether or not related to a required periodic report submitted on a Form FD-2301, shall be reported on Form FD-1932 "Adverse Drug Reaction" (except as provided in paragraph (c) of this section). Reports of adverse drug experiences may be submitted initially in the form of a written communication, but any such communication shall be followed promptly (but not necessarily within the prescribed 15 working days) by a completed Form FD-1932. A separate "Adverse Drug Reaction" form should be submitted for each patient where feasible.

(c) In lieu of Form FD-1932 the holder of an approved new animal drug application may submit:

(1) A computerized report if the information contained therein and the sequence in which it is presented are equivalent to that required by Form FD-1932 and the report is submitted in duplicate. Such reports will require initial approval by the Food and Drug Administration prior to use; and

(2) Copies of reports of reactions appearing in the published scientific literature may be submitted.

(d) Forms FD-1932 and FD-2301, with instructions for their use, may be obtained from the Food and Drug Administration, Department of Health and

Human Services, Center for Veterinary Medicine, 7500 Standish Pl., Rockville, MD 20855.

[40 FR 13807, Mar. 27, 1975, as amended at 41 FR 35844, Aug. 25, 1976; 54 FR 18280, Apr. 28, 1989; 57 FR 6475, Feb. 25, 1992]

§ 510.305 Maintenance of copies of approved applications for animal feed bearing or containing new animal drugs.

Each applicant shall maintain in a single accessible location on the premises of each establishment to which an approved medicated feed application (Form FDA 1900) or supplemental application applies either:

(a) A copy of the approved medicated feed application (Form FDA 1900) and a sample of the approved labeling; or

(b) Identification of the approved medicated feed application in a single file or in a single readable document that includes:

(1) The application number and date of its approval;

(2) The name(s) of the premix(es) and the concentration of the drug(s) contained in the premix(es);

(3) The name(s) of the approved manufacturer(s) of the premix(es);

(4) The concentration of the drug(s) in the finished medicated feed; and

(5) A sample of the approved labeling.

[41 FR 36203, Aug. 27, 1976, as amended at 51 FR 7391, Mar. 3, 1986]

Subpart E—Requirements for Specific New Animal Drugs

§ 510.410 Corticosteroids for oral, injectable, and ophthalmic use in animals; warnings and labeling requirements.

(a) The Food and Drug Administration has received reports of side effects associated with the oral, injectable, and ophthalmic use of corticosteroid animal drugs. The use of these drugs administered orally or by injection has resulted in premature parturition when administered during the last trimester of pregnancy. Premature parturition may be followed by dystocia, fetal death, retained placenta, and metritis. Additionally, corticosteroids used in dogs, rabbits, and rodents during pregnancy have produced cleft palate in offspring. Use in dogs has resulted in

§ 510.440

21 CFR Ch. I (4–1–99 Edition)

other congenital anomalies, including deformed forelegs, phocomelia, and anasarca. Drugs subject to this section are required to carry the veterinary prescription legend and are subject to the labeling requirements of § 201.105 of this chapter.

(b) In view of these potentially serious side effects, the Food and Drug Administration has concluded that the labeling on or within packaged corticosteroid-containing preparations intended for animal use shall bear conspicuously the following warning statement:

Warning: Clinical and experimental data have demonstrated that corticosteroids administered orally or by injection to animals may induce the first stage of parturition if used during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

Additionally, corticosteroids administered to dogs, rabbits, and rodents during pregnancy have resulted in cleft palate in offspring. Corticosteroids administered to dogs during pregnancy have also resulted in other congenital anomalies, including deformed forelegs, phocomelia, and anasarca.

[49 FR 48535, Dec. 13, 1984]

§ 510.440 Injectable iron preparations.

There has been an increasing interest in the use of injectable iron compounds for the prevention or treatment of iron-deficiency anemia in animals. Although some such preparations have been shown to be safe, such articles are regarded as new animal drugs within the meaning of the Federal Food, Drug, and Cosmetic Act. Accordingly, an approved new animal drug application is required prior to the marketing of such preparations within the jurisdiction of the act. In addition to the need for demonstrating the safety of such articles, the labeling of such preparations should not only recommend appropriate dosages of iron but also declare the amount (in milligrams) of available iron (Fe) per milliliter of the subject product.

§ 510.455 New animal drug requirements regarding free-choice administration in feeds.

(a) For the purpose of this section, free-choice administration of animal drugs in feeds involves feeds that are

placed in feeding or grazing areas and are not intended to be consumed fully at a single feeding or to constitute the entire diet of the animal. Such methods of administering drugs include, but are not limited to, medicated blocks (agglomerated feed compressed or rendered into a solid mass and cohesive enough to hold its form), mineral mixes, and liquid feed tank supplements (“lick tank” supplements) containing one or more animal drugs. The manufacture of medicated free-choice feeds is subject to the current good manufacturing practice regulations for medicated feeds.

(b) The Food and Drug Administration has concluded that there are questions about the safety and effectiveness of drugs when administered in free-choice feeds. Therefore, such methods of administration cause the drugs so administered to be new animal drugs, for which approved new animal drug applications (NADA’s) are required. (See § 510.3(i)). In addition, the exemption from the requirement of an approved medicated feed application provided in § 558.4 of this chapter does not apply to any free-choice medicated feed.

(c) An NADA or supplemental NADA for products for free-choice feeding submitted for approval under section 512(b) of the act shall provide for:

(1) The manufacture of a finished product for the free-choice administration of a new animal drug. Such an approval will not provide a basis upon which an application can be approved under section 512(m) of the act; or

(2) The manufacture of a Type A medicated article for use in the subsequent manufacture of a free-choice medicated feed. The approved NADA will provide a basis upon which an application can be approved under section 512(m) of the act. Data for a specific free-choice product may, if desired, be generated and submitted to the Food and Drug Administration by the manufacturer of the free-choice feed in the form of a master file which can be referenced in the NADA or supplemental NADA submitted by the new animal drug sponsor.

(d) Approval of the NADA or supplemental NADA submitted under paragraph (c) of this section will be reflected in a regulation in part 558 of this chapter published under section 512(i) of the act. The regulation will either state the formulation of the approved free-choice product or specify the specific free-choice administration products in which the drug is approved for use. If the approval is for a Type A medicated article, the regulation in part 558 of this chapter will indicate that each use of the Type A medicated article in a free-choice product must be the subject of an approved supplemental NADA.

(e) An application submitted under section 512(m) of the act to provide for manufacture of a specific free-choice feed from an approved Type A medicated article will be approved if, in addition to the information required by the medicated feed application, it includes a reference to the exact formula of the product to be manufactured as follows:

(1) The formula is the same as the one published in the new animal drug regulations; or

(2) The data in a master file have been referenced in an NADA or supplemental NADA; and

(3) Use of the Type A medicated article in the specific formulation has been approved on the basis that:

(i) The formula is the same as the one for which acceptable data have been submitted in a master file by the medicated feed applicant; or

(ii) The medicated feed applicant has written authority to reference a master file that has acceptable data for the formula in question.

(Approved by the Office of Management and Budget under control number 0910-0205)

[51 FR 19827, June 3, 1986]

Subpart F—Animal Use Exemptions From Certification and Labeling Requirements

§ 510.515 Animal feeds bearing or containing new animal drugs subject to the provisions of section 512(n) of the act.

Animal feeds that bear or contain penicillin, chlortetracycline, feed grade

zinc bacitracin, and bacitracin methylene disalicylate, with or without added suitable nutritive ingredients are exempt from the certification requirements of section 512 of the act provided they are the subject of and in compliance with regulations for their use in this subchapter E, part 558 of this chapter, or any one of the paragraphs of this section:

(a) Where indicated in paragraph (b) of this section it is manufactured with or without one, but only one, of the following ingredients in a quantity, by weight of feed, as hereinafter indicated:

(1) Arsanilic acid: Not less than 0.005 percent and not more than 0.01 percent.

(2) Sodium arsanilate: Not less than 0.005 percent and not more than 0.01 percent.

(3) 3-Nitro-4-hydroxyphenylarsonic acid: Not less than 0.0025 percent and not more than 0.0075 percent except in chicken or turkey feed which shall contain not less than 0.0025 percent and not more than 0.005 percent.

(b) It is intended for use in any one of the following conditions set forth in this paragraph:

(1) It is intended for use solely in the treatment of chronic respiratory disease (air-sac infection), infectious sinusitis, and blue comb (nonspecific infectious enteritis) in poultry and/or bacterial swine enteritis; its labeling bears adequate directions and warnings for such use; and it contains, per ton of feed, the equivalent of 100 grams of penicillin. When intended for uses specified in this paragraph, it may also contain, in the amount specified, one, but only one, of the ingredients prescribed by paragraph (a) of this section.

(2) It is intended for use solely in the treatment of chronic respiratory disease (air-sac infection) and infectious sinusitis in poultry; its labeling bears adequate directions and warnings for such use; and it contains not less than 0.1 percent para-aminobenzoic acid or the sodium or potassium salt or para-aminobenzoic acid.

(3)–(29) [Reserved]

(c) It is intended for use as follows:

§ 510.600

21 CFR Ch. I (4–1–99 Edition)

Product	Species	Use levels	Indications for use
1. Nicarbazin	Chickens	0.01 to 0.02 percent	For use in the prevention of outbreaks of coccidiosis in poultry flocks; growth promotion and feed efficiency. Do.
.....do	do	2.4 to 50 g/ton	
2. Nicarbazin	do	0.01 to 0.02 percent	
Bacitracin methylene disalicylate.	do	4 to 50 g/ton.	For use as an aid in the prevention of coccidiosis in poultry flocks; growth promotion and feed efficiency; improving pigmentation.
3. Nicarbazin	do	0.01 to 0.02 percent	
Bacitracin methylene disalicylate.	do	4 to 50 g/ton.	
3-Nitro-4-hydroxyphenylarsonic acid.	do	0.0025 to 0.005 percent.	Do.
4. Nicarbazin	do	0.01 to 0.02 percent	
Procaine penicillin	do	2.4 to 50 g/ton.	
3-Nitro-4-hydroxyphenylarsonic acid.	do	0.0025 to 0.005 percent.	Enhancement of growth and feed efficiency.
5. Chlortetracycline	Swine	10 to 50 g/ton	
Arsanilic acid	do	0.005 to 0.01 percent.	

[41 FR 8299, Feb. 25, 1976, as amended at 41 FR 11011, Mar. 15, 1976; 42 FR 18614, Apr. 8, 1977; 47 FR 42102, Sept. 24, 1982; 47 FR 51563, Nov. 16, 1982; 56 FR 41912, Aug. 23, 1991; 58 FR 30119, May 26, 1993; 61 FR 35950, July 9, 1996]

**(1) ALPHABETICAL LISTING OF SPONSORS—
Continued**

Subpart G—Sponsors of Approved Applications

§ 510.600 Names, addresses, and drug labeler codes of sponsors of approved applications.

(a) Section 512(i) of the act requires publication of names and addresses of sponsors of approved applications for new animal drugs.

(b) In this section each name and address is identified by a numerical drug labeler code. The labeler codes identify the sponsors of the new animal drug applications associated with the regulations published pursuant to section 512(i) of the act. The codes appear in the appropriate regulations and serve as a reference to the names and addresses listed in this section. The drug labeler code is established pursuant to section 510 of the act.

(c) The names, addresses, and drug labeler codes of sponsors of approved new animal drug applications are as follows:

(1) ALPHABETICAL LISTING OF SPONSORS

Firm name and address	Drug labeler code
Abbott Laboratories, North Chicago, IL 60064 ...	000074

Firm name and address	Drug labeler code
ADM Animal Health & Nutrition Div., P.O. Box 2508, Fort Wayne, IN 46801–2508	017519
Ag-Mark, Inc., P.O. Box 127, Teachey, NC 28464	024174
Agri Laboratories, Ltd., P.O. Box 3103, St. Joseph, MO 64503	057561
Agri-Tech, Inc., 4722 Broadway, Kansas City, MO 64112	017762
Carl S. Akey, Inc., P.O. Box 607, Lewisburg, OH 45338	017790
Akzo Nobel Surface Chemistry AB, Box 851, S–44485 Stenungsund, Sweden	063765
Alaco, Inc., 1500 North Wilmot Rd., suite 290–C, Tucson, AZ 85712	064146
Albion Laboratories, Inc., 101 North Main, Clearfield, UT 84015	011485
Alpharma Inc., One Executive Drive, P.O. Box 1399, Fort Lee, NJ 07024	046573
Alstoe, Ltd., Animal Health, Granary Chambers, 37–39 Burton St., Melton Mowbray, Leicestershire LE13 1AF, England	062408
Altana Inc., 60 Baylis Rd., Melville, NY 11747 ...	025463
American Cyanamid, Division of American Home Products, P.O. Box 1339, Fort Dodge, IA 50501	010042
American Pharmaceuticals Partners, Inc., 2045 North Cornell Ave., Melrose Park, IL 60160 ...	063323
Anika Therapeutics Inc., 236 West Cummings Park, Woburn, MA 01801	060865
Anthony Products Co., 5600 Peck Rd., Arcadia, CA 91006	000864
Argent Laboratories, 8702 152d Ave. NE., Redmond, WA 98052	051212
Ausa International, Inc., Rt. 8, P.O. Box 324–12, Tyler, TX 75703	059521
Balfour Guthrie & Co., Ltd., 315 North H St., Fresno, CA 93701	043728
Baxter Pharmaceutical Products, Inc., 110 Allen Rd., Liberty Corner, NJ 07938	010019
Bayer Corp., Agriculture Division, Animal Health, P.O. Box 390, Shawnee, Mission, KS 66201 ..	000859

Food and Drug Administration, HHS

§ 510.600

(1) ALPHABETICAL LISTING OF SPONSORS— Continued

Firm name and address	Drug label- er code
Biocraft Laboratories, Inc., 92 Route 46, Elmwood Park, NJ 07407	000332
Bioproducts, Inc., 320 Springside Dr., Suite 300, Fairlawn, OH 44333-2435	051359
Biopure Corp., 11 Hurley St., Cambridge, MA 02141	063075
BioScience Division of Milk Specialties Co., Illinois and Water Sts., P.O. Box 278, Dundee, IL 60118	032761
Boehringer Ingelheim Vetmedica, Inc., 2621 North Belt Highway, St. Joseph, MO 64502 ...	000010
Carnation Co., 5045 Wilshire Blvd., Los Angeles, CA 90036	047019
Chanelle Pharmaceuticals Manufacturing Ltd., Loughrea, County Galway, Ireland	061651
Chemdex, Inc., 12340 Santa Fe Dr., Lenexa, KS 66215	017287
Combe, Inc., 1101 Westchester Ave., White Plains, NY 10604	011509
ConAgra Pet Products Co., 3902 Leavenworth St., Omaha, NE 68105	021091
Contemporary Products, Inc., 3788 Elm Springs Rd., Springdale, AR 72764-6067	055462
Cooperative Research Farms, Box 69, Charlottesville, NY 12036	051267
Cross Vetpharm Group Ltd., Broomhill Rd., Tallaght, Dublin 24, Ireland	061623
Custom Feed Blenders Corp., 540 Hawkeye Ave., Fort Dodge, IA 50501	046987
Custom Feed Services Corp., 2100 N. 13th St., Norfolk, NE 68701	017473
Cyanamid Agricultural de Puerto Rico, Inc., P.O. Box 243, Manati, PR 00701	043781
Delmarva Laboratories, Inc., 2200 Wadebridge Rd., P.O. Box 525, Midlothian, VA 23113	059079
Diamond Shamrock Corp., Nutrition & Animal Health Div., 1100 Superior Ave., Cleveland, OH 44114	025001
DuPont Merck Pharmaceutical Co., DuPont Merck Plaza, MR2117, Wilmington, DE 19805	000056
Elanco Animal Health, A Division of Eli Lilly & Co., Lilly Corporate Center, Indianapolis, IN 46285	000986
Endo Pharmaceuticals, Inc., 223 Wilmington West Chester Pike, Chadds Ford, PA 19317 ..	060951
Eon Labs Manufacturing, Inc. 227-15 North Conduit Ave., Laurelton, NY 11413	000185
Equi Aid Products, Inc., 1517 West Knudsen Dr., Phoenix, AZ 85027	062240
EvSCO Pharmaceuticals, An Affiliate of IGI, Inc., Box 209, Harding Hwy., Buena, NJ 08310	017030
Farmers Feed & Supply Co., Ninth St. at Northwestern Tracks, Tipton, IA 52772	043744
Farmland Industries, Inc., Kansas City, MO 64116	021676
Farnam Companies, Inc., 301 West Osborn, Phoenix, AZ 85013-3928	017135
Feed Service Co., Inc., 303 Lundin Blvd., P.O. Box 698, Mankato, MN 56001	030841
John J. Ferrante, 11 Fairway Lane, Trumbull, CT 06611	058034
Fleming Laboratories, Inc., P.O. Box 34384, Charlotte, NC 28234	015565
Fort Dodge Animal Health, A Division of American Cyanamid Co., P.O. Box 1339, Fort Dodge, IA 50501	053501
Fort Dodge Animal Health, Division of American Home Products Corp., 800 Fifth St. NW., Fort Dodge, IA 50501	000856

(1) ALPHABETICAL LISTING OF SPONSORS— Continued

Firm name and address	Drug label- er code
Franklin Laboratories, Inc., P.O. Box 717, Fort Dodge, IA 50501	010290
Furst-McNess Co., Freeport, IL 61032	010439
Gland-O-Lac Co., 1818 Leavenworth St., Omaha, NE 68102	043735
Global Pharmaceutical Corp., Castor and Kensington Aves., Philadelphia, PA 19124	000115
Golden Sun Feeds, Inc., 111 South Fifth St., Estherville, IA 51334	021780
Michael Gordon, Inc., P.O. Box 8091, San Francisco, CA 94118	049047
Gossett Nutrition, Inc., 1676 Cascade Dr., Marion, OH 43302	050972
G. C. Hanford Manufacturing Co., P.O. Box 1017, Syracuse, NY 13201	010515
Halocarbon Laboratories, Division of Halocarbon Products Corp., 887 Kinderkamack Rd., P.O. Box 661, River Ridge, NJ 07661	012164
Happy Jack, Inc., Snow Hill, NC 28580	023851
Heinold Feeds, Inc., P.O. Box 377, Kouts, IN 46347	043727
Henwood Feed Additives, Division of Feed Specialties Co., Inc., 211 Western Rd., Box 577, Lewisburg, OH 45338	026186
Heska Corp., 1825 Sharp Point Dr., Fort Collins, CO 80525	063604
Hess & Clark, Inc., Seventh and Orange Sts., Ashland, OH 44805	050749
Dow B. Hickam, Inc., Pharmaceuticals, P.O. Box 35413, Houston, TX 77035	000514
Hoechst Roussel Vet, 30 Independence Blvd., P.O. Box 4915, Warren, NJ 07059	012799
Hubbard Milling Co., 424 North Front St., Mankato, MN 56001	012190
I. D. Russell Co. Laboratories, 1301 Iowa Ave., Longmont, CO 80501	017144
I.M.S. Inc., 13619 Industrial Rd., Omaha, NE 68137	050639
Illini Feeds, Box T, Oneida, IL 61467	037310
Inhalon Pharmaceuticals, Inc., P.O. Box 21170, Lehigh Valley, PA 18002	060307
International Nutrition, Inc., 6664 "L" St., Omaha, NE 68117	043733
Intervet, Inc., P.O. Box 318, 405 State St., Millsboro, DE 19966	057926
Ivy Laboratories, Inc., 8857 Bond Street, Overland Park, KS 66214	021641
J. C. Feed Mills, 1050 Sheffield, P.O. Box 224, Waterloo, IA 50704	039741
Jorgensen Laboratories, Inc., 1450 North Van Buren Ave., Loveland, CO 80538	045087
K. C. Pharmacal, Inc., 8345 Melrose Dr., Lenexa, KS 66214	038782
Kerber Milling Co., Box 152, 1817 E. Main St., Emmetsburg, IA 50536	029341
Koffolk, Inc., P.O. Box 675935, 14735 Las Quintas, Rancho Santa Fe, CA 92067	063271
Lambert-Kay, A Division of Carter-Wallace, Inc., P.O. Box 1001, Half Acre Rd., Cranbury, NJ 08512-0181	011615
Lemmon Co., Sellersville, PA 18960	000693
Lloyd, Inc., 604 W. Thomas Ave., Shenandoah, IA 51601	061690
Luitpold Pharmaceuticals, Inc., Animal Health Division, Shirley, NY 11967	010797
M & M Livestock Products Co., Eagle Grove, IA 50533	026282
Macleod Pharmaceuticals, Inc., 2600 Canton Ct., Fort Collins, CO 80525	058711

§ 510.600

21 CFR Ch. I (4–1–99 Edition)

(1) ALPHABETICAL LISTING OF SPONSORS—
Continued

Firm name and address	Drug label- er code
Marsam Pharmaceuticals, Inc., Bldg. 31, 24 Olney Ave., Cherry Hill, NJ 08034	000209
Mattox & Moore, Inc., 1503 East Riverside Drive, Indianapolis, IN 46207	027863
McClellan Laboratories, Inc., 19600 Sixth Ave., Lakeview, CA 92353	043738
Medicis Dermatologics, Inc., 4343 East Camelback Rd., Suite 250, Phoenix, AZ 85018–2700	099207
Med-Pharmex, Inc., 2727 Thompson Creek Rd., Pomona, CA 91767–1861	051259
Merial Ltd., 2100 Ronson Rd., Iselin, NJ 08830–3077	050604
Micro Chemical, Inc., Amarillo, TX 79105	047126
Mid-Continent Agrimarketing, Inc., 8833 Quivira Rd., Overland Park, KS 66214	059620
Monsanto Co., 800 North Lindbergh Blvd., St. Louis, MO 63167	059945
Moorman Manufacturing Co., Quincy, IL 62301	021930
Natchez Animal Supply Co., 201 John R. Junkin Dr., Natchez, MS 39120	049968
Nixon and Co., Kiewitt Plaza, Omaha, NE 68501	043729
Norbrook Laboratories, Ltd., Station Works, Newry BT35 6JP, Northern Ireland	055529
Norco Mills of Norfolk, Inc., P.O. Box 56, Norfolk, NE 68701	027190
Novartis Animal Health US, Inc.	058198
Nutra-Blend Corp., P.O. Box 485, Neosho, MO 64850	050568
NutriBasics Co., North Highway 71, P.O. Box 1014, Willmar, MN 56201	053740
Nylos Trading Co., Inc., P.O. Box 2, Route 202, Pomona, NY 10970	027454
Orion Corp. ORION–FARMOS, P.O. Box 425, SF–20101 Turku, Finland	052483
Orphan Medical, Inc., 13911 Ridgedale Dr., Suite 475, Minnetonka, MN 55305	062161
Osborn Laboratories, Inc., 2d and Oak Sts., Le Sueur, MN 56058	012487
OXIS International, Inc., 6040 N. Cutter Circle, Suite 317, Portland, OR 97217–3935	024991
Peavey Co., 730 Second Ave. South, Minneapolis, MN 55402	028459
Pegasus Laboratories, Inc., 8809 Ely Rd., Pensacola, FL 32514	055246
Pennfield Oil Co., 14040 Industrial Rd., Omaha, NE 68137	053389
Peptech Animal Health Pty, Ltd., 35–41 Waterloo Rd., North Ryde, New South Wales 2113, Australia	064288
Peter Hand Foundation, 2 East Madison St., Waukegan, IL 60085	043737
Pfizer, Inc., 235 East 42d St., New York, NY 10017	000069
Pharmacia & Upjohn Co., 7000 Portage Rd., Kalamazoo, MI 49001–0199	000009
Phoenix Pharmaceutical, Inc., 4621 Easton Rd., P.O. Box 6457 Farleigh Station, St. Joseph, MO 64506–0457	057319
Phoenix Scientific, Inc. 3915 South 48th St. Terrace, P.O. Box 6457, St. Joseph, MO 64506–0457	059130
Planalquímica Industrial Ltda., Rua das Magnolias nr. 2405, Jardim das Bandeiras, CEP 13053–120, Campinas, Sao Paulo, Brazil	060728
PM Ag Products, Inc., 1055 West 175th St., Homewood, IL 60430	036904
PM Resources, Inc., 13001 St. Charles Rock Rd., Bridgeton, MO 63044	060594

(1) ALPHABETICAL LISTING OF SPONSORS—
Continued

Firm name and address	Drug label- er code
Premier Malt Products, Inc., Milwaukee, WI 53201	032707
Protein Blenders, Inc., Box 631, Highway 218 South, Iowa City, IA 52240	033999
Purina Mills, Inc., P.O. Box 66812, St. Louis, MO 63166–6812	017800
Quali-Tech Products, Inc., 318 Lake Hazeltine Drive, Chaska, MN 55318	016968
The Rath Packing Co., P.O. Box 330, Waterloo, IA 50704	028260
Rhodia Limited, P.O. Box 46, St. Andrews Rd., Avonmouth, Bristol BS11 9YF, England, UK ..	059258
Rhone Merieux Canada, Inc., 345 Boul. Labbe Blvd., North Victoriaville, QC, G6P 1B1 Canada	047015
Rhone-Poulenc, Inc., P.O. Box 125, Black Horse Lane, Monmouth Junction, NJ 08852 ...	011526
A. H. Robins Co., P.O. Box 518, Fort Dodge, IA 50501–0518	000031
Roche Vitamins, Inc., 45 Waterview Blvd., Parsippany, NJ 07054–1298	063238
Roussel-UCLAF SA, Animal Health Division, 102 Route de Noisy, 93235 Romainville Cedex, France	012579
RSR Laboratories, Inc., 501 Fifth St., Bristol, TN 37620	058670
R. P. Scherer North America, P.O. Box 5600, Clearwater, FL 33518	011014
Schering-Plough Animal Health Corp., 1095 Morris Ave., Union, NJ 07083	000061
G. D. Searle & Co., P.O. Box 5110, Chicago, IL 60680	000014
Seeco Inc., Box 1014, North Highway 71, Willmar, MN 56201	011749
Shell Chemical Co., Division of Shell Oil Co., Animal Health, One Shell Plaza, Houston, TX 77001	011461
Sioux Biochemical, Inc., 204 Third St. NW., Sioux Center, IA 51250	063112
South St. Paul Feeds, Inc., 500 Farwell Ave., South St. Paul, MN 55075	001800
Southern Micro-Blenders, Inc., 3801 North Hawthorne St., Chattanooga, TN 37406	049685
Springfield Milling Corp., Vigorena Feeds, Springfield, MN 56087	035955
Square Deal Fortification Co., Kouts, IN 46347 ..	036108
Squire Laboratories, Inc., 100 Mill St., Revere, MA 02151	017153
Steris Laboratories, Inc., 620 North 51st Ave., Phoenix, AZ 85043–4705	000402
Sterling Winthrop, Inc., 9 Great Valley Pkwy., Malvern, PA 19355	000934
Summit Hill Laboratories, P.O. Box 535, Navesink, NJ 07752	037990
Syntex Animal Health, Division of Syntex Agribusiness, Inc., 3401 Hillview Ave., Palo Alto, CA 94304	000033
Teva Pharmaceuticals USA, 650 Cathill Rd., Sellersville, PA 18960	000093
Tevcon Ind., Inc., 8904 J St., Omaha, NE 68127	011757
Texas Vitamin Co., P.O. Box 18417, 10695 Aledo St., Dallas, TX 57218	000842
United Vaccines, A Harlan Sprague Dawley, Inc., Co., P.O. Box 4220, Madison, WI 53711 ..	058639
V.P.O., Inc., 4444 S. 76th St., Omaha, NE 68127	043743
Vet-A-Mix, Inc., P.O. Box A, Shenandoah, IA 51601	011789
Vetern, S.p.A., Viale E. Bezzi 24, 20146 Milano, Italy	055882

Food and Drug Administration, HHS

§ 510.600

(1) ALPHABETICAL LISTING OF SPONSORS— Continued

Firm name and address	Drug labeler code
Veterinary Laboratories, Inc., 12340 Santa Fe Dr., Lenexa, KS 66215	000857
Veterinary Research Associates, Inc., 20 Old Dock Rd., Yaphank, NY 11980	064408
Veterinary Service, Inc., 416 North Jefferson St., P.O. Box 2467, Modesto, CA 95354	033008
Veterinary Specialties Inc., 387 North Valley Ct., Barrington, IL 60010	062925
Walco International, Inc., 15 West Putnam, Porterville, CA 93257	049185
Waterloo Mills Co., 2050 Mitchell Ave., Waterloo, IA 50704	017139
Wayne Feed Division, Continental Grain Co., P.O. Box 459, Libertyville, IL 60048	034936
Weibel Feeds, Inc., R.R. 3, Pittsfield, IL 62363 ...	035098
Wendt Laboratories , Inc., 100 Nancy Dr., Belle Plaine, MN 56011	015579
West Agro, Inc., 11100 N. Congress Ave., Kansas City, MO 64153	033392
Westchester Veterinary Products, Inc., 180 Maroneck Ave., White Plains, NY 10601	043732
Western Chemical, Inc., 1269 Lattimore Rd., Ferndale, WA 98248	050378
Wildlife Laboratories, Inc., 1401 Duff Dr., Suite 600, Fort Collins, CO 80524	053923
Wyeth Laboratories, Division American Home Products Corp., P.O. Box 8299, Philadelphia, PA 19101	000008
Yoder Feed, Division of Yoder, Inc., Kalona, IA 52247	035369
Young's Inc., Roaring Spring, PA 16673	035393
Zema Corp., P.O. Box 12803, Research Triangle Park, Durham, NC 27709	050906

(2) NUMERICAL LISTING OF SPONSORS

Drug labeler code	Firm name and address
000008	Wyeth Laboratories, Division American Home Products Corp., P.O. Box 8299, Philadelphia, PA 19101.
000009	Pharmacia & Upjohn Co., 7000 Portage Rd., Kalamazoo, MI 49001-0199.
000010	Boehringer Ingelheim Vetmedica, Inc., 2621 North Belt Highway, St. Joseph, MO 64502.
000014	G. D. Searle & Co., P.O. Box 5110, Chicago, IL 60680.
000031	A. H. Robins Co., P.O. Box 518, Fort Dodge, IA 50501-0518.
000033	Syntex Animal Health Division of Syntex Agribusiness, Inc., 3401 Hillview Ave., Palo Alto, CA 94304.
000056	DuPont Merck Pharmaceutical Co., DuPont Merck Plaza, MR2117, Wilmington, DE 19805.
000061	Schering-Plough Animal Health Corp., 1095 Morris Ave., Union, NJ 07083.
000069	Pfizer, Inc., 235 East 42d St., New York, NY 10017.
000074	Abbott Laboratories, North Chicago, IL 60064.
000093	Teva Pharmaceuticals USA, 650 Cathill Rd., Sellersville, PA 18960.
000115	Global Pharmaceutical Corp., Castor and Kensington Aves., Philadelphia, PA 19124.

(2) NUMERICAL LISTING OF SPONSORS— Continued

Drug labeler code	Firm name and address
000185	Eon Labs Manufacturing, Inc., 227-15 North Conduit Ave., Laurelton, NY 11413.
000209	Marsam Pharmaceuticals, Inc., Bldg. 31, 24 Olney Ave., Cherry Hill, NJ 08034.
000332	Biocraft Laboratories, Inc., 92 Route 46, Elmwood Park, NJ 07407.
000402	Steris Laboratories, Inc., 620 North 51st Ave., Phoenix, AZ 85043-4705.
000514	Dow B. Hickam, Inc., Pharmaceuticals, P.O. Box 35413, Houston, TX 77035.
000693	Lemmon Co., Sellersville, PA 18960.
000842	Texas Vitamin Co., P.O. Box 18417, 10695 Aledo St., Dallas, TX 57218.
000856	Fort Dodge Animal Health, Division of American Home Products Corp., 800 Fifth St. NW., Fort Dodge, IA 50501.
000857	Veterinary Laboratories, Inc., 12340 Santa Fe Dr., Lenexa, KS 66215.
000859	Bayer Corp., Agriculture Division, Animal Health, P.O. Box 390, Shawnee Mission, KS 66201.
000864	Anthony Products Co., 5600 Peck Rd., Arcadia, CA 91006.
000934	Sterling Winthrop, Inc., 9 Great Valley Pkwy., Malvern, PA 19335.
000986	Elanco Animal Health, A Division of Eli Lilly & Co., Lilly Corporate Center, Indianapolis, IN 46285.
001800	South St. Paul Feeds, Inc., 500 Farwell Ave., South St. Paul, MN 55075.
010019	Baxter Pharmaceutical Products, Inc., 110 Allen Rd., Liberty Corner, NJ 07938-0804.
010042	American Cyanamid, Division of American Home Products, P.O. Box 1339, Fort Dodge, IA 50501.
010290	Franklin Laboratories, Inc., P.O. Box 717, Fort Dodge, IA 50501.
010290	Franklin Laboratories, P.O. Box 669, Amarillo, TX 79105.
010439	Furst-McNess Co., Freeport, IL 61032.
010515	G. C. Hanford Manufacturing Co., P.O. Box 1017, Syracuse, NY 13201.
010797	Luitpold Pharmaceuticals, Inc., Animal Health Division, Shirley, NY 11967.
011014	R. P. Scherer North America, P.O. Box 5600, Clearwater, FL 33518.
011461	Shell Chemical Co., Division of Shell Oil Co., Animal Health, One Shell Plaza, Houston, TX 77001.
011485	Albion Laboratories, Inc., 101 North Main, Clearfield, UT 84015.
011509	Combe, Inc., 1101 Westchester Ave., White Plains, NY 10604.
011526	Rhone-Poulenc, Inc., P.O. Box 125, Black Horse Lane, Monmouth Junction, NJ 08852.
011615	Lambert-Kay, A Division of Carter-Wallace, Inc., P.O. Box 1001, Half Acre Rd., Cranbury, NJ 08512-0181.
011749	Seeco Inc., Box 1014, North Highway 71, Willmar, MN 56201.
011789	Vet-A-Mix, Inc., P.O. Box A, Shenandoah, IA 51601.
012164	Halocarbon Laboratories, Division of Halocarbon Products Corp., 887 Kinderkamack Rd., P.O. Box 661, River Ridge, NJ 07661.
012190	Hubbard Milling Co., 424 North Front St., Mankato, MN 56001.

§ 510.600

21 CFR Ch. I (4–1–99 Edition)

**(2) NUMERICAL LISTING OF SPONSORS—
Continued**

Drug labeler code	Firm name and address
012487	Osborn Laboratories, Inc., 2d and Oak Sts., Le Sueur, MN 56058.
012579	Roussel-UCLAF SA, Animal Health Division, 102 Route de Noisy, 93235 Romainville Cedex, France.
012799	Hoechst Roussel Vet, 30 Independence Blvd., P.O. Box 4915, Warren, NJ 07059.
015565	Fleming Laboratories, Inc., P.O. Box 34384, Charlotte, NC 28234.
015579	Wendt Laboratories, Inc., 100 Nancy Dr., Belle Plaine, MN 56011.
016968	Quali-Tech Products, Inc., 318 Lake Hazeltine Dr., Chaska, MN 55318.
017030	Evsco Pharmaceuticals, An Affiliate of IGI, Inc., Box 209, Harding Hwy., Buena, NJ 08310.
017135	Farnam Companies, Inc., 301 West Osborn, Phoenix, AZ 85013–3928.
017139	Waterloo Mills Co., 2050 Mitchell Ave., Waterloo, IA 50704.
017144	I. D. Russell Co. Laboratories, 1301 Iowa Ave., Longmont, CO 80501.
017153	Squire Laboratories, Inc., 100 Mill St., Revere, MA 02151.
017287	Chemdex, Inc., 12340 Santa Fe Dr., Lenexa, KS 66215.
017473	Custom Feed Services Corp., 2100 N. 13th St., Norfolk, NE 68701.
017519	ADM Animal Health & Nutrition Div., P.O. Box 2508, Fort Wayne, IN 46801–2508.
017762	Agri-Tech, Inc., 4722 Broadway, Kansas City, MO 64112.
017790	Carl S. Akey, Inc., P.O. Box 607, Lewisburg, OH 45338.
017800	Purina Mills, Inc., P.O. Box 66812, St. Louis, MO 63166-6812.
021091	ConAgra Pet Products Co., 3902 Leavenworth St., Omaha, NE 68105.
021641	Ivy Laboratories, Inc., 8857 Bond Street, Overland Park, KS 66214.
021676	Farmland Industries, Inc., Kansas City, MO 64116.
021780	Golden Sun Feeds, Inc., 111 South Fifth St., Estherville, IA 51334.
021930	Moorman Manufacturing Co., Quincy, IL 62301.
023851	Happy Jack, Inc., Snow Hill, NC 28580.
024174	Ag-Mark, Inc., P.O. Box 127, Teachey, NC 28464.
024991	OXIS International, Inc., 6040 N. Cutter Circle, Suite 317 Portland, OR 97217–3935.
025001	Diamond Shamrock Corp., Nutrition & Animal Health Div., 1100 Superior Ave., Cleveland, OH 44114.
025463	Altana Inc., 60 Baylis Rd., Melville, NY 11747.
026186	Henwood Feed Additives, Division of Feed Specialties Co., Inc., 211 Western Rd., Box 577, Lewisburg, OH 45338.
026282	M & M Livestock Products Co., Eagle Grove, IA 50533.
027190	Norco Mills of Norfolk, Inc., P.O. Box 56, Norfolk, NE 68701.
027454	Nylos Trading Co., Inc., P.O. Box 2, Route 202, Pomona, NY 10970.
027863	Mattox & Moore, Inc., 1503 East Riverside Dr., Indianapolis, IN 46207.
028260	The Rath Packing Co., P.O. Box 330, Waterloo, IA 50704.

**(2) NUMERICAL LISTING OF SPONSORS—
Continued**

Drug labeler code	Firm name and address
028459	Peavey Co., 730 Second Ave. South, Minneapolis, MN 55402.
029341	Kerber Milling Co., Box 152, 1817 E. Main St., Emmetsburg, IA 50536.
030841	Feed Service Co., Inc., 303 Lundin Blvd., P.O. Box 698, Mankato, MN 56001.
032707	Premier Malt Products, Inc., Milwaukee, WI 53201.
032761	BioScience Division of Milk Specialties Co., Illinois and Water Sts., P.O. Box 278, Dundee, IL 60118.
033008	Veterinary Service, Inc., 416 North Jefferson St., P.O. Box 2467, Modesto, CA 95354.
033392	West Agro, Inc., 11100 N. Congress Ave., Kansas City, MO 64153.
033999	Protein Blenders, Inc., Box 631, Highway 218 South, Iowa City, IA 52240.
034936	Wayne Feed Division, Continental Grain Co., P.O. Box 459, Libertyville, IL 60048.
035098	Webel Feeds, Inc., R.R. 3, Pittsfield, IL 62363.
035369	Yoder Feed, Division of Yoder, Inc., Kalona, IA 52247.
035393	Young's, Inc., Roaring Spring, PA 16673.
035955	Springfield Milling Corp., Vigorena Feeds, Springfield, MN 56087.
036108	Square Deal Fortification Co., Kouts, IN 46347.
036904	PM Ag Products, Inc., 1055 West 175th St., Homewood, IL 60430.
037310	Illini Feeds, Box T, Oneida, IL 61467.
037990	Summit Hill Laboratories, P.O. Box 535, Navesink, NJ 07752.
038782	K. C. Pharmacal, Inc., 8345 Melrose Dr., Lenexa, KS 66214.
039741	J. C. Feed Mills, 1050 Sheffield, P.O. Box 224, Waterloo, IA 50704.
043727	Heinold Feeds, Inc., P.O. Box 377, Kouts, IN 46347.
043728	Balfour Guthrie & Co., Ltd., 315 North H St., Fresno, CA 93701.
043729	Nixon and Co., Kiewitt Plaza, Omaha, NE 68501.
043732	Westchester Veterinary Products, Inc., 180 Mamaroneck Ave., White Plains, NY 10601.
043733	International Nutrition, Inc., 6664 L St., Omaha, NE 68117.
043735	Gland-O-Lac Co., 1818 Leavenworth St., Omaha, NE 68102.
043737	Peter Hand Foundation, 2 East Madison St., Waukegan, IL 60085.
043738	McClellan Laboratories, Inc., 19600 Sixth Ave., Lakeview, CA 92353.
043743	V.P.O., Inc., 4444 South 76th St., Omaha, NE 68127.
043744	Farmers Feed & Supply Co., Ninth St. at Northwestern Tracks, Tipton, IA 52772.
043781	Cyanamid Agricultural de Puerto Rico, Inc., P.O. Box 243, Manati, PR 00701.
045087	Jorgensen Laboratories, Inc., 1450 North Van Buren Ave., Loveland, CO 80538.
046573	Alpharma Inc., One Executive Drive, P.O. Box 1399, Fort Lee, NJ 07024.
046987	Custom Feed Blenders Corp., 540 Hawkeye Ave., Fort Dodge, IA 50501.
047015	Rhone Merieux Canada, Inc., 345 Boul. Labbe Blvd., North Victoriaville, QC, G6P 1B1 Canada.

Food and Drug Administration, HHS

§ 510.600

(2) NUMERICAL LISTING OF SPONSORS— Continued

Drug labeler code	Firm name and address
047019	Carnation Co., 5045 Wilshire Blvd., Los Angeles, CA 90036.
047126	Micro Chemical, Inc., Amarillo, TX 79105.
049047	Michael Gordon, Inc., P.O. Box 1091, San Francisco, CA 94118.
049185	Walco International, Inc., 15 West Putnam, Porterville, CA 93257.
049685	Southern Micro-Blenders, Inc., 3801 North Hawthorne St., Chattanooga, TN 37406.
049968	Natchez Animal Supply Co., 201 John R. Junkin Dr., Natchez, MS 39120.
050378	Western Chemical, Inc., 1269 Lattimore Rd., Ferndale, WA 98248.
050568	Nutra-Blend Corp., P.O. Box 485, Neosho, MO 64850.
050604	Merial Ltd., 2100 Ronson Rd., Iselin, NJ 08830-3077.
050639	I.M.S. Inc., 13619 Industrial Rd., Omaha, NE 68137.
050749	Hess & Clark, Inc., Seventh and Orange Sts., Ashland, OH 44805.
050906	Zema Corp., P.O. Box 12803, Research Triangle Park, Durham, NC 27709.
050972	Gossett Nutrition, Inc., 1676 Cascade Dr., Marion, OH 43302.
051212	Argent Laboratories, 8702 152d Ave. NE., Redmond, WA 98052.
051259	Med-Pharmex, Inc., 2727 Thompson Creek Rd., Pomona, CA 91767-1861.
051267	Cooperative Research Farms, Box 69, Charlotteville, NY 12036.
051359	Bioproducts, Inc., 320 Springside Dr., Suite 300, Fairlawn, OH 44141.
052483	Orion Corp. ORION-FARMOS, P.O. Box 425, SF-20101 Turku, Finland.
053389	Pennfield Oil Co., 14040 Industrial Rd., Omaha, NE 68137.
053501	Fort Dodge Animal Health, A Division of American Cyanamid Co., P.O. Box 1339, Fort Dodge, IA 50501.
053740	NutriBasics Co., North Highway 71, P.O. Box 1014, Willmar, MN 56201.
053923	Wildlife Laboratories, Inc., 1401 Duff Dr., Suite 600, Fort Collins, CO 80524.
055246	Pegasus Laboratories, Inc., 8809 Ely Rd., Pensacola, FL 32514.
055462	Contemporary Products, Inc., 3788 Elm Springs Rd., Springdale, AR 72764-6067.
055529	Norbrook Laboratories, Ltd., Station Works, Newry BT35 6JP, Northern Ireland.
055882	Vetem, S.p.A., Viale E. Bezzi 24, 20146 Milano, Italy.
057319	Phoenix Pharmaceutical, Inc., 4621 Easton Rd., P.O. Box 6457 Farleigh Station, St. Joseph, MO 64506-0457.
057561	Agri Laboratories, Ltd., P.O. Box 3103, St. Joseph, MO 64503.
057926	Intervet, Inc., P.O. Box 318, 405 State St., Millsboro, DE 19966.
058034	John J. Ferrante, 11 Fairway Lane, Trumbull, CT 06611.
058198	Novartis Animal Health US, Inc., P.O. Box 18300, Greensboro, NC 27419-8300.
058639	United Vaccines, A Harlan Sprague Dawley, Inc., Co., P.O. Box 4220, Madison, WI 53711.
058670	RSR Laboratories, Inc., 501 Fifth St., Bristol, TN 37620.

(2) NUMERICAL LISTING OF SPONSORS— Continued

Drug labeler code	Firm name and address
058711	Macleod Pharmaceuticals, Inc., 2600 Canton Ct., Fort Collins, CO 80525.
059079	Delmarva Laboratories, Inc., 2200 Wadebridge Rd., P.O. Box 525, Midlothian, VA 23113.
059130	Phoenix Scientific, Inc. 3915 South 48th St. Terrace, P.O. Box 6457, St. Joseph, MO 64506-0457.
059258	Rhodia Limited, P.O. Box 46, St. Andrews Rd., Avonmouth, Bristol BS11 9YF, England, UK.
059945	Monsanto Co., 800 North Lindbergh Blvd., St. Louis, MO 63167.
059521	Ausa International, Inc., Rt. 8, P.O. Box 324-12, Tyler, TX 75703.
059620	Mid-Continent Agrimarketing, Inc., 8833 Quivira Rd., Overland Park, KS 66214.Bld., St. Louis, MO 63167.
060307	Inhalon Pharmaceuticals, Inc., P.O. Box 21170, Lehigh Valley, PA 18002.
060594	PM Resources, Inc., 13001 St. Charles Rock Rd., Bridgeton, MO 63044.
060728	Planalquimica Industrial Ltda., Rua das Magnolias nr. Jardim das Bandeiras, CEP 13053-120, Campinas, Sao Alto, Brazil.
060865	Anika Therapeutics Inc., 236 West Cummings Park, Woburn, MA 01801.
060951	Endo Pharmaceuticals, Inc., 223 Wilmington West Chester Pike, Chadds Ford, PA 19317.
061623	Cross Vetpharm Group Ltd., Broomhill Rd., Tallaght, Dublin 24, Ireland.
061651	Chanelle Pharmaceuticals Manufacturing Ltd., Loughrea, County Galway, Ireland.
061690	Lloyd, Inc., 604 W. Thomas Ave., Shendoah, IA 51601.
062161	Orphan Medical, Inc., 13911 Ridgedale Dr., Suite 475, Minnetonka, MN 55305.
062240	Equi Aid Products, Inc., 1517 West Knudsen Dr., Phoenix, AZ 85027.
062408	Alstoe, Ltd., Animal Health, Granary Chambers, 37-39 Burton St., Melton Mowbray, Leicestershire LE13 1AF, England.
062925	Veterinary Specialties Inc., 387 North Valley Ct., Barrington, IL 60010.
063075	Biopure Corp., 11 Hurley St., Cambridge, MA 02141.
063112	Sioux Biochemical, Inc., 204 Third St. NW., Sioux Center, IA 51250.
063238	Roche Vitamins, Inc., 45 Waterview Blvd., Parsippany, NJ 07054-1298.
063271	Koffolk, Inc., P.O. Box 675935, 14735 Las Quintas, Rancho Santa Fe, CA 92067.
063323	American Pharmaceuticals Partners, Inc., 2045 North Cornell Ave., Melrose Park, IL 60160.
063604	Heska Corp., 1825 Sharp Point Dr., Fort Collins, CO 80525.
063765	Akzo Nobel Surface Chemistry AB, Box 851, S-44485 Stenungsund, Sweden.
064146	Alaco, Inc., 1500 North Wilmot Rd., suite 290-C, Tucson, AZ 85712.
064288	Peptech Animal Health Pty, Ltd., 35-41 Waterloo Rd., North Ryde, New South Wales 2113, Australia.
064408	Veterinary Research Associates, Inc., 20 Old Dock Rd., Yaphank, NY 11980.

(2) NUMERICAL LISTING OF SPONSORS—
Continued

Drug labeler code	Firm name and address
099207	Medicis Dermatologics, Inc., 4343 East Camelback Rd., Suite 250, Phoenix, AZ 85018–2700.

[40 FR 13807, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 510.600, see the List of CFR Sections Affected in the Finding Aids section of this volume.

**PART 511—NEW ANIMAL DRUGS
FOR INVESTIGATIONAL USE**

AUTHORITY: 21 U.S.C. 321, 351, 352, 353, 360b, 371.

§ 511.1 New animal drugs for investigational use exempt from section 512(a) of the act.

(a) *New animal drugs for tests in vitro and in laboratory research animals.* (1) A shipment or other delivery of a new animal drug or animal feed bearing or containing a new animal drug intended solely for tests in vitro or in animals used only for laboratory research purposes shall be exempt from section 512 (a) and (m) of the act if it is labeled as follows:

Caution. Contains a new animal drug for investigational use only in laboratory research animals or for tests in vitro. Not for use in humans.

(2) The person distributing or causing the distribution of new animal drugs for tests in vitro or in animals used only for laboratory research purposes under this exemption shall use due diligence to assure that the consignee is regularly engaged in conducting such tests and that the shipment of the new animal drug will actually be used for tests in vitro or in animals used only for laboratory research.

(3) The person who introduced such shipment or who delivered the new animal drug for introduction into interstate commerce shall maintain adequate records showing the name and post office address of the expert or expert organization to whom the new animal drug is shipped and the date, quantity, and batch or code mark of each

shipment and delivery for a period of 2 years after such shipment and delivery. Upon the request of a properly authorized employee of the Department at reasonable times, he shall make such records available for inspection and copying.

(4) The exemption allowed in this paragraph shall not apply to any new animal drug intended for in vitro use in the regular course of diagnosing or treating disease, including antibacterial sensitivity discs impregnated with any new animal drug or drugs, which discs are intended for use in determining susceptibility of microorganisms to the new animal drug or drugs.

(b) *New animal drugs for clinical investigation in animals.* A shipment or other delivery of a new animal drug or an animal feed containing a new animal drug intended for clinical investigational use in animals shall be exempt from section 512(a) and (m) of the act if all the following conditions are met:

(1) The label shall bear the statements:

Caution. Contains a new animal drug for use only in investigational animals in clinical trials. Not for use in humans. Edible products of investigational animals are not to be used for food unless authorization has been granted by the U.S. Food and Drug Administration or by the U.S. Department of Agriculture.

In the case of containers too small or otherwise unable to accommodate a label with sufficient space to bear the caution statements required by paragraph (a) or (b) of this section, the statements may be included on the carton label and other labeling on or within the package from which the new animal drug is to be dispensed.

(2) The person or firm distributing or causing the distribution of the new animal drug or animal feed containing a new animal drug shall use due diligence to assure that the new animal drug or animal feed containing a new animal drug will actually be used for tests in animals and is not used in humans.

(3) The person who introduced such shipment or who delivered the new animal drug or animal feed containing a new animal drug for introduction into interstate commerce shall maintain adequate records showing the name

and post office address of the investigator to whom the new animal drug or animal feed containing a new animal drug is shipped and the date, quantity, and batch or code mark of each shipment and delivery for a period of 2 years after such shipment and delivery. Upon the request of a properly authorized employee of the Department at reasonable times, such records shall be made available for inspection and copying.

(4) Prior to shipment of the new animal drug for clinical tests in animals, the sponsor of the investigation shall submit in triplicate to the Food and Drug Administration a "Notice of Claimed Investigational Exemption for a New Animal Drug" including a signed statement containing the following information:

(i) The identity of the new animal drug.

(ii) All labeling and other pertinent information to be supplied to the investigators. When such pertinent information includes nonclinical laboratory studies, the information shall include, with respect to each nonclinical study, either a statement that the study was conducted in compliance with the requirements set forth in part 58 of this chapter, or, if the study was not conducted in compliance with such regulations, a brief statement of the reason for the noncompliance.

(iii) The name and address of each clinical investigator.

(iv) The approximate number of animals to be treated (or if not available, the amount of new animal drug to be shipped).

(v) If the new animal drug is given to food-producing animals, the statement shall contain the following additional information:

(a) A commitment that the edible products from such animals shall not be used for food without prior authorization in accordance with the provisions prescribed in this section.

(b) Approximate dates of the beginning and end of the experiment or series of experiments.

(c) The maximum daily dose(s) to be administered to a given species, the size of animal, maximum duration of administration, method(s) of adminis-

tration, and proposed withdrawal time, if any.

(vi) If a sponsor has transferred any obligations for the conduct of any clinical study to a contract research organization, a statement containing the name and address of the contract research organization, identification of the clinical study, and a listing of the obligations transferred. If all obligations governing the conduct of the study have been transferred, a general statement of this transfer—in lieu of a listing of the specific obligations transferred—may be submitted.

(5) Authorization for use of edible products derived from a treated food-producing animal may be granted under the provisions of this section and when the following specified conditions are met, except that in the case of an animal administered any unlicensed experimental veterinary biological product regulated under the viruses, serums, toxins statute (21 U.S.C., chapter V, sec. 151 *et seq.*) the product shall be exempt from the requirements of this section when U.S. Department of Agriculture approval has been obtained as provided in 9 CFR 103.2. Conditional authorization may be granted in advance of identification of the name(s) and address(es) of the clinical investigator(s) as required by paragraph (b)(4)(iii) of this section. Information required for authorization shall include, in addition to all other requirements of this section, the following:

(i) Data to show that consumption of food derived from animals treated at the maximum levels with the minimum withdrawal periods, if any, specified in accordance with paragraph (b)(4)(v)(c) of this section, will not be inconsistent with the public health; or

(ii) Data to show that food derived from animals treated at the maximum levels and with the minimum withdrawal periods, if any, specified in accordance with paragraph (b)(4)(v)(c) of this section, does not contain drug residues or metabolites.

(iii) The name and location of the packing plant where the animals will be processed, except that this requirement may be waived, on request, by the terms of the authorization.

§ 511.1

21 CFR Ch. I (4–1–99 Edition)

Authorizations granted under this paragraph do not exempt investigational animals and their products from compliance with other applicable inspection requirements. Any person who contests a refusal to grant such authorization shall have an opportunity for a regulatory hearing before the Food and Drug Administration pursuant to part 16 of this chapter.

(6) On written request of the Food and Drug Administration, the sponsor shall submit any additional information reported to or otherwise received by him with respect to the investigation deemed necessary to facilitate a determination whether there are grounds in the interest of public health for terminating the exemption.

(7) The sponsor shall assure himself that the new animal drug is shipped only to investigators who:

(i) Are qualified by scientific training and/or experience to evaluate the safety and/or effectiveness of the new animal drug.

(ii) Shall maintain complete records of the investigations, including complete records of the receipt and disposition of each shipment or delivery of the new animal drug under investigation. Copies of all records of the investigation shall be retained by the investigator for 2 years after the termination of the investigation or approval of a new animal drug application.

(iii) Shall furnish adequate and timely reports of the investigation to the sponsor.

(8) The sponsor:

(i) Shall retain all reports received from investigators for 2 years after the termination of the investigation or approval of a new animal drug application and make such reports available to a duly authorized employee of the Department for inspection at all reasonable times.

(ii) Shall provide for current monitoring of the investigation by a person qualified by scientific training and experience to evaluate information obtained from the investigation, and shall promptly investigate and report to the Food and Drug Administration and to all investigators any findings associated with use of the new animal drug that may suggest significant haz-

ards pertinent to the safety of the new animal drug.

(iii) Shall not unduly prolong distribution of the new animal drug for investigational use.

(iv) Shall not, nor shall any person acting for or on behalf of the sponsor, represent that the new animal drug is safe or effective for the purposes for which it is under investigation. This requirement is not intended to restrict the full exchange of scientific information.

(v) Shall not commercially distribute nor test-market the new animal drug until a new animal drug application is approved pursuant to section 512(c) of the act.

(9) If the shipment or other delivery of the new animal drug is imported or offered for importation into the United States for clinical investigational use in animals, it shall also meet the following conditions:

(i) The importer of all such shipments or deliveries is an agent of the foreign exporter residing in the United States or the ultimate consignee, which person has, prior to such shipments and deliveries, informed the Food and Drug Administration of his intention to import the new animal drug as sponsor in compliance with the conditions prescribed in this subdivision; or

(ii) The new animal drug is shipped directly to a scientific institution with adequate facilities and qualified personnel to conduct laboratory or clinical investigations and is intended solely for use in such institutions and which institution has submitted a statement as sponsor of the investigation.

(10) The sponsor shall submit either a claim for categorical exclusion under § 25.30 or § 25.33 of this chapter or an environmental assessment under § 25.40 of this chapter.

(c) *Withdrawal of eligibility to receive investigational-use new animal drugs.* (1) Whenever the Food and Drug Administration has information indicating that an investigator has repeatedly or deliberately failed to comply with the conditions of these exempting regulations or has submitted false information either to the sponsor of the investigation or in any required report, the Center

Food and Drug Administration, HHS

§ 511.1

for Veterinary Medicine will furnish the investigator written notice of the matter complained of in general terms and offer him an opportunity to explain the matter in an informal conference and/or in writing. If an explanation is offered but not accepted by the Center for Veterinary Medicine, the investigator shall have an opportunity for a regulatory hearing before the Food and Drug Administration pursuant to part 16 of this chapter on the question of whether the investigator is entitled to receive investigational new animal drugs.

(2) If, after evaluating all available information, including any explanation presented by the investigator, the Commissioner determines that the investigator has repeatedly or deliberately failed to comply with the conditions of the exempting regulations in this section or has repeatedly or deliberately submitted false information to the sponsor of an investigation, the Commissioner will notify the investigator and the sponsor of any investigation in which he has been named as a participant that the investigator is not entitled to receive investigational use new animal drugs with a statement of the basis for such determination.

(3) Each "Notice of Claimed Investigational Exemption for a New Animal Drug" and each approved new animal drug application containing data reported by an investigator who has been determined to be ineligible to receive investigational-use new animal drugs will be examined to determine whether he has submitted unreliable data that are essential to the continuation of the investigation or essential to the approval of any new animal drug application.

(4) If the Commissioner determines, after the unreliable data submitted by the investigator are eliminated from consideration, that the data remaining are inadequate to support a conclusion that it is reasonably safe to continue the investigation, he shall first notify the sponsor, who shall have an opportunity for a regulatory hearing before the Food and Drug Administration pursuant to part 16 of this chapter on whether the exemption should be terminated. If a danger to the public health exists, however, he shall termi-

nate the exemption forthwith and notify the sponsor of the termination. In such event the sponsor shall have an opportunity for a regulatory hearing before the Food and Drug Administration pursuant to part 16 (see 42 FR 15675, March 22, 1977) of this chapter on the question of whether the exemption should be reinstated.

(5) If the Commissioner determines, after the unreliable data submitted by the investigator are eliminated from consideration, that the data remaining are such that a new animal drug application would not have been approved, he will proceed to withdraw approval of the application in accordance with section 512(e) of the act.

(6) An investigator who has been determined to be ineligible may be reinstated as eligible to receive investigational-use new animal drugs when the Commissioner determines that he has presented adequate assurance that he will employ such new animal drugs solely in compliance with the exempting regulations in this section for investigational-use new animal drugs.

(d) *Termination of exemption.* If the Commissioner finds that:

(1) The sponsor of the investigation has failed to comply with any of the conditions for the exemption established under this section, or

(2) The continuance of the investigation is unsafe or otherwise contrary to the public interest or the drug is being or has been used for purposes other than bona fide scientific investigation, he shall first notify the sponsor and invite his immediate correction. If the conditions of the exemption are not immediately met, the sponsor shall have an opportunity for a regulatory hearing before the Food and Drug Administration pursuant to part 16 of this chapter on whether the exemption should be terminated. If the exemption is terminated the sponsor shall recall or have destroyed the unused supplies of the new animal drug.

(e) *Statements and requests.* "Notice(s) of Claimed Investigational Exemption for a New Animal Drug" and requests for authorization to use investigational animals and their products for food should be addressed to the Department of Health and Human Services, Food and Drug Administration, Center for

Pt. 514

Veterinary Medicine, 7500 Standish Pl., Rockville, MD 20855.

(f) *Contract research organizations.* (1) For purposes of this part and part 514, *contract research organization* means a person that assumes, as an independent contractor with the sponsor, one or more of the obligations of a sponsor, e.g., design of a protocol, selection or monitoring of investigations, evaluation of reports, and preparation of materials to be submitted to the Food and Drug Administration.

(2) A sponsor may transfer responsibility for any or all of the obligations set forth in this part to a contract research organization. Any such transfer shall be in writing and, if not all obligations are transferred, shall describe each of the obligations being assumed by the contract research organization. If all obligations are transferred, a general statement that all obligations have been transferred is acceptable. Any obligation not covered by the written description shall be deemed not to have been transferred.

(3) A contract research organization that assumes any obligation of a sponsor shall comply with the specific regulations in this chapter applicable to this obligation and shall be subject to the same regulatory action as a sponsor for failure to comply with any obligation assumed under these regulations. Thus, all references to *sponsor* in this part apply to a contract research organization to the extent that it assumes one or more obligations of the sponsor.

[40 FR 13823, Mar. 27, 1975, as amended at 41 FR 48268, Nov. 2, 1976; 42 FR 15675, Mar. 22, 1977; 50 FR 7517, Feb. 22, 1985; 50 FR 16668, Apr. 26, 1985; 52 FR 8847, Mar. 19, 1987; 54 FR 18280, Apr. 28, 1989; 57 FR 6475, Feb. 25, 1992; 62 FR 40599, July 29, 1997]

PART 514—NEW ANIMAL DRUG APPLICATIONS

Subpart A—General Provisions

Sec.

514.1 Applications.

514.2 Applications for animal feeds bearing or containing new animal drugs.

514.6 Amended applications.

514.7 Withdrawal of applications without prejudice.

21 CFR Ch. I (4–1–99 Edition)

514.8 Supplemental new animal drug applications.

514.9 Supplemental applications for animal feeds bearing or containing new animal drugs.

514.10 Confidentiality of data and information in an investigational new animal drug notice and a new animal drug application file for an antibiotic drug.

514.11 Confidentiality of data and information in a new animal drug application file.

514.12 Confidentiality of data and information in an investigational new animal drug notice.

514.15 Untrue statements in applications.

Subpart B—Administrative Actions on Applications

514.100 Evaluation and comment on applications.

514.105 Approval of applications.

514.106 Approval of supplemental applications.

514.110 Reasons for refusing to file applications.

514.111 Refusal to approve an application.

514.112 Return of applications for animal feeds bearing or containing new animal drugs.

514.115 Withdrawal of approval of applications.

514.116 Notice of withdrawal of approval of application.

514.117 Adequate and well-controlled studies.

514.120 Revocation of order refusing to approve an application or suspending or withdrawing approval of an application.

514.121 Service of notices and orders.

Subpart C—Hearing Procedures

514.200 Contents of notice of opportunity for a hearing.

514.201 Procedure for hearings.

Subparts D–E [Reserved]

Subpart F—Judicial Review

514.235 Judicial review.

AUTHORITY: 21 U.S.C. 351, 352, 360b, 371, 379e, 381.

SOURCE: 40 FR 13825, Mar. 27, 1975, unless otherwise noted.

Subpart A—General Provisions

§ 514.1 Applications.

(a) Applications to be filed under section 512(b) of the act shall be submitted in the form described in paragraph (b)

of this section. If any part of the application is in a foreign language, an accurate and complete English translation shall be appended to such part. Translations of literature printed in a foreign language shall be accompanied by copies of the original publication. The application must be signed by the applicant or by an authorized attorney, agent, or official. If the applicant or such authorized representative does not reside or have a place of business within the United States, the application must also furnish the name and post office address of, and must be countersigned by, an authorized attorney, agent, or official residing or maintaining a place of business within the United States. Pertinent information may be incorporated in, and will be considered as part of, an application on the basis of specific reference to such information, including information submitted under the provisions of §511.1 of this chapter, in the files of the Food and Drug Administration; however, the reference must be specific in identifying the information. Any reference to information furnished by a person other than the applicant may not be considered unless its use is authorized in a written statement signed by the person who submitted it.

(b) Applications for new animal drugs shall be submitted in triplicate and assembled in the manner prescribed by paragraph (b)(15) of this section, and shall include the following information:

(1) *Identification.* Whether the submission is an original or supplemental application; the name and the address of the applicant; the date of the application; the trade name(s) (if one has been proposed) and chemical name(s) of the new animal drug. Upon receipt, the application will be assigned a number NADA —, which shall be used for all correspondence with respect to the application.

(2) *Table of contents and summary.* The application shall be organized in a cohesive fashion, shall contain a table of contents which identifies the data and other material submitted, and shall contain a well-organized summary and evaluation of the data in the following form:

(i) Chemistry:

(a) Chemical structural formula or description for any new animal drug substance.

(b) Relationship to other chemically or pharmacologically related drugs.

(c) Description of dosage form and quantitative composition.

(ii) Scientific rationale and purpose the new animal drug is to serve:

(a) Clinical purpose.

(b) Highlights of laboratory studies: The reasons why certain types of studies were done or omitted as related to the proposed conditions of use and to information already known about this class of compounds. Emphasize any unusual or particularly significant pharmacological effects or toxicological findings.

(c) Highlights of clinical studies: The rationale of the clinical study plan showing why types of studies were done, amended, or omitted as related to laboratory studies and prior clinical experience.

(d) Conclusions: A short statement of conclusions combining the major points of effectiveness and safety as they relate to the use of the new animal drug.

(3) *Labeling.* Three copies of each piece of all labeling to be used for the article (total of 9).

(i) All labeling should be identified to show its position on, or the manner in which it is to accompany the market package.

(ii) Labeling for nonprescription new animal drugs should include adequate directions for use by the layman under all conditions of use for which the new animal drug is intended, recommended, or suggested in any of the labeling or advertising sponsored by the applicant.

(iii) Labeling for prescription veterinary drugs should bear adequate information for use under which veterinarians can use the new animal drug safely and for the purposes for which it is intended, including those purposes for which it is to be advertised or represented, in accord with §201.105 of this chapter.

(iv) All labeling for prescription or nonprescription new animal drugs shall be submitted with any necessary use restrictions prominently and conspicuously displayed.

§ 514.1

21 CFR Ch. I (4–1–99 Edition)

(v) Labeling for new animal drugs intended for use in the manufacture of medicated feeds shall include:

(a) Specimens of labeling to be used for such new animal drug with adequate directions for the manufacture and use of finished feeds for all conditions for which the new animal drug is intended, recommended, or suggested in any of the labeling, including advertising, sponsored by the applicant. Ingredient labeling may utilize collective names as provided in § 501.110 of this chapter.

(b) Representative labeling proposed to be used for Type B and Type C medicated feeds containing the new animal drug.

(vi) Draft labeling may be submitted for preliminary consideration of an application. Final printed labeling will ordinarily be required prior to approval of an application. Proposed advertising for veterinary prescription drugs may be submitted for comment or approval.

(4) *Components and composition.* A complete list of all articles used for production of the new animal drug including a full list of the composition of each article:

(i) A full list of the articles used as components of the new animal drug. This list should include all substances used in the synthesis, extraction, or other method of preparation of any new animal drug and in the preparation of the finished dosage form, regardless of whether they undergo chemical change or are removed in the process. Each component should be identified by its established name, if any, or complete chemical name, using structural formulas when necessary for specific identification. If any proprietary name is used, it should be followed by a complete quantitative statement of composition. Reasonable alternatives for any listed component may be specified.

(ii) A full statement of the composition of the new animal drug. The statement shall set forth the name and amount of each ingredient, whether active or not, contained in a stated quantity of the new animal drug in the form in which it is to be distributed (for example, amount per tablet or milliliter) and a batch formula representative of that to be employed for the manufac-

ture of the finished dosage form. All components should be included in the batch formula regardless of whether they appear in the finished product. Any calculated excess of an ingredient over the label declaration should be designated as such and percent excess shown. Reasonable variation may be specified.

(iii) If it is a new animal drug produced by fermentation:

(a) Source and type of microorganism used to produce the new animal drug.

(b) Composition of media used to produce the new animal drug.

(c) Type of precursor used, if any, to guide or enhance production of the antibiotic during fermentation.

(d) Name and composition of preservative, if any, used in the broth.

(e) A complete description of the extraction and purification processes including the names and compositions of the solvents, precipitants, ion exchange resins, emulsifiers, and all other agents used.

(f) If the new animal drug is produced by a catalytic hydrogenation process (such as tetracycline from chlortetracycline), a complete description of each chemical reaction with graphic formulas used to produce the new animal drug, including the names of the catalyst used, how it is removed, and how the new animal drug is extracted and purified.

(5) *Manufacturing methods, facilities, and controls.* A full description of the methods used in, and the facilities and controls used for, the manufacture, processing, and packing of the new animal drug. This description should include full information with respect to any new animal drug in sufficient detail to permit evaluation of the adequacy of the described methods of manufacture, processing, and packing, and the described facilities and controls to determine and preserve the identity, strength, quality, and purity of the new animal drug, and the following:

(i) If the applicant does not himself perform all the manufacturing, processing, packaging, labeling, and control operations for any new animal drug, he shall: Identify each person who will perform any part of such operations and designate the part; and provide a

Food and Drug Administration, HHS

§ 514.1

signed statement from each such person fully describing, directly or by reference, the methods, facilities, and controls he will use in his part of the operation. The statement shall include a commitment that no changes will be made without prior approval by the Food and Drug Administration, unless permitted under § 514.8.

(ii) A description of the qualifications, including educational background and experience, of the technical and professional personnel who are responsible for assuring that the new animal drug has the identity, strength, quality, and purity it purports or is represented to possess, and a statement of their responsibilities.

(iii) A description of the physical facilities including building and equipment used in manufacturing, processing, packaging, labeling, storage, and control operations.

(iv) The methods used in the synthesis, extraction, isolation, or purification of any new animal drug. When the specifications and controls applied to such new animal drugs are inadequate in themselves to determine its identity, strength, quality, and purity, the methods should be described in sufficient detail, including quantities used, times, temperature, pH, solvents, etc., to determine these characteristics. Alternative methods or variations in methods within reasonable limits that do not affect such characteristics of the new animal drug may be specified. A flow sheet and indicated equations should be submitted when needed to explain the process.

(v) Precautions to insure proper identity, strength, quality, and purity of the raw materials, whether active or not, including:

(a) The specifications for acceptance and methods of testing for each lot of raw material.

(b) A statement as to whether or not each lot of raw materials is given a serial number to identify it, and the use made of such numbers in subsequent plant operations.

(vi) The instructions used in the manufacturing, processing, packaging, and labeling of each dosage form of the new animal drug, including:

(a) The method of preparation of the master formula records and individual

batch records and the manner in which these records are used.

(b) The number of individuals checking weight or volume of each individual ingredient entering into each batch of the new animal drug.

(c) A statement as to whether or not the total weight or volume of each batch is determined at any stage of the manufacturing process subsequent to making up a batch according to the formula card and, if so, at what stage and by whom it is done.

(d) The precautions used in checking the actual package yield produced from a batch of the new animal drug with the theoretical yield. This should include a description of the accounting for such items as discards, breakage, etc., and the criteria used in accepting or rejecting batches of drugs in the event of an unexplained discrepancy.

(e) The precautions used to assure that each lot of the new animal drug is packaged with the proper label and labeling, including provisions for labeling storage and inventory control.

(f) Any special precautions used in the operations.

(vii) The analytical controls used during the various stages of the manufacturing, processing, packaging, and labeling of the new animal drug, including a detailed description of the collection of samples and the analytical procedures to which they are subjected. The analytical procedures should be capable of determining the active components within a reasonable degree of accuracy and of assuring the identity of such components.

(a) A description of practicable methods of analysis of adequate sensitivity to determine the amount of the new animal drug in the final dosage form should be included. The dosage form may be a finished pharmaceutical product, a Type A medicated article, a Type B or a Type C medicated feed, or a product for use in animal drinking water. Where two or more active ingredients are included, methods should be quantitative and specific for each active ingredient.

(b) If the article is one that is represented to be sterile, the same information with regard to the manufacturing, processing, packaging, and the collection of samples of the drug

should be given for sterility controls. Include the standards used for acceptance of each lot of the finished drug.

(viii) An explanation of the exact significance of any batch control numbers used in the manufacturing, processing, packaging, and labeling of the new animal drug, including such control numbers that may appear on the label of the finished article. State whether these numbers enable determination of the complete manufacturing history of the product. Describe any methods used to permit determination of the distribution of any batch if its recall is required.

(ix) Adequate information with respect to the characteristics of and the test methods employed for the container, closure, or other component parts of the drug package to assure their suitability for the intended use.

(x) A complete description of, and data derived from, studies of the stability of the new animal drug in the final dosage form, including information showing the suitability of the analytical methods used. A description of any additional stability studies underway or planned. Stability data for the finished dosage form of the new animal drug in the container in which it is to be marketed, including any proposed multiple dose container, and, if it is to be put into solution at the time of dispensing, for the solution prepared as directed. If the new animal drug is intended for use in the manufacture of Type C medicated feed as defined in § 558.3 of this chapter, stability data derived from studies in which representative formulations of the medicated feed articles are used. Similar data may be required for Type B medicated feeds as determined by the Food and Drug Administration on a case-by-case basis. Expiration dates shall be proposed for finished pharmaceutical dosage forms and Type A medicated articles. If the data indicate that an expiration date is needed for Type B or Type C medicated feeds, the applicant shall propose such expiration date. If no expiration date is proposed for Type B or Type C medicated feeds, the applicant shall justify its absence with data.

(xi) Additional procedures employed which are designed to prevent contamination and otherwise assure proper

control of the product. An application may be refused unless it includes adequate information showing that the methods used in, and the facilities and controls used for, the manufacturing, processing, and packaging of the new animal drug are adequate to preserve its identity, strength, quality, and purity in conformity with good manufacturing practice and identifies each establishment, showing the location of the plant conducting these operations.

(6) *Samples.* Samples of the new animal drug and articles used as components and information concerning them may be requested by the Center for Veterinary Medicine as follows:

(i) Each sample shall consist of four identical, separately packaged subdivisions, each containing at least three times the amount required to perform the laboratory test procedures described in the application to determine compliance with its control specifications for identity and assays. Each of the samples submitted shall be appropriately packaged and labeled to preserve its characteristics, to identify the material and the quantity in each subdivision of the sample, and to identify each subdivision with the name of the applicant and the new animal drug application to which it relates. Included are:

(a) A sample or samples of any reference standard and blank used in the procedures described in the application for assaying each new animal drug and other assayed components of the finished new animal drug.

(b) A representative sample or samples of each strength of the finished dosage form proposed in the application and employed in the clinical investigations and a representative sample or samples of each new animal drug from the batch(es) employed in the production of such dosage form.

(c) A representative sample or samples of finished market packages of each strength of the dosage form of the new animal drug prepared for initial marketing and, if any such sample is not from a representative commercial-scale production batch, such a sample from a representative commercial-scale production batch, and a representative sample or samples of each new animal drug from the batch(es)

employed in the production of such dosage form, provided that in the case of new animal drugs marketed in large packages the sample should contain only three times a sufficient quantity of the new animal drug to allow for performing the control tests for drug identity and assays.

(ii) The following information shall be included for the samples when requested:

(a) For each sample submitted, full information regarding its identity and the origin of any new animal drug contained therein (including a statement whether it was produced on a laboratory, pilot-plant, or full-production scale) and detailed results of all laboratory tests made to determine the identity, strength, quality, and purity of the batch represented by the sample, including assays.

(b) For any reference standard submitted, a complete description of its preparation and the results of all laboratory tests on it. If the test methods used differed from those described in the application, full details of the methods employed in obtaining the reporting results.

(7) *Analytical methods for residues.* Applications shall include a description of practicable methods for determining the quantity, if any, of the new animal drug in or on food, and any substance formed in or on food because of its use, and the proposed tolerance or withdrawal period or other use restrictions to ensure that the proposed use of this drug will be safe. When data or other adequate information establish that it is not reasonable to expect the new animal drug to become a component of food at concentrations considered unsafe, a regulatory method is not required.

(i) The kind of information required by this subdivision may include: Complete experimental protocols for determining drug residue levels in the edible products, and the length of time required for residues to be eliminated from such products following the drug's use; residue studies conducted under appropriate (consistent with the proposed usage) conditions of dosage, time, and route of administration to show levels, if any, of the drug and/or its metabolites in test animals during

and upon cessation of treatment and at intervals thereafter in order to establish a disappearance curve; if the drug is to be used in combination with other drugs, possible effects of interaction demonstrated by the appropriate disappearance curve or depletion patterns after drug withdrawal under appropriate (consistent with the proposed usage) conditions of dosage, time, and route of administration; if the drug is given in the feed or water, appropriate consumption records of the medicated feed or water and appropriate performance data in the treated animal; if the drug is to be used in more than one species, drug residue studies or appropriate metabolic studies conducted for each species that is food-producing. To provide these data, a sufficient number of birds or animals should be used at each sample interval. Appropriate use of labeled compounds (e.g. radioactive tracers), may be utilized to establish metabolism and depletion curves. Drug residue levels ordinarily should be determined in muscle, liver, kidney, and fat and where applicable, in skin, milk, and eggs (yolk and egg white). As a part of the metabolic studies, levels of the drug or metabolite should be determined in blood where feasible. Samples may be combined where necessary. Where residues are suspected or known to be present in litter from treated animals, it may be necessary to include data with respect to such residues becoming components of other agricultural commodities because of use of litter from treated animals.

(ii) A new animal drug that has the potential to contaminate human food with residues whose consumption could present a risk of cancer to people must satisfy the requirements of subpart E of part 500 of this chapter.

(8) *Evidence to establish safety and effectiveness.* (i) An application may be refused unless it contains full reports of adequate tests by all methods reasonably applicable to show whether or not the new animal drug is safe and effective for use as suggested in the proposed labeling.

(ii) An application may be refused unless it includes substantial evidence, consisting of adequate and well-controlled investigations, including field investigation, by experts qualified by

scientific training and experience to evaluate the effectiveness of the new animal drug involved, on the basis of which it could fairly and reasonably be concluded by such experts that the new animal drug will have the effect it purports or is represented to have under the conditions of use prescribed, recommended, or suggested in the proposed labeling.

(iii) An application may be refused unless it contains detailed reports of the investigations, including studies made on laboratory animals, in which the purpose, methods, and results obtained are clearly set forth of acute, subacute, and chronic toxicity, and unless it contains appropriate clinical laboratory results related to safety and efficacy. Such information should include identification of the person who conducted each investigation, a statement of where the investigations were conducted, and where the raw data are available in the application.

(iv) All information pertinent to an evaluation of the safety and effectiveness of the new animal drug received or otherwise obtained by the applicant from any source, including information derived from other investigations or commercial marketing (for example, outside the United States), or reports in the scientific literature, both favorable and unfavorable, involving the new animal drug that is the subject of the application and related new animal drugs shall be submitted. An adequate summary may be acceptable in lieu of a reprint of a published report that only supports other data submitted. Include any evaluation of the safety or effectiveness of the new animal drug that has been made by the applicant's veterinary or medical department, expert committee, or consultants.

(v) If the new animal drug is a combination of previously investigated or marketed new animal drugs, an adequate summary of preexisting information from preclinical and clinical investigation and experience with its components, including all reports received or otherwise obtained by the applicant suggesting side effects, contraindications, and ineffectiveness in use of such components, shall be submitted. Such summary should include an adequate bibliography of publica-

tions about the components and may incorporate by reference information concerning such components previously submitted to the Food and Drug Administration by the applicant; with written authorization, information may also be incorporated from the material that another applicant has on file with the Food and Drug Administration. Each ingredient designated as active in any new animal drug combination must make a contribution to the effect in the manner claimed or suggested in the labeling, and, if in the absence of express labeling claims of advantages for the combination such a product purports to be better than either component alone, it must be established that the new animal drug has that purported effectiveness.

(vi) An application shall include a complete list of the names and post office addresses of all investigators who received the new animal drug. This may be incorporated in whole or in part by reference to information submitted under the provisions of § 511.1 of this chapter.

(vii) Explain any omission of reports from any investigator to whom the investigational new animal drug has been made available. The unexplained omission of any reports of investigations made with the new animal drug by the applicant or submitted to him by an investigator or the unexplained omission of any pertinent reports of investigations or clinical experience received or otherwise obtained by the applicant from published literature or other sources that would bias an evaluation of the safety of the new animal drug or its effectiveness in use, constitutes grounds for the refusal or withdrawal of the approval of an application.

(viii) If a sponsor has transferred any obligations for the conduct of any clinical study to a contract research organization, the application is required to include a statement containing the name and address of the contract research organization, identifying the clinical study, and listing the obligations transferred. If all obligations governing the conduct of the study have been transferred, a general statement of this transfer—in lieu of a listing of the specific obligations transferred—may be submitted.

(ix) If original subject records were audited or reviewed by the sponsor in the course of monitoring any clinical study to verify the accuracy of the case reports submitted to the sponsor, a list identifying each clinical study so audited or reviewed.

(9) [Reserved]

(10) *Supplemental applications.* If it is a supplemental application, full information shall be submitted on each proposed change concerning any statement made in the approved application.

(11) *Applicant's commitment.* It is understood that the labeling and advertising for the new animal drug will prescribe, recommend, or suggest its use only under the conditions stated in the labeling which is part of this application and if the article is a prescription new animal drug, it is understood that any labeling which furnishes or purports to furnish information for use or which prescribes, recommends, or suggests a dosage for use of the new animal drug will also contain, in the same language and emphasis, information for its use including indications, effects, dosages, routes, methods, and frequency and duration of administration, any relevant hazards, contraindications, side effects, and precautions contained in the labeling which is part of this application. It is understood that all representations in this application apply to the drug produced until changes are made in conformity with § 514.8.

(12) *Additional commitments.* (i) New animal drugs as defined in § 510.3 of this chapter, intended for use in the manufacture of animal feeds in any State will be shipped only to persons who may receive such drugs in accordance with § 510.7 of this chapter.

(ii) The methods, facilities, and controls described under item 5 of this application conform to the current good manufacturing practice regulations in subchapter C of this chapter.

(iii) With respect to each nonclinical laboratory study contained in the application, either a statement that the study was conducted in compliance with the good laboratory practice regulations set forth in part 58 of this chapter, or, if the study was not conducted in compliance with such regulations, a

brief statement of the reason for the noncompliance.

(13) [Reserved]

(14) *Environmental assessment.* The applicant is required to submit either a claim for categorical exclusion under § 25.30 or § 25.33 of this chapter or an environmental assessment under § 25.40 of this chapter.

(15) *Assembling and binding the application.* Assemble and bind an original and two copies of the application as follows:

(i) Bind the original or ribbon copy of the application as copy No. 1.

(ii) Bind two identical copies as copy No. 2 and copy No. 3.

(iii) Identify each front cover with the name of the applicant, new animal drug, and the copy number.

(iv) Number each page of the application sequentially in the upper right hand corner or in another location so that the page numbers remain legible after the application has been bound, and organize the application consistent with paragraphs (b) (1) through (14) of this section. Each copy should bear the same page numbering, whether sequential in each volume or continuous and sequential throughout the application.

(v) Include complete labeling in each of the copies. It is suggested that labeling be identified by date of printing or date of preparation.

(vi) Submit separate applications for each different dosage form of the drug proposed. Repeating basic information pertinent to all dosage forms in each application is unnecessary if reference is made to the application containing such information. Include in each application information applicable to the specific dosage form, such as labeling, composition, stability data, and method of manufacture.

(vii) Submit in folders amendments, supplements, and other correspondence sent after submission of an original application. The front cover of these submissions should be identified with the name of the applicant, new animal drug, copy number, and the new animal drug application number, if known.

(c) When a new animal drug application is submitted for a new animal drug which has a stimulant, depressant, or hallucinogenic effect on the central nervous system, if it appears

§ 514.2

21 CFR Ch. I (4–1–99 Edition)

that the drug has a potential for abuse, the Commissioner shall forward that information to the Attorney General of the United States.

(d) *Minor use applications.* Applications for minor use new animal drugs:

(1) *Definitions.* For the purpose of this section:

(i) *Minor use* means the use of: (a) New animal drugs in minor animal species, or (b) new animal drugs in any animal species for the control of a disease that (1) occurs infrequently or (2) occurs in limited geographic areas.

(ii) *Minor species* means animals other than cattle, horses, swine, chickens, turkeys, dogs, and cats. Sheep are a minor species with respect to effectiveness and animal safety data collection requirements; sheep are a major species with respect to human safety data collection requirements arising from the possible presence of drug residues in food.

(2) *Animal safety, effectiveness, human food safety, and environmental considerations.* Guidelines for the preparation and submission of data to satisfy the requirements of section 512 of the act regarding animal safety, effectiveness, human food safety, and environmental considerations for new animal drugs intended for a *minor use* (as defined in paragraph (d)(1)(i) of this section) are available from the Industry Information Staff (HFV–11), Center for Veterinary Medicine, Food and Drug Administration, 7500 Standish Pl., Rockville, MD 20855.

(i) *Animal safety and effectiveness.* Where the guidelines do not specifically provide for a particular *minor use*, the Center for Veterinary Medicine, upon request, will advise interested persons on the effectiveness and animal safety data regarding the minor use that will be needed to satisfy the requirements of section 512 of the act. Where scientifically appropriate, the Center for Veterinary Medicine will allow the use of animal models and the extrapolation of data from a major species to a minor species to satisfy the requirements of the act.

(ii) *Human food safety and environmental considerations.* These guidelines do not specifically provide for a particular *minor use*. Therefore, the Center for Veterinary Medicine will, upon re-

quest, advise interested persons of the data that will be needed. Where scientifically appropriate, the Center for Veterinary Medicine will allow the extrapolation of data from a major species to a minor species to satisfy the requirements of the act.

(Approved by the Office of Management and Budget under control number 0910–0032)

[40 FR 13825, Mar. 27, 1975]

EDITORIAL NOTE: For Federal Register citations affecting § 514.1, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 514.2 Applications for animal feeds bearing or containing new animal drugs.

(a) Applications (Form FDA 1900) to be filed under section 512(m) of the act shall be completed, signed, and submitted in triplicate in the form described in paragraphs (b) and (c) of this section.

(b) Each application for a Type B or Type C medicated feed, as defined in § 558.3 of this chapter, shall include the following information:

(1) The name and address of the applicant.

(2) The registration number assigned pursuant to section 510 of the act and last date of registration of each mill.

(3) Whether the submission is an original or supplemental application.

(4) Identification of the Type A medicated article, as defined in § 558.3 of this chapter, used by generic name, potency, and manufacturer.

(5) The species of animal(s) for which the feed is intended.

(6) The form of feed to be produced, i.e., mash, meal crumbles, pellets, liquid, or other specified form.

(7) Whether the feed is a Type B or Type C medicated feed.

(8) Whether the feed is for sale or for own use (not for sale).

(9) Level of the drug(s) in the finished feed, and the amount of Type A medicated article per ton contained therein.

(10) Identification of the regulation(s) in subchapter E of this chapter on which approval relies.

(11) Labeling representative of each intended use as stated in the claim. Each generic label shall include the claim, drug level, mixing directions, feeding directions, caution and/or

Food and Drug Administration, HHS

§ 514.8

warning statements, and any other special directions required by the published regulation. This shall consist of bag labels, invoice copy, bulk labels, and placards when applicable.

(12) A commitment to establish and maintain a program of sampling and analysis consisting of an assay of the first batch manufactured, followed thereafter by two samples at periodic intervals during the calendar year. If a medicated feed contains a combination of drugs, only one of the drugs need be subject to analysis each time, provided the one tested is different from the one(s) previously tested. Reports of assays shall be kept on the premises for not less than 1 year after the date of manufacture of the medicated feed.

(13) A statement of the minimum and maximum assay value permitted from the labeled amount of the drug.

(14) Identification of the agent authorized to act on behalf of the applicant.

(15) The applicant's name, responsible individual's title and original signature, and date.

(c) Upon approval, one copy of the application will be signed by an authorized employee of the Food and Drug Administration designated by the Commissioner, and it will be returned to the applicant.

(d) Applications (Form FDA 1900) may be obtained from the Public Health Service, Consolidated Forms and Publications Distribution Center, Washington Commerce Center, 3222 Hubbard Rd., Landover, MD 20785.

(Approved by the Office of Management and Budget under control number 0910-0011)

[51 FR 7391, Mar. 3, 1986, as amended at 55 FR 14831, Apr. 19, 1990]

§ 514.6 Amended applications.

The applicant may submit an amendment to an application that is pending, including changes that may alter the conditions of use, the labeling, safety, effectiveness, identity, strength, quality, or purity of the drug or the adequacy of the manufacturing methods, facilities, and controls to preserve them, in which case the unamended application may be considered as withdrawn and the amended application may be considered resubmitted on the date on which the amendment is re-

ceived by the Food and Drug Administration. The applicant will be notified of such date.

§ 514.7 Withdrawal of applications without prejudice.

The sponsor may withdraw his pending application from consideration as a new animal drug application upon written notification to the Food and Drug Administration. Such withdrawal may be made without prejudice to a future filing. Upon resubmission, the time limitation will begin to run from the date the resubmission is received by the Food and Drug Administration. The original application will be retained by the Food and Drug Administration although it is considered withdrawn. The applicant shall be furnished a copy at cost on request.

§ 514.8 Supplemental new animal drug applications.

(a)(1) After a new animal drug application is approved, a supplemental new animal drug application may propose changes. A supplemental application may omit statements made in the approved application concerning which no change is proposed. Each supplemental application shall include up-to-date reports of any of the kinds of information required by § 510.300(a) of this chapter that has not previously been submitted. A supplemental application shall be accompanied by either a claim for categorical exclusion under § 25.30 or § 25.33 of this chapter or an environmental assessment under § 25.40 of this chapter.

(2) A supplemental new animal drug application shall be submitted for any change beyond the variations provided for in the application, including changes in the scale of production such as from pilot-plant to production batch, that may alter the conditions of use, the labeling, safety, effectiveness, identity, strength, quality, or purity of the new animal drug, or the adequacy of the manufacturing methods, facilities, or controls to preserve them.

(3) If it is a prescription drug, any mailing or promotional piece used after the drug is placed on the market is labeling requiring a supplemental application, unless:

§514.8

21 CFR Ch. I (4–1–99 Edition)

(i) The parts of the labeling furnishing directions, warnings, and information for use of the drug are the same in language and emphasis as labeling approved or permitted; and

(ii) Any other parts of the labeling are consistent with and not contrary to such approved or permitted labeling.

(4) The supplemental application shall be submitted as follows. A communication proposing a change in a new animal drug application should provide for any one of the following kinds of changes:

(i) Revision in labeling, such as updating information pertaining to effects, dosages, and side effects and contraindications, which includes information headed “side effects,” “warnings,” “precautions,” and “contraindications.”

(ii) Addition of claim.

(iii) Revision in manufacturing or control procedures; for example, changes in components, composition, method of manufacture, analytical control procedures, package or tablet size, etc.

(iv) Change in manufacturing facilities.

(v) Provision for outside firm to participate in the preparation, distribution, or packaging of a new animal drug (new distributor, packer, supplier, manufacturer, etc.); one firm per submission.

Any number of changes may be submitted at any one time; but if they fall into different categories as listed in paragraphs (a)(4) (i) through (v) of this section, the proposed changes should be covered by separate communications. Where, however, a change necessitates an overlap in categories, it should be submitted in a single communication. For example, a change in tablet potency would require other changes such as in components, composition, and labeling and should be submitted in a single communication.

(5) The following kinds of changes may be placed into effect without the approval of a supplemental application, if such change is fully described in the next periodic report required under §510.300(b)(4) of this chapter or, when such a report is not required, in a written communication to the Food and Drug Administration within 60 days of

the effective date of the change (this does not apply to a change proposed because of any mixup or any bacteriological or significant chemical, physical, or other change or deterioration in the drug or any failure of one or more distributed batches of the drug to meet its specifications):

(i) A different container size for solid oral dosage forms where container and closure are of the same materials as those provided for in the approved application.

(ii) Change in personnel not involving new facilities.

(iii) Change in equipment that does not alter the method of manufacture of a new animal drug.

(iv) Change from one commercial batch size to another without any change in manufacturing procedure.

(v) Change to more stringent specification without altering the method described in the approved application.

(vi) Inclusion of additional specifications and methods without deletion of those described in the approved application.

(vii) Alteration of specifications or methods for inactive ingredients to bring them into compliance with new or revised specifications or methods in an official compendium.

(viii) Initiation of a product identification coding system.

(ix) Addition to labeling of a reasonable expiration date where none was previously used, with related conditions of drug storage when appropriate, except when evidence shows that a significant deterioration of the drug under marketing conditions has occurred which necessitates the immediate submission of a report under §510.300(b)(1) of this chapter. The report or written communication describing such change in labeling should include stability data justifying the expiration date and recommended conditions of storage.

(x) Change from paper labels to direct printing on glass or other kinds of immediate containers without a change in text.

(6) Approval of a supplemental new animal drug application will not be required to provide for an additional distributor to distribute a drug which is the subject of an approved new animal

drug application if the conditions described below are met prior to putting such a change into effect. An order may issue refusing approval if any condition is not met or if any of the reasons for refusing or withdrawing approval, as stated in section 512(d) and (e) of the act or § 514.110 applies. For the purposes of maintaining records and making reports under the requirements of § 510.300 of this chapter, a distributor provided for under this section shall be considered an *applicant* within the meaning of § 510.300(b) of this chapter. Said conditions are:

(i) A supplemental application is furnished to the Food and Drug Administration to provide for a designated distributor.

(ii) There are no changes from the conditions of the approved application except for a different and suitable proprietary name of the new animal drug (if one is used) and the name and address of the distributor as used on the label and labeling. The name of the distributor shall be accompanied by an appropriate qualifying phrase such as “manufactured for” or “distributed by.”

(iii) A distributor’s statement is furnished to the Food and Drug Administration identifying the category of his operations (for example, wholesaler, retailer) and stating: That he will distribute the new animal drug only under the labeling provided for in the new animal drug application; that any other labeling or advertising for the drug will prescribe, recommend, or suggest its use only under the conditions stated in the labeling provided for in the application; and, if the drug is a prescription article, that he is regularly and lawfully engaged in the distribution or dispensing of prescription drugs.

(iv) Nine copies of the printed labels and other labeling to be used by the distributor are submitted, identified with the new animal drug application number.

(b) When necessary for the safety or effectiveness of the drug, a supplemental new animal drug application shall specify a period of time within which the proposed change will be made.

(c) If a material change is made in the components’ composition, manufacturing methods, facilities, or controls, or in the labeling or advertising, from the representations in an approved application for a new animal drug (except changes conforming to the conditions set forth in paragraph (a)(5) and (6) and/or paragraphs (d), (e), (f), and (g) of this section), and the drug is marketed before a supplement is approved for such change, approval of the application may be suspended or withdrawn as provided in section 512(e) of the act.

(d) Changes of the following kinds proposed in supplemental new animal drug applications should be placed into effect at the earliest possible time:

(1) The addition to package labeling, promotional labeling, and prescription drug advertising of additional warning, contraindication, side effect, and precaution information.

(2) The deletion from package labeling, promotional labeling, and drug advertising of false, misleading, or unsupported indications for use or claims for effectiveness.

(3) Changes in the methods, facilities, or controls used for the manufacture, processing, packing, or holding of the new animal drug (other than utilization of establishments not covered by the approval that is in effect) that give increased assurance that the drug will have the characteristics of identity, strength, quality, and purity which it purports or is represented to possess.

(e) The Food and Drug Administration will take no action against a new animal drug or applicant solely because changes of the kinds described in paragraph (d) of this section are placed into effect by the applicant prior to his receipt of a written notice of approval of the supplemental new animal drug application if all the following conditions are met:

(1) The supplemental new animal drug application providing a full explanation of the basis for the changes has been submitted, plainly marked on the mailing cover and on the supplement, “Special new animal drug application Supplement—changes being effected.”

(2) The applicant specifically informs the Food and Drug Administration of the date on which such changes are

§ 514.8**21 CFR Ch. I (4–1–99 Edition)**

being effected and submits to the Administration nine printed copies of any revised labeling to be placed in use, identified with the new animal drug application number.

(3) All promotional labeling and all drug advertising are promptly revised consistent with the changes made in the labeling on or within the new animal drug package.

(f) When a supplemental new animal drug application proposes changes only of the kinds described in paragraph (d) of this section, and the applicant informs the Food and Drug Administration that the changes are being put into effect, such notification will be regarded as an agreement by the applicant to an extension of the time for formal action on the application.

(g) In addition to changes as permitted by paragraphs (d) and (e) of this section, an applicant may place into effect changes proposed in a supplement to a new animal drug application that became effective prior to October 10, 1962, upon written notification from the Food and Drug Administration that such action is permitted, without approval of the supplemental application, pending the completion of the review of the effectiveness of such drug by the National Academy of Sciences-National Research Council and a determination as to whether there are grounds for refusing approval under section 512(d) of the act or for invoking section 512(e) of the act. The Food and Drug Administration will take no action against a new animal drug or an applicant solely because changes that have been permitted in a written communication are placed into effect by the applicant prior to his receipt of a written notice of approval of the supplemental new animal drug application.

(h) Except as provided in paragraphs (e) and (g) of this section, no provision of this section shall limit the authority of the Secretary or of the Commissioner to suspend or withdraw approval of a new animal drug application in accord with the provisions of section 512(e) of the act or to initiate any other regulatory proceedings with respect to a drug or applicant under provisions of the act.

(i) Changes from the conditions of an approved new animal drug application in accord with the provisions of paragraphs (d), (e), and (g) of this section are permitted on the basis of a temporary deferral of final action on the supplemental application under the provisions of section 512 (c), (d), or (e) of the act.

(j) When an applicant receives written notification from the Food and Drug Administration, under the provisions of paragraph (g) of this section, that he may place into effect changes proposed in a supplemental application without approval of the supplemental application, he may within 30 days submit a written request that the Food and Drug Administration process the supplemental application. In such case, the change shall not be put into effect until approved. Within 180 days of the receipt of such written request, the Food and Drug Administration will approve the supplemental application or furnish notice of an opportunity for a hearing under the provisions of section 512 (d) or (e), or both, of the act on a proposal to refuse approval of the supplemental application or to withdraw approval of the application and supplements thereto.

(k) A supplement to an application that became effective prior to October 10, 1962, may include a written statement to the effect that a temporary deferral of final action under the provisions of paragraph (d), (e), or (g) of this section is unacceptable to the applicant and that the applicant requests action as provided in section 512(c) of the act. Final action on such supplemental applications will be expedited in accord with applicable provisions of section 512 of the act and regulations in this subchapter E. In such cases, if the applicant places into effect any of the proposed changes prior to his receipt of a written notice of approval of the supplemental new animal drug application, such action may be regarded by the Food and Drug Administration as a basis for invoking the provisions of section 512(e)(1)(D) of the act; that is, the applicant may be furnished notice of an opportunity for a hearing on a proposal to withdraw approval of the application on the ground that the application contains an untrue statement

Food and Drug Administration, HHS

§ 514.11

of a material fact related to the changes from the conditions approved in the application.

(l) A supplemental application that contains nonclinical laboratory studies shall include, with respect to each nonclinical study, either a statement that the study was conducted in compliance with the requirements set forth in part 58 of this chapter, or, if the study was not conducted in compliance with such regulations, a brief statement of the reason for the noncompliance.

[40 FR 13825, Mar. 27, 1975, as amended at 50 FR 7517, Feb. 22, 1985; 50 FR 16668, Apr. 26, 1985; 62 FR 40600, July 29, 1997]

§ 514.9 Supplemental applications for animal feeds bearing or containing new animal drugs.

(a) After an application for an animal feed bearing or containing a new animal drug has been approved, a supplemental application may propose changes.

(b) A supplemental application shall be submitted for any change which deviates from the conditions under which the application was originally approved.

(c) Each supplemental application shall be accompanied by a fully completed Form FDA 1900 in triplicate including an explanation of the changes proposed.

[40 FR 13825, Nov. 27, 1975, as amended at 42 FR 15675, Mar. 22, 1977; 50 FR 16668, Apr. 26, 1985; 51 FR 7391, Mar. 3, 1986]

§ 514.10 Confidentiality of data and information in an investigational new animal drug notice and a new animal drug application file for an antibiotic drug.

(a) The rules established in §§ 514.11 and 514.12 of this chapter with regard to the confidentiality of an investigational new animal drug notice and a new animal drug application file shall apply to such notices and files for antibiotic drugs for new animal drug use.

(b) All records showing the Food and Drug Administration's testing of and action on a particular lot of a certifiable antibiotic drug for veterinary use are immediately available for public disclosure.

EFFECTIVE DATE NOTE: At 64 FR 403, Jan. 5, 1999, § 514.10 was removed, effective May 20, 1999.

§ 514.11 Confidentiality of data and information in a new animal drug application file.

(a) For purposes of this section the *NADA file* includes all data and information submitted with or incorporated by reference in the NADA, INAD's incorporated into the NADA, supplemental NADA's, reports under §§ 510.300 and 510.301 of this chapter, master files, and other related submissions. The availability for public disclosure of any record in the NADA file shall be handled in accordance with the provisions of this section.

(b) The existence of an NADA file will not be disclosed by the Food and Drug Administration before an approval has been published in the FEDERAL REGISTER, unless it has previously been publicly disclosed or acknowledged.

(c) If the existence of an NADA file has not been publicly disclosed or acknowledged, no data or information in the NADA file is available for public disclosure.

(d) If the existence of an NADA file has been publicly disclosed or acknowledged before an approval has been published in the FEDERAL REGISTER, no data or information contained in the file is available for public disclosure before such approval is published, but the Commissioner may, in his discretion, disclose a summary of such selected portions of the safety and effectiveness data as are appropriate for public consideration of a specific pending issue, e.g., at an open session of a Food and Drug Administration advisory committee or pursuant to an exchange of important regulatory information with a foreign government.

(e) After an approval has been published in the FEDERAL REGISTER, the following data and information in the NADA file are immediately available for public disclosure unless extraordinary circumstances are shown:

(1) All safety and effectiveness data and information previously disclosed to the public, as defined in § 20.81 of this chapter.

(2) A summary or summaries of the safety and effectiveness data and information submitted with or incorporated by reference in the NADA file. Such summaries do not constitute the full reports of investigations under section 512(b)(1) of the act (21 U.S.C. 360b(b)(1)) on which the safety or effectiveness of the drug may be approved. Such summaries shall consist of the following:

(i) For an NADA approved prior to July 1, 1975, internal agency records that describe such data and information, e.g., a summary of basis for approval or internal reviews of the data and information, after deletion of:

(a) Names and any information that would identify the investigators.

(b) Any inappropriate gratuitous comments unnecessary to an objective analysis of the data and information.

(ii) For an NADA approved on or after July 1, 1975, a summary of such data and information prepared in one of the following two alternative ways shall be publicly released when the approval is published in the FEDERAL REGISTER.

(a) The Center for Veterinary Medicine may at an appropriate time prior to approval of the NADA require the applicant to prepare a summary of such data and information, which will be reviewed and, where appropriate, revised by the Center.

(b) The Center for Veterinary Medicine may prepare its own summary of such data and information.

(3) A protocol for a test or study, unless it is shown to fall within the exemption established for trade secrets and confidential commercial information in § 20.61 of this chapter.

(4) Adverse reaction reports, product experience reports, consumer complaints, and other similar data and information, after deletion of:

(i) Names and any information that would identify the person using the product.

(ii) Names and any information that would identify any third party involved with the report, such as a physician, hospital, or other institution.

(5) A list of all active ingredients and any inactive ingredients previously disclosed to the public as defined in § 20.81 of this chapter.

(6) An assay method or other analytical method, unless it serves no regulatory or compliance purpose and is shown to fall within the exemption established in § 20.61 of this chapter.

(7) All correspondence and written summaries of oral discussions relating to the NADA, in accordance with the provisions of part 20 of this chapter.

(f) All safety and effectiveness data and information not previously disclosed to the public are available for public disclosure at the time any one of the following events occurs unless extraordinary circumstances are known:

(1) The NADA has been abandoned and no further work is being undertaken with respect to it.

(2) A final determination is made that the NADA is not approvable, and all legal appeals have been exhausted.

(3) Approval of the NADA is withdrawn, and all legal appeals have been exhausted.

(4) A final determination has been made that the animal drug is not a new animal drug.

(5) A final determination has been made that the animal drug may be marketed without submission of such safety and/or effectiveness data and information.

(g) The following data and information in an NADA file are not available for public disclosure unless they have been previously disclosed to the public as defined in § 20.81 of this chapter or they relate to a product or ingredient that has been abandoned and they no longer represent a trade secret or confidential commercial or financial information as defined in § 20.61 of this chapter:

(1) Manufacturing methods or processes, including quality control procedures.

(2) Production, sales, distribution, and similar data and information, except that any compilation of such data and information aggregated and prepared in a way that does not reveal data or information which is not available for public disclosure under this provision is available for public disclosure.

(3) Quantitative or semiquantitative formulas.

(h) For purposes of this regulation, safety and effectiveness data include

Food and Drug Administration, HHS

§ 514.100

all studies and tests of an animal drug on animals and all studies and tests on the animal drug for identity, stability, purity, potency, and bioavailability.

[40 FR 13825, Mar. 27, 1975, as amended at 42 FR 3109, Jan. 14, 1977; 42 FR 15675, Mar. 22, 1977; 54 FR 18280, Apr. 28, 1989]

§ 514.12 Confidentiality of data and information in an investigational new animal drug notice.

(a) The existence of an INAD notice will not be disclosed by the Food and Drug Administration unless it has previously been publicly disclosed or acknowledged.

(b) The availability for public disclosure of all data and information in an INAD file shall be handled in accordance with provisions established in § 514.11.

§ 514.15 Untrue statements in applications.

Among the reasons why an application for a new animal drug or animal feed bearing or containing a new animal drug may contain an untrue statement of a material fact are:

(a) Differences in:

(1) Conditions of use prescribed, recommended, or suggested by the applicant for the product from the conditions of such use stated in the application;

(2) Articles used as components of the product from those listed in the application;

(3) Composition of the product from that stated in the application;

(4) Methods used in or the facilities and controls used for the manufacture, processing, or packing of the product from such methods, facilities, and controls described in the application;

(5) Labeling from the specimens contained in the application; or

(b) The unexplained omission in whole or in part from an application or from an amendment or supplement to an application or from any record or report required under the provisions of section 512 of the act and § 510.300 or § 510.301 of this chapter of any information obtained from:

(1) Investigations as to the safety, effectiveness, identity, strength, quality, or purity of the drug, made by the applicant on the drug, or

(2) Investigations or experience with the product that is the subject of the application, or any related product, available to the applicant from any source if such information is pertinent to an evaluation of the safety, effectiveness, identity, strength, quality, or purity of the drug, when such omission would bias an evaluation of the safety or effectiveness of the product.

(c) Any nonclinical laboratory study contained in the application was not conducted in compliance with the good laboratory practice regulations as set forth in part 58 of this chapter, and the application fails to include a brief statement of the reason for the non-compliance.

[40 FR 13825, Mar. 27, 1975, as amended at 49 FR 7226, Feb. 28, 1984; 50 FR 7517, Feb. 22, 1985]

Subpart B—Administrative Actions on Applications

§ 514.100 Evaluation and comment on applications.

(a) After the filed application has been evaluated, the applicant will be furnished written comment on any apparent deficiencies in the application.

(b) When the description of the methods used in, and the facilities and controls used for, the manufacture, processing, and packing of such new animal drug appears adequate on its face, but it is not feasible to reach a conclusion as to the safety and effectiveness of the new animal drug solely from consideration of this description, the applicant may be notified that an establishment inspection is required to verify their adequacy.

(c) A request for samples of a new animal drug or any edible tissues and byproducts of animals treated with such a drug, shall specify the quantity deemed adequate to permit tests of analytical methods to determine their adequacy for regulatory purposes. The request should be made as early in the 180-day period as possible to assure timely completion. The date used for computing the 180-day limit for the purposes of section 512(c) of the act shall be moved forward 1 day for each day after the mailing date of the request until all of the requested samples

§ 514.105

21 CFR Ch. I (4–1–99 Edition)

are received. If the samples are not received within 90 days after the request, the application will be considered withdrawn without prejudice.

(d) The information contained in an application may be insufficient to determine whether a new animal drug is safe or effective in use if it fails to include (among other things) a statement showing whether such drug is to be limited to prescription sale and exempt under section 502(f) of the act from the requirement that its labeling bear adequate directions for lay use. If such drug is to be exempt, the information may also be insufficient if:

(1) The specimen labeling proposed fails to bear adequate information for professional use including indications, effects, dosages, routes, methods, and frequency and duration of administration and any relevant hazards, contraindications, side effects, and precautions under which practitioners licensed by law to administer such drug can use the drug for the purposes for which it is intended, including all purposes for which it is to be advertised, or represented, in accordance with § 201.105 of this chapter, and information concerning hazards, contraindications, side effects, and precautions relevant with respect to any uses for which such drug is to be prescribed.

(2) The application fails to show that the labeling and advertising of such drug will offer the drug for use only under those conditions for which it is offered in the labeling that is part of the application.

(3) The application fails to show that all labeling that furnishes or purports to furnish information for professional use of such drug will contain, in the same language and emphasis, the information for use including indications, effects, dosages, routes, methods, and frequency and duration of administration and any relevant warnings, hazards, contraindications, side effects, and precautions, which is contained in the labeling that is part of the application in accordance with § 201.105 of this chapter.

(e) The information contained in an application will be considered insufficient to determine whether a new animal drug is safe and effective for use when there is a refusal or failure upon

written notice to furnish inspectors authorized by the Food and Drug Administration an adequate opportunity to inspect the facilities, controls, and records pertinent to the application.

(f) On the basis of preliminary consideration of an application or supplemental application containing type-written or other draft labeling in lieu of final printed labeling, an applicant may be informed that such application is approvable when satisfactory final printed labeling identical in content to such draft copy is submitted.

(g) When an application has been found incomplete on the basis of a need for the kind of information described in § 514.6, such application shall be considered withdrawn without prejudice to future filing on the date of issuance of the letter citing the inadequacies contained in the application, unless within 30 days the sponsor chooses to avail himself of the opportunity for hearing as prescribed by § 514.111.

§ 514.105 Approval of applications.

(a) Within 180 days after an application has been filed pursuant to § 514.1, if the Commissioner determines that none of the grounds for denying approval specified in section 512(d) of the act applies:

(1) He shall forward for publication in the FEDERAL REGISTER a regulation prescribing the conditions under which the new animal drug may be used, including the name and address of the applicant; the conditions and indications for use covered by the application; any tolerance, withdrawal period, or other use restrictions; any tolerance required for the new animal drug substance or its metabolites in edible products of food-producing animals; and, if such new animal drug is intended for use in animal feed, appropriate purposes and conditions of use (including special labeling requirements) applicable to any animal feed; and such other information the Commissioner deems necessary to assure safe and effective use.

(2) He shall notify the applicant by sending him a copy of the proposed publication as described in paragraph (a)(1) of this section.

(b) Within 90 days after an application filed pursuant to § 514.2 if the Commissioner determines that none of the

grounds for denying approval specified in section 512(m)(3) of the act applies, he shall notify the applicant that it is approvable by signing and mailing to the sponsor the original copy of the Form FDA 1900.

[40 FR 13825, Mar. 27, 1975, as amended at 51 FR 7392, Mar. 3, 1986]

§ 514.106 Approval of supplemental applications.

(a) Within 180 days after a supplement to an approved application is filed pursuant to § 514.8, the Commissioner shall approve the supplemental application in accordance with procedures set forth in § 514.105(a)(1) and (2) if he/she determines that the application satisfies the requirements of applicable statutory provisions and regulations.

(b) The Commissioner will assign a supplemental application to its proper category to ensure processing of the application.

(1) *Category I.* Supplements that ordinarily do not require a reevaluation of any of the safety or effectiveness data in the parent application. Category I supplements include the following:

(i) A corporate change that alters the identity or address of the sponsor of the new animal drug application (NADA).

(ii) The sale, purchase, or construction of manufacturing facilities.

(iii) The sale or purchase of an NADA.

(iv) A change in container, container style, shape, size, or components.

(v) A change in approved labeling (color, style, format, addition, deletion, or revision of certain statements, e.g., trade name, storage, expiration dates, etc).

(vi) A change in promotional material for a prescription drug not exempted by § 514.8(a)(3)(i) and (a)(3)(ii).

(vii) Changes in manufacturing processes that do not alter the method of manufacture or change the final dosage form.

(viii) A change in bulk drug shipments.

(ix) A change in an analytical method or control procedures that do not alter the approved standards.

(x) A change in an expiration date.

(xi) Addition of an alternate manufacturer, repackager, or relabeler of the drug product.

(xii) Addition of an alternate supplier of the new drug substance.

(xiii) A change permitted in advance of approval as listed in § 514.8(d).

(xiv) Changes not requiring prior approval which are listed under § 514.8(a)(5) when submitted as supplemental applications.

(2) *Category II.* Supplements that may require a reevaluation of certain safety or effectiveness data in the parent application. Category II supplements include the following:

(i) A change in the active ingredient concentration or composition of the final product.

(ii) A change in quality, purity, strength, and identity specifications of the active or inactive ingredients.

(iii) A change in dose (amount of drug administered per dose).

(iv) A change in the treatment regimen (schedule of dosing).

(v) Addition of a new therapeutic claim to the approved uses of the product.

(vi) Addition of a new or revised animal production claim.

(vii) Addition of a new species.

(viii) A change in the prescription or over-the-counter status of a drug product.

(ix) A change in statements regarding side effects, warnings, precautions, and contraindications, except the addition of approved statements to container, package, and promotional labeling, and prescription drug advertising.

(x) A change in the drug withdrawal period prior to slaughter or in the milk discard time.

(xi) A change in the tolerance for drug residues.

(xii) A change in analytical methods for drug residues.

(xiii) A revised method of synthesis or fermentation of the new drug substance.

(xiv) Updating or changes in the manufacturing process of the new drug substance and/or final dosage form (other than a change in equipment that does not alter the method of manufacture of a new animal drug, or a change

§ 514.110

from one commercial batch size to another without any change in manufacturing procedure), or changes in the methods, facilities, or controls used for the manufacture, processing, packaging, or holding of the new animal drug (other than use of an establishment not covered by the approval that is in effect) that give increased assurance that the drug will have the characteristics of identity, strength, quality, and purity which it purports or is represented to possess.

[55 FR 46052, Nov. 1, 1990; 55 FR 49973, Dec. 3, 1990; 56 FR 12422, Mar. 25, 1991]

§ 514.110 Reasons for refusing to file applications.

(a) The date of receipt of an application for a new animal drug shall be the date on which the application shall be deemed to be filed.

(b) An application for a new animal drug shall not be considered acceptable for filing for any of the following reasons:

(1) It does not contain complete and accurate English translations of any pertinent part in a foreign language.

(2) Fewer than three copies are submitted.

(3) It is incomplete on its face in that it is not properly organized and indexed.

(4) On its face the information concerning required matter is so inadequate that the application is clearly not approvable.

(5) The new animal drug is to be manufactured, prepared, propagated, compounded, or processed in whole or in part in any State in an establishment that has not been registered or exempted from registration under the provisions of section 510 of the act.

(6) The sponsor does not reside or maintain a place of business within the United States and the application has not been countersigned by an attorney, agent, or other representative of the applicant, which representative resides in the United States and has been duly authorized to act on behalf of the applicant and to receive communications on all matters pertaining to the application.

(7) The new animal drug is a drug subject to licensing under the animal virus, serum, and toxin law of March 4,

21 CFR Ch. I (4–1–99 Edition)

1913 (37 Stat. 832; 21 U.S.C. 151 *et seq.*). Such applications will be referred to the U.S. Department of Agriculture for action.

(8) It fails to include, with respect to each nonclinical laboratory study contained in the application, either a statement that the study was conducted in compliance with the good laboratory practice regulations set forth in part 58 of this chapter, or, if the study was not conducted in compliance with such regulations, a brief statement of the reasons for the non-compliance.

(9) [Reserved]

(10) The applicant fails to submit a complete environmental assessment under § 25.40 of this chapter or fails to provide sufficient information to establish that the requested action is subject to categorical exclusion under § 25.30 or § 25.33 of this chapter.

(c) If an application is determined not to be acceptable for filing, the applicant shall be notified within 30 days of receipt of the application and shall be given the reasons therefore.

(d) If the applicant disputes the findings that his application is not acceptable for filing, he may make written request that the application be filed over protest, in which case it will be filed as of the day originally received.

[40 FR 13825, Mar. 27, 1975, as amended at 50 FR 7517, Feb. 22, 1985; 50 FR 16668, Apr. 26, 1985; 62 FR 40600, July 29, 1997]

§ 514.111 Refusal to approve an application.

(a) The Commissioner shall, within 180 days after the filing of the application, inform the applicant in writing of his intention to issue a notice of opportunity for a hearing on a proposal to refuse to approve the application, if the Commissioner determines upon the basis of the application, or upon the basis of other information before him with respect to a new animal drug, that:

(1) The reports of investigations required to be submitted pursuant to section 512(b) of the act do not include adequate tests by all methods reasonably applicable to show whether or not such drug is safe for use under the conditions prescribed, recommended, or

suggested in the proposed labeling thereof; or

(2) The results of such tests show that such drug is unsafe for use under such conditions or do not show that such drug is safe for use under such conditions; or

(3) The methods used in and the facilities and controls used for the manufacture, processing, and packing of such drug are inadequate to preserve its identity, strength, quality, and purity; or

(4) Upon the basis of the information submitted to the Food and Drug Administration as part of the application, or upon the basis of any other information before it with respect to such drug, it has insufficient information to determine whether such drug is safe for use under such conditions. In making this determination the Commissioner shall consider, among other relevant factors:

(i) The probable consumption of such drug and of any substance formed in or on food because of the use of such drug;

(ii) The cumulative effect on man or animal of such drug, taking into account any chemically or pharmacologically related substances;

(iii) Safety factors which, in the opinion of experts qualified by scientific training and experience to evaluate the safety of such drugs, are appropriate for the use of animal experimentation data; and

(iv) Whether the conditions of use prescribed, recommended, or suggested in the proposed labeling are reasonably certain to be followed in practice; or

(5) Evaluated on the basis of information submitted as part of the application and any other information before the Food and Drug Administration with respect to such drug, there is lack of substantial evidence consisting of one or more adequate and well-controlled studies by experts qualified by scientific training and experience to evaluate the effectiveness of the drug involved, on the basis of which it could fairly and reasonably be concluded by such experts that the drug will have the effect it purports to be represented to have under the conditions of use prescribed, recommended, or suggested in the labeling or proposed labeling thereof.

(6) Failure to include an appropriate proposed tolerance for residues in edible products derived from animals or a withdrawal period or other restrictions for use of such drug if any tolerance or withdrawal period or other restrictions for use are required in order to assure that the edible products derived from animals treated with such drug will be safe.

(7) Based on a fair evaluation of all material facts, the labeling is false or misleading in any particular; or

(8) Such drug induces cancer when ingested by man or animal or, after appropriate tests for evaluation of the safety of such drug, induces cancer in man or animal, except that this subparagraph shall not apply with respect to such drug if the Commissioner finds that, under the conditions of use specified in proposed labeling and reasonably certain to be followed in practice:

(i) Such drug will not adversely affect the animal for which it is intended; and

(ii) No residue of such drug will be found (by methods of examination prescribed or approved by the Commissioner by regulations) in any edible portion of such animal after slaughter or in any food yielded by, or derived from the living animals.

(9) The applicant fails to submit an adequate environmental assessment under § 25.40 of this chapter or fails to provide sufficient information to establish that the requested action is subject to categorical exclusion under § 25.30 or § 25.33 of this chapter.

(10) The drug fails to satisfy the requirements of subpart E of part 500 of this chapter.

(11) Any nonclinical laboratory study that is described in the application and that is essential to show that the drug is safe for use under the conditions prescribed, recommended, or suggested in its proposed labeling, was not conducted in compliance with the good laboratory practice regulations as set forth in part 58 of this chapter and no reason for the noncompliance is provided or, if it is, the differences between the practices used in conducting the study and the good laboratory practice regulations do not support the validity of the study.

§ 514.112

21 CFR Ch. I (4–1–99 Edition)

(b) The Commissioner shall within 90 days after the filing of the application inform the applicant in writing of his intention to issue a notice of opportunity for a hearing on a proposal to refuse to approve the application, if the Commissioner determines upon the basis of the application, or upon the basis of other information before him with respect to an animal feed bearing or containing a new animal drug that:

(1) There is not in effect a regulation established pursuant to section 512(i) of the act (identified in such application) on the basis of which such application may be approved; or

(2) Such animal feed (including the proposed use of any new animal drug therein or thereon) does not conform to an applicable regulation published pursuant to section 512(i) of the act (identified in such application), or that the purposes or conditions or indications of use prescribed, recommended, or suggested in the labeling of such feed do not conform to the applicable purposes and conditions or indications for use (including warnings) published pursuant to section 512(i) of the act or such labeling omits or fails to conform to other applicable information published pursuant to such section; or

(3) The methods used in and the facilities and controls used for the manufacturing, processing, and packaging of such animal feed are not adequate to preserve the identity, strength, quality, and purity of the new animal drug therein; or

(4) Based on a fair evaluation of all the material facts, such labeling is false or misleading in any particular.

(c) The Commissioner, as provided in § 514.200 of this chapter, shall expeditiously notify the applicant of an opportunity for a hearing on the question of whether such application is approvable, unless by the 30th day following the date of issuance of the letter informing the applicant of the intention to issue a notice of opportunity for a hearing the applicant:

(1) Withdraws the application; or

(2) Waives the opportunity for a hearing; or

(3) Agrees with the Commissioner on an additional period to precede issuance of such notice of hearing.

[40 FR 13825, Mar. 27, 1975, as amended at 43 FR 22675, May 26, 1978; 44 FR 16007, Mar. 16, 1979; 50 FR 7517, Feb. 22, 1985; 50 FR 16668, Apr. 26, 1985; 52 FR 49588, Dec. 31, 1987; 54 FR 18280, Apr. 28, 1989; 62 FR 40600, July 29, 1997; 63 FR 10770, Mar. 5, 1998]

§ 514.112 Return of applications for animal feeds bearing or containing new animal drugs.

Applications submitted pursuant to § 514.2 will be returned to the applicant if such applications are incomplete or inaccurate or do not contain an identification of the applicable regulation(s). These regulations include those published pursuant to section 512(i) of the act, and are found in part 558 of this chapter. In addition, § 510.515 of this chapter may also provide a basis on which approval of the application relies, as required by § 514.2(b)(10). All reasons for the return of the application will be made known to the applicant.

[51 FR 7392, Mar. 3, 1986]

§ 514.115 Withdrawal of approval of applications.

(a) The Secretary may suspend approval of an application approved pursuant to section 512(c) or (m)(2) of the act and give the applicant prompt notice of his action and afford the applicant the opportunity for an expedited hearing on a finding that there is an imminent hazard to the health of man or of the animals for which such new animal drug or animal feed is intended.

(b) The Commissioner shall notify in writing the person holding an application approved pursuant to section 512(c) or (m)(2) of the act and afford an opportunity for a hearing on a proposal to withdraw approval of such application if he finds:

(1) That the application contains any untrue statement of a material fact; or

(2) That the applicant has made any changes from the standpoint of safety or effectiveness beyond the variations provided for in the application unless he has supplemented the application by filing with the Secretary adequate information respecting all such changes

and unless there is in effect an approval of the supplemental application, or such changes are those for which written authorization or approval is not required as provided for in § 514.8. The supplemental application shall be treated in the same manner as the original application.

(3) That in the case of an application for use of a new animal drug approved or deemed approved pursuant to section 512(c) of the act:

(i) Experience or scientific data show that such drug is unsafe for use under the conditions of use upon the basis of which the application was approved; or

(ii) New evidence not contained in such application or not available to the Secretary until after such application was approved, or tests by new methods, or tests by methods not deemed reasonably applicable when such application was approved, evaluated together with the evidence available to the Secretary when the application was approved, shows that such drug is not shown to be safe for use under the conditions of use upon the basis of which the application was approved or that section 512 (d)(1)(H) of the act applies to such drug; or

(iii) On the basis of new information before him with respect to such drug, evaluated together with the evidence available to him when the application was approved, there is a lack of substantial evidence that such drug will have the effect it purports or is represented to have under the conditions of use prescribed, recommended, or suggested in the labeling thereof.

(4) That any nonclinical laboratory study that is described in the application and that is essential to show that the drug is safe for use under the conditions prescribed, recommended, or suggested in its proposed labeling, was not conducted in compliance with the good laboratory practice regulations as set forth in part 58 of this chapter and no reason for the noncompliance is provided or, if it is, the differences between the practices used in conducting the study and the good laboratory practice regulations do not support the validity of the study.

(c) The Commissioner may notify in writing the person holding an application approved pursuant to section

512(c) or (m)(2) of the act and afford an opportunity for a hearing on a proposal to withdraw approval of such application if he finds:

(1) That the applicant has failed to establish a system for maintaining required records, or has repeatedly or deliberately failed to maintain such records or to make required reports in accordance with a regulation or order under section 512(l)(1) or (m)(5)(A) of the act, or the applicant has refused to permit access to, or copying, or verification of, such records as required by section 512(l)(2) or (m)(5)(B) of the act; or

(2) That on the basis of new information before him evaluated together with the evidence before him when the application was approved, the methods used in, or the facilities and controls used for, the manufacture, processing, and packing of such drug or animal feed are inadequate to assure and preserve its identity, strength, quality, and purity and were not made adequate within a reasonable time after receipt of written notice from the Secretary specifying the matter complained of; or

(3) That on the basis of new information before him, evaluated together with the evidence before him when the application was approved, the labeling of such drug or animal feed, based on a fair evaluation of all material facts, is false or misleading in any particular and was not corrected within a reasonable time after receipt of written notice from the Secretary specifying the matter complained of.

(d) Approval of an application pursuant to section 512(c) or (m)(2) of the act will be withdrawn on the basis of a request for its withdrawal submitted in writing by a person holding an approved new animal drug application on the grounds that the drug subject to such application is no longer being marketed and information is included in support of this finding, provided none of the conditions cited in paragraphs (a), (b), and (c) of this section pertain to the subject drug. A written request for such withdrawal shall be construed as a waiver of the opportunity for a hearing as otherwise provided for in this section. Withdrawal of approval of an application under the

§ 514.116

21 CFR Ch. I (4–1–99 Edition)

provisions of this paragraph shall be without prejudice.

(e) On the basis of the withdrawal of approval of an application for a new animal drug approved pursuant to section 512(c) of the act, the regulation published pursuant to section 512(i) of the act covering the conditions of use of such drug as provided for in the application shall be revoked. An application providing for the manufacture of animal feeds bearing or containing such drug and approved pursuant to section 512(m)(2) of the act shall be deemed as withdrawn upon publication in the FEDERAL REGISTER of the order revoking the corresponding regulation.

[40 FR 13825, Mar. 27, 1975, as amended at 50 FR 7517, Feb. 22, 1985]

§ 514.116 Notice of withdrawal of approval of application.

When an approval of an application submitted pursuant to section 512 of the act is withdrawn by the Commissioner, he will give appropriate public notice of such action by publication in the FEDERAL REGISTER.

§ 514.117 Adequate and well-controlled studies.

(a) *Purpose.* The primary purpose of conducting adequate and well-controlled studies of a new animal drug is to distinguish the effect of the new animal drug from other influences, such as spontaneous change in the course of the disease, normal animal production performance, or biased observation. One or more adequate and well-controlled studies are required to establish, by substantial evidence, that a new animal drug is effective. The characteristics described in paragraph (b) of this section have been developed over a period of years and are generally recognized as the essentials of an adequate and well-controlled study. Well controlled, as used in the phrase adequate and well controlled, emphasizes an important aspect of adequacy. The Food and Drug Administration (FDA) considers these characteristics in determining whether a study is adequate and well controlled for purposes of section 512 of the Federal Food, Drug, and Cosmetic Act (the act) (21 U.S.C. 360b). Adequate and well-controlled studies, in addition to providing a basis for de-

termining whether a new animal drug is effective, may also be relied upon to support target animal safety. The report of an adequate and well-controlled study should provide sufficient details of study design, conduct, and analysis to allow critical evaluation and a determination of whether the characteristics of an adequate and well-controlled study are present.

(b) *Characteristics.* An adequate and well-controlled study has the following characteristics:

(1) The protocol for the study (protocol) and the report of the study results (study report) must include a clear statement of the study objective(s).

(2) The study is conducted in accordance with an appropriate standard of conduct that addresses, among other issues, study conduct, study personnel, study facilities, and study documentation. The protocol contains a statement acknowledging the applicability of, and intention to follow, a standard of conduct acceptable to FDA. The study report contains a statement describing adherence to the standard.

(3) The study is conducted with a new animal drug that is produced in accordance with appropriate manufacturing practices, which include, but are not necessarily limited to, the manufacture, processing, packaging, holding, and labeling of the new animal drug such that the critical characteristics of identity, strength, quality, purity, and physical form of the new animal drug are known, recorded, and reproducible, to permit meaningful evaluations of and comparisons with other studies conducted with the new animal drug. The physical form of a new animal drug includes the formulation and physical characterization (including delivery systems thereof, if any) of the new animal drug as presented to the animal. The protocol and study report must include an identification number which can be correlated with the specific formulation and production process used to manufacture the new animal drug used in the study.

(4) The study uses a design that permits a valid comparison with one or more controls to provide a quantitative evaluation of drug effects. The protocol and the study report must describe the

precise nature of the study design, e.g., duration of treatment periods, whether treatments are parallel, sequential, or crossover, and the determination of sample size. Within the broad range of studies conducted to support a determination of the effectiveness of a new animal drug, certain of the controls listed below would be appropriate and preferred depending on the study conducted:

(i) *Placebo concurrent control.* The new animal drug is compared with an inactive preparation designed to resemble the new animal drug as far as possible.

(ii) *Untreated concurrent control.* The new animal drug is compared with the absence of any treatment. The use of this control may be appropriate when objective measurements of effectiveness, not subject to observer bias, are available.

(iii) *Active treatment concurrent control.* The new animal drug is compared with known effective therapy. The use of this control is appropriate when the use of a placebo control or of an untreated concurrent control would unreasonably compromise the welfare of the animals. Similarity of the new animal drug and the active control drug can mean either that both drugs were effective or that neither was effective. The study report should assess the ability of the study to have detected a difference between treatments. The evaluation of the study should explain why the new animal drugs should be considered effective in the study, for example, by reference to results in previous placebo-controlled studies of the active control.

(iv) *Historical control.* The results of treatment with the new animal drug are quantitatively compared with experience historically derived from the adequately documented natural history of the disease or condition, or with a regimen (therapeutic, diagnostic, prophylactic) whose effectiveness is established, in comparable animals. Because historical control populations usually cannot be as well assessed with respect to pertinent variables as can concurrent control populations, historical control designs are usually reserved for special circumstances. Examples include studies in which the effect of the new animal drug is self-evident or stud-

ies of diseases with high and predictable mortality, or signs and symptoms of predictable duration or severity, or, in the case of prophylaxis, predictable morbidity.

(5) The study uses a method of selecting animals that provides adequate assurances that the animals are suitable for the purposes of the study. For example, the animals can reasonably be expected to have animal production characteristics typical of the class(es) of animals for which the new animal drug is intended, there is adequate assurance that the animals have the disease or condition being studied, or, in the case of prophylactic agents, evidence of susceptibility and exposure to the condition against which prophylaxis is desired has been provided. The protocol and the study report describe the method of selecting animals for the study.

(6) The study uses a method to assign a treatment or a control to each experimental unit of animals that is random and minimizes bias. Experimental units of animals are groups of animals that are comparable with respect to pertinent variables such as age, sex, class of animal, severity of disease, duration of disease, dietary regimen, level of animal production, and use of drugs or therapy other than the new animal drug. The protocol and the study report describe the method of assignment of animals to an experimental unit to account for pertinent variables and method of assignment of a treatment or a control to the experimental units. When the effect of such variables is accounted for by an appropriate design, and when, within the same animal, effects due to the test drug can be obtained free of the effects of such variables, the same animal may be used for both the test drug and the control using the controls set forth in paragraph (b)(4) of this section.

(7) The study uses methods to minimize bias on the part of observers and analysts of the data that are adequate to prevent undue influences on the results and interpretation of the study data. The protocol and study report explain the methods of observation and recording of the animal response variables and document the methods, such as “blinding” or “masking,” used in

the study for excluding or minimizing bias in the observations.

(8) The study uses methods to assess animal response that are well defined and reliable. The protocol and study report describe the methods for conducting the study, including any appropriate analytical and statistical methods, used to collect and analyze the data resulting from the conduct of the study, describe the criteria used to assess response, and, when appropriate, justify the selection of the methods to assess animal response.

(9) There is an analysis and evaluation of the results of the study in accord with the protocol adequate to assess the effects of the new animal drug. The study report evaluates the methods used to conduct, and presents and evaluates the results of, the study as to their adequacy to assess the effects of the new animal drug. This evaluation of the results of the study assesses, among other items, the comparability of treatment and control groups with respect to pertinent variables and the effects of any interim analyses performed.

(c) *Field studies.* (1) Field conditions as used in this section refers to conditions which closely approximate the conditions under which the new animal drug, if approved, is intended to be applied or administered.

(2) Studies of a new animal drug conducted under field conditions shall, consistent with generally recognized scientific principles and procedures, use an appropriate control that permits comparison, employ procedures to minimize bias, and have the characteristics generally described in paragraph (b) of this section. However, because field studies are conducted under field conditions, it is recognized that the level of control over some study conditions need not or should not be the same as the level of control in laboratory studies. While not all conditions relating to a field study need to be or should be controlled, observations of the conditions under which the new animal drug is tested shall be recorded in sufficient detail to permit evaluation of the study. Adequate and well-controlled field studies shall balance the need to control study conditions with the need to observe the true effect

of the new animal drug under closely approximated actual use conditions.

(d) *Waiver.* The Director of the Center for Veterinary Medicine (the Director) may, on the Director's own initiative or on the petition of an interested person, waive in whole or in part any of the criteria in paragraph (b) of this section with respect to a specific study. A petition for a waiver is required to set forth clearly and concisely the specific criteria from which waiver is sought, why the criteria are not reasonably applicable to the particular study, what alternative procedures, if any, are to be, or have been employed, and what results have been obtained. The petition is also required to state why the studies so conducted will yield, or have yielded, substantial evidence of effectiveness, notwithstanding nonconformance with the criteria for which waiver is requested.

(e) *Uncontrolled studies.* Uncontrolled studies or partially controlled studies are not acceptable as the sole basis for the approval of claims of effectiveness or target animal safety. Such studies, carefully conducted and documented, may provide corroborative support of adequate and well-controlled studies regarding effectiveness and may yield valuable data regarding safety of the new animal drug. Such studies will be considered on their merits in light of the characteristics listed here. Isolated case reports, random experience, and reports lacking the details which permit scientific evaluation will not be considered.

[63 FR 10770, Mar. 5, 1998]

§ 514.120 Revocation of order refusing to approve an application or suspending or withdrawing approval of an application.

The Commissioner, upon his own initiative or upon request of an applicant stating reasonable grounds therefor and if he finds that the facts so require, may issue an order approving an application that previously has had its approval refused, suspended, or withdrawn.

§ 514.121 Service of notices and orders.

All notices and orders under this subchapter E and section 512 of the act

pertaining to new animal drug applications shall be served:

(a) In person by any officer or employee of the Department designated by the Commissioner; or

(b) By mailing the order by certified mail addressed to the applicant or respondent at his last known address in the records of the Food and Drug Administration.

Subpart C—Hearing Procedures

§ 514.200 Contents of notice of opportunity for a hearing.

(a) The notice to the applicant of opportunity for a hearing on a proposal by the Commissioner to refuse to approve an application or to withdraw the approval of an application will specify the grounds upon which he proposes to issue his order. On request of the applicant, the Commissioner will explain the reasons for his action. The notice of opportunity for a hearing will be published in the FEDERAL REGISTER and will specify that the applicant has 30 days after issuance of the notice within which he is required to file a written appearance electing whether:

(1) To avail himself of the opportunity for a hearing; or

(2) Not to avail himself of the opportunity for a hearing.

(b) If the applicant fails to file a written appearance in answer to the notice of opportunity for hearing, his failure will be construed as an election not to avail himself of the opportunity for the hearing, and the Commissioner without further notice may enter a final order.

(c) If the applicant elects to avail himself of the opportunity for a hearing, he is required to file a written appearance requesting the hearing within 30 days after the publication of the notice, giving the reason why the application should not be refused or should not be withdrawn, together with a well-organized and full-factual analysis of the clinical and other investigational data he is prepared to prove in support of his opposition to the Commissioner's proposal. A request for a hearing may not rest upon mere allegations or denials, but must set forth specific facts showing there is a genuine and substantial issue of fact that re-

quires a hearing. When it clearly appears from the data in the application and from the reasons and a factual analysis in the request for the hearing that no genuine and substantial issue of fact precludes the refusal to approve the application or the withdrawal of approval of the application (for example, no adequate and well-controlled clinical investigations to support the claims of effectiveness have been identified), the Commissioner will enter an order on this data, stating his findings and conclusions. If a hearing is requested and is justified by the applicant's response to the notice of opportunity for a hearing, the issues will be defined, an Administrative Law Judge will be named, and he shall issue a written notice of the time and place at which the hearing will commence. In the case of denial of approval, such time shall be not more than 90 days after the expiration of such 30 days unless the Administrative Law Judge and the applicant otherwise agree; and, in the case of withdrawal of approval, such time shall be as soon as practicable.

(d) The hearing will be open to the public; however, if the Commissioner finds that portions of the application which serve as a basis for the hearing contain information concerning a method or process entitled to protection as a trade secret, the part of the hearing involving such portions will not be public, unless the respondent so specifies in his appearance.

[40 FR 13825, Mar. 27, 1975, as amended at 43 FR 1941, Jan. 13, 1978]

§ 514.201 Procedure for hearings.

Hearings relating to new animal drugs under section 512 (d), (e), (m)(3), and (m)(4) of the act shall be governed by part 12 of this chapter.

[42 FR 4717, Jan. 25, 1977, as amended at 42 FR 10980, Feb. 25, 1977; 42 FR 15675, Mar. 22, 1977]

Subparts D–E [Reserved]

Subpart F—Judicial Review

§ 514.235 Judicial review.

(a) The transcript and record shall be certified by the Commissioner. In any

Pt. 520**21 CFR Ch. I (4–1–99 Edition)**

case in which the Commissioner enters an order without a hearing pursuant to §314.200(g) of this chapter, the request(s) for hearing together with the data and information submitted and the Commissioner's findings and conclusions shall be included in the record certified by the Commissioner.

(b) Judicial review of an order withdrawing approval of a new drug application, whether or not a hearing has been held, may be sought by a manufacturer or distributor of an identical, related, or similar drug product, as defined in §310.6 of this chapter, in a United States court of appeals pursuant to section 505(h) of the act.

[42 FR 4717, Jan. 25, 1977]

**PART 520—ORAL DOSAGE FORM
NEW ANIMAL DRUGS**

Sec.

520.23 Acepromazine maleate tablets.
520.44 Acetazolamide sodium soluble powder.
520.45 Albendazole oral dosage forms.
520.45a Albendazole suspension.
520.45b Albendazole paste.
520.48 Altrenogest solution.
520.62 Aminopentamide hydrogen sulphate tablets.
520.82 Aminopropazine fumarate oral dosage forms.
520.82a Aminopropazine fumarate tablets.
520.82b Aminopropazine fumarate, neomycin sulfate tablets.
520.88 Amoxicillin oral dosage forms.
520.88a Amoxicillin trihydrate film-coated tablets.
520.88b Amoxicillin trihydrate for oral suspension.
520.88c Amoxicillin trihydrate oral suspension.
520.88d Amoxicillin trihydrate soluble powder.
520.88e Amoxicillin trihydrate boluses.
520.88f Amoxicillin trihydrate tablets.
520.88g Amoxicillin trihydrate and clavulanate potassium film-coated tablets.
520.88h Amoxicillin trihydrate and clavulanate potassium for oral suspension.
520.90 Ampicillin oral dosage forms.
520.90a Ampicillin capsules.
520.90b Ampicillin trihydrate tablets.
520.90c Ampicillin trihydrate capsules.
520.90d Ampicillin trihydrate for oral suspension.
520.90e Ampicillin trihydrate soluble powder.
520.90f Ampicillin trihydrate boluses.

520.100 Amprolium oral dosage forms.
520.100a Amprolium drinking water.
520.100b Amprolium drench.
520.100c Amprolium crumbles.
520.110 Apramycin sulfate soluble powder.
520.154 Bacitracin oral dosage forms.
520.154a Soluble bacitracin methylene disalicylate.
520.154b Soluble bacitracin methylene disalicylate and streptomycin sulfate oral powder.
520.154c Bacitracin zinc soluble powder.
520.182 Bicyclohexylammonium fumagillin.
520.222 Bunamidine hydrochloride.
520.246 Butorphanol tartrate tablets.
520.260 *n*-Butyl chloride capsules.
520.300 Cambendazole oral dosage forms.
520.300a Cambendazole suspension.
520.300b Cambendazole pellets.
520.300c Cambendazole paste.
520.309 Carprofen caplets.
520.310 Caramiphen ethanedisulfonate and ammonium chloride tablets.
520.312 Carnidazole tablets.
520.314 Cefadroxil tablets.
520.315 Cefadroxil powder for oral suspension.
520.390 Chloramphenicol oral dosage forms.
520.390a Chloramphenicol tablets.
520.390b Chloramphenicol capsules.
520.390c Chloramphenicol palmitate oral suspension.
520.420 Chlorothiazide tablets and boluses.
520.434 Chlorphenesin carbamate tablets.
520.445 Chlortetracycline oral dosage forms.
520.445a Chlortetracycline bisulfate/sulfamethazine bisulfate soluble powder.
520.445b Chlortetracycline powder (chlortetracycline hydrochloride or chlortetracycline bisulfate).
520.445c Chlortetracycline tablets and boluses.
520.446 Clindamycin hydrochloride capsules.
520.447 Clindamycin hydrochloride liquid.
520.452 Clenbuterol syrup.
520.455 Clomipramine hydrochloride tablets.
520.462 Clorsulon drench.
520.530 Cythioate oral liquid.
520.531 Cythioate tablets.
520.534 Decoquate.
520.540 Dexamethasone oral dosage forms.
520.540a Dexamethasone powder.
520.540b Dexamethasone tablets and boluses.
520.540c Dexamethasone chewable tablets.
520.550 Dextrose/glycine/electrolyte.
520.563 Diatrizoate meglumine and diatrizoate sodium oral solution.
520.580 Dichlorophene and toluene capsules.
520.581 Dichlorophene tablets.
520.600 Dichlorvos.
520.608 Dicloxacillin sodium monohydrate capsules.
520.620 Diethylcarbamazine oral dosage forms.

Food and Drug Administration, HHS

Pt. 520

- 520.622 Diethylcarbamazine citrate oral dosage forms.
- 520.622a Diethylcarbamazine citrate tablets.
- 520.622b Diethylcarbamazine citrate syrup.
- 520.622c Diethylcarbamazine citrate chewable tablets.
- 520.622d Diethylcarbamazine citrate capsules.
- 520.623 Diethylcarbamazine citrate, oxbendazole chewable tablets.
- 520.645 Difloxacin.
- 520.763 Dithiazanine iodide oral dosage forms.
- 520.763a Dithiazanine iodide tablets.
- 520.763b Dithiazanine iodide powder.
- 520.763c Dithiazanine iodide and piperazine citrate suspension.
- 520.784 Doxylamine succinate tablets.
- 520.804 Enalapril tablets.
- 520.812 Enrofloxacin tablets.
- 520.813 Enrofloxacin oral solution.
- 520.816 Epsiprantel tablets.
- 520.823 Erythromycin phosphate.
- 520.863 Ethylisobutrazine hydrochloride tablets.
- 520.870 Etodolac.
- 520.903 Febantel oral dosage forms.
- 520.903a Febantel paste.
- 520.903b Febantel suspension.
- 520.903c [Reserved]
- 520.903d Febantel-praziquantel paste.
- 520.903e Febantel tablets.
- 520.905 Fenbendazole oral dosage forms.
- 520.905a Fenbendazole suspension.
- 520.905b Fenbendazole granules.
- 520.905c Fenbendazole paste.
- 520.905d Fenbendazole powder.
- 520.905e Fenbendazole blocks.
- 520.960 Flumethasone tablets.
- 520.970 Flunixin oral dosage forms.
- 520.970a Flunixin meglumine granules.
- 520.970b Flunixin meglumine paste.
- 520.1010 Furosemide oral dosage forms.
- 520.1010a Furosemide tablets or boluses.
- 520.1010b Furosemide powder.
- 520.1010c Furosemide syrup.
- 520.1044 Gentamicin sulfate oral dosage forms.
- 520.1044a Gentamicin sulfate oral solution.
- 520.1044b Gentamicin sulfate pig pump oral solution.
- 520.1044c Gentamicin sulfate soluble powder.
- 520.1100 Griseofulvin.
- 520.1120 Haloxon oral dosage forms.
- 520.1120a Haloxon drench.
- 520.1120b Haloxon boluses.
- 520.1130 Hetacillin oral dosage forms.
- 520.1130a Hetacillin potassium capsules.
- 520.1130b Hetacillin potassium oral suspension.
- 520.1130c Hetacillin potassium tablets.
- 520.1157 Iodinated casein tablets.
- 520.1158 Iodochlorhydroxyquin boluses.
- 520.1182 Iron dextran oral suspension.
- 520.1192 Ivermectin paste.
- 520.1193 Ivermectin tablets and chewables.
- 520.1194 Ivermectin drench.
- 520.1195 Ivermectin liquid.
- 520.1196 Ivermectin and pyrantel pamoate chewable tablet.
- 520.1197 Ivermectin sustained-release bolus.
- 520.1204 Kanamycin sulfate, aminopentamide hydrogen sulfate, pectin, bismuth subcarbonate, activated attapulgit suspension.
- 520.1205 Kanamycin sulfate, pectin, bismuth subcarbonate, activated attapulgit tablets.
- 520.1242 Levamisole hydrochloride oral dosage forms.
- 520.1242a Levamisole hydrochloride drench and drinking water.
- 520.1242b Levamisole hydrochloride tablet or oblet (bolus).
- 520.1242c Levamisole hydrochloride and piperazine dihydrochloride.
- 520.1242d Levamisole resinate.
- 520.1242e Levamisole hydrochloride effervescent tablets.
- 520.1242f Levamisole hydrochloride gel.
- 520.1242g Levamisole resinate and famphur paste.
- 520.1263 Lincomycin hydrochloride monohydrate oral dosage forms.
- 520.1263a Lincomycin hydrochloride monohydrate tablets and sirup.
- 520.1263b Lincomycin hydrochloride monohydrate and spectinomycin sulfate tetrahydrate soluble powder.
- 520.1263c Lincomycin hydrochloride soluble powder.
- 520.1284 Sodium liothyronine tablets.
- 520.1288 Lufenuron tablets.
- 520.1289 Lufenuron suspension.
- 520.1320 Mebendazole oral.
- 520.1326 Mebendazole and trichlorfon oral dosage forms.
- 520.1326a Mebendazole and trichlorfon powder.
- 520.1326b Mebendazole and trichlorfon paste.
- 520.1330 Meclofenamic acid granules.
- 520.1331 Meclofenamic acid tablets.
- 520.1341 Megestrol acetate tablets.
- 520.1380 Methocarbamol tablets.
- 520.1408 Methylprednisolone tablets.
- 520.1409 Methylprednisolone, aspirin tablets.
- 520.1422 Metoserpate hydrochloride.
- 520.1430 Mibolerone.
- 520.1445 Milbemycin oxime tablets.
- 520.1446 Milbemycin oxime/lufenuron tablets.
- 520.1448 Monensin oral dosage forms.
- 520.1448a Monensin blocks.
- 520.1450 Morantel tartrate oral dosage forms.
- 520.1450a Morantel tartrate bolus.
- 520.1450b Morantel tartrate cartridge.
- 520.1450c Morantel tartrate sustained-release trilaminar cylinder/sheet.
- 520.1451 Moxidectin.
- 520.1452 Moxidectin gel.
- 520.1468 Naproxen granules.

Pt. 520**21 CFR Ch. I (4–1–99 Edition)**

- 520.1484 Neomycin sulfate soluble powder.
520.1485 Neomycin sulfate oral solution.
520.1616 Orbifloxacin tablets.
520.1628 Oxfendazole powder and pellets.
520.1629 Oxfendazole paste.
520.1630 Oxfendazole suspension.
520.1631 Oxfendazole and trichlorfon paste.
520.1638 Oxibendazole paste.
520.1640 Oxibendazole suspension.
520.1660 Oxytetracycline.
520.1660a Oxytetracycline and carbomycin in combination.
520.1660b Oxytetracycline hydrochloride capsules.
520.1660c Oxytetracycline hydrochloride tablets/boluses.
520.1660d Oxytetracycline hydrochloride soluble powder.
520.1696 Penicillin oral dosage forms.
520.1696a Buffered penicillin powder, penicillin powder with buffered aqueous diluent.
520.1696b Penicillin G potassium in drinking water.
520.1696c Penicillin V potassium for oral solution.
520.1696d Penicillin V potassium tablets.
520.1720 Phenylbutazone oral dosage forms.
520.1720a Phenylbutazone tablets and boluses.
520.1720b Phenylbutazone granules.
520.1720c Phenylbutazone paste.
520.1720d Phenylbutazone gel.
520.1802 Piperazine-carbon disulfide complex oral dosage forms.
520.1802a Piperazine-carbon disulfide complex suspension.
520.1802b Piperazine-carbon disulfide complex boluses.
520.1802c Piperazine-carbon disulfide complex with phenothiazine suspension.
520.1803 Piperazine citrate capsules.
520.1804 Piperazine phosphate capsules.
520.1805 Piperazine phosphate with thenium closylate tablets.
520.1806 Piperazine monohydrochloride liquid.
520.1840 Poloxalene.
520.1846 Polyoxyethylene (23) lauryl ether blocks.
520.1870 Praziquantel tablets.
520.1871 Praziquantel/pyrantel pamoate tablets.
520.1872 Praziquantel, pyrantel pamoate, and febantel tablets.
520.1880 Prednisolone tablets.
520.1900 Primidone tablets.
520.1920 Prochlorperazine, isopropamide sustained release capsules.
520.1921 Prochlorperazine, isopropamide, with neomycin sustained-release capsules.
520.1962 Promazine hydrochloride.
520.2002 Propiopromazine hydrochloride.
520.2041 Pyrantel pamoate chewable tablets.
520.2042 Pyrantel pamoate tablets.
520.2043 Pyrantel pamoate suspension.
520.2044 Pyrantel pamoate paste.
520.2045 Pyrantel tartrate powder; pyrantel tartrate pellets.
520.2087 Roxarsone soluble powder.
520.2088 Roxarsone tablets.
520.2089 Roxarsone liquid.
520.2095 Sarafloxacin soluble powder.
520.2098 Selegiline hydrochloride tablets.
520.2100 Selenium, vitamin E capsules.
520.2122 Spectinomycin dihydrochloride oral solution.
520.2123 Spectinomycin dihydrochloride pentahydrate oral dosage forms.
520.2123a Spectinomycin dihydrochloride pentahydrate tablets.
520.2123b Spectinomycin dihydrochloride pentahydrate soluble powder.
520.2150 Stanozolol oral dosage forms.
520.2150a Stanozolol tablets.
520.2150b Stanozolol chewable tablets.
520.2158 Streptomycin/dihydrostreptomycin oral dosage forms.
520.2158a Streptomycin sulfate oral solution.
520.2158b Dihydrostreptomycin tablets.
520.2158c Dihydrostreptomycin oral suspension.
520.2160 Styrylpyridinium, diethylcarbazine oral dosage forms.
520.2170 Sulfabromomethazine sodium boluses.
520.2184 Sodium sulfachloropyrazine monohydrate.
520.2200 Sulfachlorpyridazine oral dosage forms.
520.2200a Sulfachlorpyridazine bolus.
520.2200b Sulfachlorpyridazine medicated milk and drinking water.
520.2200c Sulfachlorpyridazine tablets.
520.2220 Sulfadimethoxine oral dosage forms.
520.2220a Sulfadimethoxine oral solution and soluble powder.
520.2220b Sulfadimethoxine tablets and boluses.
520.2220c Sulfadimethoxine oral suspension.
520.2220d Sulfadimethoxine-ormetoprim tablets.
520.2240 Sulfaethoxypyridazine.
520.2240a Sulfaethoxypyridazine drinking water.
520.2240b Sulfaethoxypyridazine tablets.
520.2260 Sulfamethazine oral dosage forms.
520.2260a Sulfamethazine oblets and boluses.
520.2260b Sulfamethazine sustained-release boluses.
520.2260c Sulfamethazine sustained-release tablets.
520.2261 Sulfamethazine sodium oral dosage forms.
520.2261a Sulfamethazine sodium drinking water solution.
520.2261b Sulfamethazine sodium soluble powder.
520.2280 Sulfamethizole and methenamine mandelate tablets.

Food and Drug Administration, HHS

§ 520.44

520.2320 Sulfanitran and aklomide in combination.
520.2325 Sulfaquinoxaline oral dosage forms.
520.2325a Sulfaquinoxaline drinking water.
520.2325b Sulfaquinoxaline drench.
520.2330 Sulfisoxazole tablets.
520.2345 Tetracycline oral dosage forms.
520.2345a Tetracycline hydrochloride capsules.
520.2345b Tetracycline tablets.
520.2345c Tetracycline boluses.
520.2345d Tetracycline hydrochloride soluble powder.
520.2345e Tetracycline oral liquid.
520.2345f Tetracycline phosphate complex and sodium novobiocin capsules.
520.2345g Tetracycline hydrochloride and sodium novobiocin tablets.
520.2345h Tetracycline hydrochloride, sodium novobiocin, and prednisolone tablets.
520.2362 Thienium closylate tablets.
520.2380 Thiabendazole oral dosage forms.
520.2380a Thiabendazole top dressing and mineral protein feed block.
520.2380b Thiabendazole drench or oral paste.
520.2380c Thiabendazole bolus.
520.2380d Thiabendazole, piperazine citrate suspension.
520.2380e Thiabendazole with trichlorfon.
520.2380f Thiabendazole, piperazine phosphate powder.
520.2455 Tiamulin soluble powder.
520.2456 Tiamulin liquid concentrate.
520.2460 Ticarbodine oral dosage forms.
520.2460a Ticarbodine tablets.
520.2460b Ticarbodine capsules.
520.2473 Tioxidazole oral dosage forms.
520.2473a Tioxidazole granules.
520.2473b Tioxidazole paste.
520.2481 Triamcinolone acetonide tablets.
520.2482 Triamcinolone acetonide oral powder.
520.2520 Trichlorfon oral dosage forms.
520.2520a Trichlorfon oral.
520.2520b Trichlorfon and atropine.
520.2520e Trichlorofon boluses.
520.2520f Trichlorofon granules.
520.2520g Trichlorfon, phenothiazine, and piperazine dihydrochloride powder.
520.2582 Triflupromazine hydrochloride tablets.
520.2604 Trimeprazine tartrate and prednisolone tablets.
520.2605 Trimeprazine tartrate and prednisolone capsules.
520.2610 Trimethoprim and sulfadiazine tablets.
520.2611 Trimethoprim and sulfadiazine oral paste.
520.2612 Trimethoprim and sulfadiazine oral suspension.
520.2613 Trimethoprim and sulfadiazine powder.
520.2640 Tylosin.

AUTHORITY: 21 U.S.C. 360b.

SOURCE: 40 FR 13838, Mar. 27, 1975, unless otherwise noted.

§ 520.23 Acepromazine maleate tablets.

(a) *Sponsors.* See drug labeler codes in § 510.600(c) of this chapter for identification of sponsors as follows:

(1) For No. 000856, use of 5-, 10-, or 25-milligram tablets as in paragraph (b) of this section.

(2) For No. 000010, use of 10- or 25-milligram tablets as in paragraph (c) of this section.

(b) *Conditions of use.* It is used in dogs and cats as follows:

(1) *Indications for use.* It is used in dogs and cats as a tranquilizer.

(2) *Amount.* Dogs: 0.25 to 1.0 milligram per pound of body weight; Cats: 0.5 to 1.0 milligram per pound of body weight.

(3) *Limitations.* The drug is administered orally. Dosage may be repeated as required. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(c) *Conditions of use.* It is used in dogs as follows:

(1) *Indications for use.* It is used in dogs as an aid in tranquilization and as a preanesthetic agent.

(2) *Amount.* Dogs: 0.25 to 1.0 milligram per pound of body weight.

(3) *Limitations.* The drug is administered orally. Dosage may be repeated as required. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 44443, Sept. 4, 1981, as amended at 49 FR 49091, Dec. 18, 1984; 52 FR 666, Jan. 8, 1987; 53 FR 40727, Oct. 18, 1988; 56 FR 37473, Aug. 7, 1991; 62 FR 35075, June 30, 1997]

§ 520.44 Acetazolamide sodium soluble powder.

(a) *Specifications.* The drug is in a powder form containing acetazolamide sodium, USP equivalent to 25 percent acetazolamide activity.

(b) *Sponsor.* See No. 010042 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in dogs as an aid in the treatment of mild congestive heart failure and for rapid reduction of intraocular pressure.¹

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as

Continued

§ 520.45

(2) It is administered orally at a dosage level of 5 to 15 milligrams per pound of body weight daily.¹

(3) For use only by or on the order of a licensed veterinarian.¹

§ 520.45 Albendazole oral dosage forms.

§ 520.45a Albendazole suspension.

(a)(1) *Specifications.* The product contains 11.36 percent albendazole.

(2) *Sponsor.* See No. 000069 in § 510.600 of this chapter.

(3) *Related tolerances.* See § 556.34 of this chapter.

(4) *Conditions of use in cattle*—(i) *Amount.* 4.54 milligrams per pound of body weight (10 milligrams per kilogram).

(ii) *Indications for use.* For removal and control of the following internal parasites of cattle: Adult liver flukes (*Fasciola hepatica*); heads and segments of tapeworms (*Moniezia benedeni*, *M. expansa*); adult and 4th stage larvae of stomach worms (brown stomach worms including 4th stage inhibited larvae (*Ostertagia ostertagi*), barberpole worm (*Haemonchus contortus*, *H. placei*), small stomach worm (*Trichostrongylus axei*)); adult and 4th stage larvae of intestinal worms (thread-necked intestinal worm (*Nematodirus spathiger*, *N. helvetianus*), small intestinal worm (*Cooperia punctata* and *C. oncophora*)); adult stages of intestinal worms (hookworm (*Bunostomum phlebotomum*), bankrupt worm (*Trichostrongylus colubriformis*), nodular worm (*Oesophagostomum radiatum*)); adult and 4th stage larvae of lungworms (*Dictyocaulus viviparus*).

(iii) *Limitations.* Administer as a single oral dose using dosing gun or dosing syringe. Do not slaughter within 27 days of last treatment. Do not use in female dairy cattle of breeding age: Do not administer to female cattle during first 45 days of pregnancy or for 45 days after removal of bulls. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(b)(1) *Specifications.* The product contains 4.55 or 11.36 percent albendazole.

¹specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

21 CFR Ch. I (4–1–99 Edition)

(2) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(3) *Related tolerances.* See § 556.34 of this chapter.

(4) *Conditions of use in sheep*—(i) *Amount.* 7.5 milligrams per kilogram of body weight (3.4 milligrams per pound).

(ii) *Indications for use.* For removal and control of the following internal parasites of sheep: Adult liver flukes (*Fasciola hepatica*, *Fascioloides magna*); heads and segments of common tapeworms (*Moniezia expansa*) and fringed tapeworm (*Thysanosoma actinioides*); adult and fourth stage larvae of stomach worms (brown stomach worm (*Ostertagia circumcincta* and *Marshallagia marshalli*), barberpole worm (*Haemonchus contortus*), small stomach worm (*Trichostrongylus axei*)); adult and fourth stage larvae of intestinal worms (thread-necked intestinal worm (*Nematodirus spathiger* and *N. filicollis*), Cooper's worm (*Cooperia oncophora*), bankrupt worm (*Trichostrongylus colubriformis*), nodular worm (*Oesophagostomum columbianum*), and large-mouth bowel worm (*Chabertia ovina*)); adult and larval stages of lungworms (*Dictyocaulus filaria*).

(iii) *Limitations.* Administer as a single oral dose using dosing gun or dosing syringe. Do not slaughter within 7 days of last treatment. Do not administer to ewes during first 30 days of pregnancy or for 30 days after removal of rams. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[54 FR 25115, June 13, 1989, as amended at 56 FR 50653, Oct. 8, 1991; 59 FR 65711, Dec. 21, 1994; 60 FR 55658, Nov. 2, 1995; 61 FR 4875, Feb. 9, 1996; 64 FR 1504, Jan. 11, 1999]

§ 520.45b Albendazole paste.

(a) *Specifications.* The product contains 30 percent albendazole.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.34 of this chapter.

(d) *Conditions of use in cattle*—(1) *Amount.* Equivalent to 4.54 milligrams per 1 pound of body weight (10 milligrams per kilogram).

(2) *Indications for use.* For removal and control of the following internal parasites of cattle: adult liver flukes (*Fasciola hepatica*); heads and segments

Food and Drug Administration, HHS

§ 520.82a

of tapeworms (*Moniezia benedeni*, *M. expansa*); adult and 4th stage larvae of stomach worms (brown stomach worms including 4th stage inhibited larvae (*Ostertagia ostertagi*); barberpole worm (*Haemonchus contortus*, *H. placei*); small stomach worm (*Trichostrongylus axei*); adult and 4th stages larvae of intestinal worms (thread-necked intestinal worm (*Nematodirus spathiger*, *N. helvetianus*); small intestinal worm (*Cooperia punctata* and *C. oncophora*); adult stages of intestinal worms (hookworm (*Bunostomum phlebotomum*); bankrupt worm (*Trichostrongylus colubriformis*), nodular worm (*Oesophagostomum radiatum*)); adult and 4th stage larvae of lungworms (*Dictyocaulus viviparus*).

(3) *Limitations.* Administer as a single oral dose. Do not slaughter within 27 days of last treatment. Do not use in female dairy cattle of breeding age. Do not administer to female cattle during first 45 days of pregnancy or for 45 days after removal of bulls. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[54 FR 51385, Dec. 15, 1989, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55658, Nov. 2, 1995]

§ 520.48 Altrenogest solution.

(a) *Specifications.* Each milliliter of altrenogest solution contains 2.2 milligrams of altrenogest.

(b) *Sponsor.* See No. 012579 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* Administer orally at the rate of 1 milliliter per 110 pounds body weight (0.044 milligram per kilogram body weight). Give one dose daily for 15 consecutive days.

(2) *Indications for use.* For suppression of estrus in mares.

(3) *Limitations.* For oral use in horses only; avoid contact with the skin. Do not administer to horses intended for use as food. The drug is contraindicated for use in mares having a previous or current history of uterine inflammation (i.e., acute, subacute, or chronic endometritis). Natural or synthetic gestagen therapy may exacerbate existing low-grade or smoldering uterine inflammation into a fulminating uterine infection in some in-

stances. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 40887, Sept. 12, 1983, as amended at 55 FR 26431, June 28, 1990]

§ 520.62 Aminopentamide hydrogen sulphate tablets.

(a) *Chemical name.* 4-(Dimethylamino)-2,2-diphenylvaleramide hydrogen sulfate.

(b) *Specifications.* Each tablet contains 0.2 milligram of the drug.

(c) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) It is intended for use in dogs and cats only for the treatment of vomiting and/or diarrhea, nausea, acute abdominal visceral spasm, pylorospasm, or hypertrophic gastritis.

NOTE: Not for use in animals with glaucoma because of the occurrence of mydriasis.

(2) Dosage is administered by oral tablet every 8 to 12 hours, as follows:

Weight of animal in pounds	Dosage in milligrams
Up to 10	0.1
11 to 20	0.2
21 to 50	0.3
51 to 100	0.4
Over 100	0.5

Dosage may be gradually increased up to a maximum of five times the suggested dosage. Oral administration of tablets may be preceded by subcutaneous or intramuscular use of the injectable form of the drug.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 53 FR 27851, July 25, 1988]

§ 520.82 Aminopropazine fumarate oral dosage forms.

§ 520.82a Aminopropazine fumarate tablets.

(a) *Specifications.* The drug is in tablet form. Each tablet contains aminopropazine fumarate equivalent to 25 milligrams of aminopropazine base.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs and cats for reducing excessive smooth muscle contractions,

§ 520.82b

such as occur in urethral spasms associated with urolithiasis.¹

(2) It is administered at a dosage level of 1 to 2 milligrams per pound of body weight. The dosage can be repeated every 12 hours, as indicated.¹

(3) Not for use in animals intended for food purposes.

(4) For use only by or on the order of a licensed veterinarian.¹

[40 FR 13838, Mar. 27, 1975, as amended at 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61624, Nov. 19, 1997]

§ 520.82b Aminopropazine fumarate, neomycin sulfate tablets.

(a) *Specifications.* The drug is in tablet form. Each tablet contains both aminopropazine fumarate equivalent to 25 milligrams of aminopropazine base and neomycin sulfate equivalent to 50 milligrams of neomycin base.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs to control bacterial diarrhea caused by organisms susceptible to neomycin and to reduce smooth muscle contractions.¹

(2) It is administered at a dosage level of one to two tablets per 10 pounds of body weight twice daily for 3 days.¹

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13838, Mar. 27, 1975, as amended at 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61624, Nov. 19, 1997]

§ 520.88 Amoxicillin oral dosage forms.

§ 520.88a Amoxicillin trihydrate film-coated tablets.

(a) *Specifications.* Each tablet contains amoxicillin trihydrate equivalent to 50, 100, 150, 200, or 400 milligrams of amoxicillin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Dogs—(i) Amount.* 5 milligrams per pound of body weight, twice a day.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

21 CFR Ch. I (4–1–99 Edition)

(ii) *Indications for use.* Treatment of infections of the respiratory tract (tonsillitis, tracheobronchitis), genitourinary tract (cystitis), gastrointestinal tract (bacterial gastroenteritis), and soft tissues (abscesses, lacerations, wounds), caused by susceptible strains of *Staphylococcus aureus*, *Streptococcus* spp., *Escherichia coli*, *Proteus mirabilis*, and bacterial dermatitis caused by *S. aureus*, *Streptococcus* spp., and *P. mirabilis*.

(iii) *Limitations.* Administer for 5 to 7 days or 48 hours after all symptoms have subsided. If no improvement is seen in 5 days, review diagnosis and change therapy. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats—(i) Amount.* 50 milligrams (5 to 10 milligrams per pound of body weight) once a day.

(ii) *Indications for use.* Treatment of infections caused by susceptible organisms as follows: upper respiratory tract due to *S. aureus*, *Streptococcus* spp., and *E. coli*; genitourinary tract (cystitis) due to *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*; gastrointestinal tract due to *E. coli*; and skin and soft tissue (abscesses, lacerations, and wounds) due to *S. aureus*, *Streptococcus* spp., *E. coli*, and *Pasteurella multocida*.

(iii) *Limitations.* Administer for 5 to 7 days or 48 hours after all symptoms have subsided. If no improvement is seen in 5 days, review diagnosis and change therapy. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37319, Aug. 18, 1992, as amended at 60 FR 55658, Nov. 2, 1995]

§ 520.88b Amoxicillin trihydrate for oral suspension.

(a) *Specifications.* When reconstituted, each milliliter contains amoxicillin trihydrate equivalent to 50 milligrams of amoxicillin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(1) *Conditions of use—(i) Dogs—(A) Amount.* 5 milligrams per pound of body weight twice daily.

(B) *Indications for use.* Treatment of infections caused by susceptible strains of organisms as follows: respiratory tract (tonsillitis, tracheobronchitis)

Food and Drug Administration, HHS

§ 520.88d

caused by *Staphylococcus aureus*, *Streptococcus* spp., *Escherichia coli*, and *Proteus mirabilis*; genitourinary tract (cystitis) caused by *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*; gastrointestinal tract (bacterial gastroenteritis) caused by *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*; bacterial dermatitis caused by *S. aureus*, *Streptococcus* spp., and *P. mirabilis*; and soft tissues (abscesses, lacerations, and wounds) caused by *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*.

(C) *Limitations*. Use for 5 to 7 days or 48 hours after all symptoms have subsided. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(ii) *Cats*—(A) *Amount*. 50 milligrams (5 to 10 milligrams per pound) once daily.

(B) *Indications for use*. Treatment of infections caused by susceptible strains of organisms as follows: upper respiratory tract due to *Staphylococcus* spp., *Streptococcus* spp., *Hemophilus* spp., *E. coli*, *Pasteurella* spp., and *P. mirabilis*; genitourinary tract (cystitis) due to *S. aureus*, *Streptococcus* spp., *E. coli*, *P. mirabilis*, and *Corynebacterium* spp.; gastrointestinal tract due to *E. coli*, *Proteus* spp., *Staphylococcus* spp., and *Streptococcus* spp.; skin and soft tissue (abscesses, lacerations, and wounds) due to *Staphylococcus* spp., *Streptococcus* spp., *E. coli*, and *Pasteurella multocida*.

(C) *Limitations*. Use for 5 to 7 days or 48 hours after all symptoms have subsided. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

(c) *Sponsor*. See Nos. 000031 and 000093 in § 510.600(c) of this chapter.

(1) *Conditions of use. Dogs*—(i) *Amount*. 5 milligrams per pound of body weight twice daily.

(ii) *Indications for use*. Treatment of bacterial dermatitis due to *S. aureus*, *Streptococcus* spp., *Staphylococcus* spp., and *E. coli*, and soft tissue infections (abscesses, wounds, lacerations) due to *S. aureus*, *Streptococcus* spp., *E. coli*, *P. mirabilis* and *Staphylococcus* spp.

(iii) *Limitations*. Use for 5 to 7 days. Continue for 48 hours after all symptoms have subsided. If no improvement

is seen in 5 days, review diagnosis and change therapy. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[57 FR 37319, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992, as amended at 60 FR 55658, Nov. 2, 1995; 62 FR 13302, Mar. 20, 1997]

§ 520.88c Amoxicillin trihydrate oral suspension.

(a) *Specifications*. Each 0.8-milliliter dose contains amoxicillin trihydrate equivalent to 40 milligrams of amoxicillin.

(b) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.510 of this chapter.

(d) *Conditions of use. Swine*—(1) *Amount*. 40 milligrams orally, twice a day using a dosing pump.

(2) *Indications for use*. Treatment of baby pigs under 10 pounds for porcine colibacillosis caused by *Escherichia coli* susceptible to amoxicillin.

(3) *Limitations*. Treat animals for 48 hours after all symptoms have subsided but not beyond 5 days. Do not slaughter during treatment or for 15 days after latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37319, Aug. 18, 1992, as amended at 60 FR 55658, Nov. 2, 1995]

§ 520.88d Amoxicillin trihydrate soluble powder.

(a) *Specifications*. Each gram contains amoxicillin trihydrate equivalent to 115.4 milligrams of amoxicillin.

(b) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.38 of this chapter.

(d) *Conditions of use. Preruminating calves including veal calves*—(1) *Amount*. 400 milligrams per 100 pounds of body weight twice daily.

(2) *Indications for use*. Treatment of bacterial enteritis when due to susceptible *Escherichia coli* in preruminating calves including veal calves.

(3) *Limitations*. Administer by drench or by mixing in milk. Treatment should be continued for 48 hours after all symptoms have subsided but not to exceed 5 days. For use in preruminating calves including veal

§ 520.88e

calves only, not for use in other animals which are raised for food production. Do not slaughter animals during treatment or for 20 days after the latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37319, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992; 58 FR 18304, Apr. 8, 1993, as amended at 60 FR 55658, Nov. 2, 1995; 62 FR 5525, Feb. 6, 1997]

§ 520.88e Amoxicillin trihydrate boluses.

(a) *Specifications.* Each bolus contains the equivalent of 400 milligrams of amoxicillin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.38 of this chapter.

(d) *Conditions of use. Preruminating calves including veal calves—*(1) *Amount.* 400 milligrams per 100 pounds of body weight twice daily.

(2) *Indications for use.* Treatment of bacterial enteritis when due to susceptible *Escherichia coli* in preruminating calves including veal calves.

(3) *Limitations.* For oral use in preruminating calves including veal calves only, not for use in other animals which are raised for food production. Treatment should be continued for 48 hours after all symptoms have subsided but not to exceed 5 days. Do not slaughter animals during treatment or for 20 days after the latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37320, Aug. 18, 1992, as amended at 60 FR 55659, Nov. 2, 1995; 62 FR 5526, Feb. 6, 1997]

§ 520.88f Amoxicillin trihydrate tablets.

(a) *Specifications.* Each tablet contains amoxicillin trihydrate equivalent to 50, 100, 200, or 400 milligrams of amoxicillin.

(b) *Sponsor.* See Nos. 000031 or 000093 in § 510.600(c) of this chapter.

(c) *Conditions of use—*(1) *Dogs—*(i) *Amount.* 5 milligrams per pound of body weight twice a day.

(ii) *Indications for use.* Treatment of bacterial dermatitis due to *Staphylococcus aureus*, *Streptococcus* spp., *Staphylococcus* spp., and *Escherichia*

21 CFR Ch. I (4–1–99 Edition)

coli; and soft tissue infections (abscesses, wounds, lacerations) due to *S. aureus*, *Streptococcus* spp., *E. coli*, *Proteus mirabilis*, and *Staphylococcus* spp.

(iii) *Limitations.* Use for 5 to 7 days or 48 hours after all symptoms have subsided. If no improvement is seen in 5 days, review diagnosis and change therapy. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[57 FR 37320, Aug. 18, 1992, as amended at 62 FR 13302, Mar. 20, 1997]

§ 520.88g Amoxicillin trihydrate and clavulanate potassium film-coated tablets.

(a) *Specifications.* Each tablet contains amoxicillin trihydrate and clavulanate potassium, equivalent to either 50 milligrams of amoxicillin and 12.5 milligrams clavulanic acid, or 100 milligrams of amoxicillin and 25 milligrams clavulanic acid, or 200 milligrams amoxicillin and 50 milligrams clavulanic acid or 300 milligrams amoxicillin and 75 milligrams clavulanic acid.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use—*(1) *Dogs—*(i) *Amount.* 6.25 milligrams (equivalent to 5 milligrams amoxicillin and 1.25 milligrams clavulanic acid) per pound of body weight twice daily.

(ii) *Indications for use.* Treatment of skin and soft tissue infections such as wounds, abscesses, cellulitis, superficial/juvenile and deep pyoderma due to susceptible strains of beta-lactamase (penicillinase) *Staphylococcus aureus*, nonbeta-lactamase *S. aureus*, *Staphylococcus* spp., *Streptococcus* spp., and *Escherichia coli*. Treatment of periodontal infections due to susceptible strains of aerobic and anaerobic bacteria.

(iii) *Limitations.* Wounds, abscesses, cellulitis, and superficial/juvenile pyoderma: Treat for 5 to 7 days or for 48 hours after all signs have subsided. If no improvement is seen after 5 days of treatment, discontinue therapy and reevaluate diagnosis. Deep pyoderma may require treatment for 21 days; do not treat for more than 30 days. Not for use in dogs maintained for breeding. Federal law restricts this drug to use

Food and Drug Administration, HHS

§ 520.90a

by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Amount*. 62.5 milligrams (1 milliliter) (50 milligrams amoxicillin and 12.5 milligrams clavulanic acid) twice daily.

(ii) *Indications for use*. Treatment of skin and soft tissue infections, such as wounds, abscesses and cellulitis/dermatitis due to susceptible strains of beta-lactamase (penicillinase) producing *S. aureus*, nonbeta-lactamase producing *S. aureus*, *Staphylococcus* spp., *Streptococcus* spp., *E. coli*, and *Pasteurella* spp. Also, treatment of urinary tract infections (cystitis) due to susceptible strains of *E. coli*.

(iii) *Limitations*. Skin and soft tissue infections: abscesses, cellulitis/dermatitis should be treated for 5 to 7 days or for 48 hours after all signs have subsided. If no response is seen after 3 days of treatment, therapy should be discontinued and diagnosis reevaluated. Urinary tract infections may require treatment for 10 to 14 days or longer. The maximum duration of treatment should not exceed 30 days. Safety of use in pregnant or breeding animals has not been established. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37320, Aug. 18, 1992, as amended at 60 FR 55659, Nov. 2, 1995; 63 FR 13121, Mar. 18, 1998]

§ 520.88h Amoxicillin trihydrate and clavulanate potassium for oral suspension.

(a) *Specifications*. When reconstituted, each milliliter contains amoxicillin trihydrate equivalent to 50 milligrams of amoxicillin with clavulanate potassium equivalent to 12.5 milligrams of clavulanic acid.

(b) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount*. 6.25 milligrams (equivalent to 5 milligrams amoxicillin and 1.25 milligrams clavulanic acid) per pound of body weight twice daily.

(ii) *Indications for use*. Treatment of skin and soft tissue infections such as wounds, abscesses, cellulitis, superficial/juvenile and deep pyoderma due to susceptible strains of beta-lactamase (penicillinase) producing *Staphylococcus aureus*, nonbeta-

lactamase *Staphylococcus aureus*, *Staphylococcus* spp., *Streptococcus* spp., and *Escherichia coli*. Treatment of periodontal infections due to susceptible strains of aerobic and anaerobic bacteria.

(iii) *Limitations*. Administer for 5 to 7 days or 48 hours after all symptoms subsided. Deep pyoderma may require 21 days, not to exceed 30 days. If no improvement is seen in 5 days, discontinue therapy and reevaluate the case. Not for use in dogs maintained for breeding. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Amount*. 62.5 milligrams (1 milliliter) (50 milligrams of amoxicillin and 12.5 milligrams clavulanic acid) twice daily.

(ii) *Indications for use*. Treatment of feline skin and soft tissue infections, such as wounds, abscesses and cellulitis/dermatitis due to susceptible strains of beta-lactamase (penicillinase) producing *S. aureus*, nonbeta-lactamase *S. aureus*, *Staphylococcus* spp., *Streptococcus* spp., *E. coli*, *Pasteurella multocida*, and *Pasteurella* spp.

(iii) *Limitations*. Administer 48 hours after all symptoms have subsided. If no improvement is seen after 3 days of treatment, discontinue therapy and reevaluate diagnosis. Maximum duration of treatment should not exceed 30 days. Not for use in cats maintained for breeding. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37320, Aug. 18, 1992, as amended at 60 FR 55659, Nov. 2, 1995; 63 FR 13121, Mar. 18, 1998]

§ 520.90 Ampicillin oral dosage forms.

§ 520.90a Ampicillin capsules.

(a) *Specifications*. Each capsule contains 125 milligrams or 250 milligrams of ampicillin.

(b) *Sponsor*. See No. 000008 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount*. 5 to 10 milligrams per pound of body weight, e.g., one 125 mg capsule per 14 to 25 pounds, given 2 to 4 times daily; for animals weighing 6 to 14 pounds, one capsule twice daily.

§ 520.90b

21 CFR Ch. I (4–1–99 Edition)

(ii) *Indications for use.* Treatment of urinary tract infections (cystitis) due to *Proteus* spp., hemolytic and non-hemolytic streptococci, beta hemolytic streptococci, and *Escherichia coli*. In upper respiratory tract infections tracheobronchitis (kennel cough), tonsillitis due to alpha and beta hemolytic streptococci, hemolytic positive staphylococci, *E. coli*, and *Proteus* spp. In infections associated with abscesses, lacerations, and wounds due to *Staphylococcus* spp. and *Streptococcus* spp.

(iii) *Limitations.* Bacteriologic studies to determine the causative organisms and their susceptibility to ampicillin should be performed. Use of the drug is contraindicated in animals with a history of an allergic reaction to any of the penicillins. Ampicillin is contraindicated in infections caused by penicillinase-producing organisms. Not for use in animals which are raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Amount.* 125 milligrams twice daily; in more acute conditions three times daily.

(ii) *Indications for use.* Treatment of respiratory tract infections (bacterial pneumonia) due to alpha and beta hemolytic streptococci, hemolytic positive staphylococci, *E. coli*, and *Proteus* spp. In infections associated with abscesses, lacerations, and wounds due to *Staphylococcus* spp. and *Streptococcus* spp.

(iii) *Limitations.* Bacteriologic studies to determine the causative organisms and their susceptibility to ampicillin should be performed. Use of the drug is contraindicated in animals with a history of an allergic reaction to any of the penicillins. Ampicillin is contraindicated in infections caused by penicillinase-producing organisms. Not for use in animals which are raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37321, Aug. 18, 1992]

§ 520.90b Ampicillin trihydrate tablets.

(a) *Specifications.* Each tablet contains ampicillin trihydrate equivalent to 50 or 100 milligrams of ampicillin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* *Dogs*—(1) *Amount.* 5 milligrams per pound of body weight, at 8-hour intervals, 1 to 2 hours prior to feeding, to be continued 36 to 48 hours after all symptoms have subsided. If no improvement is seen within 5 days, stop treatment, reevaluate diagnosis, and change therapy.

(2) *Indications for use.* Oral treatment of infections caused by susceptible organisms as follows: Upper respiratory infections, tonsillitis, and bronchitis due to *Streptococcus* spp., *Staphylococcus* spp., *Escherichia coli*, *Proteus mirabilis*, and *Pasteurella* spp., urinary tract infections (cystitis) due to *Streptococcus* spp., *Staphylococcus* spp., *E. coli*, *P. mirabilis*, and *Enterococcus* spp.; gastrointestinal infections due to *Staphylococcus* spp., *Streptococcus* spp., *Enterococcus* spp., and *E. coli*; infections associated with abscesses, lacerations, and wounds caused by *Staphylococcus* spp., and *Streptococcus* spp.

(3) *Limitations.* Not for use in animals which have shown hypersensitivity to penicillin or for infections caused by penicillinase-producing organisms. Not for use in animals which are raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37321, Aug. 18, 1992, as amended at 60 FR 55659, Nov. 2, 1995]

§ 520.90c Ampicillin trihydrate capsules.

(a) *Specifications.* Each capsule contains ampicillin trihydrate equivalent to 125, 250, or 500 milligrams of ampicillin.

(b) *Sponsor.* See No. 055529 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount.* 5 to 10 milligrams per pound of body weight two or three times daily. In severe or acute conditions, 10 milligrams per pound of body weight, three times daily. Administer 1 to 2 hours prior to feeding.

(ii) *Indications for use.* Treatment against strains of gram-negative and gram-positive organisms sensitive to ampicillin and associated with respiratory tract infections (tracheobronchitis and tonsillitis); urinary tract infections (cystitis); bacterial gastroenteritis; generalized infections (septicemia) associated with

abscesses, lacerations, and wounds; and bacterial dermatitis.

(iii) *Limitations*. The drug may be given as an emergency measure; however, in vitro sensitivity tests on samples collected prior to treatment should be made. Ampicillin is contraindicated for use in infections caused by penicillinase-producing organisms and for use in animals known to be allergic to any of the penicillins. Not for use in animals raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Amount*. 10 to 30 milligrams per pound of body weight or three times daily. Administer 1 to 2 hours prior to feeding.

(ii) *Indications for use*. Treatment against strains of gram-negative and gram-positive organisms sensitive to ampicillin and associated with respiratory tract infections (bacterial pneumonia); urinary tract infections (cystitis); and generalized infections (septicemia) associated with abscesses, lacerations, and wounds.

(iii) *Limitations*. The drug may be given as an emergency measure; however, in vitro sensitivity tests on samples collected prior to treatment should be made. Ampicillin is contraindicated for use in infections caused by penicillinase-producing organisms and for use in animals known to be allergic to any of the penicillins. Not for use in animals raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37321, Aug. 18, 1992, as amended at 58 FR 61016, Nov. 19, 1993]

§ 520.90d Ampicillin trihydrate for oral suspension.

(a) *Specifications*. When reconstituted as directed, each milliliter contains ampicillin trihydrate equivalent to 25 milligrams of ampicillin.

(b) *Sponsor*. See No. 055529 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount*. 5 to 10 milligrams per pound of body weight orally, 2 or 3 times daily, 1 to 2 hours prior to feeding. In severe or acute conditions, 10 milligrams per pound of body weight 3 times daily.

(ii) *Indications for use*. Treatment of respiratory tract infections (tracheobronchitis and tonsillitis) due to *Escherichia coli*, *Pseudomonas* spp., *Proteus* spp., *Staphylococcus* spp., and *Streptococcus* spp.; urinary tract infections (cystitis) due to *E. coli*, *Staphylococcus* spp., *Streptococcus* spp., and *Proteus* spp.; bacterial gastroenteritis due to *E. coli*; generalized infections (septicemia) associated with abscesses, lacerations, and wounds, due to *Staphylococcus* spp. and *Streptococcus* spp.; bacterial dermatitis due to *Staphylococcus* spp., *Streptococcus* spp., *Proteus* spp., and *Pseudomonas* spp.

(iii) *Limitations*. Duration of treatment is usually 3 to 5 days. Continue treatment 48 hours after the animal's temperature has returned to normal and all other signs of infection have subsided. If no response is obtained within 3 to 5 days, reevaluate diagnosis and treatment. Appropriate laboratory tests should be conducted, including in vitro culturing and susceptibility tests on samples collected prior to treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Amount*. 10 to 30 milligrams per pound of body weight orally, 2 or 3 times daily, 1 to 2 hours prior to feeding.

(ii) *Indications for use*. Treatment of respiratory tract infections (bacterial pneumonia) due to *Staphylococcus* spp., *Streptococcus* spp., *E. coli*, and *Proteus* spp.; urinary tract infections (cystitis) due to *E. coli*, *Staphylococcus* spp., *Streptococcus* spp., *Proteus* spp., and *Corynebacterium* spp.; generalized infections (septicemia) associated with abscesses, lacerations, and wounds, due to *Staphylococcus* spp., *Streptococcus* spp., *Bacillus* spp., and *Pasteurella* spp.

(iii) *Limitations*. Duration of treatment is usually 3 to 5 days. Continue treatment 48 hours after the animal's temperature has returned to normal and all other signs of infection have subsided. If no response is obtained within 3 to 5 days, reevaluate diagnosis and treatment. Appropriate laboratory tests should be conducted, including in vitro culturing and susceptibility tests on samples collected prior to treatment. Federal law restricts this drug

§ 520.90e

to use by or on the order of a licensed veterinarian.

[57 FR 37321, Aug. 18, 1992, as amended at 58 FR 61016, Nov. 19, 1993]

§ 520.90e Ampicillin trihydrate soluble powder.

(a) *Specifications.* Each gram contains ampicillin trihydrate equivalent to 88.2 milligrams of ampicillin.

(b) *Sponsor.* See No. 055529 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.40 of this chapter.

(d) *Conditions of use. Swine*—(1) *Amount.* 5 milligrams of ampicillin per pound of body weight twice daily, orally by gavage or in drinking water for up to 5 days.

(2) *Indications for use.* Oral treatment of porcine colibacillosis (*Escherichia coli*) and salmonellosis (*Salmonella* spp.) infections in swine up to 75 pounds of body weight, and bacterial pneumonia caused by *Pasteurella multocida*, *Staphylococcus* spp., *Streptococcus* spp., and *Salmonella* spp.

(3) *Limitations.* For use in swine only. Not for use in other animals which are raised for food production. Treated swine must not be slaughtered for food during treatment and for 24 hours following the last treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37322, Aug. 18, 1992, as amended at 58 FR 61016, Nov. 19, 1993]

§ 520.90f Ampicillin trihydrate boluses.

(a) *Specifications.* Each bolus contains ampicillin trihydrate equivalent to 400 milligrams of ampicillin.

(b) *Sponsor.* See No. 055529 in § 510.600(c) of this chapter for use as in paragraph (d)(1), 000069 for use as in paragraph (d)(2).

(c) *Related tolerances.* See § 556.40 of this chapter.

(d) *Conditions of use. Nonruminating calves*—(1) *Amount.* 5 milligrams per pound of body weight twice daily for up to 5 days.

(i) *Indications for use.* Oral treatment of colibacillosis caused by *Escherichia coli*, bacterial enteritis caused by *Salmonella* spp., and bacterial pneumonia caused by *Pasteurella* spp.

(ii) *Limitations.* Treated calves must not be slaughtered for food during

21 CFR Ch. I (4–1–99 Edition)

treatment and for 15 days after the last treatment. Not for use in other animals raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Amount.* 5 milligrams per pound of body weight twice daily not to exceed 4 days.

(i) *Indications for use.* Oral treatment of bacterial enteritis (colibacillosis) caused by *E. coli*.

(ii) *Limitations.* Treated calves must not be slaughtered for food during treatment and for 7 days after the last treatment. Not for use in other animals raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37322, Aug. 18, 1992, as amended at 58 FR 61016, Nov. 19, 1993; 60 FR 55659, Nov. 2, 1995]

§ 520.100 Amprolium oral dosage forms.

§ 520.100a Amprolium drinking water.

(a) *Chemical name.* 1-(4-Amino-2-n-propyl-5-pyrimidinylmethyl)-2-picolinium chloride hydrochloride.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.50 of this chapter.

(d) *Conditions of use.* It is used in drinking water as follows:

(1) *Chickens and turkeys*—(i) *Amount.* 20 percent soluble powder.

(ii) *Indications for use.* Treatment of coccidiosis.

(iii) *Limitations.* Administer at the 0.012 percent level in drinking water as soon as coccidiosis is diagnosed and continue for from 3 to 5 days (in severe outbreaks, give amprolium at the 0.024 percent level); continue with 0.006 percent amprolium-medicated water for an additional 1 to 2 weeks; no other source of drinking water should be available to the birds during this time; as sole source of amprolium.

(2) *Calves*—(i) *Amount.* 9.6 percent solution or 20 percent soluble powder.

(a) *Indications for use.* As an aid in the treatment of coccidiosis caused by *Eimeria bovis* and *E. zurnii*.

(b) *Limitations.* Add 16 fluid ounces of the 9.6 percent solution to each 100 gallons of drinking water; or 4 ounces of the soluble powder to each 50 gallons of

Food and Drug Administration, HHS

§ 520.100c

drinking water; at the usual rate of water consumption, this will provide an intake of approximately 10 milligrams per kilogram (2.2 pounds) of body weight; offer this solution as the only source of water for 5 days; for a satisfactory diagnosis, a microscopic examination of the feces should be done by a veterinarian or diagnostic laboratory before treatment; when treating outbreaks, the drug should be administered promptly after diagnosis is determined; withdraw 24 hours before slaughter.

(ii) *Amount.* 9.6 percent solution or 20 percent soluble powder.

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii*.

(b) *Limitations.* Add 8 fluid ounces of the 9.6 percent solution or 4 ounces of the 20 percent soluble powder to each 100 gallons of drinking water; at the usual rate of water consumption, this will provide an intake of approximately 5 milligrams per kilogram (2.2 pounds) of body weight; offer this solution as the only source of water for 21 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard; withdraw 24 hours before slaughter.

[40 FR 13838, Mar. 27, 1975, as amended at 62 FR 63270, Nov. 28, 1997]

§ 520.100b Amprolium drench.

(a) *Chemical name.* 1-(4-Amino-2-n-propyl - 5 - pyrimidinylmethyl) - 2 - picolinium chloride hydrochloride.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.50 of this chapter.

(d) *Conditions of use.* It is used for calves as follows:

(1) *Amount.* 9.6 percent solution or 20 percent soluble powder.

(i) *Indications for use.* As an aid in the treatment of coccidiosis caused by *Eimeria bovis* and *E. zurnii*.

(ii) *Limitations.* Add 3 fluid ounces of the 9.6 percent solution to 1 pint of water or 3 ounces of the 20 percent soluble powder to each quart of water and with a dose syringe administer 1 fluid ounce of this solution for each 100 pounds of body weight; this will provide a dose of approximately 10 milligrams per kilogram (2.2 pounds) of

body weight; administer daily for 5 days; for a satisfactory diagnosis, a microscopic examination of the feces should be done by a veterinarian or diagnostic laboratory before treatment; when treating outbreaks, the drug should be administered promptly after diagnosis is determined; withdraw 24 hours before slaughter.

(2) *Amount.* 9.6 percent solution or 20 percent soluble powder.

(i) *Indications for use.* As an aid in the prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii*.

(ii) *Limitations.* Add 1½ fluid ounces of the 9.6 percent solution to 1 pint of water or 1½ ounces of the 20 percent soluble powder to each quart of water and with a dose syringe administer 1 fluid ounce of this solution for each 100 pounds of body weight; this will provide a dose of approximately 5 milligrams per kilogram (2.2 pounds) of body weight; administer daily for 21 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard; withdraw 24 hours before slaughter.

[40 FR 13838, Mar. 27, 1975, as amended at 62 FR 63270, Nov. 28, 1997]

§ 520.100c Amprolium crumbles.

(a) *Specifications.* Amprolium crumbles contain 1.25 percent amprolium.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.50 of this chapter.

(d) *Conditions of use.* It is top-dressed on or thoroughly mixed in the daily feed ration of calves as follows:

(1) *Amount.* 1.6 ounces of crumbles per 250 pounds of body weight per day (5 milligrams per kilogram of body weight).

(i) *Indications for use.* As an aid in the prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii*.

(ii) *Limitations.* Administer for 21 consecutive days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard. Withdraw 24 hours before slaughter. Use as sole source of amprolium.

(2) *Amount.* 3.2 ounces of crumbles per 250 pounds of body weight per day (10 milligrams per kilogram of body weight).

§ 520.110

(i) *Indications for use.* As an aid in the treatment of coccidiosis caused by *Eimeria bovis* and *E. zurnii*.

(ii) *Limitations.* Administer for 5 consecutive days. For satisfactory diagnosis, a microscopic fecal examination should be done by a veterinarian or diagnostic laboratory before treatment. When treating outbreaks, the drug should be administered promptly after diagnosis is determined. Withdraw 24 hours before slaughter. Use as sole source of amprolium.

[42 FR 41855, Aug. 19, 1977, as amended at 62 FR 63270, Nov. 28, 1997]

§ 520.110 Apramycin sulfate soluble powder.

(a) *Specifications.* A water soluble powder used to make a medicated drinking water containing apramycin sulfate equivalent to 0.375 gram of apramycin activity per gallon of drinking water.

(b) *Sponsor.* See No. 000986 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.52 of this chapter.

(d) *Conditions of use.* (1) In swine for control of porcine colibacillosis (weanling pig scours) caused by strains of *E. coli* sensitive to apramycin.

(2) It is administered for 7 days in drinking water at the rate of 12.5 milligrams of apramycin per kilogram (5.7 milligrams per pound) of body weight per day. Swine will normally consume 1 gallon per day of medicated water containing 375 milligrams of apramycin for each 66 pounds of body weight. Water consumption should be monitored to determine that the required amount of apramycin is being consumed. The drug concentration should be adjusted according to water consumption which varies depending on ambient temperature, humidity, and other factors.

(3) Prepare fresh medicated water daily.

(4) Do not slaughter treated swine for 28 days following treatment

[47 FR 15771, Apr. 13, 1982, as amended at 49 FR 19642, May 9, 1984; 53 FR 37753, Sept. 28, 1988]

21 CFR Ch. I (4–1–99 Edition)

§ 520.154 Bacitracin oral dosage forms.

§ 520.154a Soluble bacitracin methylene disalicylate.

(a) *Specifications.* Each pound of soluble powder contains the equivalent of 50 grams of bacitracin activity for use as in paragraph (d)(1) or (d)(2) of this section, or the equivalent of 200 grams of bacitracin activity for use as in paragraph (d) of this section.

(b) *Sponsor.* See No. 046573 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.70 of this chapter.

(d) *Conditions of use*—(1) *Growing turkeys*—(i) *Amount.* 400 milligrams per gallon in drinking water.

(ii) *Indications for use.* Aid in the control of transmissible enteritis complicated by organisms susceptible to bacitracin methylene disalicylate.

(iii) *Limitations.* Prepare a fresh solution daily.

(2) *Broiler and replacement chickens*—(i) *Amount.* 100 milligrams per gallon in drinking water.

(A) *Indications for use.* Aid in the prevention of necrotic enteritis caused by *Clostridium perfringens* susceptible to bacitracin methylene disalicylate.

(B) *Limitations.* Prepare a fresh solution daily.

(ii) *Amount.* 200 to 400 milligrams per gallon in drinking water.

(A) *Indications for use.* Aid in the control of necrotic enteritis caused by *C. perfringens* susceptible to bacitracin methylene disalicylate.

(B) *Limitations.* Prepare a fresh solution daily.

(3) *Swine*—(i) *Amount.* 1 gram per gallon in drinking water.

(ii) *Indications for use.* Treatment of swine dysentery associated with *Treponema hyodysenteriae*. Administer continuously for 7 days or until signs of dysentery disappear.

(iii) *Limitations.* Prepare a fresh solution daily. Treatment not to exceed 14 days. If symptoms persist after 4 to 5 days consult a veterinarian. Not to be given to swine that weigh more than 250 pounds.

(4) *Growing quail*—(i) *Amount.* 400 milligrams per gallon in drinking water.

(ii) *Indications for use.* For prevention of ulcerative enteritis due to *Clostridium colinum* susceptible to bacitracin methylene disalicylate.

(iii) *Limitations.* Prepare fresh solution daily. Use as sole source of drinking water.

[57 FR 37322, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992, as amended at 63 FR 38474, July 17, 1998; 64 FR 13068, Mar. 17, 1999]

§ 520.154b Soluble bacitracin methylene disalicylate and streptomycin sulfate oral powder.

(a) *Specifications.* Each gram contains 200 units of soluble bacitracin methylene disalicylate, streptomycin sulfate equivalent to 20 milligrams of streptomycin, and 850 milligrams of carob flour.

(b) *Sponsor.* See No. 062925 in § 510.600(c) of this chapter.

(c) *Conditions of use.* *Dogs*—(1) *Amount.* 1 level teaspoonful per 10 pounds of body weight three times daily, mixed in a small quantity of liquid or feed.

(2) *Indications for use.* Treatment of bacterial enteritis caused by pathogens susceptible to bacitracin and streptomycin such as *Escherichia coli*, *Proteus* spp., *Staphylococcus* spp., and *Streptococcus* spp., and for the symptomatic treatment of associated diarrhea.

(3) *Limitations.* If no improvement is noted in 2 to 3 days, diagnosis should be reevaluated. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37322, Aug. 18, 1992, as amended at 61 FR 66581, Dec. 18, 1996]

§ 520.154c Bacitracin zinc soluble powder.

(a) *Specifications.* Each pound contains the equivalent of not less than 5 grams of bacitracin.

(b) *Sponsor.* See No. 010042 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.70 of this chapter.

(d) *Conditions of use.* (1) *Broiler chickens*—(i) *Amount.* 100 milligrams per gallon in drinking water.

(A) *Indications for use.* Prevention of necrotic enteritis caused by *Clostridium perfringens* susceptible to bacitracin zinc.

(B) *Limitations.* Prepare a fresh solution daily.

(ii) *Amount.* 200 to 400 milligrams per gallon in drinking water.

(A) *Indications for use.* Control of necrotic enteritis caused by *Clostridium perfringens* susceptible to bacitracin zinc.

(B) *Limitations.* Prepare a fresh solution daily.

(2) *Growing quail*—(i) *Amount.* 500 milligrams per gallon in drinking water for 5 days followed by 165 milligrams per gallon in drinking water for 10 days.

(ii) *Indications for use.* Control of ulcerative enteritis caused by *Clostridium* spp. susceptible to bacitracin zinc.

(iii) *Limitations.* Prepare a fresh solution daily.

[57 FR 37322, Aug. 18, 1992]

§ 520.182 Bicyclohexylammonium fumagillin.

(a) *Specifications.* The drug is a soluble powder containing bicyclohexylammonium fumagillin and appropriate phosphate buffers.

(b) *Sponsor.* See No. 059620 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used for the prevention of nosema in honey bees.¹

(2) It is administered usually in a 2:1 sugar sirup containing a concentration of from 75 to 100 milligrams of fumagillin activity per gallon of sugar sirup.¹

(3) Colonies used for package production should be fed medicated sirup as a principal food supply for a month prior to stocking nuclei or shaking packages for market.¹

(4) The medicated sirup should not be fed immediately before or during the honey flow.

[40 FR 13838, Mar. 27, 1975, as amended at 42 FR 65151, Dec. 30, 1977; 56 FR 43699, Sept. 4, 1991; 58 FR 5608, Jan. 22, 1993]

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

§ 520.222

21 CFR Ch. I (4–1–99 Edition)

§ 520.222 Bunamidine hydrochloride.

(a) *Chemical name.* *N,N*-Dibutyl-4-(hexyloxy)-1-naphthamidine hydrochloride.

(b) *Specifications.* The drug is an oral tablet containing 100, 200, or 400 milligrams of bunamidine hydrochloride.

(c) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) The drug is intended for oral administration to dogs for the treatment of the tapeworms *Dipylidium caninum*, *Taenia pisiformis*, and *Echinococcus granulosus*, and to cats for the treatment of the tapeworms *Dipylidium caninum* and *Taenia taeniaeformis*.

(2) It is administered to cats and dogs at the rate of 25 to 50 milligrams per kilogram of body weight. The drug should be given on an empty stomach and food should not be given for 3 hours following treatment.

(3) Tablets should not be crushed, mixed with food, or dissolved in liquid. Repeat treatments should not be given within 14 days. The drug should not be given to male dogs within 28 days prior to their use for breeding. Do not administer to dogs or cats having known heart conditions.

(4) For use only by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 42 FR 13018, Mar. 8, 1977; 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61624, Nov. 19, 1997]

§ 520.246 Butorphanol tartrate tablets.

(a) *Specifications.* Each tablet contains 1, 5, or 10 milligrams of butorphanol base activity as butorphanol tartrate.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* The drug is used for the treatment of dogs as follows:

(1) *Amount.* 0.25 milligram of butorphanol base activity per pound of body weight.

(2) *Indications for use.* For the relief of chronic nonproductive cough associated with tracheo-bronchitis, tracheitis, tonsillitis, laryngitis, and pharyngitis associated with inflammatory conditions of the upper respiratory tract.

(3) *Limitations.* For oral use in dogs only. Repeat at intervals of 6 to 12

hours as required. If necessary, increase dose to a maximum of 0.5 milligram per pound of body weight. Treatment should not normally be required for longer than 7 days. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[47 FR 14702, Apr. 6, 1982, as amended at 53 FR 27851, July 25, 1988]

§ 520.260 *n*-Butyl chloride capsules.

(a)(1) *Specifications.* *n*-Butyl chloride capsules, veterinary contain 272 milligrams or 816 milligrams of *n*-butyl chloride in each capsule.

(2) *Sponsor.* See No. 021091 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) It is used for the removal of ascarids (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum*, *Ancylostoma braziliense*, and *Uncinaria stenocephala*) from dogs and of the ascarid (*Toxocara cati*) and hookworm (*Ancylostoma tubaeforme*) from cats.

(ii)(a) Animals should not be fed for 18 to 24 hours before being given the drug. Puppies and kittens should be wormed at 6 weeks of age. However, if heavily infested, they may be wormed at 4 or 5 weeks of age. Administration of the drug should be followed in ½ to 1 hour with a teaspoonful to a tablespoonful of milk of magnesia or 1 or 2 milk of magnesia tablets. Normal rations may be resumed 4 to 8 hours after treatment. Puppies and kittens should be given a repeat treatment in a week or 10 days. After that they should be treated every 2 months (or as symptoms reappear) until a year old. When the puppy or kitten is a year old, one treatment every 3 to 6 months is sufficient.

(b) For dogs or cats that have been wormed regularly, treatment every 3 to 6 months will be sufficient. If a dog or cat has not been wormed previously and has the symptoms of large roundworms a dose should be given and repeated in 10 days. Removal of hookworms may require 3 or 4 doses at 10-day intervals.

(c) Puppies, dogs, cats, or kittens weighing 1 to 3 pounds should be given 2 capsules per dose which contain 272 milligrams of *n*-butyl chloride each. Such animals weighing 4 to 5 pounds

should be given 3 such capsules. Animals weighing 6 to 7 pounds should be given 4 such capsules and animals weighing 8 to 9 pounds should be given 5 such capsules. Animals weighing 10 to 20 pounds should be given 3 capsules which contain 816 milligrams of *n*-butyl chloride each, animals weighing 20 to 40 pounds should be given 4 such capsules and animals weighing over 40 pounds should be given 5 such capsules with the maximum dosage being 5 capsules, each of which contains 816 milligrams of *n*-butyl chloride.

(iii) A veterinarian should be consulted before using in severely debilitated dogs or cats and also prior to repeated use in cases which present signs of persistent parasitism.

(b)(1) *Specifications.* *n*-Butyl chloride capsules contain 221, 442, 884, or 1,768 milligrams or 4.42 grams of *n*-butyl chloride in each capsule.¹

(2) *Sponsors.* See No. 023851 in § 510.600(c) of this chapter for 221, 442, 884, or 1,768 milligram or 4.42 gram capsules; No. 000115 or 038782 for 884 or 1,768 milligram or 4.42 gram capsules; and No. 000069 for 221 milligram capsules.

(3) *Conditions of use.* (i) It is used for the removal of ascarids (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum*, *Ancylostoma braziliense*, and *Uncinaria stenocephala*) from dogs.¹

(ii)(a) Dogs should not be fed for 18 to 24 hours before being given the drug. Administration of the drug should be followed in ½ to 1 hour with a mild cathartic. Normal feeding may be resumed 4 to 8 hours after treatment. Animals subject to reinfection may be retreated in 2 weeks.¹

(b) The drug is administered orally to dogs. Capsules containing 221 milligrams of *n*-butyl chloride are administered to dogs weighing under 5 pounds at a dosage level of 1 capsule per 1¼ pound of body weight. Capsules containing 442 milligrams of *n*-butyl chloride are administered to dogs weighing under 5 pounds at a dosage level of 1 capsule per 2½ pounds body weight. Capsules containing 884 milligrams of

n-butyl chloride are administered to dogs as follows: Weighing under 5 pounds, 1 capsule; weighing 5 to 10 pounds, 2 capsules; weighing 10 to 20 pounds, 3 capsules; weighing 20 to 40 pounds, 4 capsules; over 40 pounds, 5 capsules. Capsules containing 1,768 milligrams of *n*-butyl chloride are administered at a dosage level of 1 capsule per dog weighing 5 to 10 pounds. Capsules containing 4.42 grams of *n*-butyl chloride are administered at a dosage level of 1 capsule per dog weighing 40 pounds or over.¹

(iii) A veterinarian should be consulted before using in severely debilitated dogs.¹

(c)(1) *Specifications.* *n*-Butyl chloride capsules, veterinary contain 884 or 1,768 milligrams or 4.42 grams of *n*-butyl chloride in each capsule.

(2) *Sponsor.* See No. 000115 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) It is used for the removal of ascarids (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum*, *Ancylostoma braziliense*, and *Uncinaria stenocephala*) from dogs.

(ii)(a) Dogs should not be fed for 18 to 24 hours before being given the drug. Administration of the drug should be followed in ½ to 1 hour with a mild cathartic. Normal rations may be resumed 4 to 8 hours after treatment.

(b) The drug is administered orally to dogs. Capsules containing 884 milligrams of *n*-butyl chloride are administered to dogs as follows: weighing under 5 pounds, 1 capsule; weighing 5–10 pounds, 2 capsules; weighing 10–20 pounds, 3 capsules; weighing 20–40 pounds, 4 capsules; over 40 pounds, 5 capsules. Capsules containing 1,768 milligrams of *n*-butyl chloride are administered at a dosage level of 1 capsule per dogs to dogs weighing 5–10 pounds and 2 capsules per dog to dogs weighing 20–40 pounds. Capsules containing 4.42 grams of *n*-butyl chloride are administered at dosage level of 1 capsule per dog to dogs weighing 40 pounds or over.

(iii) A veterinarian should be consulted before using in severely debilitated dogs.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter.

[40 FR 13838, Mar. 27, 1975, as amended at 40 FR 39858, Aug. 29, 1975; 44 FR 10059, Feb. 16, 1979; 54 FR 38515, Sept. 19, 1989; 55 FR 24556, June 18, 1990; 64 FR 15684, Apr. 1, 1999]

§ 520.300

21 CFR Ch. I (4–1–99 Edition)

§ 520.300 Cambendazole oral dosage forms.

§ 520.300a Cambendazole suspension.

(a) *Specifications.* Each fluid ounce contains 0.9 gram of cambendazole.

(b) *Sponsor.* No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in horses for the control of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); small strongyles (*Trichonema*, *Poteriostomum*, *Cylicobrachytus*, *Craterostomum*, *Oesophagodontus*); roundworms (*Parascaris*); pinworms (*Oxyuris*); and threadworms (*Strongyloides*).

(2) It is administered by stomach tube or as a drench at a dose of 0.9 gram of cambendazole per 100 pounds of body weight (20 milligrams per kilogram).

(3) For animals maintained on premises where reinfection is likely to occur, re-treatments may be necessary. For most effective results, re-treat in 6 to 8 weeks.

(4) Not for use in horses intended for food.

(5) Caution: Do not administer to pregnant mares during first 3 months of pregnancy.

(6) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975. Redesignated at 41 FR 1276, Jan. 7, 1976, and amended at 42 FR 3838, Jan. 21, 1977; 62 FR 63270, Nov. 28, 1997]

§ 520.300b Cambendazole pellets.

(a) *Specifications.* The drug is in feed pellets containing 5.3 percent cambendazole.

(b) *Sponsor.* No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in horses for the control of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); small strongyles (*Trichonema*, *Poteriostomum*, *Cylicobrachytus*, *Craterostomum*, *Oesophagodontus*); roundworms (*Parascaris*); pinworms (*Oxyuris*); and threadworms (*Strongyloides*).

(2) Administer 20 milligrams cambendazole per kilogram body weight (6 ounces per 1,000 pounds) by mixing with normal grain ration given at one feeding. Doses for individual

horses should be mixed and fed separately to assure that each horse will consume the correct amount.

(3) For animals maintained on premises where reinfection is likely to occur, re-treatments may be necessary. For most effective results, re-treat in 6 to 8 weeks.

(4) Not for use in horses intended for food.

(5) Caution: Do not administer to pregnant mares during first 3 months of pregnancy.

(6) Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[41 FR 1276, Jan. 7, 1976, as amended at 42 FR 3838, Jan. 21, 1977; 62 FR 63270, Nov. 28, 1997]

§ 520.300c Cambendazole paste.

(a) *Specifications.* The drug is a paste containing 45 percent cambendazole.

(b) *Sponsor.* No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in horses for the control of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); small strongyles (*Trichonema*, *Poteriostomum*, *Cylicobrachytus*, *Craterostomum*, *Oesophagodontus*); roundworms (*Parascaris*); pinworms (*Oxyuris*); and threadworms (*Strongyloides*).

(2) Administer 20 milligrams cambendazole per kilogram body weight (5 grams per 550 pounds (250 kilograms)) by depositing the paste on the back of the tongue using a dosing gun.

(3) For animals maintained on premises where reinfection is likely to occur, re-treatments may be necessary. For most effective results, re-treat in 6 to 8 weeks.

(4) Not for use in horses intended for food.

(5) Caution: Do not administer to pregnant mares during first 3 months of pregnancy.

(6) Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[41 FR 1276, Jan. 7, 1976, as amended at 42 FR 3838, Jan. 21, 1977; 62 FR 63270, Nov. 28, 1997]

§ 520.309 Carprofen caplets.

(a) *Specification.* Each caplet contains 25, 75, or 100 milligrams of carprofen.

Food and Drug Administration, HHS

§ 520.314

(b) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount*. 1 milligram per pound of body weight twice daily. Caplets are scored and dosage should be calculated in half-caplet increments.

(2) *Indications for use*. For the relief of pain and inflammation in dogs. Carprofen has been shown to be clinically effective for the relief of signs associated with osteoarthritis in dogs.

(3) *Limitations*. The safe use of carprofen in pregnant dogs, dogs used for breeding purposes, or in lactating bitches has not been established. As a class, cyclo-oxygenase inhibitory non-steroidal anti-inflammatory drugs (NSAID's) may be associated with gastrointestinal and renal toxicity. Patients at greatest risk for renal toxicity are those on concomitant diuretic therapy, or those with renal, cardiovascular, and/or hepatic dysfunction. Because many NSAID's possess the potential to induce gastrointestinal ulceration, avoid or closely monitor concomitant use of carprofen with other anti-inflammatory drugs, such as corticosteroids and NSAID's. Carprofen treatment was not associated with renal toxicity or gastrointestinal ulceration in safety studies of up to 10 times the dose in dogs. Do not use in dogs with bleeding disorders (e.g., Von Willebrand's disease). Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[61 FR 66581, Dec. 18, 1996]

§ 520.310 Caramiphen ethanedisulfonate and ammonium chloride tablets.

(a) *Specifications*. Each tablet contains 10 milligrams of caramiphen ethanedisulfonate and 80 milligrams of ammonium chloride.¹

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount*. One tablet per 15 to 30 pounds of body weight every 4 to 6 hours.¹

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(2) *Indications for use*. For relief of cough.¹

[43 FR 55385, Nov. 28, 1978]

§ 520.312 Carnidazole tablets.

(a) *Specifications*. Each tablet contains 10 milligrams of carnidazole.

(b) *Sponsor*. See 053923 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. Adult pigeons: 1 tablet (10 milligrams); newly weaned pigeons: ½ tablet (5 milligrams).

(2) *Indications for use*. For treating trichomoniasis (canker) in ornamental and homing pigeons.

(3) *Limitations*. Not for use in pigeons intended for human food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism or when severely ill birds do not respond to treatment.

[54 FR 32336, Aug. 7, 1989]

§ 520.314 Cefadroxil tablets.

(a) *Specifications*. 50-, 100-, and 200-milligram tablets for dogs and cats; 1 gram tablet for dogs.

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) For use in dogs as follows:

(i) *Indications for use*. For the treatment of skin and soft tissue infections including cellulitis, pyoderma, dermatitis, wound infections, and abscesses due to susceptible strains of *Staphylococcus aureus*. For the treatment of genitourinary tract infections (cystitis) due to susceptible strains of *Escherichia coli*, *Proteus mirabilis*, and *Staphylococcus aureus*.

(ii) *Amount*. Ten milligrams per pound of body weight twice daily.

(iii) *Limitations*. The drug is administered orally. For skin and soft tissue infections, treatment should be continued for a minimum of 3 days. For genitourinary tract infections, treatment should be continued for a minimum of 7 days. Continue treatment at least 48 hours after the dog has become afebrile or asymptomatic. If no response is seen after 3 days of treatment, therapy should be discontinued and the case re-evaluated. Do not treat for more than 30 days. Safety for use in pregnant

§ 520.315

bitches and stud dogs has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) For use in cats as follows:

(i) *Indications for use.* For the treatment of skin and soft tissue infections including abscesses, wound infections, cellulitis, and dermatitis caused by susceptible strains of *Pasteurella multocida*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, and *Streptococcus* spp.

(ii) *Amount.* Ten milligrams per pound of body weight once daily.

(iii) *Limitations.* The drug is administered orally. Continue treatment at least 48 hours after the cat has become afebrile or asymptomatic. If no response is seen after 3 days of treatment, therapy should be discontinued and the case reevaluated. Do not treat for more than 21 days. Safety for use in pregnant cats and breeding male cats has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[47 FR 41105, Sept. 17, 1982, as amended at 49 FR 43052, Oct. 26, 1984; 51 FR 4165, Feb. 3, 1986; 52 FR 11989, Apr. 14, 1987; 53 FR 27851, July 25, 1988]

§ 520.315 Cefadroxil powder for oral suspension.

(a) *Specifications.* Cefadroxil powder is reconstituted to form a 50 milligram-milliliter aqueous suspension.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) For use in dogs as follows:

(i) *Indications for use.* For treating genitourinary tract infections (cystitis) caused by susceptible strains of *Escherichia coli*, *Proteus mirabilis*, and *Staphylococcus aureus*; and skin and soft tissue infections including cellulitis, pyoderma, dermatitis, wound infections, and abscesses caused by susceptible strains of *Staphylococcus aureus*.

(ii) *Amount.* 10 milligrams per pound of body weight, twice daily.

(2) For use in cats as follows:

(i) *Indications for use.* For treating skin and soft tissue infections including abscesses, wound infections, cellulitis, and dermatitis caused by susceptible strains of *Pasteurella*

21 CFR Ch. I (4–1–99 Edition)

multocida, *Staphylococcus aureus*, *Staphylococcus epidermidis*, and *Streptococcus* spp.

(ii) *Amount.* 10 milligrams per pound of body weight, once daily.

(3) *Limitations.* Discard unused portion of reconstituted product after 14 days. Treatment should continue for 48 hours after animal is afebrile or asymptomatic. If no response after 3 days, discontinue treatment and re-evaluate therapy. Not for use in animals raised for food production. Safe use in pregnant or breeding animals has not been established. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[53 FR 27344, July 20, 1988]

§ 520.390 Chloramphenicol oral dosage forms.

§ 520.390a Chloramphenicol tablets.

(a)(1) *Specifications.* Each tablet contains 100, 250, or 500 milligrams, 1 or 2.5 grams of chloramphenicol.

(2) *Sponsor.* In § 510.600(c) of this chapter: No. 000010 for 100-, 250-, and 500-milligram and 1-gram tablets; No. 000856 for 100-, 250-, and 500-milligram tablets; No. 017030 for 100-milligram tablets; No. 000010 for 100-, 250-, and 500-milligram and 1- and 2.5-gram tablets; No. 000069 for 250-milligram tablets.

(3) *Conditions of use.* *Dogs*—(i) *Amount.* 25 milligrams per pound of body weight every 6 hours.

(ii) *Indications for use.* Oral treatment of bacterial pulmonary infections, bacterial infections of the urinary tract, bacterial enteritis, and bacterial infections associated with canine distemper caused by susceptible organisms.

(iii) *Limitations.* Laboratory tests should be conducted, including in vitro culturing and susceptibility tests on samples collected prior to treatment. If no response to chloramphenicol therapy is obtained in 3 to 5 days, discontinue its use and review diagnosis. Not for animals which are raised for food production. Chloramphenicol products must not be used in meat-, egg-, or milk-producing animals. The length of time that residues persist in milk or tissues has not been determined. Because of potential antagonism, chloramphenicol should not be

Food and Drug Administration, HHS

§ 520.420

administered simultaneously with penicillin or streptomycin. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Specifications*. Each tablet contains 50, 100, 250, or 500 milligrams, or 1 gram of chloramphenicol.

(2) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(3) *Conditions of use*. *Dogs*—(i) *Amount*. 25 milligrams per pound of body weight every 6 hours.

(ii) *Indications for use*. Oral treatment of bacterial gastroenteritis associated with bacterial diarrhea, bacterial pulmonary infections, and bacterial infections of the urinary tract caused by susceptible organisms.

(iii) *Limitations*. Laboratory tests should be conducted, including in vitro culturing and susceptibility tests on samples collected prior to treatment. If no response is obtained in 3 to 5 days, discontinue use and reevaluate diagnosis. Not for animals that are raised for food production. Chloramphenicol products should not be administered in conjunction with or 2 hours prior to the induction of general anesthesia with pentobarbital because of prolonged recovery. Chloramphenicol should not be administered to dogs maintained for breeding purposes. Because of potential antagonism, chloramphenicol should not be administered simultaneously with penicillin or streptomycin. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37323, Aug. 18, 1992, as amended at 62 FR 35076, June 30, 1997]

§ 520.390b Chloramphenicol capsules.

(a) *Specifications*. Each capsule contains 50, 100, 250, or 500 milligrams of chloramphenicol.

(b) *Sponsor*. (1) For chloramphenicol capsules containing 50, 100, 250, or 500 milligrams of chloramphenicol see Nos. 000069, 000185, and 027454 in § 510.600(c) of this chapter.

(2) For chloramphenicol capsules containing 100 or 250 milligrams of chloramphenicol see No. 058034 in § 510.600(c) of this chapter.

(c) *Conditions of use*. *Dogs*—(1) *Amount*. 25 milligrams per pound of body weight every 6 hours.

(2) *Indications for use*. Oral treatment of bacterial pulmonary infections, bacterial infections of the urinary tract, bacterial enteritis, and bacterial infections associated with canine distemper caused by susceptible organisms.

(3) *Limitations*. Laboratory tests should be conducted including in vitro culturing and susceptibility tests on samples collected prior to treatment. This product must not be used in meat-, egg-, or milk-producing animals. The length of time that residues persist in milk or tissues has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37323, Aug. 18, 1992, as amended at 63 FR 5255, Feb. 2, 1998]

§ 520.390c Chloramphenicol palmitate oral suspension.

(a) *Specifications*. Each milliliter contains chloramphenicol palmitate equivalent to 30 milligrams of chloramphenicol.

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*. *Dogs*—(1) *Amount*. 25 milligrams per pound of body weight every 6 hours. If no response is obtained in 3 to 5 days, discontinue use and reevaluate diagnosis.

(2) *Indications for use*. Treatment of bacterial pulmonary infections, infections of the urinary tract, enteritis, and infections associated with canine distemper that are caused by organisms susceptible to chloramphenicol.

(3) *Limitations*. Not for use in animals that are raised for food production. Must not be used in meat-, egg-, or milk-producing animals. The length of time that residues persist in milk or tissues has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37323, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992]

§ 520.420 Chlorothiazide tablets and boluses.

(a)(1) *Specifications*. Each tablet contains 0.25 gram of chlorothiazide.

(2) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(3) *Conditions of use*—(i) *Amount*. Usual dosage is 5 to 10 milligrams per

§ 520.434

pound of body weight two or three times daily.¹

(ii) *Indications for use.* For use in dogs for treatment of congestive heart failure and renal edema.¹

(iii) *Limitations.* (a) Dosage must be adjusted to meet the changing needs of the individual animal. In mild and responsive cases, it is suggested that a dose of 5 milligrams per pound of body weight be administered two or three times daily. In moderately edematous and moderately responsive animals, a dose of 7.5 to 10 milligrams per pound of body weight may be administered three times daily. Severe conditions may require higher doses. Certain animals may respond adequately to intermittent therapy; in these cases, the drug may be administered either every other day or for 3 to 5 days each week.

(b) Animals should be regularly and carefully observed for early signs of fluid and electrolyte imbalance. Take appropriate countermeasures if this should occur. In some dogs, hypochloremic alkalosis may occur (that is, excretion of chloride in relation to sodium is excessive; the plasma bicarbonate level increases and alkalosis results). Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

(b)(1) *Specifications.* Each bolus contains 2 grams of chlorothiazide.

(2) *Sponsor.* See No. 000006 in § 510.600(c) of this chapter.

(3) *Conditions of use*—(i) *Amount.* 2 grams once or twice daily for 3 or 4 days.¹

(ii) *Indications for use.* For use in cattle as an aid in reduction of postparturient udder edema.¹

(iii) *Limitations.* Animals should be regularly and carefully observed for early signs of fluid and electrolyte imbalance. Take appropriate countermeasures if this should occur. Milk taken from dairy animals during treatment and for 72 hours (six milkings) after latest treatment must not be used for food. Federal law restricts this

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

21 CFR Ch. I (4–1–99 Edition)

drug to use by or on the order of a licensed veterinarian.¹

[43 FR 39085, Sept. 1, 1978, as amended at 62 FR 63270, Nov. 28, 1997]

§ 520.434 Chlorphenesin carbamate tablets.

(a) *Specifications.* Each tablet contains 400 milligrams of chlorphenesin carbamate.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount.* 50 milligrams per pound of body weight on first day; 25 milligrams per pound of body weight each following day. Divide total daily dose into 2 or 3 equal doses—administer at 12- or 8-hour intervals.

(2) *Indications for use.* For use as an adjunct to therapy of acute inflammatory and traumatic conditions of skeletal muscles. The drug provides relief of the signs of discomfort associated with myositis, muscle sprains, traumatic injuries, stifle injuries—especially when administered before or after surgery—and intervertebral disc syndrome (can be used concurrently with adrenal corticosteroids).

(3) *Limitations.* Not recommended for pregnant animals or those with a known hepatic dysfunction. Periodic liver function studies are recommended for animals on prolonged treatment. If no response is evident within 5 days of the beginning of treatment, the diagnosis should be redetermined and appropriate therapy instituted. Not recommended for use with general anesthetics other than the barbiturates. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 16009, Mar. 16, 1979]

§ 520.445 Chlortetracycline oral dosage forms.

§ 520.445a Chlortetracycline bisulfate/sulfamethazine bisulfate soluble powder.

(a) *Specifications.* Each pound contains chlortetracycline bisulfate equivalent to 102.4 grams of chlortetracycline hydrochloride with sulfamethazine bisulfate equivalent to 102.4 grams of sulfamethazine.

Food and Drug Administration, HHS

§ 520.445b

(b) *Sponsor*. See No. 010042 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See §§ 556.150 and 556.670 of this chapter.

(d) *Conditions of use*. *Swine*—Used in drinking water as follows:

(1) *Amount*. 250 milligrams of chlortetracycline with 250 milligrams of sulfamethazine per gallon.

(2) *Indications for use*. Prevention and treatment of bacterial enteritis; aid in the reduction of the incidence of cervical abscesses; aid in the maintenance of weight gains in the presence of bacterial enteritis and atrophic rhinitis.

(3) *Limitations*. Not to be used for more than 28 consecutive days; withdraw 15 days before slaughter; as sole source of chlortetracycline and sulfonamide.

[57 FR 37323, Aug. 18, 1992]

§ 520.445b Chlortetracycline powder (chlortetracycline hydrochloride or chlortetracycline bisulfate).

(a) *Specifications*. Chlortetracycline powder contains not less than 15 milligrams per gram chlortetracycline hydrochloride, or chlortetracycline bisulfate equivalent to 25.6, 64 or 102.4 grams per pound (56.4, 141 or 225.6 milligrams per gram) chlortetracycline hydrochloride.

(b) *Sponsors*. See No. 053389 in § 510.600(c) of this chapter for conditions of use as in paragraphs (d)(1)(i)(A) and (d)(2)(i)(A) of this section; No. 010042 for conditions of use as in paragraph (d)(4) of this section; No. 000010 for conditions of use as in paragraphs (d)(4)(i)(A) and (B) and (d)(4)(ii) through (iv) of this section; Nos. 017519 and 059130 for conditions of use as in paragraphs (d)(4)(i)(A) and (B) and (d)(4)(ii) and (iii) of this section.

(c) *Related tolerances*. See § 556.150 of this chapter.

(d) *Conditions of use*. (1) Use as chlortetracycline hydrochloride in drinking water as follows:

(i) *Swine*—(A) *Amount*. Ten milligrams per pound of body weight daily in divided doses.

(1) *Indications for use*. Control and treatment of bacterial enteritis (scours) caused by *Escherichia coli* and bacterial pneumonia associated with *Pasteurella* spp., *Actinobacillus*

pleuropneumoniae (*Hemophilus* spp.), and *Klebsiella* spp.

(2) *Limitations*. Prepare a fresh solution twice daily; as sole source of chlortetracycline; administer for not more than 5 days; do not slaughter animals for food within 5 days of treatment.

(B) [Reserved]

(ii) [Reserved]

(2) Use as chlortetracycline hydrochloride in a drench or drinking water as follows:

(i) *Calves*—(A) *Amount*. Ten milligrams per pound of body weight daily in divided doses.

(1) Control and treatment of bacterial enteritis (scours) caused by *E. coli* and bacterial pneumonia (shipping fever) associated with *Pasteurella* spp., *A. pleuropneumoniae* (*Hemophilus* spp.), and *Klebsiella* spp.

(2) *Limitations*. Prepare fresh solution daily; as sole source of chlortetracycline; administer for not more than 5 days; do not slaughter animals for food within 24 hours of treatment; do not administer this product with milk or milk replacers; administer 1 hour before or 2 hours after feeding milk or milk replacers; a withdrawal period has not been established in preruminating calves; do not use in calves to be processed for veal.

(B) [Reserved]

(ii) [Reserved]

(3) [Reserved]

(4) The following uses of chlortetracycline hydrochloride or chlortetracycline bisulfate in drinking water or drench were reviewed by the National Academy of Sciences/National Research Council (NAS/NRC) and found effective:

(i) *Chickens*—(A) *Amount*. 200 to 400 milligrams per gallon.

(1) *Indications for use*. Control of infectious synovitis caused by *Mycoplasma synoviae*.

(2) *Limitations*. Prepare fresh solution daily; as sole source of chlortetracycline; do not use for more than 14 days; do not slaughter animals for food within 24 hours of treatment; do not use in laying chickens.

(B) *Amount*. 400 to 800 milligrams per gallon.

(1) *Indications for use*. Control of chronic respiratory disease and air-sac

§ 520.445c

infections caused by *M. gallisepticum* and *E. coli*.

(2) *Limitations*. Prepare fresh solution daily; as sole source of chlortetracycline; do not use for more than 14 days; do not slaughter animals for food within 24 hours of treatment; do not use in laying chickens.

(C) *Amount*. One thousand milligrams per gallon.

(1) *Indications for use*. Control of mortality due to fowl cholera caused by *Pasteurella multocida* susceptible to chlortetracycline.

(2) *Limitations*. See paragraph (d)(4)(i)(A)(2) of this section.

(ii) *Growing turkeys*—(A) *Amount*. 400 milligrams per gallon.

(1) *Indications for use*. Control of infectious synovitis caused by *M. synoviae*.

(2) *Limitations*. Prepare fresh solution daily; as sole source of chlortetracycline; do not use for more than 14 days; do not slaughter animals for food within 24 hours of treatment.

(B) *Amount*. 25 milligrams per pound of body weight daily.

(1) *Indications for use*. Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis).

(2) *Limitations*. Prepare fresh solution daily; as sole source of chlortetracycline; do not use for more than 14 days; do not slaughter animals for food within 24 hours of treatment.

(iii) *Swine*—(A) *Amount*. 10 milligrams per pound body weight daily in divided doses.

(B) *Indications for use*. Control and treatment of bacterial enteritis (scours) caused by *E. coli* and *Salmonella* spp. and bacterial pneumonia associated with *Pasteurella* spp., *Actinobacillus pleuropneumoniae* (*Hemophilus* spp.), and *Klebsiella* spp.

(C) *Limitations*. Prepare fresh solution daily; as sole source of chlortetracycline; do not use for more than 5 days; for 012286, 053389, and 054273 do not slaughter animals for food within 5 days of treatment; for 010042 do not slaughter animals for food within 24 hours of treatment.

(iv) *Calves, beef cattle, and nonlactating dairy cattle*—(A) *Amount*. 10 milligrams per pound daily in divided doses.

21 CFR Ch. I (4–1–99 Edition)

(B) *Indications for use*. Control and treatment of bacterial enteritis (scours) caused by *E. coli* and *Salmonella* spp. and bacterial pneumonia (shipping fever complex) associated with *Pasteurella* spp., *A. pleuropneumoniae* (*Hemophilus* spp.), and *Klebsiella* spp.

(C) *Limitations*. Prepare fresh solution daily; use as a drench; as sole source of chlortetracycline; do not use for more than 5 days; do not slaughter animals for food within 24 hours of treatment; do not use in lactating cattle; do not administer this product with milk or milk replacers; administer 1 hour before or 2 hours after feeding milk or milk replacers; a withdrawal period has not been established in prurminating calves; do not use in calves to be processed for veal.

[57 FR 37324, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992; 58 FR 61015, Nov. 19, 1993; 59 FR 39439, Aug. 3, 1994; 60 FR 26827, May 19, 1995; 60 FR 47052, Sept. 11, 1995; 62 FR 27691, May 21, 1997; 62 FR 35076, June 30, 1997; 62 FR 60656, Nov. 12, 1997]

§ 520.445c Chlortetracycline tablets and boluses.

(a) *Specifications*. Each tablet/bolus contains 25, 250, or 500 milligrams of chlortetracycline hydrochloride.

(b) *Sponsors*. See No. 000010 in § 510.600(c) of this chapter for the 250-milligram chlortetracycline hydrochloride bolus; see No. 010042 for the 25-milligram tablet and the 500 milligram bolus.

(c) *Related tolerances*. See § 556.150 of this chapter.

(d) *National Academy of Sciences/National Research Council NAS/NRC status*. The conditions of use specified in this section were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified in § 514.111 of this chapter but may require bioequivalency and safety information.

(e) *Conditions of use*. *Calves*—(1) *Amount*. One 250 milligram bolus per 50 pounds of body weight twice a day for 3 to 5 days.

(i) *Indications for use*. Treatment of bacterial enteritis (scours) caused by *Escherichia coli* and bacterial pneumonia associated with *Pasteurella* spp., *Klebsiella* spp., and *Hemophilus* spp.

(ii) *Limitations.* Administer bolus directly by mouth or crush and dissolve in milk or water for drenching or bucket feeding; if no improvement is noted after 3 days of treatment, consult a veterinarian; do not use for more than 5 days; do not administer within 24 hours of slaughter.

(2) *Amount.* One 25 milligram tablet for each 5 pounds of body weight every 12 hours daily for 3 to 5 days.

(i) *Indications for use.* Control and treatment of bacterial enteritis (scours) caused by *E. coli* and *Salmonella* spp. and bacterial pneumonia associated with *Pasteurella* spp., *Hemophilus* spp., and *Klebsiella* spp., susceptible to chlortetracycline.

(ii) *Limitations.* Administer tablet directly by mouth or crush and dissolve in water for drenching; if no improvement is noted after 3 days of treatment, consult a veterinarian; do not use for more than 5 days; when feeding milk or milk replacer, administration 1 hour before or 2 hours after feeding; do not administer within 24 hours of slaughter.

(3) *Amount.* One 500 milligram bolus per 100 pounds of body weight twice a day for 3 to 5 days.

(i) *Indications for use.* Treatment of bacterial enteritis (scours) caused by *E. coli* and *Salmonella* spp., and bacterial pneumonia associated with *Pasteurella* spp., *Hemophilus* spp., and *Klebsiella* spp., susceptible to chlortetracycline.

(ii) *Limitations.* Administer directly by mouth or crush and dissolve in water for drenching; if no improvement is noted after 3 days of treatment, consult a veterinarian; do not use for more than 5 days; do not administer within 24 hours of slaughter.

[57 FR 37325, Aug. 18, 1992]

§ 520.446 Clindamycin hydrochloride capsules.

(a) *Specifications.* Each capsule contains the equivalent of 25, 75, or 150 milligrams of clindamycin as the hydrated hydrochloride salt.

(b) *Sponsor.* No 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs—(1) Amount.* Wounds, abscesses, and dental infections: 2.5 milligrams per pound of body weight every 12 hours for a maximum of 28 days. Osteomyelitis: 5.0

milligrams per pound of body weight every 12 hours for a minimum of 28 days.

(2) *Indications for use.* For use in dogs for treatment of soft tissue infections (wounds and abscesses), dental infections, and osteomyelitis caused by susceptible strains of *Staphylococcus aureus*, soft tissue infections (deep wounds and abscesses), dental infections, and osteomyelitis caused by or associated with susceptible strains of *Bacteroides fragilis*, *Bacteroides melaninogenicus*, *Fusobacterium necrophorum*, and *Clostridium perfringens*.

(3) *Limitations.* Wound infections, abscesses, and dental infections: Do not use more than 4 days if no improvement of acute infection is observed. Osteomyelitis: Do not use for more than 28 consecutive days if no improvement is observed. Because of potential adverse gastrointestinal effects, do not administer to rabbits, hamsters, guinea pigs, and horses. Use with caution in animals receiving neuromuscular blocking agents, because clindamycin may potentiate their action. Prescribe with caution in atopic animals. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[49 FR 23341, June 6, 1984, as amended at 51 FR 34960, Oct. 1, 1986; 54 FR 47767, Nov. 17, 1989]

§ 520.447 Clindamycin hydrochloride liquid.

(a) *Specifications.* Each milliliter of 8.64 percent alcoholic solution contains the equivalent of 25 milligrams of clindamycin (as the hydrochloride).

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter for use as in paragraphs (c) and (d) of this section. See No. 059130 for use as in paragraph (c) of this section.

(c) *Conditions of use in dogs—(1) Amount.* Wounds, abscesses, and dental infections: 2.5 milligrams per pound of body weight every 12 hours for a maximum of 28 days. Osteomyelitis: 5.0 milligrams per pound of body weight every 12 hours for a minimum of 28 days.

(2) *Indications for use.* For use in dogs for treatment of soft tissue infections

§ 520.452

(wounds and abscesses), dental infections, and osteomyelitis caused by susceptible strains of *Staphylococcus aureus* and for soft tissue infections (deep wounds and abscesses), dental infections, and osteomyelitis caused by or associated with susceptible strains of *Bacteroides fragilis*, *Bacteroides melaninogenicus*, *Fusobacterium necrophorum*, and *Clostridium perfringens*.

(3) *Limitations*. Wound infections, abscesses, and dental infections: Do not use for more than 4 days if no improvement of acute infection is observed. Osteomyelitis: Do not use for more than 28 consecutive days if no improvement is observed. Because of potential adverse gastrointestinal effects, do not administer to rabbits, hamsters, guinea pigs, horses, chinchillas, or ruminating animals. Use with caution in animals receiving neuromuscular blocking agents, because clindamycin may potentiate their action. Prescribe with caution in atopic animals. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d) *Conditions of use in cats*—(1) *Amount*. 5.0 to 10.0 milligrams per pound of body weight every 24 hours for a maximum of 14 days (11 to 22 milligrams per kilogram of body weight per day).

(2) *Indications for use*. Aerobic bacteria: Treatment of soft tissue infections (wounds and abscesses) and dental infections caused by or associated with susceptible strains of *Staphylococcus aureus*, *S. intermedius*, and *Streptococcus spp.* Anaerobic bacteria: Treatment of soft tissue infections (deep wounds and abscesses) and dental infections caused by or associated with susceptible strains of *Clostridium perfringens* and *Bacteroides fragilis*.

(3) *Limitations*. Wound infections, abscesses, and dental infections: Do not use for more than 4 days if no improvement of acute infection is observed. Because of potential adverse gastrointestinal effects, do not administer to rabbits, hamsters, guinea pigs, horses, chinchillas, or ruminating animals. Use with caution in animals receiving neuromuscular blocking agents, because clindamycin may potentiate their action. Prescribe with caution in

21 CFR Ch. I (4–1–99 Edition)

atopic animals. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 21239, May 23, 1985, as amended at 51 FR 34960, Oct. 1, 1986; 54 FR 47766, Nov. 17, 1989; 61 FR 59003, Nov. 20, 1996; 62 FR 46669, Sept. 4, 1997]

§ 520.452 Clenbuterol syrup.

(a) *Specifications*. Each milliliter contains 72.5 micrograms of clenbuterol hydrochloride.

(b) *Sponsor*. See 000010 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Horses*—(i) *Amount*. Administer orally twice a day (b.i.d.). Initial dose is 0.5 milliliter per 100 pounds body weight (0.8 micrograms per kilogram) for 3 days (6 treatments). If no improvement, administer 1 milliliter per 100 pounds (1.6 micrograms per kilogram) for 3 days (6 treatments). If no improvement, administer 1.5 milliliters per 100 pounds (2.4 micrograms per kilogram) for 3 days (6 treatments). If no improvement, administer 2.0 milliliters per 100 pounds (3.2 micrograms per kilogram) for 3 days (6 treatments). If no improvement, horse is non-responder to clenbuterol and treatment should be discontinued.

(ii) *Indications for use*. Indicated for the management of horses affected with airway obstruction, such as occurs in chronic obstructive pulmonary disease (COPD).

(iii) *Limitations*. Treat at effective dose for 30 days. At the end of the 30-day treatment period, drug should be withdrawn. If signs return, the 30-day treatment period may be repeated. If repeating treatment, the step-wise dosage schedule should be repeated. The effect of this drug on breeding stallions and brood mares has not been determined. Treatment starting with dosages higher than the initial dose is not recommended. Federal law prohibits the extralabel use of this drug in food animals. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[63 FR 41419, Aug. 4, 1998]

§ 520.455 Clomipramine hydrochloride tablets.

(a) *Specifications.* Each tablet contains 20, 40, or 80 milligrams of clomipramine hydrochloride.

(b) *Sponsor.* See No. 058198 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* 2 to 4 milligrams of clomipramine hydrochloride per kilogram (0.9 to 1.8 milligrams per pound) of body weight per day, administered as a single daily dose or divided twice daily.

(2) *Indications for use.* For use as part of a comprehensive behavioral management program to treat separation anxiety in dogs greater than 6 months of age.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[64 FR 1762, Jan. 12, 1999]

§ 520.462 Clorsulon drench.

(a) *Specifications.* The drug is a suspension containing 8.5 percent clorsulon (85 milligrams per milliliter).

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use. Cattle—(1) Amount.* One-quarter fluid ounce per 200 pounds of body weight (7 milligrams per kilogram or 3.2 milligrams per pound of body weight).

(2) *Indications for use.* For the treatment of immature and adult liver fluke (*Fasciola hepatica*) infestations in cattle.

(3) *Limitations.* Using dose syringe, deposit drench over back of tongue. Do not treat cattle within 8 days of slaughter. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[50 FR 10221, Mar. 14, 1985, as amended at 62 FR 63270, Nov. 28, 1997]

§ 520.530 Cythioate oral liquid.

(a) *Specifications.* Each milliliter contains 15 milligrams of cythioate.

(b) *Sponsor.* See Nos. 000859 and 010042 in § 510.600 of this chapter.

(c) *Special considerations.* Cythioate is a cholinesterase inhibitor. Do not use this product in animals simultaneously

with or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, insecticides, pesticides, or chemicals.

(d) *Conditions of use—(1) Amount.* 15 milligrams cythioate per 10 pounds of body weight every third day or twice a week.

(2) *Indications for use.* Dogs, for control of fleas.

(3) *Limitations.* For oral use in dogs only. Do not use in greyhounds or in animals that are pregnant, sick, under stress, or recovering from surgery. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[49 FR 5614, Feb. 14, 1984]

§ 520.531 Cythioate tablets.

(a) [Reserved]

(b) *Sponsors.* See No. 000859 in § 510.600(c) of this chapter for use of 30- and 90-milligram (mg) tablets and see No. 010042 in § 510.600(c) of this chapter for use of 30-mg tablet.

(c) *Special considerations.* Cythioate is a cholinesterase inhibitor. Do not use this product in animals simultaneously with or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, insecticides, pesticides, or chemicals.

(d) *Conditions of use—(1) Amount.* 30 milligrams cythioate per 20 pounds of body weight every third day or twice a week.

(2) *Indications for use.* Dogs, for control of fleas.

(3) *Limitations.* For oral use in dogs only. Do not use in greyhounds or in animals that are pregnant, sick, under stress, or recovering from surgery. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[49 FR 5615, Feb. 14, 1984, as amended at 59 FR 26942, May 25, 1994]

§ 520.534 Decoquinat.

(a) *Specifications.* The drug is a powder containing 0.8 percent decoquinat.

(b) *Sponsor.* See No. 046573 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.170 of this chapter.

(d) *Conditions of use. Replacement calves—(1) Amount.* Feed 22.7 milligrams per 100 pounds of body weight (0.5 milligram per kilogram) per day.

§ 520.540

21 CFR Ch. I (4–1–99 Edition)

(2) *Indications for use.* For the prevention of coccidiosis in ruminating and nonruminating calves, including veal calves, caused by *Eimeria bovis* and *E. zuernii*.

(3) *Limitations.* Feed in whole milk at the rate of 22.7 milligrams per 100 pounds body weight daily (0.5 milligram per kilogram) for at least 28 days.

[64 FR 10103, Mar. 2, 1999]

§ 520.540 Dexamethasone oral dosage forms.

§ 520.540a Dexamethasone powder.

(a) *Specifications.* Dexamethasone powder is packaged in packets containing 10 milligrams of dexamethasone.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) Dexamethasone powder is indicated in cases where cattle and horses require additional steroid therapy following its parenteral administration. The drug is used as supportive therapy for management or inflammatory conditions such as acute arthritic lameness, and for various stress conditions where corticosteroids are required while the animal is being treated for a specific condition.

(2) The drug is administered at a dosage level of 5 to 10 milligrams per animal the first day then 5 milligrams per day as required by drench or by sprinkling on a small amount of feed.

(3) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975; 41 FR 9149, Mar. 3, 1976; 52 FR 7832, Mar. 13, 1987]

§ 520.540b Dexamethasone tablets and boluses.

(a)(1) *Specifications.* Each bolus is half-scored and contains 10 milligrams of dexamethasone.

(2) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) Dexamethasone bolus is indicated in cases where cattle and horses require additional steroid therapy following its parenteral administration. The drug may be used as supportive therapy for management of inflammatory conditions such as acute arthritic lamenesses, and for various stress conditions where corticosteroids are required while the animal is being treated for a specific condition.

(ii) Administered orally, 5 to 10 milligrams for the first day, then 5 milligrams per day as required.

(iii) Do not use in viral infections during the viremic stage. With bacterial infections, appropriate antibacterial therapy should be used.

(iv) Do not use in animals with chronic nephritis and hypercorticalism (Cushingoid syndrome), except for emergency therapy.

(v) Clinical and experimental data have demonstrated that corticosteroids administered orally or by injection to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(vi) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Specifications.* Each tablet contains 0.25 milligram of dexamethasone.¹

(2) *Sponsors.* See Nos. 000061 and 050604 in § 510.600(c) of this chapter.

(3) *Conditions of use—(i) Amount.* Dogs: Administer orally at 0.25 to 1.25 milligrams per day for up to 7 days. Cats: 0.125 to 0.5 milligram per day for up to 7 days.¹

(ii) *Indications for use.* In treatment of dogs and cats as an anti-inflammatory agent.¹

(iii) *Limitations.* (a) Clinical and experimental data have demonstrated that corticosteroids administered orally or by injection to animals may induce the first stage of parturition when administered during the last trimester of pregnancy; and they may precipitate

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter.

Food and Drug Administration, HHS

§ 520.550

premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(b) Do not use in viral infections. Anti-inflammatory action of corticosteroids may mask signs of infections. Do not use in animals with tuberculosis, chronic nephritis, cushingoid syndrome, or peptic ulcers, except for emergency therapy.¹

(c) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 26273, June 23, 1975, as amended at 44 FR 7130, Feb. 6, 1979; 50 FR 49372, Dec. 2, 1985; 52 FR 7832, Mar. 13, 1987; 55 FR 8461, Mar. 8, 1990]

§ 520.540c Dexamethasone chewable tablets.

(a) *Specifications.* Each half-scored tablet contains 0.25 milligram of dexamethasone.¹

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 0.25 to 1.25 milligrams per day.¹

(2) *Indications for use.* Supportive therapy in nonspecific dermatosis and inflammatory conditions in dogs.¹

(3) *Limitations.* (i) Administer by free-choice feeding or crumble over food. Administer 0.25 to 1.25 milligrams daily in single or two divided doses until response is noted or 7 days have elapsed. When response is attained, dosage should be gradually reduced by 0.125 milligram per day until maintenance level is achieved.

(ii) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy; and they may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(iii) Do not use in viral infections. Anti-inflammatory action of corticosteroids may mask signs of infection. Do not use in animals with tuberculosis, chronic nephritis, cushingoid syndrome, or peptic ulcers, except for emergency therapy.¹

(iv) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 7130, Feb. 6, 1979, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.550 Dextrose/glycine/electrolyte.

(a) *Specifications.* The product is distributed in packets each of which contains the following ingredients: sodium chloride 8.82 grams, potassium phosphate 4.20 grams, citric acid anhydrous 0.5 gram, potassium citrate 0.12 gram, aminoacetic acid (glycine) 6.36 grams, and dextrose 44.0 grams.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) Dextrose/glycine/electrolyte is indicated for use in the control of dehydration associated with diarrhea (scours) in calves. It is used as an early treatment at the first signs of scouring. It may also be used as followup treatment following intravenous fluid therapy.

(2) Dissolve each packet in two quarts of warm water and administer to each calf as follows:

(i) *Scouring and/or dehydrated calves.* Feed 2 quarts of solution, twice daily for 2 days (four feedings). No milk or milk replacer should be fed during this period. For the next four feedings (days 3 and 4), use 1 quart of solution together with 1 quart of milk replacer. Thereafter, feed as normal.

(ii) *Newly purchased calves.* Feed 2 quarts of solution instead of milk as the first feed upon arrival. For the next scheduled feeding, use 1 quart of solution mixed together with 1 quart of milk or milk replacer. Thereafter, feed as normal.

(3) The product should not be used in animals with severe dehydration (down, comatose, or in a state of shock). Such animals need intravenous therapy. Oral therapy in these cases is too slow. Animals which cannot drink after initial intravenous therapy may need to be dosed with a stomach tube or esophageal tube. Adequate colostrum intake during the first 12 hours is essential for healthy, vigorous calves. Antibacterial therapy is often indicated in bacterial scours due to *E. coli* and/or *Salmonella*. The product does not contain antibacterial agents. A veterinarian should be consulted in severely

§ 520.563

scouring calves or cases requiring antibacterial therapy. The product is not nutritionally complete if administered by itself for long periods of time. It should not be administered beyond the recommended treatment period without the addition of milk or milk replacer.

[48 FR 38606, Aug. 25, 1983, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.563 Diatrizoate meglumine and diatrizoate sodium oral solution.

(a) *Specifications.* Diatrizoate meglumine oral solution is a water soluble radiopaque medium containing 66 percent diatrizoate meglumine and 10 percent diatrizoate sodium.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is indicated for radiography of the gastrointestinal tract in dogs and cats.

(2) It is administered orally at a dosage level of 0.5 to 1.0 milliliter per pound of body weight by gavage or stomach tube. It is administered rectally at a dosage level of 0.5 to 1.0 milliliter per pound of body weight diluted with 1 part of the drug to 5 parts of water.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 12993, Mar. 9, 1979, as amended at 50 FR 41489, Oct. 11, 1985]

§ 520.580 Dichlorophene and toluene capsules.

(a) *Specifications.* Each soft gelatin capsule contains 50 milligrams of dichlorophene and 60 milligrams of toluene or multiples thereof.¹

(b) *Sponsor.* (1) For single dose only, see 000010, 000061, 000115, 000842, 011615, 017135, 023851, 049968, 050906, and 058670 in § 510.600(c) of this chapter.

(2) For single and multiple dose, see 000010, 000061, and 038782 in § 510.600(c) of this chapter.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

21 CFR Ch. I (4–1–99 Edition)

(c) *Required statement.* Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism, and before administering to weak or debilitated animals.

(d) *Conditions of use*—(1) *Amount.* (i) Single dose of 100 milligrams of dichlorophene and 120 milligrams of toluene per pound of body weight.¹

(ii) Divided dose of 100 milligrams of dichlorophene and 120 milligrams of toluene per 5 pounds of body weight (20 and 24 milligrams per pound) daily for 6 days.¹

(2) *Indications for use.* It is used for the removal of ascarids (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) and as an aid in removing tapeworms (*Taenia pisiformis*, *Dipylidium caninum*, and *Echinococcus granulosus*) from dogs and cats.¹

(3) *Limitations.* Withhold solid foods and milk for at least 12 hours prior to medication and for 4 hours afterward. Repeat treatment in 2 to 4 weeks in animals subject to reinfection.¹

[45 FR 10332, Feb. 15, 1980]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 520.580, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 520.581 Dichlorophene tablets.

(a) *Specifications.* Each tablet contains 1 gram of dichlorophene.

(b) *Sponsor.* See 023851 in § 510.600(c) of this chapter.

(c) *Required statement.* Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism, and before administering to weak or debilitated animals.

(d) *Conditions of use.* *Dogs*—(1) *Amount.* Single dose of 1 tablet (1 gram of dichlorophene) for each 10 pounds of body weight.

(2) *Indications for use.* It is used as an aid in the removal of tapeworms (*Taenia pisiformis* and *Dipylidium caninum*).

(3) *Limitations.* Withhold solid foods and milk for at least 12 hours prior to medication and for 4 hours afterward.

[45 FR 10333, Feb. 15, 1980]

Food and Drug Administration, HHS

§ 520.600

§ 520.600 Dichlorvos.

(a) *Chemical name.* 2,2-Dichlorovinyl dimethyl phosphate.

(b) [Reserved]

(c) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(d) *Related tolerances.* See § 556.180 of this chapter.

(e) *Conditions of use in swine.* (1) It is recommended for the removal and control of sexually mature (adult), sexually immature and/or 4th stage larvae of the whipworm (*Trichuris suis*), nodular worms (*Oesophagostomum* spp.), large round-worm (*Ascaris suum*), and the mature thick stomach worm (*Ascarops strongylina*) occurring in the lumen of the gastrointestinal tract of pigs, boars, and open or bred gilts and sows.

(2) The preparation should be added to the indicated amount of feed as set forth in paragraph (e)(2) of this section and administered shortly after mixing, as follows:

Weight of animal in pounds	Pounds of feed to be mixed with each 0.08 ounce of dichlorvos	Pounds of mixed feed to be administered to each pig as a single treatment	Number of pigs to be treated per 0.08 ounce of dichlorvos
20-30	4	0.33	12
31-40	5	0.56	9
41-60	6	1.00	6
61-80	5	1.00	5
81-100	4	1.00	4
Adult Gilts, Sows, and Boars	16	4.00	4

(3) Do not use this product on animals either simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. The preparation should be mixed thoroughly with the feed on a clean, impervious surface. Do not allow swine access to feed other than that containing the preparation until treatment is complete. Do not treat pigs with signs of scours until these signs subside or are alleviated by proper medication. Resume normal feeding schedule afterwards. Swine may be retreated in 4 to 5 weeks.

(f) *Conditions of use in dogs.* (1) For removal of *Toxocara canis* and *Toxascaris leonina* (roundworms), *Ancylostoma caninum* and *Uncinaria stenocephala*

(hookworms), and *Trichuris vulpis* (whipworm) residing in the lumen of the gastrointestinal tract.

(2) The drug is in capsule form for direct administration and in pellet form for administration in about one-third of the regular canned dog food ration or in ground meat. Dogs may be treated with any combination of capsules and/or pellets so that the animal receives a single dose equaling 12 to 15 milligrams of the active ingredient per pound of body weight. One-half of the single recommended dosage may be given, and the other half may be administered 8 to 24 hours later. This split dosage schedule should be used in animals which are very old, heavily parasitized, anemic, or otherwise debilitated. The drug should not be used in dogs weighing less than 2 pounds.

(3) In some dogs, efficacy against *Trichurias vulpis* (whipworm) may be erratic. Dogs that do not develop a negative stool for *Trichuris vulpis* ova 10 to 14 days following initial treatment should be re-treated. If a negative stool is not obtained in 10 to 14 days following re-treatment, alternate means of therapy should be considered.

(4) Do not use in dogs infected with *Dirofilaria immitis*.

(5) Do not use with other anthelmintics, taeniocides, antifilarial agents, muscle relaxants, or tranquilizers.

(6) The drug is a cholinesterase inhibitor. Not for use simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(7) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(g) *Conditions of use in horses when administered in grain.* (1) It is recommended for the removal and control of bots (*Gastrophilus intestinalis*, *G. nalis*), large strongyles (*Strongylus vulgaris*, *S. equinus*, *S. edentatus*), small strongyles (of the genera *Cyathostomum*, *Cylicocercus*, *Cylicocylus*, *Cylicodontophorus*, *Triodontophorus*, *Poteriostomum*, *Gyalocephalus*), pinworms (*Oxyuris equi*), and large roundworm (*Parascaris equorum*) in horses including ponies and

mules. Not for use in foals (sucklings and young weanlings).

(2) For a satisfactory diagnosis, a microscopic fecal examination should be performed by a veterinarian or a diagnostic laboratory prior to worming.

(3) It is administered in the grain portion of the ration at a dosage of 14.2 milligrams to 18.5 milligrams per pound of body weight as a single dose. It may be administered at one-half of the single recommended dosage and repeated 8 to 12 hours later in the treatment of very aged, emaciated or debilitated subjects or those reluctant to consume medicated feed. In suspected cases of severe ascarid infection sufficient to cause concern over mechanical blockage of the intestinal tract, the split dosage should be utilized.

(4) Do not use in horses which are severely debilitated, suffering from diarrhea or severe constipation, infectious disease, toxemia or colic. Do not administer in conjunction with or within 1 week of administration of muscle relaxant drugs, phenothiazine derived tranquilizers or central nervous system depressant drugs. Horses should not be subjected to insecticide treatment for 5 days prior to or after treating with the drug. Do not administer to horses afflicted with chronic alveolar emphysema (heaves) or related respiratory conditions. The product is a cholinesterase inhibitor and should not be used simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

(5) Do not use in animals other than horses, ponies, and mules. Do not use in horses, ponies, and mules intended for food purposes. Do not allow fowl access to feed containing this preparation or to fecal excrement from treated animals.

(h) *Conditions of use in horses when administered orally by syringe.* (1) It is recommended for the removal and control of first, second, and third instar bots (*Gastrophilus intestinalis* and *G. nasalis*), sexually mature and sexually immature (4th stage) ascarids (*Parascaris equorum*) in horses and foals.

(2) The product is in the form of a gel which is administered directly from a syringe onto the horse's tongue. The product is administered at a dosage

level of 20 milligrams of dichlorvos per kilogram of body weight for the removal of bots and ascarids. The same dosage level is repeated every 21 to 28 days for the control of bots and ascarids. For the control of bots only, the repeat dosage is 10 milligrams per kilogram of body weight every 21 to 28 days during bot fly season.

(3) Do not use this product in animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides or chemicals. Do not administer in conjunction with or within 1 week of administration of muscle-relaxant drugs, phenothiazine derived tranquilizers, or central nervous system depressants.

(4) Do not use in horses which are severely debilitated or suffering from diarrhea or severe constipation, infectious disease, toxemia, or colic. Do not administer to horses affected with chronic alveolar emphysema (heaves) or other respiratory conditions.

(5) Do not use in horses intended for food purposes.

(6) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(i) *Conditions of use in cats and puppies.* (1) It is indicated for the removal and control of roundworms (*Toxocara canis*, *Toxocara cati*, *Toxascaris leonina*) and hookworms (*Ancylostoma caninum*, *Ancylostoma tubaeforme*, *Uncinaria stenocephala*) occurring in the intestinal tracts of cats and puppies.

(2) The drug is in tablet form and is administered orally at a dosage level of 5 mg of the active ingredient per pound of body weight.

(3) Do not administer to puppies or cats showing signs of constipation, mechanical blockage of the intestinal tract, impaired liver function, or to animals recently exposed to or showing signs of infectious disease. The drug is a cholinesterase inhibitor and should not be used simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(4) Do not use in animals under 10 days of age or under 1 pound of body weight.

Food and Drug Administration, HHS

§ 520.622a

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 48 FR 40704, Sept. 9, 1983; 51 FR 28546, Aug. 8, 1986; 62 FR 35076, June 30, 1997]

§ 520.608 Dicloxacillin sodium monohydrate capsules.

(a) *Specifications.* Each capsule contains dicloxacillin sodium monohydrate equivalent to 50, 100, 200, or 500 milligrams of dicloxacillin.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use. Dogs—*(1) *Amount.* 5 to 10 milligrams per pound of body weight, three times daily. In severe cases, up to 25 milligrams per pound of body weight three times daily.

(2) *Indications for use.* Treatment of pyoderma (pyogenic dermatitis) due to penicillinase-producing staphylococci sensitive to the drug.

(3) *Limitations.* For the treatment of dogs only. Continue treatment for 24 to 48 hours after the animal has become afebrile or asymptomatic. Administer 1 to 2 hours before feeding to ensure maximum absorption. Not for use in animals which are raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37325, Aug. 18, 1992]

§ 520.620 Diethylcarbamazine oral dosage forms.

§ 520.622 Diethylcarbamazine citrate oral dosage forms.

§ 520.622a Diethylcarbamazine citrate tablets.

(a) *Sponsors.* (1) See 015579 in § 510.600(c) of this chapter for use of 50, 200, and 400 milligram tablets for prevention of heartworm disease in dogs and as an aid in the treatment of ascarid infections in dogs and cats.

(2) See 053501 in § 510.600(c) of this chapter for use of 100, 200, and 300 milligram tablets for prevention of heartworm disease in dogs and as an aid in the treatment of ascarid infections in dogs.

(3) See 050604 in § 510.600(c) of this chapter for use of 50, 100, 200, 300, or 400

milligram tablets for prevention of heartworm disease in dogs, as an aid in the control of ascarid infections in dogs, and as an aid in the treatment of ascarid infections in dogs and cats.

(4) See 017030 in § 510.600(c) of this chapter for use of 50, 100, 200, 300, and 400 milligram tablets for prevention of heartworm disease in dogs and as an aid in the treatment of ascarid infections in dogs and cats.

(5) See 000081 in § 510.600(c) of this chapter for use of 60, 120, or 180 milligram tablets for prevention of heartworm disease in dogs, as an aid in the control of ascarid infections in dogs, and as an aid in the treatment of ascarid infections in dogs and cats.

(6) See No. 000010 in § 510.600(c) of this chapter for use of 50, 100, 200, 300, or 400 milligram tablets for prevention of heartworm disease in dogs, as an aid in the control of ascarid infections in dogs, and as an aid in the treatment of ascarid infections in dogs and cats.

(b) *Conditions of use—*(1) *Dosage/indications for use.* (i) Three milligrams per pound of body weight daily for prevention of heartworm disease (*Dirofilaria immitis*) in dogs.

(ii) Three milligrams per pound of body weight daily as an aid in the control of ascarid infections (*Toxocara canis*) in dogs.

(iii) Twenty-five to 50 milligrams per pound of body weight as an aid in the treatment of ascarid infections in dogs (*Toxocara canis*) and cats (*Toxocara canis* and *Toxascaris leonina*).

(2) *Limitations.* Administer orally either pulverized and given in feed or water or directly by mouth. For the treatment of ascarid infections, repeat in 10 to 20 days to remove immature worms that may enter the intestine from the lungs after the first dose. Do not treat dogs with established heartworm infections until they have been converted to a negative status by the use of adulticidal and microfilaricidal drugs. Inadvertent administration to heartworm-infected dogs may cause adverse reactions because of pulmonary occlusion. Overdosage may cause emesis. For prevention of heartworm disease in heartworm-endemic areas, administration of the drug should start at the beginning of mosquito activity and be continued daily throughout the

§ 520.622b

mosquito season and for approximately a month thereafter. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 23230, Apr. 24, 1981, as amended at 46 FR 41038, Aug. 14, 1981; 46 FR 46315, Sept. 18, 1981; 46 FR 61653, Dec. 18, 1981; 47 FR 10805, Mar. 12, 1982; 47 FR 14150, Apr. 2, 1982; 50 FR 41489, Oct. 11, 1985; 50 FR 49372, Dec. 2, 1985; 53 FR 40056, Oct. 13, 1988; 53 FR 40727, Oct. 18, 1988; 55 FR 8461, Mar. 8, 1990; 61 FR 34728, July 3, 1996; 62 FR 35076, June 30, 1997]

§ 520.622b Diethylcarbamazine citrate syrup.

(a)(1) *Specifications.* Each milliliter of syrup contains 60 milligrams of diethylcarbamazine citrate.

(2) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) The drug is indicated for use in dogs for the prevention of infection with *Dirofilaria immitis* and *T. canis* and *T. leonina*. It is also indicated for treatment of ascarid infections of *T. canis* and *T. leonina* in dogs and *T. cati* in cats.

(ii) For prevention of heartworm and ascarid infections in dogs, the drug may be added to the daily diet at a dosage rate of 3.0 milligrams per pound of body weight per day or given directly by mouth at the same dosage rate. For treatment of ascarid infections in dogs and cats, the drug is administered at a dosage level of 25 to 50 milligrams per pound of body weight preferably administered immediately after feeding.

(iii) Older dogs should be proven negative for the presence of *Dirofilaria immitis* infection before administration of the drug. Those with proven infection of *Dirofilaria immitis* should be rendered negative using adulticidal and microfilaricidal drugs before administration of this drug.

(iv) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Specifications.* Each milliliter of syrup contains 60 milligrams of diethylcarbamazine citrate.

(2) *Sponsors.* (i) See No. 017030 for use as in paragraphs (b)(3)(ii)(a) and (b)(3)(ii)(c) of this section.

(ii) See No. 017030 for use as in paragraphs (b)(3)(ii) (a) and (c) of this section.

(3) *Conditions of use—(i) Amount.* 3 milligrams per pound of body weight

21 CFR Ch. I (4–1–99 Edition)

per day for prevention of heartworm disease and as an aid in control of large roundworms; 25 to 50 milligrams per pound of body weight as an aid in treatment of ascarid infections.

(ii) *Indications for use.* (a) For prevention of heartworm disease (*Dirofilaria immitis*) in dogs.

(b) As an aid in control of large roundworms (*T. canis*) in dogs.

(c) As an aid in treatment of ascarid infections in dogs (*T. canis*) and cats (*T. canis* and *T. leonina*).

(iii) *Limitations.* The drug may be placed on the daily ration or given directly by mouth. For treatment of ascarid infections, a repeat dose should be given in 10 to 20 days to remove immature worms which may enter the intestine from the lungs after the first dose. Older dogs should be proven negative for presence of *Dirofilaria immitis* infections before administering the drug. Dogs with established heartworm infections should not receive the drug until they have been converted to a negative status by the use of adulticidal and microfilaricidal drugs. Inadvertent administration to heartworm-infected dogs may cause adverse reactions due to pulmonary occlusion. Overdosage may cause emesis. For prevention of heartworm disease in heartworm-endemic areas, administration of the drug should start 1 month before the mosquito season and be continued daily throughout the mosquito season and for 2 months thereafter. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(c)(1) *Specifications.* Each milliliter of syrup contains 60 milligrams of diethylcarbamazine citrate.

(2) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) The drug is used in dogs between 4 weeks and 8 months of age for the removal of ascarids (*Toxocara canis*) and in animals over 4 weeks of age for the prevention of heartworm disease (*Dirofilaria immitis*).

(ii) The drug is administered (a) for removal of ascarids at a dosage of 50 milligrams per pound of body weight divided into two equal doses and administered 8 to 12 hours apart (morning and night), orally or mixed with either dry or wet food, and (b) for prevention

of heartworm disease at a dosage of 3 milligrams per pound of body weight daily, orally or in food, in heartworm endemic areas, from the beginning of mosquito activity, during the mosquito season, and for 2 months following the end thereof.

(iii) Dogs older than 8 months of age may be infected with *Dirofilaria immitis*. Use of the drug is contraindicated in dogs with active *D. immitis* infections.

(iv) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 41 FR 28265, July 9, 1976; 44 FR 3967, Jan. 19, 1979; 47 FR 14150, Apr. 2, 1982; 47 FR 35186, Aug. 13, 1982; 49 FR 33997, Aug. 28, 1984; 50 FR 41489, Oct. 11, 1985; 53 FR 47027, Oct. 18, 1988; 61 FR 34728, July 3, 1996; 62 FR 35076, June 30, 1997; 62 FR 38906, July 21, 1997]

§ 520.622c Diethylcarbamazine citrate chewable tablets.

(a) *Specifications.* Each chewable tablet contains 30, 45, 60, 120, 150, or 180 milligrams of diethylcarbamazine citrate.

(b) *Sponsors.* See drug listing nos. in § 510.600(c) of this chapter for identification of sponsors as follows:

(1) For 015579, use of 30 or 120 milligram tablets as in paragraph (c)(2)(i) of this section.

(2) For 000069, use of 60, 120, or 180 milligram tablets as in paragraph (c)(2)(ii) of this section.

(3) For 061690, use of 45 or 150 milligram tablets as in paragraph (c)(2)(iii) of this section.

(4) For 050604, use of 60-, 120-, or 180-milligram tablets as in paragraph (c)(2)(i) of this section.

(5) For 000061, use of 60-milligram tablets as in paragraph (c)(2)(i) of this section.

(6) For 000010, use of 30, 60, 120, or 180 milligram tablets as in paragraph (c)(2)(i) of this section.

(7) [Reserved]

(c) *Conditions of use*—(1) *Amount.* 3 milligrams per pound of body weight per day for prevention of heartworm disease and control of ascarids; 25 to 50 milligrams per pound of body weight as an aid in treatment of ascarid infections.

(2) *Indications for use.* (i) For prevention of heartworm disease (*Dirofilaria immitis*) in dogs; as an aid in control of

ascarids (*Toxocara canis*) in dogs; as an aid in treatment of ascarid (*Toxocara canis* and *Toxascaris leonina*) infections in dogs and cats.

(ii) For prevention of infection with *Dirofilaria immitis* (heartworm disease) in dogs; as an aid in treatment of ascarid (*Toxocara canis* and *Toxascaris leonina*) infections in dogs.

(iii) For prevention of heartworm disease (*Dirofilaria immitis*) in dogs.

(3) *Limitations.* Tablets are administered orally or pulverized and given in the feed. For treatment of ascarid infections, a repeat dose should be given in 10 to 20 days to remove immature worms which may enter the intestine from the lungs after the first dose. Dogs with established heartworm infections should not receive the drug until they have been converted to a negative status by the use of adulticidal and microfilaricidal drugs. Inadvertent administration to heartworm-infected dogs may cause adverse reactions due to pulmonary occlusion. Overdosage may cause emesis. For prevention of heartworm disease in heartworm-endemic areas, administration of the drug should start at the beginning of mosquito activity and be continued daily throughout the mosquito season and for approximately a month thereafter. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 6941, Feb. 17, 1978]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 520.622c, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 520.622d Diethylcarbamazine citrate capsules.

(a)(1) *Specifications.* Each capsule contains either 12.5, 50, 200, or 400 milligrams of diethylcarbamazine citrate.

(2) *Sponsor.* See 011014 in § 510.600(c) of this chapter.

(3) *Conditions of use*—(i) *Amount/indications for use.* 3 milligrams per pound of body weight daily for prevention of heartworm disease (*Dirofilaria immitis*) in dogs; 25 to 50 milligrams per pound of body weight in a single dose as an aid in the treatment of ascarid infections in dogs (*Toxocara canis* and *Toxascaris leonina*).

§ 520.623

(ii) *Limitations.* Administer orally directly or added to the daily ration. For ascarid infections, repeat treatment in 10 to 20 days to remove immature worms that may enter the intestine from the lungs after the first dose. Do not treat dogs with established heartworm infections until they have been converted to a negative status by the use of adulticidal and microfilaricidal drugs. Inadvertent administration to heartworm-infected dogs may cause adverse reactions due to pulmonary occlusion or shock. Overdosage may cause emesis. For prevention of heartworm disease in heartworm-endemic areas, administration of the drug should begin 1 month before and continue 2 months after the mosquito season. Dogs receiving prophylactic therapy should be examined every 6 months for the presence of microfilariae. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Specifications.* Each capsule contains either 12.5, 50, 200, or 400 milligrams of diethylcarbamazine citrate.

(2) *Sponsor.* See 023851 in § 510.600(c) of this chapter.

(3) *Conditions of use*—(i) *Amount/indications for use.* 3 milligrams per pound of body weight daily for prevention of heartworm disease (*Dirofilaria immitis*) in dogs.

(ii) *Limitations.* Capsules may be administered to the dog directly or added to the daily ration. For oral administration only. Do not treat dogs with established heartworm infections until they have been converted to a negative status by the use of adulticidal and microfilaricidal drugs. Inadvertent administration to heartworm infected dogs may cause adverse reactions due to pulmonary occlusion or shock. For prevention of heartworm disease in heartworm-endemic areas, administration of the drug should begin 1 month before and continue 2 months after the mosquito season. Dogs receiving prophylactic therapy should be examined every 6 months for the presence of microfilariae. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[47 FR 26377, June 18, 1982, as amended at 48 FR 46979, Oct. 17, 1983; 49 FR 5099, Feb. 10, 1984]

21 CFR Ch. I (4–1–99 Edition)

§ 520.623 Diethylcarbamazine citrate, oxibendazole chewable tablets.

(a) *Specifications.* Each tablet contains either 60, 120, or 180 milligrams of diethylcarbamazine citrate with 45, 91, or 136 milligrams of oxibendazole, respectively.

(b) *Sponsor.* See 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount.* Administer orally to dogs at a dosage level of 6.6 milligrams of diethylcarbamazine citrate per kilogram of body weight (3 milligrams per pound of body weight) and 5.0 milligrams of oxibendazole per kilogram of body weight (2.27 milligrams per pound of body weight).

(2) *Indications for use.* For prevention of infection with *Dirofilaria immitis* (heartworm disease) and *Ancylostoma caninum* (hookworm infection) and for removal and control of *Trichuris vulpis* (whipworm infection) and mature and immature stages of intestinal *Toxocara canis* (ascarid infection).

(3) *Limitations.* Orally administer daily during heartworm season. For free-choice feeding or broken and placed on or mixed with feed. Do not use in dogs that may harbor adult heartworms. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 28768, July 16, 1985, as amended at 53 FR 45759, Nov. 14, 1988; 54 FR 3776, Jan. 26, 1989; 54 FR 6804, Feb. 14, 1989; 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.645 Difloxacin.

(a) *Specifications.* Each tablet contains 11.4, 45.4, or 136 milligrams (mg) of difloxacin hydrochloride.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Dogs*—(i) *Amount.* 5 to 10 mg per kilogram (2.3 to 4.6 mg/pound) of body weight.

(ii) *Indications for use.* For management of diseases in dogs associated with bacteria susceptible to difloxacin.

(iii) *Limitations.* Use once a day for 2 to 3 days beyond cessation of clinical signs of disease up to a maximum of 30 days. Federal law prohibits the extra-label use of this drug in food-producing animals. Federal law restricts this

Food and Drug Administration, HHS

§ 520.763b

drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[63 FR 8123, Feb. 18, 1998]

§ 520.763 Dithiazanine iodide oral dosage forms.

§ 520.763a Dithiazanine iodide tablets.

(a) *Chemical name.* 3-Ethyl-2-[5-(3-ethyl-2-benzothiazolinyldene)-1,3-pentadienyl]-benzothiazolium iodide.

(b) *Specifications.* Dithiazanine iodide tablets contain 10 milligrams, 50 milligrams, 100 milligrams, or 200 milligrams of dithiazanine iodide in each tablet.

(c) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) The tablets are administered orally to dogs immediately after feeding using the following dosage schedule for various parasite infestations:

	Milligrams per pound of body weight	Length of treatment—days
Large roundworms (<i>Toxocara canis</i> , <i>Toxascaris leonina</i>)	10	3–5
Hookworms (<i>Ancylostoma caninum</i> , <i>Uncinaria stenocephala</i>)	10	7
Whipworms (<i>Trichuris vulpis</i>)	10	
Strongyloides (<i>Strongyloides canis</i> , <i>Strongyloides stercoralis</i>)	10	10–12
Heartworm microfilariae (<i>Dirofilaria immitis</i>)	3–5	7–10

Note: Treatment with dithiazanine iodide for heartworm microfilariae should follow 6 weeks after therapy for adult worms.

(2) The drug is contraindicated in animals sensitive to dithiazanine iodide and should be used cautiously, if at all, in dogs with reduced renal function.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(e) Use for treating dogs for large roundworms, hookworms, whipworms, and strongyloides as provided for in this section has been NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111

of this chapter, but may require bioequivalency and safety information.

[40 FR 13838, Mar. 27, 1975, as amended at 47 FR 51564, Nov. 16, 1982; 48 FR 32342, July 15, 1983; 53 FR 40727, Oct. 18, 1988; 62 FR 35076, June 30, 1997]

§ 520.763b Dithiazanine iodide powder.

(a) *Chemical name.* 3-Ethyl-2-[5-(3-ethyl-2-benzothiazolinyldene)-1,3-pentadienyl]-benzothiazolium iodide.

(b) *Specifications.* Dithiazanine iodide powder contains 200 milligrams of dithiazanine iodide per level standard tablespoon.

(c) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) Dithiazanine iodide powder is administered to dogs by mixing the proper dosage in the dog's food, using the following dosage schedule for various parasite infestations:

	Milligrams per pound of body weight	Length of treatment—days
Large roundworms (<i>Toxocara canis</i> , <i>Toxascaris leonina</i>)	10	3–5
Hookworms (<i>Ancylostoma caninum</i> , <i>Uncinaria stenocephala</i>)	10	7
Whipworms (<i>Trichuris vulpis</i>)	10	7
Strongyloides (<i>Strongyloides canis</i> , <i>Strongyloides stercoralis</i>)	10	10–12
Heartworm microfilariae (<i>Dirofilaria immitis</i>)	3–5	7–10

Note: Treatment with dithiazanine iodide for heartworm microfilariae should follow 6 weeks after therapy for adult worms.

(2) The drug is contraindicated in animals sensitive to dithiazanine iodide and should be used cautiously, if at all, in dogs with reduced renal function.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(e) Use for treating dogs for large roundworms, hookworms, whipworms, and strongyloides as provided for in this section has been NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111

§ 520.763c

21 CFR Ch. I (4–1–99 Edition)

of this chapter, but may require bioequivalency and safety information.

[40 FR 13838, Mar. 27, 1975, as amended at 47 FR 51564, Nov. 16, 1982; 48 FR 32342, July 15, 1983; 53 FR 40727, Oct. 18, 1988; 62 FR 35076, June 30, 1997]

§ 520.763c Dithiazanine iodide and piperazine citrate suspension.

(a) *Specifications.* Each milliliter of the drug contains 69 milligrams of dithiazanine iodide and 83 milligrams of piperazine base (as piperazine citrate).

(b) *Sponsor.* See 000010 in § 510.600(c) of this chapter.

(c) *NAS/NRC status.* The conditions of use are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use—(1) Amount.* 1 ounce (30 milliliters) per 100 pounds of body weight for the first 500 pounds; $\frac{3}{4}$ ounce for each 100 pounds thereafter, up to 1,200 pounds; $10\frac{1}{4}$ ounces to animals over 1,200 pounds.

(2) *Indications for use.* For control of large roundworms, *Parascaris equorum*; small strongyles; large strongyles, *Strongylus vulgaris*; and pinworms, *Oxyuris equi*.

(3) *Limitations.* Administer by drench or mixed with the daily ration as a single dose. Treatment is recommended in spring and fall. In a heavily infested environment, treatment may be repeated every 30 days. Not for use in horses intended for food purposes. Severely debilitated animals should not be wormed except on the advice of a veterinarian. If the drug is for administration by stomach tube, it shall be labeled: “Federal law restricts this drug to use by or on the order of a licensed veterinarian.”

[47 FR 52696, Nov. 23, 1982, as amended at 48 FR 32342, July 15, 1983; 53 FR 40727, Oct. 18, 1988; 62 FR 35076, June 30, 1997]

§ 520.784 Doxylamine succinate tablets.

(a) *Specifications.* The drug is in tablet form and contains doxylamine succinate as the active drug ingredient.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in conditions in which antihistaminic therapy may be expected to alleviate some signs of disease in horses, dogs, and cats.¹

(2) It is administered orally to horses at a dosage level of 1 to 2 milligrams per pound of body weight per day divided into 3 or 4 equal doses. It is administered orally to dogs and cats at a dosage level of 2 to 3 milligrams per pound of body weight per day divided into 3 or 4 equal doses.¹

(3) Not for use in horses intended for food.¹

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13838, Mar. 27, 1975, as amended at 42 FR 60140, Nov. 25, 1977; 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61624, Nov. 19, 1997]

§ 520.804 Enalapril tablets.

(a) *Specifications.* Each tablet contains either 1.0, 2.5, 5.0, 10.0, or 20.0 milligrams of enalapril maleate.

(b) *Sponsor.* See 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Dogs—(i) Amount.* 0.5 to 1.0 milligram of enalapril maleate per kilogram of body weight per day.

(ii) *Indications for use.* Treatment of mild, moderate, and severe (modified New York Heart Association Class II, III, IV) heart failure in dogs.

(iii) *Limitations.* Use 0.5 milligram per kilogram once daily. In the absence of adequate clinical response within a 2-week period, use may be increased to twice daily (a total of 1.0 milligram per kilogram). Enalapril maleate is administered as conjunctive therapy with furosemide and digoxin in the treatment of dilated cardiomyopathy and furosemide with or without digoxin in the treatment of chronic valvular disease. The safety of enalapril for use in breeding dogs has not been established. Use in pregnant bitches is not recommended. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter.

(2) [Reserved]

[59 FR 17694, Apr. 14, 1994, as amended at 62 FR 63270, Nov. 28, 1997]

§ 520.812 Enrofloxacin tablets.

(a) *Specifications.* Each tablet contains either 5.7, 22.7, or 68.0 milligrams of enrofloxacin.

(b) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Amount.* 5 to 20 milligrams per kilogram (2.27 to 9.07 milligrams per pound) of body weight.

(2) *Indications for use.* Dogs and cats for management of diseases associated with bacteria susceptible to enrofloxacin.

(3) *Limitations.* Administer orally as a single dose or divided into 2 equal doses at 12 hour intervals, daily. Administer for at least 2 to 3 days beyond cessation of clinical symptoms, for a maximum of 30 days. Safety in breeding or pregnant cats has not been established. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[54 FR 3444, Jan. 24, 1989, as amended at 55 FR 43327, Oct. 29, 1990; 62 FR 38906, July 21, 1997]

§ 520.813 Enrofloxacin oral solution.

(a) *Specifications.* Each milliliter of concentrate solution contains 32.3 milligrams of enrofloxacin.

(b) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.228 of this chapter.

(d) *Conditions of use.* It is used in drinking water as follows:

(1) *Chickens and turkeys*—(i) *Amount.* 25 to 50 parts per million of enrofloxacin in drinking water.

(ii) *Indications.* Chickens: Control of mortality associated with *Escherichia coli* susceptible to enrofloxacin. Turkeys: Control of mortality associated with *E. coli* and *Pasteurella multocida* (fowl cholera) susceptible to enrofloxacin.

(iii) *Limitations.* Do not use in laying hens producing eggs for human consumption. Administer medicated water continuously as sole source of drinking water for 3 to 7 days. Prepare fresh stock solution daily. Effects on the reproductive function of turkeys have

not been determined. Treated animals must not be slaughtered for food within 2 days of the last treatment. Individuals with a history of hypersensitivity to quinolones should avoid exposure to this product. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[61 FR 56893, Nov. 5, 1996]

§ 520.816 Epsiprantel tablets.

(a) *Specifications.* Each tablet contains either 12.5, 25, 50, or 100 milligrams of epsiprantel.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount.* 2.5 milligrams per pound of body weight.

(ii) *Indications for use.* Removal of canine cestodes *Dipylidium caninum* and *Taenia pisiformis*.

(2) *Cats*—(i) *Amount.* 1.25 milligrams per pound of body weight.

(ii) *Indications for use.* Removal of feline cestodes *D. caninum* and *T. taeniaeformis*.

(3) *Limitations.* For oral use only as a single dose. Do not use in animals less than 7 weeks of age. Safety of use in pregnant or breeding animals has not been established. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[54 FR 50615, Dec. 8, 1989, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.823 Erythromycin phosphate.

(a) *Specifications.* Erythromycin phosphate is the phosphate salt of the antibiotic substance produced by the growth of *Streptomyces erythreus* or the same antibiotic substance produced by any other means. One gram of erythromycin phosphate is equivalent to 0.89 gram of erythromycin master standard.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.230 of this chapter.

(d) *Conditions of use.* It is used in drinking water as follows:

(1) *Broiler and replacement chickens*—(i) *Amount.* 0.500 gram per gallon.

§ 520.863

21 CFR Ch. I (4–1–99 Edition)

(ii) *Indications for use.* As an aid in the control of chronic respiratory disease due to *Mycoplasma gallisepticum* susceptible to erythromycin.

(iii) *Limitations.* Administer for 5 days; do not use in replacement pullets over 16 weeks of age; do not use in chickens producing eggs for human consumption; to assure effectiveness, treated birds must consume enough medicated water to provide a therapeutic dosage; solutions older than 3 days should not be used; withdraw 1 day before slaughter.

(2) *Replacement chickens and chicken breeders*—(i) *Amount.* 0.500 gram per gallon.

(ii) *Indications for use.* As an aid in the control of infectious coryza due to *Hemophilus gallinarum* susceptible to erythromycin.

(iii) *Limitations.* Administer for 7 days; do not use in replacement pullets over 16 weeks of age; do not use in chickens producing eggs for human consumption; to assure effectiveness, treated birds must consume enough medicated water to provide a therapeutic dosage; solutions older than 3 days should not be used; withdraw 1 day before slaughter.

(3) *Growing turkeys*—(i) *Amount.* 0.500 gram per gallon.

(ii) *Indications for use.* As an aid in the control of blue comb (nonspecific infectious enteritis) caused by organisms susceptible to erythromycin.

(iii) *Limitations.* Administer for 7 days; do not use in turkeys producing eggs for human consumption; to assure effectiveness, treated birds must consume enough medicated water to provide a therapeutic dosage; solutions older than 3 days should not be used; withdraw 1 day before slaughter.

[40 FR 13838, Mar. 27, 1975, as amended at 45 FR 56798, Aug. 26, 1980]

§ 520.863 Ethylisobutrazine hydrochloride tablets.

(a) *Specifications.* Each tablet contains either 10 milligrams or 50 milligrams of ethylisobutrazine hydrochloride.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is administered orally to dogs as a tranquilizer.¹

(2) It is administered once daily at a dosage level of 2 to 5 milligrams of ethylisobutrazine hydrochloride per pound of body weight.¹

(3) It is not to be used in conjunction with organophosphates and/or procaine hydrochloride because phenothiazine may potentiate the toxicity of organophosphates and the activity of procaine hydrochloride.¹

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13838, Mar. 27, 1975, as amended at 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61624, Nov. 19, 1997]

§ 520.870 Etodolac.

(a) *Specifications.* Each tablet contains 150 or 300 milligrams (mg) of etodolac.

(b) *Sponsor.* See 053501 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Dogs*—(i) *Amount.* 10 to 15 mg per kilogram (4.5 to 6.8 mg/pound) of body weight per day.

(ii) *Indications for use.* For the management of pain and inflammation associated with osteoarthritis in dogs.

(iii) *Limitations.* Use once-a-day. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[63 FR 51300, Sept. 25, 1998]

§ 520.903 Febantel oral dosage forms.

§ 520.903a Febantel paste.

(a) *Chemical name.* Dimethyl [[2-[(methoxyacetyl)amino]-4-(phenylthio)phenyl] carbonimidoyl]bis [carbamate].

(b) *Specifications.* The drug is a paste containing 45.5 percent febantel.

(c) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use*—(1) *Amount*. Six milligrams per kilogram (2.73 milligrams per pound) of body weight in horses.

(2) *Indications for use*. For removal of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); ascarids (*Parascaris equorum*—sexually mature and immature); pinworms (*Oxyuris equi*—adult and 4th stage larva); and the various small strongyles in horses, foals, and ponies.

(3) *Limitations*. (i) The paste may be administered on the base of the tongue or well mixed into a portion of the normal grain ration.

(ii) [Reserved]

(iii) For animals maintained on premises where reinfection is likely to occur, retreatment may be necessary. For most effective results, retreat in 6 to 8 weeks.

(iv) Not for use in horses intended for food.

(v) Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[43 FR 8797, Mar. 3, 1978; 43 FR 12311, Mar. 24, 1978, as amended at 43 FR 60882, Dec. 29, 1978. Redesignated at 45 FR 8587, Feb. 8, 1980]

§ 520.903b Febantel suspension.

(a) *Specifications*. The suspension contains 9.3 percent (2.75 grams per ounce) febantel.

(b) *Sponsor*. See 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 3 milliliters per 100 pounds body weight or 1 fluid ounce per 1000 pounds (6 milligrams per kilogram body weight).

(2) *Indications for use*. For removal of ascarids (*Parascaris equorum*—adult and sexually immature), pinworms (*Oxyuris equi*—adult and 4th stage larvae), large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*), and the various small strongyles in horses, breeding stallions and mares, pregnant mares, foals, and ponies.

(3) *Limitations*. Administer by stomach tube or drench, or by mixing well into a portion of the normal grain ration. For animals maintained on premises where reinfection is likely to occur, retreatment may be necessary. For most effective results, retreat in 6 to 8 weeks. Not for use in horses intended for food. Federal law restricts

this drug to use by or on the order of a licensed veterinarian.

(d) *Special considerations*. Febantel suspension may be used in combination with trichlorfon oral liquid in accordance with the provisions of § 520.2520c, this section, and the following conditions:

(1) Combine 1 part febantel suspension with 5 parts trichlorfon liquid.

(2) Allow animal to consume a portion of daily grain ration; administer mixture by stomach tube at rate of 18 milliliters per 100 pounds of body weight.

[45 FR 8587, Feb. 8, 1980]

§ 520.903c [Reserved]

§ 520.903d Febantel-praziquantel paste.

(a) *Specifications*. Each gram of paste contains 34 milligrams of febantel and 3.4 milligrams of praziquantel.

(b) *Sponsor*. See No. 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*—(i) Dogs and cats (over 6 months of age): 10 milligrams of febantel and 1 milligram of praziquantel per kilogram of body weight (1 gram of paste per 7.5 pounds body weight) administered by mouth or in the food once daily for 3 days.

(ii) Puppies and kittens (less than 6 months of age): 15 milligrams of febantel and 1.5 milligrams of praziquantel per kilogram of body weight (1 gram of paste per 5 pounds body weight) administered by mouth on a full stomach once daily for 3 days.

(2) *Indications for use*. (i) Dogs and puppies: For removal of hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*), whipworms (*Trichuris vulpis*), ascarids (*Toxocara canis* and *Toxascaris leonina*), and tapeworms (*Dipylidium caninum* and *Taenia pisiformis*).

(ii) Cats and kittens: For removal of hookworms (*Ancylostoma tubaeforme*), ascarids (*Toxocara cati*) and tapeworms (*Dipylidium caninum* and *Taenia taeniaeformis*).

(3) *Limitations*. Do not use in pregnant animals. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(4) *Special considerations*. Consider alternative therapy or use with caution

§ 520.903e

in animals with pre-existing liver or kidney dysfunction.

[50 FR 19167, May 7, 1985, as amended at 53 FR 48533, Dec. 1, 1988; 56 FR 50813, Oct. 9, 1991]

§ 520.903e Febantel tablets.

(a) *Specifications.* Each scored tablet contains 27.2 milligrams of febantel for use in dogs, puppies, cats, and kittens or 163.3 milligrams of febantel for use in dogs, puppies, and cats.

(b) *Sponsor.* See 000859 in § 510.600(c)(2) of this chapter.

(c) *Conditions of use*—(1) *Amount*—(i) *Dogs and cats.* Ten milligrams per kilogram body weight. Administer once daily for 3 consecutive days.

(ii) *Puppies and kittens fewer than 6 months of age.* Fifteen milligrams per kilogram body weight. Administer once daily for 3 consecutive days.

(2) *Indications for use.* (i) For removal of hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*), ascarids (*Toxocara canis* and *Toxascaris leonina*) and whipworms (*Trichuris vulpis*) in dogs and puppies.

(ii) For removal of hookworms (*Ancylostoma tubaeforme*) and ascarids (*Toxocara cati*) in cats and kittens.

(3) *Limitations.* Do not use in pregnant animals. Consider alternative therapy or use with caution in animals with preexisting liver or kidney dysfunction. Administer to puppies and kittens on a full stomach. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[56 FR 50655, Oct. 8, 1991]

§ 520.905 Fenbendazole oral dosage forms.

§ 520.905a Fenbendazole suspension.

(a) *Specifications.* The drug is a suspension containing 10 percent (100 milligrams per milliliter) fenbendazole.

(b) *Sponsor.* See No. 012799 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.275 of this chapter.

(d) *Conditions of use*—(1) *Horses*—(i) *Amount.* 5 milligrams per kilogram (2.3 milligrams per pound) for the control of large strongyles, small strongyles, and pinworms; 10 milligrams per kilogram for the control of ascarids.

21 CFR Ch. I (4–1–99 Edition)

(ii) *Indications for use.* For the control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), small strongyles (*Cyathostomum* spp., *Cylicocyclus* spp., *Cylicostephanus* spp., *Triodontophorus* spp.), pinworms (*Oxyuris equi*), and ascarids (*Parascaris equorum*) in horses.

(iii) *Limitations.* Administer orally by dose syringe or suitable plastic syringe. Do not use in horses intended for food. Consult a veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) *Cattle including dairy cows of breeding age*—(i) *Amount.* Administer orally 5 milligrams per kilogram of body weight (2.3 milligrams per pound).

(ii) *Indications for use.* For the removal and control of lungworm (*Dictyocaulus viviparus*); stomach worm (adults)—brown stomach worm (*Ostertagia ostertagi*); stomach worms (adults and 4th-stage larvae)—barberpole worm (*Haemonchus contortus* and *H. placei*) and small stomach worm (*Trichostrongylus axei*); intestinal worms (adults and 4th-stage larvae)—hookworm (*Bunostomum phlebotomum*), threadnecked intestinal worm (*Nematodirus helvetianus*), small intestinal worm (*Cooperia punctata* and *C. oncophora*), bankrupt worm (*Trichostrongylus colubriformis*), and nodular worm (*Oesophagostomum radiatum*).

(iii) *Limitations.* Retreatment may be needed after 4 to 6 weeks. Cattle must not be slaughtered within 8 days following last treatment. Consult a veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(3) *Beef cattle*—(i) *Amount.* Administer orally 10 milligrams per kilogram of body weight.

(ii) For the removal and control of stomach worm (4th-stage inhibited larvae/type II ostertagiasis), *Ostertagia ostertagi*, and tapeworm, *Moniezia benedeni*.

(iii) *Limitations.* Retreatment may be needed after 4 to 6 weeks. Cattle must not be slaughtered within 8 days following last treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(4) *Goats*—(i) *Amount*. Administer orally 5 milligrams per kilogram of body weight (2.3 milligrams per pound).

(ii) *Indications for use*. For the removal and control of stomach and intestinal worms *Haemonchus contortus* and *Ostertagia circumcincta*.

(iii) *Limitations*. Retreatment may be needed after 4 to 6 weeks. Goats must not be slaughtered for food within 6 days following last treatment. Do not use in lactating goats. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(e) *Special considerations*. Fenbendazole suspension 10 percent and approved forms of trichlorfon, when used concomitantly for treating the indications provided in paragraph (d) of this section and for treating infections of stomach bot as provided in § 520.2520, have been shown to be compatible and not to interfere with one another.

[42 FR 59069, Nov. 15, 1977; 43 FR 12311, Mar. 24, 1978. Redesignated at 44 FR 1375, Jan. 5, 1979, and amended at 46 FR 29464, June 2, 1981; 47 FR 15327, Apr. 9, 1982; 48 FR 42809, Sept. 20, 1983; 49 FR 1983, Jan. 17, 1984; 53 FR 40058, Oct. 13, 1988; 59 FR 26943, May 25, 1994; 61 FR 29478, June 11, 1996; 63 FR 63983, Nov. 18, 1998]

§ 520.905b Fenbendazole granules.

(a) *Specifications*. The drug is in granular form containing 22 percent (222 milligrams per gram) fenbendazole.

(b) *Sponsor*. See No. 012799 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Horses*—(i) *Amount*. 5 milligrams per kilogram (2.3 milligrams per pound) for the control of large strongyles, small strongyles, and pinworms; 10 milligrams per kilogram for the control of ascarids.

(ii) *Indications for use*. For the control of infections of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), small strongyles, pinworms (*Oxyuris equi*), and ascarids (*Parascaris equorum*).

(iii) *Limitations*. Sprinkle the appropriate amount of drug on a small amount of the usual grain ration. Prepare for each horse individually. Withholding feed or water is not necessary. Retreat in 6 to 8 weeks if required. Do not use in horses intended for food. Consult your veterinarian for assist-

ance in the diagnosis, treatment, and control of parasitism.

(2) *Dogs*—(i) *Amount*. 50 milligrams per kilogram (22.7 milligrams per pound) daily for 3 consecutive days.

(ii) *Indications for use*. For the removal of ascarids (*Toxocara canis*, *Toxascaris leonina*), hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*), whipworms (*Trichuris vulpis*), and tapeworms (*Taenia pisiformis*).

(iii) *Limitations*. Mix the appropriate amount of drug with a small amount of the usual food; dry dog food may require slight moistening to facilitate mixing. Medicated food must be fully consumed. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(3) *Zoo and wildlife animals*—(i) *Amount*. Ten milligrams per kilogram per day for 3 days.

(ii) *Indications for use*. For control of internal parasites of *Felidae* and *Ursidae* as follows:

(A) Lion (*Panthera leo*) and Tiger (*Panthera tigris*): Ascarid (*Toxocara cati*, *Toxascaris leonina*), Hookworm (*Ancylostoma* spp.).

(B) Cheetah (*Acinonyx jubatus*): Ascarid (*Toxocara cati*, *Toxascaris leonina*).

(C) Puma (*Felis concolor*), Panther (*Panthera* spp.), Leopard (*Panthera pardus*), Jaguar (*Panthera onca*): Ascarid (*Toxocara cati*, *Toxascaris leonina*), Hookworm (*Ancylostoma* spp.), Tapeworm (*Taenia hydatigena*, *T. krabbei*, *T. taeniaeformis*).

(D) Black Bear (*Ursus americanus*): Ascarid (*Baylisascaris transfuga*, *Toxascaris leonina*), Hookworm (*Ancylostoma caninum*), Tapeworm (*Taenia hydatigena*, *T. krabbei*).

(E) Polar Bear (*Ursus maritimus*) and Grizzly Bear (*Ursus horribilis*): Ascarid (*Baylisascaris transfuga*, *Toxascaris leonina*).

(iii) *Limitations*. Top dress or mix with a small portion of food. Must be fully consumed prior to feeding. Federal law restricts this drug to use by or on the order of a licensed veterinarian. Do not use 14 days before or during the hunting season.

[44 FR 1375, Jan. 5, 1979, as amended at 47 FR 15327, Apr. 9, 1982; 48 FR 50528, Nov. 2, 1983; 59 FR 35252, July 11, 1994]

§ 520.905c

21 CFR Ch. I (4–1–99 Edition)

§ 520.905c Fenbendazole paste.

(a) *Specifications.* The product is an aqueous paste containing 10 percent fenbendazole.

(b) *Sponsor.* See No. 012799 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.275 of this chapter.

(d) *Conditions of use—(1) Horses—(i)(a) Amount.* 2.3 milligrams per pound of body weight (one 2.5-gram fenbendazole syringe for a 1,100-pound horse). For foals and weanlings (less than 18 months of age), 4.6 milligrams per pound of body weight (one 2.5-gram fenbendazole syringe for each 550 pounds of body weight).

(b) *Indications for use.* For control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), small strongyles, pinworms (*Oxyuris equi*), and ascarids (*Parascaris equorum*) in horses.

(c) *Limitations.* Retreatment at intervals of 6 to 8 weeks may be required. Do not use in horses intended for food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(ii)(a) *Amount.* 4.6 milligrams per pound of body weight (one 2.5-gram fenbendazole syringe for a 550-pound horse) daily for 5 days.

(b) *Indications for use.* For control of arteritis caused by the fourth stage larvae of *Strongylus vulgaris*.

(c) *Limitations.* Treatment should be initiated in the spring and repeated in 6 months. Do not use in horses intended for food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of fourth stage larvae of *S. vulgaris*.

(iii)(a) *Amount.* 4.6 milligrams per pound of body weight (10 milligrams per kilogram) daily for 5 consecutive days.

(b) *Indications for use.* For treatment of encysted mucosal cyathostome (small strongyle) larvae including early third stage (hypobiotic), late third stage, and fourth stage larvae in horses.

(c) *Limitations.* (Consult your veterinarian for assistance in the diagnosis, treatment, and control of encysted mucosal cyathostomes). Do not use in horses intended for food.

(2) *Beef and dairy cattle—(i) Amount.* Administer orally 5 milligrams per kilogram of body weight (2.3 milligrams per pound).

(ii) *Indications for use.* For the removal and control of lungworm (*Dictyocaulus viviparus*), barberpole worm (*Haemonchus contortus*), brown stomach worm (*Ostertagia ostertagi*), small stomach worm (*Trichostrongylus axei*), hookworm (*Bunostomum phlebotomum*), thread-necked intestinal worm (*Nematodirus helvetianus*), small intestinal worms (*Cooperia punctata* and *C. oncophora*), bankrupt worm (*Trichostrongylus colubriformis*), and nodular worm (*Oesophagostomum radiatum*).

(iii) *Limitations.* Re-treatment may be needed after 4 to 6 weeks. Cattle must not be slaughtered within 8 days following last treatment. Consult a veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(e) *Special considerations.* Fenbendazole paste 10 percent may be used concomitantly with approved forms of trichlorfon for the indications provided in paragraph (d)(1)(i) of this section and for treating infections of stomach bots as provided in § 520.2520.

[46 FR 32018, June 19, 1981, as amended at 47 FR 15327, Apr. 9, 1982; 49 FR 8433, Mar. 7, 1984; 50 FR 26358, June 26, 1985; 61 FR 29478, June 11, 1996; 63 FR 31624, June 10, 1998]

§ 520.905d Fenbendazole powder.

(a) *Specifications.* (1) Each 2-ounce packet contains 2.27 grams (4 percent) of fenbendazole plus other inert ingredients.

(2) Each 4-ounce packet contains 1.7 grams (1.5 percent) of fenbendazole plus other inert ingredients.

(b) *Sponsors.* (1) See No. 012799 in § 510.600(c) of this chapter for use of the 4-percent product.

(2) See No. 017800 in § 510.600(c) of this chapter for use of the 1.5-percent product.

(c) *Related tolerances.* See § 556.275 of this chapter.

(d) *Conditions of use.* It is administered to swine as follows:

(1) *Amount.* 3 milligrams fenbendazole per kilogram body weight per day (1.36 milligrams per pound per day).

(2) *Indications for use.* For removal and control of large roundworms (*Ascaris suum*); lungworms (*Metastrongylus apri*); nodular worms (*Oesophagostomum dentatum*, *O. quadrispinulatum*); small stomach worms (*Hyostrongylus rubidus*); whipworms (*Trichuris suis*); and kidneyworms (*Stephanurus dentatus*—mature and immature).

(3) *Limitations.* Thoroughly mix the contents of the packet(s) with swine ration and administer according to label directions. Feed as sole ration for 3 consecutive days. Can be fed to pregnant sows. No prior withdrawal of feed or water is necessary. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[49 FR 18090, Apr. 27, 1984, as amended at 49 FR 20485, May 15, 1984]

§ 520.905e Fenbendazole blocks.

(a) *Specifications.* (1) Each pound of molasses block contains 750 milligrams of fenbendazole.

(2) Each pound of protein block contains 750 milligrams of fenbendazole.

(b) *Sponsor.* See 012799 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.275 of this chapter.

(d) *Conditions of use*—(1) *Amount.* 0.1 pound of block per 100 pounds of body weight per day for 3 days. Total dose for the 3-day period is 2.27 milligrams of fenbendazole per pound of body weight for mature cattle.

(2) *Indications for use.* For removal and control of infections of lungworms (*Dictyocaulus viviparus*) and gastrointestinal roundworms (*Haemonchus contortus*, *Ostertagia ostertagi*, *Trichostrongylus axei*, *Bunostomum phlebotomum*, *Nematodirus helvetianus*, *Cooperia oncophora* and *C. punctata*, *Trichostrongylus colubriformis*, and *Oesophagostomum radiatum*) in beef cattle.

(3) *Limitations.* Administer free choice of beef cattle on pasture that have become accustomed to nonmedicated block feeding during an adaptation period of 12 to 19 days. Molasses block: Cattle must not be slaughtered within 11 days following last treatment. Protein block: Cattle must not be slaugh-

tered within 16 days following last treatment; do not use in dairy cattle of breeding age. Animals maintained under conditions of constant worm exposure may require retreatment within 6 to 8 weeks. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[51 FR 41783, Nov. 19, 1986, as amended at 54 FR 20787, May 15, 1989]

§ 520.960 Flumethasone tablets.

(a) *Specifications.* Each tablet contains 0.0625 milligram of flumethasone.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* (i) *Dogs:* Administer orally from 0.0625 to 0.25 milligram daily in divided doses.

(ii) *Cats:* Administer orally from 0.03125 to 0.125 milligram daily in divided doses.

(2) *Indications for use.* (i) *Dogs:* It is used for musculoskeletal conditions due to inflammation of muscles or joints and accessory structures, where permanent structural changes do not exist, such as arthritis, the disc syndrome, and myositis.

(ii) *Dogs and cats:* It is used in certain acute and chronic dermatoses of varying etiology to help control the pruritus, irritation, and inflammation associated with these conditions.

(3) *Limitations.* Do not use in viral infections. Anti-inflammatory action of corticosteroids may mask signs of infection. Do not use in animals with tuberculosis, chronic nephritis, cushingoid syndrome, or where peptic ulcers occur, except for emergency therapy. Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 7131, Feb. 6, 1979, as amended at 61 FR 5506, Feb. 13, 1996]

§ 520.970

§ 520.970 Flunixin oral dosage forms.

§ 520.970a Flunixin meglumine granules.

(a) *Specifications.* Each 10-gram packet contains flunixin meglumine equivalent to 250 milligrams of flunixin.

(b) *Sponsor.* No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 0.5 milligram of flunixin per pound of body weight (one packet per 500 pounds) per day.

(2) *Indications for use.* For alleviation of inflammation and pain associated with musculoskeletal disorders in the horse.

(3) *Limitations.* Administer daily dose for up to 5 days by sprinkling on small amount of feed. The effect of this drug on pregnancy has not been determined. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 36381, June 22, 1979. Redesignated at 50 FR 38114, Sept. 20, 1985, and amended at 52 FR 7832, Mar. 13, 1987]

§ 520.970b Flunixin meglumine paste.

(a) *Specifications.* Each 30-gram syringe contains flunixin meglumine equivalent to 1,500 milligrams of flunixin.

(b) *Sponsor.* No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use. Horses*—(1) *Amount.* 0.5 milligram of flunixin per pound of body weight daily.

(2) *Indications for use.* For alleviation of inflammation and pain associated with musculoskeletal disorders.

(3) *Limitations.* For oral use only. Treatment should not exceed 5 consecutive days. The effect of this drug on pregnancy has not been determined. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 38114, Sept. 20, 1985, as amended at 52 FR 7832, Mar. 13, 1987]

21 CFR Ch. I (4–1–99 Edition)

§ 520.1010 Furosemide oral dosage forms.

§ 520.1010a Furosemide tablets or boluses.

(a) *Specifications.* Each tablet contains 12.5 or 50 milligrams of furosemide. Each bolus contains 2 grams of furosemide.

(b) *Sponsor.* See No. 012799 in § 510.600(c) of this chapter for conditions of use provided for in paragraphs (c) (1) and (2) of this section; see No. 000010 in § 510.600(c) of this chapter for use in dogs as provided for in paragraph (c)(1) of this section; see No. 000093 in § 510.600(c) of this chapter for use of a 12.5- and 50-milligram tablet for conditions of use provided for in paragraph (c)(3) of this section.

(c) *Conditions of use.* It is used as follows:

(1) *Dogs and cats*—(i) *Amount.* 1 to 2 milligrams per pound of body weight, once or twice daily, with a 6- to 8-hour interval between successive daily doses.

(ii) *Indications for use.* It is used for the treatment of edema (pulmonary congestion, ascites) associated with cardiac insufficiency and acute non-inflammatory tissue edema.

(iii) *Limitations.* The dosage should be adjusted to the animal's response. In severe edematous or refractory cases, the dosage may be doubled or increased by increments of 1 milligram per pound of body weight to establish the effective dose. This dose should be administered once or twice daily on an intermittent schedule. Diuretic therapy should be discontinued after reduction of edema, or when necessary, maintained after determining a programmed dosage schedule to prevent recurrence. The drug, if given in excessive amounts or over extended periods of time may result in dehydration and/or electrolyte imbalance. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cattle*—(i) *Amount.* 1 to 2 milligrams per pound of body weight, or one 2-gram bolus per animal, per day.

(ii) *Indications for use.* The drug is used for the treatment of physiological parturient edema of the mammary gland and associated structures.

(iii) *Limitations.* Treatment not to exceed 48 hours post-parturition. For oral use only. When treatment is initiated with an injectable, it is followed by a 12-hour interval, and maintained by oral administration. Milk taken during treatment and for 48 hours (four milkings) after the last treatment must not be used for food. Cattle must not be slaughtered for food within 48 hours following last treatment. The drug, if given in excessive amounts or over extended periods of time, may result in dehydration and/or electrolyte imbalance. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(3) *Dogs*—(i) *Amount.* 1 to 2 milligrams per pound of body weight, once or twice daily, with a 6- to 8-hour interval between successive daily doses.

(ii) *Indications for use.* It is used for the treatment of edema (pulmonary congestion, ascites) associated with cardiac insufficiency.

(iii) *Limitations.* The dosage should be adjusted to the animal's response. In severe cases, the dosage may be doubled or increased by increments of 1 milligram per pound of body weight to establish the effective dose. This dose should be administered once or twice daily on an intermittent schedule. Diuretic therapy should be discontinued after reduction of edema, or when necessary, maintained after determining a programmed dosage schedule to prevent recurrence. The drug, if given in excessive amounts or over extended periods of time, may result in dehydration and/or electrolyte imbalance. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 52446, Nov. 30, 1976. Redesignated at 43 FR 14647, Apr. 7, 1978, and amended at 48 FR 40517, Sept. 8, 1983; 48 FR 49841, Oct. 28, 1983; 53 FR 40727, Oct. 18, 1988; 56 FR 37473, Aug. 7, 1991; 62 FR 6724, Feb. 13, 1997; 62 FR 35076, June 30, 1997]

§ 520.1010b Furosemide powder.

(a) *Specifications.* Furosemide powder is packaged in packets containing 2 grams of furosemide plus other inert ingredients.

(b) *Sponsor.* See No. 012799 in § 510.600(c) of this chapter.

(c) *Conditions of use.* It is administered to dairy cattle alone, as a "top

dressing" upon a small amount of feed or as a drench.

(1) *Amount.* 1 to 2 milligrams per pound of body weight but not to exceed one packet per animal, per day.

(2) *Indications for use.* The drug is used for the treatment of physiological parturient edema of the mammary gland and associated structures.

(3) *Limitations.* Treatment not to exceed 48 hours post-parturition. For oral use only. The individual dose is one packet administered once daily; when treatment is initiated with an injectable, at least a 12-hour interval must follow before oral administration. Milk taken during treatment and for 48 hours (four milkings) after the last treatment must not be used for food. Cattle must not be slaughtered for food within 48 hours following last treatment. The drug, if given in excessive amounts or over extended periods of time, may result in dehydration and electrolyte imbalance. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 14647, Apr. 7, 1978, as amended at 47 FR 15327, Apr. 9, 1982]

§ 520.1010c Furosemide syrup.

(a) *Specifications.* Each milliliter of syrup contains 10 milligrams of furosemide.

(b) *Sponsor.* See No. 012799 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 1 to 2 milliliters orally (10 to 20 milligrams) per 10 pounds of body weight (approximately 2.5 to 5 milligrams per kilogram), once or twice daily, with a 6- to 8-hour interval between successive daily doses.

(2) *Indications for use.* It is used in dogs for the treatment of edema (pulmonary congestion, ascites) associated with cardiac insufficiency and acute noninflammatory tissue edema.

(3) *Limitations.* The dosage should be adjusted to the animal's response. In severe edematous or refractory cases, the dosage may be doubled or increased by increments of 1 milligram per pound of body weight to establish the effective dose. This dose should be administered once or twice daily on an intermittent schedule. Diuretic therapy should be discontinued after reduction

§ 520.1044

of edema or, when necessary, maintained after determining a programmed dosage schedule to prevent recurrence. If given in excessive amounts or over extended periods of time, the drug may result in dehydration and/or electrolyte imbalance. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 11177, Mar. 17, 1978, as amended at 47 FR 15327, Apr. 9, 1982]

§ 520.1044 Gentamicin sulfate oral dosage forms.

§ 520.1044a Gentamicin sulfate oral solution.

(a) *Specifications.* Each milliliter of aqueous solution contains gentamicin sulfate equivalent to 50 milligrams of gentamicin.

(b) *Sponsor.* See Nos. 000061 and 051259 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.300 of this chapter.

(d) *Conditions of use*—(1) *Amount.* Colibacillosis: 1 milliliter per 2 gallons of drinking water for 3 consecutive days, to provide 0.5 milligram/pound/day; swine dysentery: 1 milliliter per 1 gallon of drinking water for 3 consecutive days, to provide 1.0 milligram/pound/day.

(2) *Indications for use.* In weanling swine for control and treatment of colibacillosis caused by strains of *E. coli* sensitive to gentamicin, and in swine for control and treatment of swine dysentery associated with *Treponema hyodysenteriae*.

(3) *Limitations.* For use in swine drinking water only. Do not store or offer medicated drinking water in rusty containers since the drug is quickly destroyed in such containers. Medicated drinking water should be prepared daily and be the sole source of drinking water for 3 consecutive days. Treatment may be repeated if dysentery recurs. Do not slaughter treated swine for food for at least 3 days following treatment.

[48 FR 10302, Mar. 11, 1983. Redesignated at 49 FR 572, Jan. 5, 1984, and amended at 49 FR 14332, Apr. 11, 1984; 52 FR 7832, Mar. 13, 1987; 62 FR 34169, June 25, 1997]

21 CFR Ch. I (4–1–99 Edition)

§ 520.1044b Gentamicin sulfate pig pump oral solution.

(a) *Specifications.* Each milliliter of pig pump oral solution contains gentamicin sulfate equivalent to 4.35 milligrams of gentamicin.

(b) *Sponsor.* See Nos. 000061 and 059130 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.300 of this chapter.

(d) *Conditions of use*—(1) *Amount.* Administer 1.15 milliliters of pig pump oral solution (5 milligrams of gentamicin) orally per pig one time.

(2) *Indications for use.* In neonatal swine 1 to 3 days of age for control and treatment of colibacillosis caused by strains of *E. coli* sensitive to gentamicin.

(3) *Limitations.* For use in neonatal swine only. Do not slaughter treated swine for food for at least 14 days following treatment.

[49 FR 572, Jan. 5, 1984, as amended at 52 FR 7832, Mar. 13, 1987; 62 FR 29011, May 29, 1997]

§ 520.1044c Gentamicin sulfate soluble powder.

(a) *Specifications.* Each gram of gentamicin sulfate soluble powder contains gentamicin sulfate equivalent to 16.7, 66.7, or 333.3 milligrams of gentamicin.

(b) *Sponsor.* See Nos. 000061 and 057561 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.300 of this chapter.

(d) *Conditions of use*—(1) *Amount.* Colibacillosis: gentamicin sulfate equivalent to 25 milligrams of gentamicin per gallon of drinking water for 3 consecutive days, to provide 0.5 milligram per pound of body weight per day; swine dysentery: gentamicin sulfate equivalent to 50 milligrams of gentamicin per gallon of drinking water for 3 consecutive days, to provide 1 milligram per pound of body weight per day.

(2) *Indications for use.* In weanling swine for control and treatment of colibacillosis caused by strains of *E. coli* sensitive to gentamicin, and in swine for control and treatment of swine dysentery associated with *Treponema hyodysenteriae*.

(3) *Limitations.* For use in swine drinking water only. Do not store or offer medicated drinking water in

Food and Drug Administration, HHS

§ 520.1120a

rusty containers since the drug is quickly destroyed in such containers. Medicated drinking water should be prepared daily and be the sole source of drinking water for 3 consecutive days. Treatment may be repeated if dysentery recurs. Do not slaughter treated swine for food for at least 10 days following treatment.

[49 FR 29778, July 24, 1984, as amended at 52 FR 7832, Mar. 13, 1987; 52 FR 48675, Dec. 24, 1987; 62 FR 29013, May 29, 1997]

§ 520.1100 Griseofulvin.

(a) *Chemical name.* 7-Chloro-2',4,6-trimethoxy-6'-methylspiro [benzofuran-2(3H), 1'-[2]-cyclohexene]-3,4'-dione.

(b) *Specifications.* Complies with U.S.P. for griseofulvin microsize.

(c) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) As a soluble powder for horses, it is administered as a drench or as a top dressing on feed. It is used for equine ringworm infection caused by *Trichophyton equinum* or *Microsporum gypseum*. Administer for not less than 10 days a daily dose as follows: Adults, 2.5 grams; yearlings, 1.25 to 2.5 grams; and foals, 1.25 grams. Not for use in horses intended for food. For use only by or on the order of a licensed veterinarian.

(2)(i) Boluses containing 2.5 grams of griseofulvin are used in horses for treating ringworm infection caused by *Trichophyton equinum*. It is administered to adult horses at a level of one bolus per day, to yearlings at one-half to one bolus per day, and to foals at one-half bolus per day. All three dosage levels should be administered for a period of not less than 10 days. In responsive cases, treatment should be continued until all infected areas are proven negative by appropriate culture. Not for use in horses intended for food.

(ii) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(3) Dogs and cats: (i) *Amount.* 125- and 500-milligram tablets administered orally as follows:

(a) Daily (single or divided) dose:

Body weight (pounds)	Dosage (milligrams)
Up to 6	62.5
6 to 18	125

Body weight (pounds)	Dosage (milligrams)
18 to 36	250
36 to 48	375
48 to 75	500

(b) Weekly (single) dose: If experience indicates that treatment is more effective for the drug given in large doses, administer at intervals of 7 to 10 days, a dose equal to 10 milligrams/pound of body weight x body weight x number of days between treatments. Dosage should be adjusted according to response. Administer additional dose after the animal is free of infection.

(ii) *Indications for use.* For treatment of fungal infections of the skin, hair, and claws caused by *Trichophyton mentagrophytes*, *T. rubrum*, *T. schoenleini*, *T. sulphurem*, *T. verrucosum*, *T. interdigitale*, *Epidermophyton floccosum*, *Microsporum gypseum*, *M. canis*, *M. audouini*.

(iii) *Limitations.* For satisfactory diagnosis, a microscopic tissue examination or culture is recommended prior to treatment. Treatment should be continued for 3 to 4 weeks in skin and hair infections, and up to 4 months for infections involving nails or claws. Clipping of hair, nails, and claws to help remove any remaining viable fungi is indicated. Safety for use of griseofulvin for pregnant animals has not been established. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 41 FR 42948, Sept. 29, 1976; 43 FR 28458, June 30, 1978; 52 FR 7832, Mar. 13, 1987; 54 FR 30205, July 19, 1989]

§ 520.1120 Haloxon oral dosage forms.

§ 520.1120a Haloxon drench.

(a) *Chemical name.* 3-Chloro-7-hydroxy-4-methylcoumarin bis (2-chloroethyl) phosphate.

(b) *Specifications.* Haloxon assay of not less than 96 percent by infrared spectrum at 8.62 microns.

(c) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(d) *Special considerations.* Do not use any drug, insecticide, pesticide, or other chemical having cholinesterase-

§ 520.1120b

inhibiting activity either simultaneously or within a few days before or after treatment with haloxon.

(e) *Related tolerances.* See § 556.310 of this chapter.

(f) *Conditions of use.* It is used as a drench as follows:

(1) *Cattle*—(i) *Amount.* 141.5 grams per packet.

(ii) *Indications for use.* Control of gastrointestinal roundworms of the genera *Haemonchus*, *Ostertagia*, *Trichostrongylus*, and *Cooperia*.

(iii) *Limitations.* (a) Dissolve each packet in 32 fluid ounces of water and administer as follows:

Weight of animal (pounds)	Dose (fluid ounces)
Up to 100	½
100 to 150	¾
150 to 200	1
200 to 300	1½
300 to 450	2
450 to 700	3
700 to 1,000	4
1,000 to 1,200	5
Over 1,200	6

(b) Do not treat within 1 week of slaughter; do not treat dairy animals of breeding age; animals should be retreated in 3 to 4 weeks.

[40 FR 13838, Mar. 27, 1975, as amended at 45 FR 10333, Feb. 15, 1980; 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61624, Nov. 19, 1997]

§ 520.1120b Haloxon boluses.

(a) *Chemical name.* 3-Chloro-7-hydroxy-4-methylcoumarin bis (2-chloroethyl) phosphate.

(b) *Specifications.* Each bolus contains 10.1 grams of haloxon.

(c) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(d) *Related tolerances.* See § 556.310 of this chapter.

(e) *Conditions of use.* (1) Haloxon bolus is an anthelmintic used in cattle for the control of gastrointestinal roundworms of the genera *Haemonchus*, *Ostertagia*, *Trichostrongylus* and *Cooperia*.

(2) It is administered by giving one bolus per approximately 500 pounds body weight (35 to 50 milligrams per kilogram of body weight).

(3) For most effective results, retreat animals in 3 to 4 weeks. If rein-

21 CFR Ch. I (4–1–99 Edition)

fection is likely to occur, additional retreatments may be necessary.

(4) Do not use any drug, pesticide or other chemical having cholinesterase inhibiting activity either simultaneously or within a few days before or after treatment with haloxon.

(5) Do not treat animals within one week of slaughter.

(6) Do not treat dairy animals of breeding age or older.

[40 FR 13838, Mar. 27, 1975, as amended at 44 FR 61591, Oct. 29, 1979; 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 520.1130 Hetacillin oral dosage forms.

§ 520.1130a Hetacillin potassium capsules.

(a) *Specifications.* Each capsule contains hetacillin potassium equivalent to 50, 100, or 200 milligrams of ampicillin.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) *Dogs*—(i) *Amount.* 5 milligrams per pound of body weight, twice daily. In severe infections, up to three times daily, or up to 10 milligrams per pound of body weight twice daily. For stubborn urinary tract infections, up to 20 milligrams per pound of body weight twice daily.

(ii) *Indications for use.* Treatment against strains of organisms sensitive to hetacillin potassium and associated with respiratory tract infections, urinary tract infections, gastrointestinal infections, skin infections, soft tissue infections, and postsurgical infections.

(iii) *Limitations.* For use in dogs and cats only. Continue treatment for 48 to 72 hours after the animal has become afebrile or asymptomatic. Administer 1 to 2 hours prior to feeding to ensure maximum absorption. In stubborn infections, therapy may be required for several weeks. Not for use in animals raised for food production. Federal law restricts this drug to use only by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Amount.* Administer 50 milligrams twice daily.

(ii) *Indications for use.* Treatment against strains of organisms sensitive to hetacillin potassium and associated with respiratory tract infections, urinary tract infections, gastrointestinal

Food and Drug Administration, HHS

§ 520.1130c

infections, skin infections, soft tissue infections, and postsurgical infections.

(3) *Limitations.* For use in dogs and cats only. Continue treatment for 48 to 72 hours after the animal has become afebrile or asymptomatic. Administer in a fasting state to ensure maximum absorption. In stubborn infections, therapy may be required for several weeks. Not for use in animals raised for food production. Federal law restricts this drug to use only by or on the order of a licensed veterinarian.

[57 FR 37325, Aug. 18, 1992]

§ 520.1130b Hetacillin potassium oral suspension.

(a) *Specifications.* Each milliliter contains hetacillin potassium equivalent to 50 milligrams of ampicillin.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount.* 5 milligrams per pound of body weight twice daily. In severe infections, up to three times daily, or up to 10 milligrams per pound of body weight twice daily. For stubborn urinary tract infections, up to 20 milligrams per pound of body weight twice daily.

(ii) *Indications for use.* Treatment against strains of organisms susceptible to hetacillin potassium and associated with respiratory tract infections, urinary tract infections, gastrointestinal infections, skin infections, soft-tissue infections, and postsurgical infections.

(iii) *Limitations.* For use in dogs only. Not for use in animals raised for food production. Continue treatment 48 to 72 hours after the animal has become afebrile or asymptomatic. Administer 1 to 2 hours prior to feeding to ensure maximum absorption. In stubborn infections, therapy may be required for several weeks. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Amount.* 50 milligrams twice daily.

(ii) *Indications for use.* Treatment against strains of organisms susceptible to hetacillin potassium and associated with respiratory tract infections, urinary tract infections, gastrointestinal infections, skin infections, soft-tissue infections, and postsurgical infections.

(iii) *Limitations.* For use in cats only. Not for use in animals raised for food production. Continue treatment 48 to 72 hours after the animal has become afebrile or asymptomatic. Administer 1 to 2 hours prior to feeding to ensure maximum absorption. In stubborn infections, therapy may be required for several weeks. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37326, Aug. 18, 1992]

§ 520.1130c Hetacillin potassium tablets.

(a) *Specifications.* Each tablet contains hetacillin potassium equivalent to 50, 100, or 200 milligrams of ampicillin.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount.* 5 milligrams per pound of body weight twice daily. In severe infections, up to three times daily, or up to 10 milligrams per pound of body weight twice daily. For stubborn urinary tract infections, up to 20 milligrams per pound of body weight twice daily.

(ii) *Indications for use.* Oral treatment against strains of organisms sensitive to hetacillin potassium and associated with respiratory tract infections, urinary tract infections, gastrointestinal infections, skin infections, soft tissue infections, and postsurgical infections.

(iii) *Limitations.* For use in dogs and cats only. Continue treatment for 48 to 72 hours after the animal has become afebrile or asymptomatic. Administer 1 to 2 hours prior to feeding to ensure maximum absorption. In stubborn infections, therapy may be required for several weeks. Not for use in animals which are raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Amount.* 50 milligrams twice daily.

(ii) *Indications for use.* Treatment against strains of organisms sensitive to hetacillin potassium and associated with respiratory tract infections, urinary tract infections, gastrointestinal infections, skin infections, soft tissue infections, and postsurgical infections.

(iii) *Limitations.* For use in dogs and cats only. Continue treatment for 48 to

§ 520.1157

72 hours after the animal has become afebrile or asymptomatic. Administer 1 to 2 hours prior to feeding to ensure maximum absorption. In stubborn infections, therapy may be required for several weeks. Not for use in animals which are raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37326, Aug. 18, 1992]

§ 520.1157 Iodinated casein tablets.

(a) *Specifications.* Each 1-gram tablet contains 25 milligrams of iodinated casein.

(b) *Sponsor.* See No. 017762 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* $\frac{1}{2}$ to 1 tablet per 10 pounds of body weight (equivalent to 0.5 to 2.5 milligrams of iodinated casein per pound of body weight).

(2) *Indications for use.* For dogs for apparent decreased thyroid activity where the signs are alopecia, scaliness of the skin surface, loss of hair, seborrhea, thickening of the skin, hyperpigmentation, and lethargy.

(3) *Limitations.* If no response is observed in 30 to 45 days, the drug should be withdrawn and the diagnosis reconsidered. Do not use in the presence of cardiac disease, ischemia, adrenal insufficiency, or nephrosis. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[49 FR 22469, May 30, 1984]

§ 520.1158 Iodochlorhydroxyquin boluses.

(a) *Specifications.* Each bolus contains 10 grams of iodochlorhydroxyquin.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 1 bolus (10 grams) daily for a 1,000-pound horse.

(2) *Indications for use.* For treatment of equine diarrhea.

(3) *Limitations.* For horses only; not to be administered to food-producing animals. Do not administer to horses intended for use as food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 8054, Feb. 25, 1983, as amended at 50 FR 41489, Oct. 11, 1985]

21 CFR Ch. I (4–1–99 Edition)

§ 520.1182 Iron dextran oral suspension.

(a) *Specifications.* Each 1.8 milliliter contains 100 milligrams of elemental iron as ferric hydroxide in complex with a low molecular weight dextran and 0.2 percent phenol as a preservative.

(b) *Sponsor.* See 017800 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 100 milligrams of elemental iron to each pig.

(2) *Indications for use.* Prevention of iron deficiency anemia in baby pigs.

(3) *Limitations.* Treat each pig within 24 hours of farrowing. Administer 1.8 milliliters orally by automatic dose dispenser.

[45 FR 75199, Nov. 14, 1980]

§ 520.1192 Ivermectin paste.

(a) *Specifications*—(1) *Horses.* Paste contains 1.87 percent ivermectin.

(2) *Cattle.* Paste contains 0.153 percent ivermectin.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Horses*—(i) *Amount.* 200 micrograms per kilogram (91 micrograms per pound) of body weight.

(ii) *Indications for use.* It is used in horses for the treatment and control of large strongyles (adult) (*Strongylus equinus*), (adult and arterial larval stages) (*Strongylus vulgaris*), (adult and migrating tissue stages) (*Strongylus edentatus*), (adult) (*Triodontophorus* spp.); small strongyles, including those resistant to some benzimidazole class compounds (adult and fourth stage larvae) (*Cyathostomum* spp., *Cylicocyclus* spp., *Cylicodontophorus* spp., *Cylicostephanus* spp.); pinworms (adult and fourth stage larvae) (*Oxyuris equi*); ascarids (third- and fourth-stage larvae and adults) (*Parascaris equorum*); hairworms (adult) (*Trichostrongylus axei*); large mouth stomach worms (adult) (*Habronema muscae*); stomach bots (oral and gastric stages) (*Gastrophilus* spp.); lungworms (adults and fourth stage larvae) (*Dictyocaulus arnfieldi*); intestinal threadworms (adults) (*Strongyloides westeri*); summer sores caused by *Habronema* and

Draschia spp. cutaneous third stage larvae; and dermatitis caused by neck threadworm microfilariae (*Onchocerca* spp.).

(iii) *Limitations*. For oral use only. Do not use in horses intended for food purposes. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) *Cattle*—(i) *Amount*. 23 milligrams per 250 pounds of body weight.

(ii) *Indications for use*. It is used in cattle for the treatment and control of gastrointestinal roundworms (adults and fourth-stage larvae) (*Ostertagia ostertagi* (including inhibited forms), *O. lyrata*, *Haemonchus placei*, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia oncophora*, *C. punctata*, *Nematodirus helvetianus*, *Bunostomum phlebotomum*, *Strongyloides papillosus* (adults only), *Oesophagostomum radiatum*, *Trichuris ovis* (adults only)); lungworms (adults and fourth-stage larvae) (*Dictyocaulus viviparus*); grubs (first, second, and third instars) (*Hypoderma bovis*, *H. lineatum*); and sucking lice (*Linognathus vituli*, *Haematopinus eurysternus*).

(iii) *Limitations*. For oral use only. Do not treat cattle within 24 days of slaughter. Because withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[49 FR 22275, May 29, 1984, as amended at 50 FR 27819, July 8, 1985; 51 FR 44449, Dec. 10, 1986; 53 FR 51273, Dec. 21, 1988; 62 FR 63270, Nov. 28, 1997]

§ 520.1193 Ivermectin tablets and chewables.

(a) *Specifications*—(1) *Dogs*. Each tablet or chewable contains 68, 136, or 272 micrograms of ivermectin.

(2) *Cats*. Each chewable contains 55 or 165 micrograms of ivermectin.

(b) *Sponsor*. See 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount*. 6.0 micrograms per kilogram body weight (2.72 micrograms per pound), minimum. For dogs up to 25 pounds, 68 micrograms; dogs 26 to 50 pounds, 136 micrograms; dogs 51 to 100 pounds, 272 micrograms; dogs over 100 pounds, a combination of the appro-

priate tablets. The drug is administered at monthly dosing intervals.

(2) *Indications for use*. To prevent canine heartworm disease by eliminating the tissue stage of heartworm larvae (*Dirofilaria immitis*) for 1 month (30 days) after infection.

(3) *Limitations*. Use once-a-month. Recommended for dogs 6 weeks of age and older. Initial use within 1 month after first exposure to mosquitoes. Final use within 1 month after last exposure to mosquitoes. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d) *Conditions of use in cats*—(1) *Amount*. Up to 2.3 kilograms (up to 5 pounds), 55 micrograms; 2.3 to 6.8 kilograms (5 to 15 pounds), 165 micrograms; over 6.8 kilograms (15 pounds), a combination of the appropriate chewables (recommended minimum dose of 24 micrograms of ivermectin per kilogram of body weight (10.9 micrograms per pound)).

(2) *Indications for use*. To prevent feline heartworm disease by eliminating the tissue stage of heartworm larvae (*Dirofilaria immitis*) for a month (30 days) after infection, and for removal and control of adult and immature (L4) hookworms *Ancylostoma tubaeforme* and *A. braziliense*.

(3) *Limitations*. For use in cats 6 weeks of age and older. Administer once a month. The initial dose must be given within a month after cats first exposure to mosquitoes. The final dose must be given within a month after the cats last exposure to mosquitoes. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[52 FR 11042, Apr. 7, 1987, as amended at 54 FR 32337, Aug. 7, 1989; 61 FR 39868, July 31, 1996; 62 FR 5319, Feb. 5, 1997; 62 FR 63270, Nov. 28, 1997]

§ 520.1194 Ivermectin drench.

(a) *Specifications*. Each milliliter of 0.08 percent (weight per volume) micellar solution contains 0.08 milligram of ivermectin.

(b) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.344 of this chapter.

(d) *Conditions of use*—(1) *Amount*. 3.0 milliliters (2.4 milligrams of ivermectin) per 26 pounds of body

§ 520.1195

weight (or 200 micrograms per kilogram of body weight).

(2) *Indications for use.* It is used in sheep for treatment and control of the adult and fourth-stage larvae of the following parasites of sheep: gastrointestinal roundworms (*Haemonchus contortus*, *H. placei* (adults only), *Ostertagia circumcincta*, *Trichostrongylus axei*, *T. Colubriformis*, *Cooperia oncophora* (adults only), *C. curticei*, *Oesophagostomum columbianum*, *O. venulosum* (adults only), *Nematodirus battus*, *N. spathiger*, *Strongyloides papillosus* (adults only), *Chabertia ovina* (adults only), *Trichuris ovis* (adults only)), lungworms (*Dictyocaulus filaria*); and all larval stages of the nasal bot *Oestrus ovis*.

(3) *Limitations.* It is used as a drench in sheep only. Do not treat sheep within 11 days of slaughter. Do not use in other animal species as severe adverse reactions, including fatalities in dogs, may result. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[53 FR 27958, July 26, 1988, as amended at 62 FR 63270, Nov. 28, 1997]

§ 520.1195 Ivermectin liquid.

(a) *Specifications.* Each milliliter contains 10 milligrams of ivermectin.

(b) *Sponsor.* See Nos. 050604 and 059130 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* 200 micrograms per kilogram of body weight as a single dose.

(2) *Indications for use.* It is used in horses for the treatment and control of large strongyles (adult) (*Strongylus equinus*), (adult and arterial larval stages) (*Strongylus vulgaris*), (adult and migrating tissue stages) (*Strongylus edentatus*), (adult) (*Triodontophorus* spp.); small strongyles, including those resistant to some benzimidazole class compounds (adult and fourth stage larvae) (*Cyathostomum* spp., *Cylicocyclus* spp., *Cylicodontophorus* spp., *Cylicostephanus* spp.); pinworms (adult and fourth stage larvae) (*Oxyuris equi*); ascarids (third- and fourth-stage larvae and adults) (*Parascaris equorum*); hairworms (adult) (*Trichostrongylus axei*); large mouth stomach worms (adult) (*Habronema muscae*); stomach bots (oral and gastric stages) (*Gastrophilus* spp.); lungworms (adults

21 CFR Ch. I (4–1–99 Edition)

and fourth stage larvae) (*Dictyocaulus arnfieldi*); intestinal threadworms (adults) (*Strongyloides westeri*); summer sores caused by *Habronema* and *Draschia* spp. cutaneous third stage larvae; and dermatitis caused by neck threadworm microfilariae (*Onchocerca* spp.).

(3) *Limitations.* Administer by stomach tube or as an oral drench. Do not use in horses intended for food purposes. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[52 FR 34637, Sept. 14, 1987, as amended at 53 FR 51273, Dec. 21, 1988; 62 FR 63270, Nov. 28, 1997; 63 FR 38474, July 17, 1998]

§ 520.1196 Ivermectin and pyrantel pamoate chewable tablet.

(a) *Specifications.* Each chewable tablet contains either 68 micrograms (µg) of ivermectin and 57 milligrams (mg) of pyrantel (as pamoate salt), or 136 µg and 114 mg, or 272 µg and 227 mg, respectively.

(b) *Sponsor.* See 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Dogs—(i) Amount.* A minimum of 6 µg of ivermectin and 5 mg of pyrantel (as pamoate salt) per kilogram (2.72 µg and 2.27 mg per pound) of body weight.

(ii) *Indications for use.* To prevent canine heartworm disease by eliminating the tissue larval stages of *Dirofilaria immitis* for up to a month (30 days) after infection and treatment and control of adult ascarids *Toxocara canis* and *Toxascaris leonina*, and adult hookworms *Ancylostoma caninum*, *A. braziliense*, and *Uncinaria stenocephala*.

(iii) *Limitations.* Use monthly. Recommended for dogs 6 weeks of age and older. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[58 FR 8542, Feb. 16, 1993, as amended at 61 FR 15186, Apr. 5, 1996; 61 FR 59004, Nov. 20, 1996; 62 FR 63270, Nov. 28, 1997]

§ 520.1197 Ivermectin sustained-release bolus.

(a) *Specifications.* Each sustained-release bolus contains 1.72 grams of ivermectin.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.344 of this chapter.

(d) *Conditions of use in ruminating calves*—(1) *Amount.* Administer one bolus per calf weighing at least 275 pounds (lb) (125 kilograms (kg)) and not more than 660 lb (300 kg) on the day of administration.

(2) *Indications.* For treatment and control, throughout the grazing season (approximately 135 days), of gastrointestinal roundworms *Haemonchus placei*, *Ostertagia ostertagi* (including inhibited fourth-stage larvae), *Trichostrongylus axei*, *T. colubriformis*, *Cooperia* spp., *Nematodirus helvetianus*, *Bunostomum* spp., *phlebotomum*, *Oesophagostomum radiatum*; lungworms *Dictyocaulus viviparus*; grubs *Hypoderma* spp.; sucking lice *Linognathus vituli*, *Solenopotes capillatus*; mange mites *Psoroptes ovis*, *Sarcoptes scabiei*, and ticks *Amblyomma americanum*.

(3) *Limitations.* The bolus was specifically designed for use in cattle; do not use in other animal species. Calves must be ruminating and older than 12 weeks of age. Do not administer to calves weighing less than 275 lb (125 kg). Do not administer a damaged bolus. Because a milk withdrawal time has not been established, do not use in female dairy cattle of breeding age. Do not slaughter cattle within 180 days of treatment. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[61 FR 67452, Dec. 23, 1996, as amended at 62 FR 63270, Nov. 28, 1997]

§ 520.1204 Kanamycin sulfate, aminopentamide hydrogen sulfate, pectin, bismuth subcarbonate, activated attapulgitic suspension.

(a) *Specifications.* Each five milliliters of suspension of the drug contains: 100 milligrams of kanamycin as the sulfate, 0.033 milligram of aminopentamide hydrogen sulfate, 25 milligrams of pectin, 250 milligrams of bismuth subcarbonate, and 500 milligrams of activated attapulgitic.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is administered orally to dogs for the symptomatic relief of acute bacterial diarrhea caused by kanamycin-susceptible organisms.

(2) The drug is recommended for use at the rate of one teaspoonful (5 milliliters) of suspension per 20 pounds of body weight every 8 hours. Animals weighing under 10 pounds should be given one-half the above amount every 8 hours. The initial dose should be twice the amount of a single dose. Maximum dosage should not exceed three times the recommended dose.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 53 FR 27851, July 25, 1988; 56 FR 8710, Mar. 1, 1991; 64 FR 403, Jan. 5, 1999]

EFFECTIVE DATE NOTE: At 64 FR 403, Jan. 5, 1999, § 520.1204 was amended in paragraph (a) by removing the phrase “(the kanamycin used conforms to the standards of identity, strength, quality, and purity prescribed by § 444.30 of this chapter)”, effective May 20, 1999.

§ 520.1205 Kanamycin sulfate, pectin, bismuth subcarbonate, activated attapulgitic tablets.

(a) *Specifications.* Each tablet contains 100 milligrams of kanamycin (as the sulfate), 25 milligrams of pectin, 250 milligrams of bismuth subcarbonate, and 500 milligrams of activated attapulgitic.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* One tablet per 44 kilograms (20 pounds) of body weight every 8 hours. Maximum dose 3 tablets every 8 hours. For animals under 22 kilograms (10 pounds) ½ tablet every 8 hours. The initial loading dose should be twice the amount of a single dose.

(2) *Indications for use.* For the treatment of bacterial enteritis caused by organisms susceptible to kanamycin and the symptomatic relief of associated diarrhea in dogs.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[56 FR 8710, Mar. 1, 1991]

§ 520.1242 Levamisole hydrochloride oral dosage forms.

§ 520.1242a Levamisole hydrochloride drench and drinking water.

(a) *Specifications.* Each package contains either 9.075, 11.7, 18.15, 46.8, or

§ 520.1242a

21 CFR Ch. I (4–1–99 Edition)

544.5 grams of levamisole hydrochloride.

(b) *Sponsors.* Approval for sponsors in 21 CFR 510.600(c) for use as in paragraph (d) of this section as follows:

(1) See 043781 for use of 46.8 gram package as in paragraph (d)(1) of this section, for 11.7 and 46.8 gram packages as in paragraph (d)(2) of this section, and for 9.075 and 18.15 gram packages as in paragraph (d)(3) of this section.

(2) See 000061 for use of 46.8 and 544.5 gram packages as in paragraph (d)(1) of this section, for 11.7, 46.8, and 544.5 gram packages as in paragraph (d)(2) of this section, and for 18.15 gram package as in paragraph (d)(3) of this section.

(3) See 057561 for use of 46.8 and 544.5 gram packages as in paragraphs (d)(1) and (d)(2) of this section.

(c) *Related tolerances.* See § 556.350 of this chapter.

(d) *Conditions of use.* It is used as an anthelmintic at 0.365 gram per 100 pounds of body weight as follows:

(1) *Cattle*—(i) *Amount.* As a single oral dose drench using 46.8 or 544.5 gram packet.

(ii) *Indications for use.* Anthelmintic effective against the following nematode infections: Stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*), and lungworms (*Dictyocaulus*).

(iii) *Limitations.* Conditions of constant helminth exposure may require retreatment within 2 to 4 weeks after the first treatment. Do not slaughter for food within 48 hours of treatment. Not for use in dairy animals of breeding age. Consult your veterinarian before using in severely debilitated animals. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Prepare solutions for use as follows:

(a) Dissolve contents of 46.8 gram package in water to provide 1 quart (32 fluid ounces) of drench solution and administer as a drench at 1/4 ounce per 100 pounds of body weight as a single oral dose.

(b) Dissolve contents of 46.8 gram package in water to provide 8.75 fluid ounces of concentrate solution and administer as a drench at 2 milliliters per

100 pounds of body weight as a single oral dose by syringe.

(c) Dissolve contents of 544.5 gram package in 3 liters of water and administer as a drench at 2 milliliters per 100 pounds of body weight as a single oral dose.

(2) *Sheep*—(i) *Amount.* As a single oral dose drench using 11.7, 46.8, or 544.5 gram packet.

(a) *Indications for use.* Anthelmintic effective against the following nematode infections: Stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*, *Chabertia*), and lungworms (*Dictyocaulus*).

(b) *Limitations.* Dissolve in 1 gallon (128 fluid ounces) of water and administer as a single drench at 1 ounce (0.365 gram) per 100 pounds of body weight; conditions of constant helminth exposure may require re-treatment within 2 to 4 weeks after the first treatment; do not slaughter for food within 72 hours of treatment; consult veterinarian before using in severely debilitated animals.

(ii) *Amount.* 11.7 grams per packet.

(a) *Indications for use.* See paragraph (f)(2)(i)(a) of this section.

(b) *Limitations.* Dissolve in 1 quart (32 fluid ounces) of water and administer as a single drench at 1 ounce (0.365 gram) per 100 pounds of body weight or dissolve 1 packet in 10.9 fluid ounces of water and administer as a single drench at 1 cubic centimeter (0.036 gram) per 10 pounds of body weight; conditions of constant helminth exposure may require re-treatment within 2 to 4 weeks after the first treatment; do not slaughter for food within 72 hours of treatment; consult veterinarian before using in severely debilitated animals.

(iii) *Limitations.* Conditions of constant helminth exposure may require retreatment within 2 to 4 weeks after the first treatment. Do not slaughter for food within 72 hours of treatment. Consult your veterinarian before using in severely debilitated animals. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Prepare solutions for use as follows:

(a) Dissolve contents of 11.7 gram package in 1 quart (32 ounces) of water and administer as a drench at 1 ounce per 100 pounds of body weight, or dissolve in 10.9 fluid ounces of water and administer as a drench at 1 milliliter per 10 pounds of body weight as a single oral dose.

(b) Dissolve contents of 46.8 gram package in 128 fluid ounces (1 gallon) of water and administer as a drench at 1 ounce per 100 pounds of body weight as a single oral dose.

(c) Dissolve contents of 544.5 gram package in 3 liters of water and administer as a drench at 2 milliliters per 100 pounds of body weight as a single oral dose.

(3) *Swine*—(i) *Amount*. 9.075 or 18.15 grams per bottle.

(ii) *Indications for use*. Anthelmintic effective against the following nematode infections: Large roundworms (*Ascaris suum*), nodular worms (*Oesophagostomum* spp.), intestinal thread worms (*Strongyloides ransomi*) and lungworms (*Metastrongylus* spp.).

(iii) *Limitations*. Dissolve in water to provide 9.075 grams per 250 milliliters or 18.15 grams per 500 milliliters. Add 10 milliliters (2 teaspoons) of this concentrate solution to each gallon of drinking water. Allow 1 gallon of medicated drinking water for each 100 pounds of body weight of pigs to be treated. No other source of water should be offered. After pigs have consumed medicated water, resume use of regular water. Pigs maintained under conditions of constant exposure to worms may require retreatment within 4 to 5 weeks after the first treatment. Consult your veterinarian before administering to sick swine. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Do not administer within 72 hours of slaughter for food.

[40 FR 13838, Mar. 27, 1975, as amended at 43 FR 11176, Mar. 17, 1978; 44 FR 59508, Oct. 16, 1979; 44 FR 72586, Dec. 14, 1979; 46 FR 25085, May 5, 1981; 52 FR 20598, June 2, 1987; 54 FR 18280, Apr. 28, 1989; 62 FR 61625, Nov. 19, 1997; 63 FR 59713, Nov. 5, 1998]

§ 520.1242b Levamisole hydrochloride tablet or oblet (bolus).

(a) *Chemical name*. (-)-2,3,5,6-Tetrahydro-6-phenylimidazo [2,1-*b*] thiazole monohydrochloride.

(b) *Specifications*. Assay of not less than 98 percent by nonaqueous titration with 0.1 *N* potassium isopropoxide; 1 isomer minimum 95 percent pure by optical rotation.

(c) *Sponsor*. See Nos. 000061 and 043781 in § 510.600(c) of this chapter.

(d) *Required labeling*. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(e) *Related tolerances*. See § 556.350 of this chapter.

(f) *Conditions of use*. (1) It is used in an oblet for cattle as follows:

(i) *Amount*. 2.19 grams per oblet.

(ii) *Indications for use*. Anthelmintic effective against the following nematode infections: Stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*), and lungworms (*Dictyocaulus*).

(iii) *Limitations*. Administer as a single dose as follows: 250 to 450 pounds, ½ oblet; 450 to 750 pounds, 1 oblet; and 750 to 1,050 pounds, 1½ oblets; conditions of constant helminth exposure may require re-treatment within 2 to 4 weeks after the first treatment; do not slaughter for food within 48 hours of treatment; not for use in dairy animals of breeding age; consult veterinarian before using in severely debilitated animals.

(2) It is used in a tablet for sheep as follows:

(i) *Amount*. 0.184 gram per tablet.

(ii) *Indications for use*. Anthelmintic effective against the following nematode infections: Stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*, *Chabertia*), and lungworms (*Dictyocaulus*).

(iii) *Limitations*. Administer one tablet for each 50 pounds of body weight;

§ 520.1242c

conditions of constant helminth exposure may require re-treatment within 2 to 4 weeks after the first treatment; do not slaughter for food within 72 hours of treatment; consult a veterinarian before using in severely debilitated animals.

[40 FR 13838, Mar. 27, 1975, as amended at 44 FR 59507, Oct. 16, 1979; 62 FR 61625, Nov. 19, 1997]

§ 520.1242c Levamisole hydrochloride and piperazine dihydrochloride.

(a) *Specifications.* (1) The drug is an aqueous solution which contains in each fluid ounce 0.36 gram of levamisole hydrochloride and piperazine dihydrochloride equivalent to 3.98 grams of piperazine base.

(2) The drug is a soluble powder which when reconstituted with water contains in each fluid ounce 0.45 gram of levamisole hydrochloride and piperazine dihydrochloride equivalent to 5.0 grams of piperazine base.

(b) *Sponsor.* See No. 043781 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use.* It is used as a drench for horses as follows:

(1) *Indications for use.* An anthelmintic effective against infections of large strongyles (*Strongylus vulgaris*, *S. edentatus*), small strongyles (*Cylicocercus* spp., *Cylicocylclus* spp., *Cylicodontophorus* spp., *Cylicostephanus* spp., *Cylicotetrapedon* spp.), ascarids (*Parascaris equorum*), and pinworms (*Oxyuris equi*).

(2) *Limitations.* Aqueous solution: administer by stomach tube or drench 1 fluid ounce per 100 pounds of body weight. Reconstituted soluble powder: administer by stomach tube 1 fluid ounce per 125 pounds of body weight. If reinfection occurs, re-treat animals at 6- to 8-week intervals. Do not treat animals intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 32831, Aug. 5, 1975, as amended at 41 FR 48731, Nov. 5, 1976; 43 FR 11176 Mar. 17, 1978]

§ 520.1242d Levamisole resinate.

(a) *Specifications.* The drug is levamisole adsorbed on a resin, in a concentration equivalent to 10 percent levamisole hydrochloride. Each 2.05-

21 CFR Ch. I (4–1–99 Edition)

ounce (58.1 gram) packet contains levamisole equivalent to 5.806 grams of levamisole hydrochloride.

(b) *Sponsor.* See No. 043781 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.350 of this chapter.

(d) *Conditions of use.* In swine it is used as follows:

(1) *Amount.* The equivalent of 8 milligrams per kilogram of body weight, as a single dose, mixed in the animal's ration.

(2) *Indications for use.* For the removal of and control of the following nematode infections: large roundworms (*Ascaris suum*), nodular worms (*Oesophagostomum* spp.), lungworms (*Metastrongylus* spp.), intestinal threadworms (*Strongyloides ransomi*), and swine kidney worms (*Stephanurus dentatum*).

(3) *Limitations.* For pigs from weaning to market weight, mix one 58.1-gram packet of levamisole resinate containing the equivalent of 10-percent levamisole hydrochloride in 40 pounds of feed and administer 1 pound of medicated feed per 40 pounds of body weight as sole ration. For breeding swine, mix 1 packet of the 10-percent resinate in 16 pounds of feed and administer 1 pound of medicated feed per 100 pounds of body weight as sole ration. Administer as single doses. Withhold regular feed overnight and administer medicated feed the following morning. Do not withhold water during fasting. Do not treat within 72 hours of slaughter. Salivation or muzzle foam may be observed. The reaction will disappear a short time after feeding. If pigs are infected with mature lungworms, coughing and vomiting may be observed. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[43 FR 18171, Apr. 28, 1978, as amended at 45 FR 3574, Jan. 18, 1980]

§ 520.1242e Levamisole hydrochloride effervescent tablets.

(a) *Specifications.* Each tablet contains 907 milligrams of levamisole hydrochloride.

(b) *Sponsor.* See No. 043781 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.350 of this chapter.

Food and Drug Administration, HHS

§ 520.1242g

(d) *Conditions of use.* It is used for swine as follows:

(1) *Amount.* The equivalent of 8 milligrams of levamisole hydrochloride per kilogram of body weight, as a single dose.

(2) *Indications for use.* See § 520.1242a(f)(3)(ii).

(3) *Limitations.* Withholding water from pigs before treatment is not necessary. Add one tablet for each 2½ gallons of water; mix thoroughly. Allow 1 gallon of medicated water for each 100 pounds body weight of pigs to be treated. No other source of water should be offered. After pigs have consumed medicated water, resume use of regular water. Pigs maintained under conditions of constant worm exposure may require re-treatment within 4 to 5 weeks. Consult your veterinarian before administering to sick swine. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Do not administer within 72 hours of slaughter for food.

[45 FR 6087, Jan. 25, 1980]

§ 520.1242f Levamisole hydrochloride gel.

(a) *Specifications.* The drug is a gel containing 11.5 percent levamisole hydrochloride.

(b) *Sponsor.* See No. 043781 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.350 of this chapter.

(d) *Conditions of use—(1) Cattle—(i) Amount.* Eight milligrams of levamisole hydrochloride per kilogram of body weight, as a single oral dose.

(ii) *Indications for use.* Anthelmintic effective against the following nematode infections: Stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*), and lungworms (*Dictyocaulus*).

(iii) *Limitations.* Conditions of constant helminth exposure may require re-treatment within 2 to 4 weeks after the first treatment; do not administer to cattle within 6 days of slaughter for food; do not administer to dairy animals of breeding age; consult veterinarian before using in severely debilitated animals.

(2) *Breeding swine—(i) Amount.* Eight milligrams per kilogram of body weight (3.6 milligrams per pound) as a single oral dose.

(ii) *Conditions of use.* For treating breeding swine infected with the following nematodes: Large roundworms (*Ascaris suum*), nodular worms (*Oesophagostomum* spp.), lungworms (*Metastrongylus* spp.), intestinal threadworms (*Strongyloides ransomi*), and kidney worms (*Stephanurus dentatus*).

(iii) *Limitations.* May require retreatment in 4 to 5 weeks. Do not use within 11 days of slaughter for food. Consult your veterinarian for assistance before using in severely debilitated animals and in the diagnosis, treatment, and control of parasitism.

[47 FR 22517, May 25, 1982; 47 FR 30242, July 13, 1982, as amended at 48 FR 11429, Mar. 18, 1983; 51 FR 29215, Aug. 15, 1986]

§ 520.1242g Levamisole resinate and famphur paste.

(a) *Chemical name of famphur.* O, O-Dimethyl O-[p-(dimethylsulfamoyl)phenyl]phosphorothioate.

(b) *Specifications.* The drug is a paste containing 11.6 percent levamisole resinate (50 percent potency) and 23.6 percent famphur.

(c) *Sponsor.* See 000061 in § 510.600(c) of this chapter.

(d) *Special considerations.* Do not use any cholinesterase-inhibiting drugs, pesticides, insecticides, or chemicals on cattle simultaneously or within a few days before or after treatment with this product.

(e) *Related tolerances.* See § 556.350 of this chapter for levamisole and § 556.273 of this chapter for famphur.

(f) *Conditions of use in cattle—(1) Amount.* 8 milligrams of levamisole hydrochloride (equivalent) and 30 milligrams of famphur activity per kilogram of body weight.

(2) *Indications for use.* For treatment of cattle infected with the following parasites: Stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*), lungworms (*Dictyocaulus*), cattle grubs

§ 520.1263

(*Hypoderma*), biting lice (*Bovicola*), and sucking lice (*Linognathus*, *Solenoptes*).

(3) *Limitations*. Drug is not effective against lice eggs. Conditions of constant helminth and ectoparasitic exposure may require retreatment within 2 to 4 weeks after first treatment. Do not administer to cattle within 19 days of slaughter. Do not administer to dairy animals of breeding age. Do not use in calves less than 3 months old, or in debilitated animals. Do not treat Brahman bulls. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[53 FR 23757, June 24, 1988, as amended at 54 FR 1353, Jan. 13, 1989; 57 FR 7652, Mar. 4, 1992; 62 FR 55160, Oct. 23, 1997; 62 FR 61625, Nov. 19, 1997]

§ 520.1263 Lincomycin hydrochloride monohydrate oral dosage forms.

§ 520.1263a Lincomycin hydrochloride monohydrate tablets and sirup.

(a) *Specifications*. The sirup contains lincomycin hydrochloride equivalent to either 25 milligrams or 50 milligrams of lincomycin.

(b) *Sponsor*. See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) The drug is indicated in infections caused by gram-positive organisms which are sensitive to its action, particularly streptococci and staphylococci.

(2) It is administered orally to dogs and cats at a dosage level of 10 mgs per pound of body weight every 12 hours, or 7 mgs per pound of body weight every 8 hours. Treatment may be continued for periods as long as 12 days if clinical judgment indicates.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 44 FR 7130, Feb. 6, 1979; 64 FR 403, Jan. 5, 1999]

EFFECTIVE DATE NOTE: At 64 FR 403, Jan. 5, 1999, § 520.1263a was amended by revising paragraph (a), effective May 20, 1999. For the convenience of the user, the superseded text is set forth as follows:

§ 520.1263a Lincomycin hydrochloride monohydrate tablets and sirup.

(a) *Specifications*. The lincomycin hydrochloride monohydrate of the tablet meets the specifications prescribed by § 453.30(a)(1)

21 CFR Ch. I (4–1–99 Edition)

of this chapter. The sirup meets the specifications prescribed by § 453.130b(a)(1) of this chapter.

* * * * *

§ 520.1263b Lincomycin hydrochloride monohydrate and spectinomycin sulfate tetrahydrate soluble powder.

(a) *Specifications*. The spectinomycin sulfate tetrahydrate used in manufacturing the drug is the antibiotic substance produced by the growth of *Streptomyces spectabilis* or the same antibiotic substance produced by any other means. The quantity of total antibiotic activity cited in this section refers to the equivalent weight of the base activity of the drugs. Lincomycin hydrochloride monohydrate and spectinomycin sulfate tetrahydrate are present in the drug in the ratio of 1 to 2 on the basis of equivalency of lincomycin base to equivalency of spectinomycin base.

(b) *Sponsor*. See No. 000009 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See §§ 556.600 and 556.360 of this chapter.

(d) *Conditions of use*. (1) It is administered in the drinking water of chickens up to 7 days of age as an aid in the control of airsacculitis caused by either *Mycoplasma synoviae* or *Mycoplasma gallisepticum* susceptible to lincomycin-spectinomycin and complicated chronic respiratory disease (air sac infection) caused by *Escherichia coli* and *M. gallisepticum* susceptible to lincomycin-spectinomycin.

(2) For aid in the control of these conditions it is administered in the drinking water at a level of 2 grams of antibiotic activity per gallon of water as the sole source of water for the first 5 to 7 days of life.

[40 FR 13838, Mar. 27, 1975, as amended at 41 FR 1891, Jan. 13, 1976; 64 FR 403, Jan. 5, 1999]

EFFECTIVE DATE NOTE: At 64 FR 403, Jan. 5, 1999, § 520.1263b was amended by removing the first sentence from paragraph (a), effective May 20, 1999. For the convenience of the user, the superseded text is set forth as follows:

Food and Drug Administration, HHS

§ 520.1288

§ 520.1263b Lincomycin hydrochloride monohydrate and spectinomycin sulfate tetrahydrate soluble powder.

(a) * * * The lincomycin hydrochloride monohydrate meets the specifications prescribed by § 453.30(a)(1) of this chapter. * * *

* * * * *

§ 520.1263c Lincomycin hydrochloride soluble powder.

(a) *Specifications.* Each 40-gram packet (1.41 ounce) contains lincomycin hydrochloride equivalent to 16 grams of lincomycin. Each 80-gram packet (2.82 ounces) contains lincomycin hydrochloride equivalent to 32 grams of lincomycin. The 40-gram measuring device contains lincomycin hydrochloride equivalent to 16 grams of lincomycin (the measuring device is packaged with a 32-ounce jar).

(b) *Sponsors.* Approval for use of 40- and 80-gram packet to Nos. 000009 and 017144 in § 510.600(c) of this chapter. Approval for use of 40- and 80-gram packet and 32-ounce jar to No. 051259 in § 510.600(c) of this chapter.

(c) *Tolerances.* See § 556.360 of this chapter.

(d) *Conditions of use*—(1) It is used in drinking water for swine as follows:

(i) *Amount.* 250 milligrams per gallon.

(A) *Dosage.* 3.8 milligrams per pound of body weight per day.

(B) *Indications for use.* Treatment of swine dysentery (bloody scours).

(C) *Limitations.* Discard medicated drinking water if not used within 2 days. Prepare fresh stock solution daily. Do not use for more than 10 days. If clinical signs of disease have not improved within 6 days, discontinue treatment and reevaluate diagnosis. Not for use in swine weighing more than 250 pounds.

(ii) [Reserved]

(2) It is used in drinking water for broiler chickens as follows:

(i) *Amount.* 64 milligrams per gallon.

(A) *Indications for use.* For the control of necrotic enteritis caused by *Clostridium perfringens* susceptible to lincomycin.

(B) *Limitations.* Discard medicated drinking water if not used within 2 days. Prepare fresh stock solution daily. Administer for 7 consecutive days. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants ac-

cess to water containing lincomycin. Not for use in layer and breeder chickens.

(ii) [Reserved]

[48 FR 3966, Jan. 28, 1983, as amended at 55 FR 3209, Jan. 31, 1990; 60 FR 14217, Mar. 16, 1995; 62 FR 65020, Dec. 10, 1997; 64 FR 13341, Mar. 18, 1999; 64 FR 13508, Mar. 19, 1999]

§ 520.1284 Sodium liothyronine tablets.

(a) *Specifications.* Sodium liothyronine tablets consist of tablets intended for oral administration which contain liothyronine at 60 or 120 micrograms per tablet, as the sodium salt.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is indicated in cases of hypothyroidism in dogs.

(2) It is administered orally to dogs at levels up to 12.8 micrograms per kilogram of body weight per day. Dosage should be adjusted according to the severity of the condition and the response of the patient. Dosage at the total replacement level (12.8µg per kilogram of body weight) should be considered for initiating therapy and then titrated downward for optimum maintenance effect. Twice daily administration is recommended.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.1288 Lufenuron tablets.

(a) *Specifications*—(1) *Dogs.* Each tablet contains either 45, 90, 204.9, or 409.8 milligrams (mg) of lufenuron.

(2) *Cats.* Each tablet contains either 90, 135, 204.9 or 270 mg of lufenuron.

(b) *Sponsor.* See No. 058198 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount.* Minimum of 10 mg of lufenuron per kilogram (4.5 mg per pound (lb)) of body weight.

(2) *Indications for use.* For use in dogs and puppies, 6 weeks of age and older, for the prevention and control of flea populations. Lufenuron controls flea populations by preventing the development of flea eggs and does not kill

§ 520.1289

adult fleas. Concurrent use of insecticides may be necessary for adequate control of adult fleas.

(3) *Limitations.* Administer tablet(s) after or in conjunction with a full meal to ensure adequate absorption. Administer tablet(s) once a month. All dogs and cats in a household should be treated to achieve maximum efficacy.

(d) *Conditions of use in cats—(1) Amount.* Minimum of 30 mg of lufenuron per kilogram (13.6 mg/lb) of body weight.

(2) *Indications for use.* For use in cats and kittens, 6 weeks of age and older, for the control of flea populations. Lufenuron controls flea populations by preventing the development of flea eggs and does not kill adult fleas. Concurrent use of insecticides may be necessary for adequate control of adult fleas.

(3) *Limitations.* Administer tablet(s) after or in conjunction with a full meal to ensure adequate absorption. Administer tablet(s) once a month. All dogs and cats in a household should be treated to achieve maximum efficacy.

[63 FR 52968, Oct. 2, 1998]

§ 520.1289 Lufenuron suspension.

(a) *Specifications.* Each individual dose pack contains either 135 or 270 milligrams of lufenuron.

(b) *Sponsor.* See No. 058198 in § 510.600(c) of this chapter.

(c) *Conditions of use in cats—(1) Amount.* Minimum of 13.6 milligrams per pound of body weight (30 milligrams per kilogram). Recommended dose of 135 milligrams for up to 10 pounds of body weight or 270 milligrams for 11 to 20 pounds. Cats over 20 pounds are provided the appropriate combination of packs.

(2) *Indications for use.* For control of flea populations.

(3) *Limitations.* For oral use in cats 6 weeks of age or older, once a month, mixed with food. Administer in conjunction with a full meal to ensure adequate absorption. Treat all cats in the household to ensure maximum benefits. Because the drug has no effect on adult fleas, the concurrent use of insecticides that kill adults may be nec-

21 CFR Ch. I (4–1–99 Edition)

essary depending on the severity of the infestation.

[60 FR 20402, Apr. 26, 1995, as amended at 62 FR 8371, Feb. 25, 1997]

§ 520.1320 Mebendazole oral.

(a) *Chemical name.* Methyl 5-benzoyl-benzimidazole-2-carbamate.

(b) *Specifications.* As oral powder: Each gram contains either 40 or 166.7 milligrams of mebendazole. As oral paste: Each gram contains 200 milligrams of mebendazole. As oral suspension: Each milliliter contains 33.3 milligrams of mebendazole.

(c) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(d) *Conditions of use—(1) Horses—(i) Amount.* 1 gram of mebendazole per 250 pounds of body weight per dose, as an oral powder, paste or suspension.

(ii) *Indications for use.* It is used in horses for treatment of infections caused by large roundworms (*Parascaris equorum*); large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*); small strongyles; and mature and immature (4th larval stage pinworms (*Oxyuris equi*)).

(iii) *Limitations—(a) Oral powder.* The drug is given by sprinkling directly on the grain portion of the ration or dissolving in 2 to 4 pints of water and administering by stomach tube. The drug is compatible with carbon disulfide, which can be used concurrently for both control (*Gastrophilus spp.*). Routine cautions regarding the use of carbon disulfide must be observed. Do not administer to horses intended for use as food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b) *Oral paste.* The drug is given by dosing gun (syringe), inserting the tip of the gun at the interdental space in the horse's mouth and depositing the paste on the animal's tongue. The hand is placed under the animal's jaw, and the head is raised to assure that the paste is deposited on the roof of the mouth. Not for use in horses intended for food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(c) *Oral suspension.* The drug is administered by stomach tube. Not for horses intended for food use. Federal

Food and Drug Administration, HHS

§ 520.1330

law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Dogs*—(i) *Amount*. One hundred milligrams of mebendazole per 10 pounds of body weight, once daily for 3 days, as an oral powder.

(ii) *Indications for use*. The drug is used for treatment of infections of roundworms (*Toxocara canis*), hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*), whipworms (*Trichuris vulpis*), and tapeworms (*Taenia pisiformis*).

(iii) *Limitations*. Administer as an oral powder by mixing with a small quantity of food, preferably before the regular meal. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[42 FR 61255, Dec. 2, 1977, as amended at 43 FR 35686, Aug. 11, 1978; 45 FR 3574, Jan. 18, 1980; 46 FR 47218, Sept. 25, 1981; 46 FR 53658, Oct. 30, 1981; 62 FR 61625, Nov. 19, 1997]

§ 520.1326 Mebendazole and trichlorfon oral dosage forms.

§ 520.1326a Mebendazole and trichlorfon powder.

(a) *Specifications*. Each gram of powder contains 83.3 milligrams of mebendazole and 375.0 milligrams of trichlorfon.

(b) *Sponsor*. See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*. *Horses*—(1) *Amount*. 8.8 milligrams of mebendazole and 40 milligrams of trichlorfon per kilogram of body weight.

(2) *Indications for use*. It is used in horses for the treatment of infections of bots (*Gastrophilus intestinalis* and *G. nasalis*), large roundworms (*Parascaris equorum*), large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), small strongyles, and pinworms (*Oxyuris equi*).

(3) *Limitations*. Administer orally as an individual dose by stomach tube or thoroughly mixed in the ground grain portion of the ration to be consumed in one feeding. Discard treated feed not consumed. Do not administer more than once every 30 days. Do not treat sick or debilitated animals, foals under 4 months of age, or mares in the last month of pregnancy. Trichlorfon is a cholinesterase inhibitor. Do not administer simultaneously or within a few

days before or after treatment with, or exposure to, cholinesterase-inhibiting drugs, pesticides or chemicals. Do not administer intravenous anesthetics, especially muscle relaxants, concurrently. Not for horses intended for food use. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 10759, Feb. 19, 1980, as amended at 46 FR 52330, Oct. 27, 1981. Redesignated at 51 FR 13212, Apr. 18, 1986, as amended at 62 FR 61625, Nov. 19, 1997]

§ 520.1326b Mebendazole and trichlorfon paste.

(a) *Specifications*. Each gram of paste contains 100 milligrams of mebendazole and 454 milligrams of trichlorfon.

(b) *Sponsor*. See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 8.8 milligrams of mebendazole and 40 milligrams of trichlorfon per kilogram of body weight.

(2) *Indications for use*. It is used in horses for treatment of infections of bots (*Gastrophilus intestinalis* and *G. nasalis*), large roundworms (*Parascaris equorum*), large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), small strongyles, and pinworms (*Oxyuris equi*).

(3) *Limitations*. Do not administer more than once every 30 days. Do not treat sick or debilitated animals, foals under 4 months of age, or mares in the last month of pregnancy. Trichlorfon is a cholinesterase inhibitor. Do not administer simultaneously or within a few days before or after treatment with, or exposure to, cholinesterase-inhibiting drugs, pesticides, or chemicals. Do not administer intravenous anesthetics, especially muscle relaxants, concurrently. Not for use in horses intended for food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[51 FR 13212, Apr. 18, 1986, as amended at 62 FR 61625, Nov. 19, 1997]

§ 520.1330 Meclofenamic acid granules.

(a) *Chemical name*. N-(2,6-Dichloromethyl) anthranilic acid.

(b) *Specifications*. The drug is in granular form containing 5 percent meclofenamic acid.

§ 520.1331

21 CFR Ch. I (4–1–99 Edition)

(c) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) The drug is used in horses for the treatment of acute or chronic inflammatory diseases involving the musculoskeletal system.

(2) It is administered orally at a dosage of 1 milligram per pound of body weight (1 gram per 1,000 pounds) once daily for 5 to 7 days by addition to the daily grain ration.

(3) Treatment beyond the initial 5- to 7-day period may be indicated. A maintenance dosage level should be individualized for each animal.

(4) This drug should not be administered to horses with active gastrointestinal, hepatic, or renal disease.

(5) Not for use in horses intended for food.

(6) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 5632, Feb. 9, 1976, as amended at 53 FR 23390, June 22, 1988]

§ 520.1331 Meclofenamic acid tablets.

(a) *Specifications.* Each tablet contains either 10 or 20 milligrams of meclufenamic acid.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs—*(1) *Amount.* 1.1 milligrams per kilogram (0.5 milligram per pound) daily for 5 to 7 days.

(2) *Indications for use.* For the relief of signs and symptoms of chronic inflammatory disease involving the musculoskeletal system.

(3) *Limitations.* For oral use only. Should not be administered to animals with congestive heart failure or active gastrointestinal, hepatic, or renal disease. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 43385, Oct. 25, 1985, as amended at 53 FR 23390, June 22, 1988]

§ 520.1341 Megestrol acetate tablets.

(a) *Specifications.* Each tablet contains 5 or 20 milligrams of megestrol acetate.

(b) *Sponsor.* No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in female dogs for the postpone-

ment of estrus and the alleviation of false pregnancy.

(2) It is administered orally, intact, or crushed and mixed with food as follows:

(i) For the postponement of estrus by proestrus treatment, 1 milligram per pound of body weight per day for 8 days.

(ii) For the postponement of estrus by anestrus treatment, 0.25 milligram per pound of body weight per day for 32 days.

(iii) For alleviation of false pregnancy, 1 milligram per pound of body weight per day for 8 days.

(3) Full dosage regimen must be completed to produce the desired effect.

(4) Examination of vaginal smears is recommended to confirm detection of proestrus.

(5) Do not administer for more than two consecutive treatments.

(6) Once therapy is started, the animal should be confined for 3 to 8 days or until cessation of bleeding, since dogs in proestrus accept a male.

(7) Do not use prior to or during first estrus cycle.

(8) Do not use in pregnant animals.

(9) Do not use in the presence of a disease of the reproductive system or with mammary tumors.

(10) Should estrus occur within 30 days after cessation of treatment, mating should be prevented.

(11) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 52 FR 7832, Mar. 13, 1987]

§ 520.1380 Methocarbamol tablets.

(a) *Chemical name.* 3-(O-Methoxyphenoxy)-1,2-propanediol 1-carbamate.

(b) *Specifications.* Each tablet contains 500 milligrams of methocarbamol.

(c) *Sponsor.* See No. 000031 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) The drug is administered to dogs and cats as an adjunct to therapy for acute inflammatory and traumatic conditions of the skeletal muscles in order to reduce muscular spasms.

(2) Dosage is based upon severity of symptoms and response noted. The usual initial dose in 60 milligrams per

pound of body weight in two or three equally divided doses followed by 30 to 60 milligrams per pound of body weight each following day, usually not to exceed 14 to 21 days.

(3) For use only by or on the order of a licensed veterinarian.

§ 520.1408 Methylprednisolone tablets.

(a) *Specifications.* Each table contains 1, 2, or 4 milligrams of methylprednisolone.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter for use of 1- and 4-milligram tablets; see No. 000010 for use of 1- and 2-milligram tablets.

(c) *NAS/NRC status.* The conditions of use have been NAS/NRC reviewed and found effective. NADA's for approval of drugs for these conditions of use need not include effectiveness data specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Special consideration.* (1) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(2) Systemic therapy with methylprednisolone is contraindicated in animals with arrested tuberculosis, peptic ulcer, acute psychoses, or cushingoid syndrome. The presence of active tuberculosis, diabetes, osteoporosis, chronic psychotic reactions, predisposition to thrombophlebitis, hypertension, congestive heart failure, or renal insufficiency necessitates carefully controlled use of corticosteroids. Some of these conditions occur only rarely in dogs and cats but should be kept in mind.

(3) Anti-inflammatory action of corticosteroids may mask signs of infection.

(e) *Conditions of use—(1) Amount.* Dogs and cats: 5 to 15 pounds, 2 milligrams; 15 to 40 pounds, 2 to 4 milligrams; 40 to 80 pounds, 4 to 8 milligrams.

(2) *Indications for use.* For use in dogs and cats as an anti-inflammatory agent.

(3) *Limitations.* Administer total daily dose orally in equally divided doses 6 to 10 hours apart until response is noted or 7 days have elapsed. When response is attained, dosage should be gradually reduced until maintenance level is achieved. Hazardous for human use. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[47 FR 52697, Nov. 23, 1982, as amended at 49 FR 20810, May 17, 1984; 50 FR 32844, Aug. 15, 1985; 53 FR 40727, Oct. 18, 1988; 62 FR 35076, June 30, 1997]

§ 520.1409 Methylprednisolone, aspirin tablets.

(a) *Specifications.* Each tablet contains 0.5 milligram of methylprednisolone and 300 milligrams of aspirin.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *NAS/NRC status.* The conditions of use have been NAS/NRC reviewed and found effective. New animal drug applications for approval of drugs for these conditions of use need not include effectiveness data specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Special considerations.* (1) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(2) Systemic therapy with methylprednisolone is contraindicated in animals with tuberculosis, chronic nephritis, peptic ulcer, or Cushingoid syndrome. The presence of diabetes mellitus, osteoporosis, predisposition to thrombophlebitis, hypertension, congestive heart failure, or renal insufficiency necessitates carefully controlled use of corticosteroids.

(3) Anti-inflammatory action of corticosteroids may mask signs of infection.

(e) *Conditions of use—(1) Amount.* Dogs under 15 pounds, ¼ to 1 tablet daily; 15 to 60 pounds, 1 to 2 tablets daily; 60 pounds and over, 2 tablets daily.

§ 520.1422

(2) *Indications for use.* As an anti-inflammatory and analgesic agent in dogs.

(3) *Limitations.* Administer total daily dose in divided doses 6 to 10 hours apart, with a light feeding. When response is attained, dosage should be gradually reduced until maintenance level is achieved. Do not administer to cats. Do not overdose. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 21566, May 13, 1983]

§ 520.1422 Metoserpate hydrochloride.

(a) *Chemical name.* Methyl-*o*-methyl-18-epireserpate hydrochloride.

(b) *Sponsor.* See No. 000003 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.410 of this chapter.

(d) *Conditions of use.* It is used in drinking water for replacement chickens as follows:

(1) *Amount.* 568.5 milligrams per gallon (0.015 percent).

(i) *Indications for use.* As a tranquilizer for flock treatment of chickens prior to handling.

(ii) *Limitations.* To be used one time as a treatment for replacement chickens up to 16 weeks of age; usual drinking water should be withheld prior to treatment to provide adequate consumption of medicated drinking water; not for use in laying chickens; chickens slaughtered within 72 hours following treatment must not be used for food.

(2) *Amount.* 2 to 4 milligrams per 2.2 pounds of body weight.

(i) *Indications for use.* As an aid in control of hysteria.

(ii) *Limitations.* To be used as a treatment for replacement chickens up to 16 weeks of age; usual drinking water should be withheld prior to treatment to provide adequate consumption of medicated drinking water; the drug should be administered at a dosage level of 4 milligrams per 2.2 pounds of body weight followed by 2 treatments at 4-day intervals of 2 milligrams per 2.2 pounds of body weight; not for use in laying chickens; chickens slaughtered within 72 hours following treatment must not be used for food.

21 CFR Ch. I (4–1–99 Edition)

§ 520.1430 Mibolerone.

(a) *Specifications.* Each milliliter contains 100 micrograms of mibolerone.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 30 micrograms for animals weighing 1 to 25 pounds; 60 micrograms for animals weighing 26 to 50 pounds; 120 micrograms for animals weighing 51 to 100 pounds; 180 micrograms for animals weighing over 100 pounds, German Shepherds, or German Shepherd mix.

(2) *Indications for use.* For the prevention of estrus (heat) in adult female dogs not intended primarily for breeding purposes.

(3) *Limitations.* Administer daily, orally or in a small amount of food, at least 30 days before expected initiation of heat, and continue daily as long as desired, but not for more than 24 months. Mibolerone should not be used in bitches before the first estrous period. It is not intended for animals being used primarily for breeding purposes. Use orally in adult female dogs only. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 15625, Apr. 14, 1978]

§ 520.1445 Milbemycin oxime tablets.

(a) *Specifications*—(1) *Dogs.* Each tablet contains 2.3, 5.75, 11.5, or 23.0 milligrams of milbemycin oxime.

(2) *Cats.* Each tablet contains 5.75, 11.5, or 23.0 milligrams of milbemycin oxime.

(b) *Sponsor.* See 058198 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Dogs and puppies*—(i) *Amount.* For hookworm, roundworm, and whipworm, use 0.23 milligram per pound of body weight (0.5 milligram per kilogram). For heartworm, use 0.05 milligram per pound of body weight (0.1 milligram per kilogram).

(ii) *Indications for use.* For prevention of heartworm disease caused by *Dirofilaria immitis*, control of hookworm infections caused by *Ancylostoma caninum*, and removal and control of adult roundworm infections caused by *Toxocara canis* and *Toxascaris leonina* and whipworm infections caused by

Food and Drug Administration, HHS

§ 520.1448a

Trichuris vulpis in dogs and in puppies 4 weeks of age or greater and 2 pounds of body weight or greater.

(iii) *Limitations*. Do not use in puppies less than 4 weeks of age and less than 2 pounds of body weight. Administer once a month. First dose given within 1 month after first exposure to mosquitoes and continue regular use until at least 1 month after end of mosquito season. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats and kittens*—(i) *Amount*. 0.91 milligram per pound of body weight (2.0 milligrams per kilogram).

(ii) *Indications for use*. For prevention of heartworm disease caused by *Dirofilaria immitis* and the removal of adult *Toxocara cati* (roundworm) and *Ancylostoma tubaeforme* (hookworm) infections in cats 6 weeks of age or greater and 1.5 pounds body weight or greater.

(iii) *Limitations*. Do not use in kittens less than 6 weeks of age or 1.5 pounds body weight. Administer once a month. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[55 FR 25301, June 21, 1990, as amended at 55 FR 49888, Dec. 3, 1990; 58 FR 5608, Jan. 22, 1993; 60 FR 50097, Sept. 28, 1995; 61 FR 43654, Aug. 26, 1996; 63 FR 29352, May 29, 1998; 63 FR 41189, Aug. 3, 1998]

§ 520.1446 Milbemycin oxime/lufenuron tablets.

(a) *Specifications*. Tablets containing: 2.3 milligrams milbemycin oxime/46 milligrams lufenuron, 5.75 milligrams/115 milligrams, 11.5 milligrams/230 milligrams, and 23 milligrams/460 milligrams.

(b) *Sponsor*. See No. 058198 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Dogs*—(i) *Amount*. 0.5 milligrams of milbemycin and 10 milligrams of lufenuron per kilogram of body weight.

(ii) *Indications for use*. For use in dogs and puppies for the prevention of heartworm disease caused by *Dirofilaria immitis*, for prevention and control of flea populations, control of adult *Ancylostoma caninum* (hookworm), and removal and control of adult *Toxocara canis*, *Toxascaris leonina* (roundworm),

and *Trichuris vulpis* (whipworm) infections. Lufenuron controls flea populations by preventing the development of flea eggs and does not kill adult fleas. Concurrent use of insecticides may be necessary for adequate control of adult fleas.

(iii) *Limitations*. Administer tablets once a month, preferably on the same date each time. All dogs in a household should be treated to achieve maximum efficacy. Do not use in dogs less than 4 weeks of age and less than 2 pounds body weight. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[62 FR 28629, May 27, 1997, as amended at 63 FR 41190, Aug. 3, 1998]

§ 520.1448 Monensin oral dosage forms.

Monensin, as the base or the sodium salt, contains a minimum of 90 percent monensin activity derived from monensin A and a minimum of 95 percent derived from monensin A plus B. Using thin layer chromatography, the R_f value must be comparable to a reference standard (the R_f value is the distance the spots travel from the starting line divided by the distance the solvent front travels from the starting line). The loss on drying is not more than 10 percent when dried in vacuum at 60 °C for 2 hours.

[55 FR 3586, Feb. 2, 1990]

§ 520.1448a Monensin blocks.

(a)(1) *Specifications*. Each pound of protein-mineral block contains 400 milligrams of monensin (0.088 percent) as monensin sodium.

(2) *Sponsor*. See 036904 in § 510.600(c) of this chapter.

(3) *Related tolerances*. See § 556.420 of this chapter.

(4) *Conditions of use*—(i) *Amount*. 80 to 200 milligrams of monensin (0.2 to 0.5 pound of block) per head per day.

(ii) *Indications for use*. Increased rate of weight gain.

(iii) *Limitations*. Block to be fed free choice to pasture cattle (slaughter, stocker, feeder, and dairy and beef replacement heifers). Provide at least 1 block per 5 head of cattle. Feed blocks continuously. Do not feed salt or minerals containing salt. Do not allow

§ 520.1450

horses or other equines access to formulations containing monensin (ingestion of monensin by equines has been fatal). The effectiveness of this block in cull cows and bulls has not been established.

(b) [Reserved]

(c)(1) *Specifications.* Each pound of protein block contains 175 milligrams of monensin (0.038 percent) as monensin sodium.

(2) *Sponsor.* See 021676 in § 510.600(c) of this chapter.

(3) *Related tolerances.* See § 556.420 of this chapter.

(4) *Conditions of use*—(i) *Amount.* 40 to 200 milligrams of monensin (0.25 to 1.13 pounds or 4 to 18 ounces of block) per head per day.

(ii) *Indications for use.* Increased rate of weight gain.

(iii) *Limitations.* Blocks to be fed free choice to pasture cattle (slaughter, stocker, and feeder). Provide at least 1 block per 4 head of cattle. Do not allow cattle access to salt or mineral while being fed this product. Ingestion by cattle of monensin at levels of 600 milligrams per head per day and higher has been fatal. Do not allow horses or other equines access to formulations containing monensin (ingestion of monensin by equines has been fatal). Block's effectiveness in cull cows and bulls has not been established.

(d)(1) *Specifications.* Each pound of block contains 400 milligrams of monensin (0.088 percent) as monensin sodium.

(2) *Sponsor.* See 051267 in § 510.600(c) of this chapter.

(3) *Related tolerances.* See § 556.420 of this chapter.

(4) *Conditions of use*—(i) *Amount.* 50 to 200 milligrams of monensin (2 to 8 ounces of block) per head per day.

(ii) *Indications for use.* Pasture cattle: Increased rate of weight gain.

(iii) *Limitations.* Blocks to be fed free choice to pasture cattle (slaughter, stocker, feeder, and dairy and beef replacement heifers). Provide at least one block per five head of cattle. Feed blocks continuously. Do not feed salt or mineral supplements in addition to the blocks. Ingestion by cattle of monensin at levels of 600 milligrams per head per day and higher has been fatal. Do not allow horses or other equines access to

21 CFR Ch. I (4–1–99 Edition)

formulations containing monensin (ingestion of monensin by equines has been fatal). The effectiveness of this block in cull cows and bulls has not been established.

[46 FR 19466, Mar. 31, 1981]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 520.1448a, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 520.1450 Morantel tartrate oral dosage forms.

§ 520.1450a Morantel tartrate bolus.

(a) *Specifications.* Each bolus contains 2.2 grams morantel tartrate equivalent to 1.3 grams of morantel base.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.425 of this chapter.

(d) *Conditions of use*—(1) *Amount.* One bolus per 500 pounds of body weight (4.4 milligrams per pound of body weight) as a single oral dose. Boluses may be divided in half for more accurate dosing as follows: up to 325 pounds, ½ bolus; 326 to 600 pounds, 1 bolus; 601 to 900 pounds, 1½ boluses; and 901 to 1,200 pounds, 2 boluses.

(2) *Indications for use.* For removal and control of mature gastrointestinal nematode infections of cattle including stomach worms (*Haemonchus* spp., *Ostertagia* spp., *Trichostrongylus* spp.), worms of the small intestine (*Cooperia* spp., *Trichostrongylus* spp., *Nematodirus* spp.), and worms of the large intestine (*Oesophagostomum radiatum*).

(3) *Limitations.* Conditions of constant worm exposure may require retreatment in 2 to 4 weeks. Consult your veterinarian before administering to severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism. Do not treat within 14 days of slaughter.

[46 FR 50949, Oct. 16, 1981. Redesignated at 49 FR 47831, Dec. 7, 1984, and amended at 51 FR 9005, Mar. 17, 1986]

§ 520.1450b Morantel tartrate cartridge.

(a) *Specifications.* The drug product consists of a stainless-steel cylinder having both ends closed with polyethylene diffusing discs and containing a morantel tartrate paste. The paste

Food and Drug Administration, HHS

§ 520.1452

contains 22.7 grams of morantel tartrate equivalent to 13.5 grams of morantel base.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.425 of this chapter.

(d) *Conditions of use—(1) Amount.* Grazing cattle: Administer 1 cartridge to each animal at the start of the grazing season.

(2) *Indications for use.* For control of the adult stage of the following gastrointestinal nematode infections in weaned calves and yearling cattle weighing a minimum of 200 pounds: *Ostertagia* spp., *Trichostrongylus axei*, *Cooperia* spp., and *Oesophagostomum radiatum*.

(3) *Limitations.* Administer orally with the dosing gun to all cattle that will be grazing the same pasture. Effectiveness of the drug product is dependent upon continuous control of the gastrointestinal parasites for approximately 90 days following administration. Therefore, treated cattle should not be moved to pastures grazed in the same grazing season/calendar year by untreated cattle. Do not administer to cattle within 106 days of slaughter. Consult your veterinarian before administering to severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.

[49 FR 47831, Dec. 7, 1984, as amended at 51 FR 23415, June 27, 1986; 51 FR 41081, Nov. 13, 1986]

§ 520.1450c Morantel tartrate sustained-release trilaminar cylinder/sheet.

(a) *Specifications.* The drug product consists of a trilaminar, perforated, plastic sheet formed into a cylinder having plastic plugs in its ends. The core lamina contains 19.8 grams of morantel tartrate equivalent to 11.8 grams of morantel base.

(b) *Sponsor.* See 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.425 of this chapter.

(d) *Conditions of use—(1) Amount.* Grazing cattle: Administer 1 cartridge to each animal at the start of the grazing season.

(2) *Indications for use.* For control of the adult stage of the following gastrointestinal nematode infections in weaned calves and yearling cattle weighing a minimum of 200 pounds: *Ostertagia* spp., *Trichostrongylus axei*, *Cooperia* spp., and *Oesophagostomum radiatum*.

(3) *Limitations.* Administer orally with the dosing gun to all cattle that will be grazing the same pasture. Effectiveness of the drug product is dependent upon continuous control of the gastrointestinal parasites for approximately 90 days following administration. Therefore, treated cattle should not be moved to pastures grazed in the same grazing season/calendar year by untreated cattle. Do not administer to cattle within 102 days of slaughter. Consult your veterinarian before administering to severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.

[56 FR 13396, Apr. 2, 1991]

§ 520.1451 Moxidectin.

(a) *Specifications.* Each tablet contains 30, 68, or 136 micrograms of moxidectin.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use—(1) Amount.* 3 micrograms per kilogram (1.36 micrograms per pound) of body weight.—

(2) *Indications for use.* To prevent infection by the canine heartworm *Dirofilaria immitis* and the subsequent development of canine heartworm disease.

(3) *Limitations.* Use once-a-month in dogs at 8 weeks of age or older. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[62 FR 37713, July 15, 1997]

§ 520.1452 Moxidectin gel.

(a) *Specifications.* The gel contains 2 percent moxidectin (20 milligrams per milliliter).

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) [Reserved]

§ 520.1468

(d) *Conditions of use*—(1) *Amount*. 0.4 milligram moxidectin per kilogram (2.2 pounds) of body weight.

(2) *Indications for use*. Horses and ponies for treatment and control of large strongyles (*Strongylus vulgaris* (adults and L4/L5 arterial stages), *S. edentatus* (adult and tissue stages), *Triodontophorus brevicauda* (adults), *T. serratus* (adults)); small strongyles (*Cyathostomum* spp. (adults), *Cylicocycclus* spp. (adults), *Cylicostephanus* spp. (adults), *Gyalocephalus capitatus* (adults), undifferentiated luminal larvae); encysted cyathostomes (late L3 and L4 mucosal cyathostome larvae); ascarids (*Parascaris equorum* (adults and L4 larval stages)); pinworms (*Oxyuris equi* (adults and L4 larval stages)), hairworms (*Trichostrongylus axei* (adults)), large-mouth stomach worms (*Habronema muscae* (adults)), and horse stomach bots (*Gasterophilus intestinalis* (2nd and 3rd instars)). One dose also suppresses small strongyle egg production for 84 days.

(3) *Limitations*. For horses and ponies including breeding mares and stallions. Not for use in horses and ponies intended for food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[62 FR 42902, Aug. 11, 1997]

§ 520.1468 Naproxen granules.

(a) *Specifications*. Naproxen granules contain 50 percent naproxen.

(b) *Sponsor*. No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Horses*. The drug is used for the relief of inflammation and associated pain and lameness exhibited with arthritis, as well as myositis and other soft tissue diseases of the musculoskeletal system of the horse.

(2)(i) For oral maintenance therapy following initial intravenous dosage, administer 10 milligrams naproxen per kilogram of animal body weight twice daily as top dressing in the animal's feed for up to 14 consecutive days. The initial intravenous dosage is 5 milligrams per kilogram of body weight.

(ii) For oral dosage only, administer 10 milligrams naproxen per kilogram of animal body weight twice daily as a

21 CFR Ch. I (4–1–99 Edition)

top dressing in the animal's feed for up to 14 consecutive days.

(3) Not for use in horses intended for food.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 14188, Apr. 2, 1976, as amended at 51 FR 24525, July 7, 1986; 61 FR 5506, Feb. 13, 1996]

§ 520.1484 Neomycin sulfate soluble powder.

(a) *Specifications*. The drug contains 20.3 grams of neomycin sulfate per ounce which is equivalent to 14.2 grams of neomycin base.

(b) *Sponsors*. See Nos. 000009, 000069, 046573, 050604, and 051259 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 10 milligrams of neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days.

(2) *Indications for use*. For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep, and goats.

(3) *Limitations*. Add to drinking water or milk; not for use in liquid supplements. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days. Discontinue treatment prior to slaughter as follows: Cattle (not for use in veal calves), 1 day; sheep, 2 days; swine and goats, 3 days.

[57 FR 19085, May 4, 1992, as amended at 57 FR 26604, June 15, 1992; 60 FR 14217, Mar. 16, 1995; 61 FR 31027, June 19, 1996; 61 FR 31399, June 20, 1996; 62 FR 55160, Oct. 23, 1997; 63 FR 17329, Apr. 9, 1998]

§ 520.1485 Neomycin sulfate oral solution.

(a) *Specifications*. Each milliliter contains 200 milligrams of neomycin sulfate (equivalent to 140 milligrams of neomycin base).

(b) *Sponsors*. See Nos. 000009, 050604, and 059130 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.430 of this chapter.

(d) *Conditions of use*—(1) *Amount*. 10 milligrams of neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days.

(2) *Indications for use*. For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin in cattle (excluding veal calves), swine, sheep, and goats.

(3) *Limitations*. Administer undiluted or in drinking water. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days. Discontinue treatment prior to slaughter as follows: 1 day for cattle, 2 days for sheep, and 3 days for swine and goats.

[58 FR 38972, July 21, 1993, as amended at 60 FR 3079, Jan. 13, 1995; 61 FR 31398, June 20, 1996; 62 FR 60657, Nov. 12, 1997; 63 FR 45944, Aug. 28, 1998]

§ 520.1616 Orbifloxacin tablets.

(a) *Specifications*. Each tablet contains 5.7, 22.7, or 68 milligrams of orbifloxacin.

(b) *Sponsor*. See No. 000061 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Dogs and cats*—(i) *Amount*. 2.5 to 7.5 milligrams per kilogram body weight.

(ii) *Indications for use*. For management of diseases associated with bacteria susceptible to orbifloxacin.

(iii) *Limitations*. Administer orally once daily for 2 to 3 days beyond cessation of clinical signs for up to a maximum of 30 days. If no improvement is seen within 5 days, diagnosis should be reevaluated and a different course of therapy considered. Orbifloxacin is contraindicated in immature dogs and cats during the rapid growth phase. Orbifloxacin and other quinolones have been shown to cause arthropathy in immature animals of most species. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[62 FR 29012, May 29, 1997, as amended at 62 FR 61627, Nov. 19, 1997]

§ 520.1628 Oxfendazole powder and pellets.

(a) *Specifications*—(1) *Powder for suspension*. Each gram of powder contains 7.57 percent oxfendazole.

(2) *Pellets*. Each gram of pellets contains 6.49 percent oxfendazole.

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 10 milligrams per kilogram of body weight.

(2) *Indications for use*. The drug is used in horses for removal of the following gastrointestinal worms: Large roundworms (*Parascaris equorum*), mature and immature pinworms (*Oxyuris equi*), large strongyles (*Strongylus edentatus*, *Strongylus vulgaris*, and *Strongylus equinus*), and small strongyles.

(3) *Limitations*—(i) *Powder for suspension*. For gravity administration via stomach tube or for positive administration via stomach tube and dose syringe. Discard unused portions of suspension after 24 hours. Mix drug according to directions prior to use. Administer drug with caution to sick or debilitated horses. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(ii) *Pellets*. The drug is given by sprinkling on the grain portion of the ration. Withholding feed or water prior to administration is not necessary. Administer drug with caution to sick or debilitated horses. Not for use in horses intended for food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[44 FR 35211, June 19, 1979, as amended at 46 FR 26301, May 12, 1981; 46 FR 60570, Dec. 11, 1981; 49 FR 28549, July 13, 1984; 61 FR 5506, Feb. 13, 1996]

§ 520.1629 Oxfendazole paste.

(a)(1) *Specifications*. Each gram of paste contains 0.375 gram oxfendazole (37.5 percent).

(2) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(3) *Conditions of use*—(i) *Amount*. 10 milligrams per kilogram (2.2 pounds) of body weight.

§ 520.1630

21 CFR Ch. I (4–1–99 Edition)

(ii) *Indications for use.* The drug is used in horses for removal of the following gastrointestinal worms: Large roundworms (*Parascaris equorum*), mature and 4th stage larvae pinworms (*Oxyuris equi*), large strongyles (*Strongylus edentatus*, *S. vulgaris*, and *S. equinus*), and small strongyles.

(iii) *Limitations.* Horses maintained on premises where reinfection is likely to occur should be retreated in 6 to 8 weeks. Withholding feed or water prior to use is unnecessary. Administer drug with caution to sick or debilitated horses. Not for use in horses intended for food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(b)(1) *Specifications.* Each gram of paste contains 185 milligrams of oxfendazole (18.5 percent).

(2) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(3) *Related tolerances.* See § 556.495 of this chapter.

(4) *Conditions of use*—(i) *Amount.* 4.5 milligrams per kilogram of body weight (2.05 milligrams per pound).

(ii) *Indications for use.* The drug is used in cattle for the removal and control of the following worms: lungworms (*Dictyocaulus viviparus*—adult, L4); stomach worms: barberpole worms (*Haemonchus contortus* and *H. placei*—adult), small stomach worms (*Trichostrongylus axei*—adult), brown stomach worms (*Ostertagia ostertagi*—adult, L4, inhibited L4); intestinal worms; nodular worms (*Oesophagostomum radiatum*—adult), hookworms (*Bunostomum phlebotomum*—adult), small intestinal worms (*Cooperia punctata*, *C. oncophora*, and *C. mcmasteri*—adult, L4); and tapeworms (*Moniezia benedeni*—adult).

(iii) *Limitations.* For use in cattle only. Treatment may be repeated in 4 to 6 weeks. Cattle must not be slaughtered until 11 days after treatment. Do not use in female dairy cattle of breeding age. Consult a veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[49 FR 38250, Sept. 28, 1984, as amended at 58 FR 39443, July 23, 1993; 61 FR 5506, Feb. 13, 1996]

§ 520.1630 Oxfendazole suspension.

(a) *Specifications.* Each milliliter contains 90.6 or 225.0 milligrams oxfendazole (9.06 or 22.5 percent).

(b) *Sponsor.* See 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.495 of this chapter.

(d) *Conditions of use*—(1) *Horses* (9.06 percent suspension only).

(i) *Amount.* 10 milligrams per kilogram (2.2 pounds) of body weight.

(ii) *Indications for use.* For removal of large roundworms (*Parascaris equorum*), mature and 4th stage larvae pinworms (*Oxyuris equi*), large strongyles (*Strongylus edentatus*, *S. vulgaris*, and *S. equinus*), and small strongyles.

(iii) *Limitations.* Administer 9.06 percent suspension by stomach tube or dose syringe. Horses maintained on premises where reinfection is likely to occur should be retreated in 6 to 8 weeks. Withholding feed or water prior to use is unnecessary. Administer drug with caution to sick or debilitated horses. Do not use in horses intended for food. If administered by stomach tube: Federal law restricts this drug to use by or on the order of a licensed veterinarian. If administered by dose syringe only: Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) *Cattle.* (9.06 or 22.5 percent suspension). (i) *Amount.* 4.5 milligrams per kilogram of body weight (2.05 milligrams per pound).

(ii) *Indications for use.* For the removal and control of: lungworms (*Dictyocaulus viviparus*—adult, L4); stomach worms: barberpole worms (*Haemonchus contortus* and *H. placei*—adult), small stomach worms (*Trichostrongylus axei*—adult), brown stomach worms (*Ostertagia ostertagi*—adult, L4, inhibited L4); intestinal worms; nodular worms (*Oesophagostomum radiatum*—adult), hookworms (*Bunostomum phlebotomum*—adult), small intestinal worms (*Cooperia punctata*, *C. oncophora*, and *C. mcmasteri*—adult, L4), and tapeworms (*Moniezia benedeni*—adult).

(iii) *Limitations.* For use in cattle only. Administer 9.06 percent suspension orally only with a dose syringe, and 22.5 percent suspension either orally with a dose syringe or

intraruminally with a rumen injector. Treatment may be repeated in 4 to 6 weeks. Cattle must not be slaughtered until 7 days after treatment. Do not use in lactating dairy cattle. For use of 9.06 percent suspension orally: Consult a veterinarian for assistance in the diagnosis, treatment, and control of parasitism. For use of 22.5 percent suspension orally or intraruminally: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[55 FR 46943, Nov. 8, 1990, as amended at 56 FR 8710, Mar. 1, 1991; 61 FR 5506, Feb. 13, 1996]

§ 520.1631 Oxfendazole and trichlorfon paste.

(a) *Specifications.* Each gram of paste contains 28.5 milligrams oxfendazole and 454.5 milligrams trichlorfon.

(b) *Sponsor.* See 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 2.5 milligrams of oxfendazole and 40 milligrams of trichlorfon per kilogram of body weight.

(2) *Indications for use.* The drug is used in horses for removal of bots (*Gasterophilus intestinalis*, 2nd and 3rd instars; *G. nasalis*, 3rd instar) and the following gastrointestinal worms: Large roundworms (*Parascaris equorum*), pinworms (*Oxyuris equi*), adult and 4th stage larvae; large strongyles (*Strongylus edentatus*, *S. vulgaris*, and *S. equinus*); and small strongyles.

(3) *Limitations.* Horses maintained on premises where reinfection is likely to occur should be retreated in 6 to 8 weeks. Withholding feed or water before use is unnecessary. Administer with caution to sick or debilitated horses. Not for use in horses intended for food. Do not administer to mares during the last month of pregnancy. Trichlorfon is a cholinesterase inhibitor. Do not use this product in animals simultaneously with, or within a few days before or after treatment with or exposure to, cholinesterase-inhibiting drugs, pesticides, or chemicals. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[50 FR 50291, Dec. 10, 1985, as amended at 61 FR 5506, Feb. 13, 1996]

§ 520.1638 Oxibendazole paste.

(a) *Specifications.* The paste contains 22.7 percent oxibendazole.

(b) *Sponsor.* See 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use in horses*—(1) *Amount.* For uses other than for threadworms (*Strongyloides westeri*), 10 milligrams of oxibendazole per kilogram of body weight; for threadworms (*Strongyloides westeri*), 15 milligrams per kilogram.

(2) *Indications for use.* For removal and control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*); small strongyles (genera *Cylicostephanus*, *Cylicocyclus*, *Cyathostomum*, *Triodontophorus*, *Cylicodontophorus*, and *Gyalocephalus*); large roundworms (*Parascaris equorum*); pinworms (*Oxyuris equi*) including various larval stages; and threadworms (*Strongyloides westeri*).

(3) *Limitations.* Administer orally by syringe. Horses maintained on premises where reinfection is likely to occur should be re-treated in 6 to 8 weeks. Not for use in horses intended for food. Consult a veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[46 FR 50948, Oct. 16, 1981, as amended at 47 FR 36418, Aug. 20, 1982; 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.1640 Oxibendazole suspension.

(a) *Specifications.* The suspension contains 10 percent oxibendazole.

(b) *Sponsor.* See 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use in horses*—(1) *Amount.* For use other than threadworms (*Strongyloides westeri*), 10 milligrams of oxibendazole per kilogram of body weight; for threadworms, 15 milligrams per kilogram of body weight.

(2) *Indications for use.* For removal and control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*); small strongyles (species of the genera *Cylicostephanus*, *Cylicocyclus*, *Cyathostomum*, *Triodontophorus*, *Cylicodontophorus*, and *Gyalocephalus*); large roundworms (*Parascaris equorum*); pinworms (*Oxyuris equi*) including various larval stages; and threadworms (*Strongyloides westeri*).

§ 520.1660

21 CFR Ch. I (4–1–99 Edition)

(3) *Limitations.* Administer by stomach tube in 3 to 4 pints of warm water, or by top dressing or mixing into a portion of the normal grain ration. Prepare individual doses to ensure that each animal receives the correct amount. Horses maintained on premises where reinfection is likely to occur should be re-treated in 6 to 8 weeks. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 78119, Nov. 28, 1980, as amended at 47 FR 39812, Sept. 10, 1982; 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.1660 Oxytetracycline.

§ 520.1660a Oxytetracycline and carbomycin in combination.

(a) *Specifications.* (1) Oxytetracycline: The antibiotic substance produced by growth of *Streptomyces rimosus* or the same antibiotic substance produced by any other means.

(2) Carbomycin: The antibiotic substance produced by growth of *Streptomyces halstedii* or the same antibiotic substance produced by any other means.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Special considerations.* The quantities of oxytetracycline in paragraph (e) of this section refer to the activity of oxytetracycline hydrochloride and the quantities of carbomycin listed refer to the activity of an appropriate standard.

(d) *Related tolerances.* See §§ 556.110 and 556.500 of this chapter.

(e) *Conditions of use.* It is used as oxytetracycline hydrochloride plus carbomycin base in drinking water of chickens as follows:

(1) *Amount.* 1.0 gram of oxytetracycline and 1.0 gram carbomycin per gallon.

(2) *Indications for use.* As an aid in the prevention and treatment of complicated chronic respiratory disease (air-sac infection) caused by *Mycoplasma gallisepticum* and secondary bacterial organisms associated with chronic respiratory disease such as *E. coli*.

(3) *Limitations.* Administer for not more than 5 days; not for use in chick-

ens producing eggs for human consumption; withdraw 24 hours before slaughter.

§ 520.1660b Oxytetracycline hydrochloride capsules.

(a) *Specifications.* The drug is in capsule form with each capsule containing 125 or 250 milligrams of oxytetracycline hydrochloride. Oxytetracycline is the antibiotic substance produced by growth of *Streptomyces rimosus* or the same antibiotic substance produced by any other means.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in dogs and cats for the treatment of bacterial pneumonia caused by *Brucella bronchiseptica*, tonsillitis caused by *Streptococcus hemolyticus*, bacterial enteritis caused by *Escherichia coli*, urinary tract infections caused by *Escherichia coli*, and wound infections caused by *Staphylococcus aureus*.¹

(2) The drug is administered orally to dogs and cats at a dosage level of 25–50 milligrams per pound of body weight per day in divided doses at 12-hour intervals. The drug can be used for continuation of compatible antibiotic therapy following parenteral oxytetracycline administration where rapidly attained, sustained antibiotic blood levels are required. The duration of treatment required to obtain favorable response will depend to some extent on the severity and degree of involvement and the susceptibility of the infectious agent. Clinical response to antibiotic therapy usually occurs within 48 to 72 hours. If improvement is not observed within that period, the diagnosis and course of treatment should be reconsidered. To assure adequate treatment, administration of the drug should continue for at least 48 hours following favorable clinical response.¹

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

§ 520.1660c Oxytetracycline hydrochloride tablets/boluses.

(a) *Specifications.* Each tablet or bolus contains 250, 500, or 1,000 milligrams of oxytetracycline hydrochloride.

(b) *Sponsors.* For sponsors in § 510.600(c) of this chapter: See 000010 for use of 500 and 1,000 milligram boluses. See 000069 for use of 250 and 500 milligram tablets.

(c) *Tolerances.* See § 556.500 of this chapter.

(d) *Conditions of use in beef and dairy cattle—*(1)(i) *Amount.* 250 milligrams per 100 pounds of body weight every 12 hours (5 milligrams per pound of body weight daily in two doses).

(ii) *Indications for use.* For control of bacterial enteritis caused by *Salmonella typhimurium* and *Escherichia coli* (colibacillosis) and bacterial pneumonia (shipping fever complex, pasteurellosis) caused by *Pasteurella multocida*.

(2)(i) *Amount.* 500 milligrams per 100 pound of body weight every 12 hours (10 milligrams per pound of body weight daily in two doses).

(ii) *Indications for use.* For treatment of bacterial enteritis caused by *Salmonella typhimurium* and *Escherichia coli* (colibacillosis) and bacterial pneumonia (shipping fever complex, pasteurellosis) caused by *Pasteurella multocida*.

(3) *Limitations.* Dosage should continue until the animal returns to normal and for 24 hours to 48 hours after symptoms have subsided. Treatment should not exceed 4 consecutive days. Do not exceed 500 milligrams per 100 pounds of body weight every 12 hours (10 milligrams per pound daily). For sponsor 000069: Discontinue treatment 7 days prior to slaughter. Not for use in lactating dairy cattle.

[46 FR 32440, June 23, 1981, as amended at 50 FR 1045, Jan. 9, 1985; 63 FR 70334, Dec. 21, 1998]

§ 520.1660d Oxytetracycline hydrochloride soluble powder.

(a) *Specifications.* The drug is a soluble powder distributed in packets or pails having several concentrations of oxytetracycline hydrochloride (independent of the various net weights) as follows:

(1) Each 18.14 grams of powder contains 1 gram of oxytetracycline hydrochloride (OTC HCl) (packets: 4, 6.4, and 16 oz.);

(2) Each 4.43 grams of powder contains 1 gram of OTC HCl (packets: 4 and 16 oz.);

(3) Each 1.32 grams of powder contains 1 gram of OTC HCl (packets: 2.39, 4.78, and 9.55 oz.; jars: 2.25 lbs.; and pails: 4.5 lbs.);

(4) Each 2.73 grams of powder contains 1 gram of OTC HCl (packets: 2.46 and 9.87 oz; pail: 3.09 lb).

(5) Each 4.2 grams of powder contains 1 gram of OTC HCl (packets: 3.8 and 15.2 oz; pails: 4.74 and 23.7 lb).

(6) Each 1.32 grams of powder contains 1 gram of OTC HCl (packet: 4.78 oz.).

(7) Each 18.1 grams of powder contains 1 gram of OTC HCl (pails: 2 and 5 lb), each 272.2 grams (9.6 oz) of powder contains 204.8 grams of OTC HCl, each 907.2 grams (2 lb) of powder contains 686 grams of OTC HCl, each 2.26 kilograms (5 lb) of powder contains 1,715 grams of OTC HCl.

(8) Each 135.5-gram packet (4.78 ounce) contains 102.4 grams of OTC HCl. Each 677.5-gram packet (23.9 ounce) contains 512 grams of OTC HCl.

(9) Each 2.73 grams of powder contains 1 gram of OTC HCl (packets: 9.87 and 19.75 oz; pails: 5 lb).

(b) *Sponsor.* See sponsor numbers in § 510.600(c) of this chapter as follows:

(1) No. 000069 for use of OTC HCl concentrations in paragraphs (a)(1), (a)(2), and (a)(3) of this section in chickens, turkeys, swine, cattle, sheep, and honey bees.

(2) No. 017144 for use of OTC HCl concentration in paragraph (a)(4) of this section in chickens, turkeys, and swine.

(3) No. 000010 for use of OTC HCl concentration in paragraph (a)(5) of this section in turkeys and chickens.

(4) No. 057561 for use of OTC HCl concentration in paragraph (a)(6) of this section in chickens, turkeys, and swine.

(5) No. 059130 for use of OTC HCl concentration in paragraph (a)(7) of this section in chickens, turkeys, swine, cattle, and sheep.

(6) No. 053389 for use of OTC HCl concentrations in paragraph (a)(8) of this

§ 520.1660d

21 CFR Ch. I (4–1–99 Edition)

section in chickens, turkeys, swine, cattle, and sheep.

(7) No. 050604 for use of OTC HCl concentration in paragraph (a)(9) of this section in chickens, turkeys, and swine.

(c) *Related tolerances.* See § 556.500 of this chapter.

(d) *Conditions of use.* (1) It is used in drinking water as follows:

(i) *Chickens—(A)(1) Amount per gallon.* 200 to 400 milligrams.

(2) *Indications for use.* Control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to oxytetracycline.

(3) *Limitations.* Prepare a fresh solution daily. Administer 7 to 14 days. Not to be used for more than 14 consecutive days. Use as sole source of drinking water. Do not use in birds producing eggs for human consumption.

(B)(1) *Amount per gallon.* 400 to 800 milligrams.

(2) *Indications for use.* Control of chronic respiratory disease (CRD) and air sac infections caused by *Mycoplasma gallisepticum* and *E. coli* susceptible to oxytetracycline; control of fowl cholera caused by *Pasteurella multocida* susceptible to oxytetracycline.

(3) *Limitations.* Prepare a fresh solution daily. Administer 7 to 14 days. Not to be used for more than 14 consecutive days. Use as sole source of drinking water. Do not use in birds producing eggs for human consumption.

(ii) *Turkeys—(A)(1) Amount per gallon.* 200 to 400 milligrams.

(2) *Indications for use.* Control of hexamitiasis caused by *Hexamita meleagridis* susceptible to oxytetracycline.

(3) *Limitations.* Prepare a fresh solution daily. Administer 7 to 14 days. Not to be used for more than 14 consecutive days. Use as sole source of drinking water. Do not use in birds producing eggs for human consumption. Withdraw 5 days prior to slaughter those products sponsored by Nos. 000069, 017144, 057561, and 059130 in § 510.600(c) of this chapter. Withdraw 4 days prior to slaughter those products sponsored by No. 000010. Zero-day withdrawal for those products sponsored by No. 053389.

(B)(1) *Amount per gallon.* 400 milligrams.

(2) *Indications for use.* Control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to oxytetracycline.

(3) *Limitations.* Prepare a fresh solution daily. Administer 7 to 14 days. Not to be used for more than 14 consecutive days. Use as sole source of drinking water. Do not use in birds producing eggs for human consumption. Withdraw 5 days prior to slaughter those products sponsored by Nos. 000069, 017144, 057561, and 059130 in § 510.600(c) of this chapter. Withdraw 4 days prior to slaughter those products sponsored by No. 000010. Zero-day withdrawal for those products sponsored by No. 053389.

(C)(1) *Amount.* 25 milligrams per pound of body weight.

(2) *Indications for use.* Growing turkeys. Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline.

(3) *Limitations.* Prepare a fresh solution daily. Administer 7 to 14 days. Not to be used for more than 14 consecutive days. Use as sole source of drinking water. Do not use in birds producing eggs for human consumption. Withdraw 5 days prior to slaughter those products sponsored by Nos. 000069, 017144, 057561, and 059130 in § 510.600(c) of this chapter. Withdraw 4 days prior to slaughter those products sponsored by No. 000010. Zero-day withdrawal for those products sponsored by No. 053389.

(iii) *Swine—(A) Amount.* 10 milligrams per pound of body weight daily.

(B) *Indications for use.* Control and treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to oxytetracycline. For breeding swine: Control and treatment of leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by *Leptospira pomona* susceptible to oxytetracycline.

(C) *Limitations.* Prepare a fresh solution daily. Use as sole source of OTC. Administer up to 14 days; do not use for more than 14 consecutive days; withdraw 5 days prior to slaughter those products sponsored by Nos. 000069 and 059130. Administer up to 5 days; do not use for more than 5 consecutive days;

Food and Drug Administration, HHS

§ 520.1696b

withdraw 13 days prior to slaughter those products sponsored by Nos. 017144 and 057561.

(iv) *Calves, beef cattle, and nonlactating dairy cattle*—(A) *Amount*. 10 milligrams per pound of body weight daily.

(B) *Indications for use*. Control and treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia (shipping fever complex) caused by *P. multocida* susceptible to oxytetracycline.

(C) *Limitations*. Prepare a fresh solution daily. Administer up to 14 days. Do not use for more than 14 consecutive days. Use as sole source of oxytetracycline. Do not administer this product with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. Withdraw 5 days prior to slaughter. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

(v) *Sheep*—(A) *Amount*. 10 milligrams per pound of body weight daily.

(B) *Indications for use*. Control and treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia (shipping fever complex) caused by *P. multocida* susceptible to oxytetracycline.

(C) *Limitations*. Prepare a fresh solution daily. Administer up to 14 days. Do not use for more than 14 consecutive days. Use as sole source of oxytetracycline. Withdraw 5 days prior to slaughter.

(2) It is used in the food of honey bees as follows:

(i) *Amount*. 200 milligrams per colony, administered via either a 1:1 sugar syrup (equal parts of sugar and water weight to weight) or dusting with a powdered sugar mixture.

(ii) *Indications for use*. Control and treatment of American and European foul brood caused by *Bacillus larvae* susceptible to oxytetracycline.

(iii) *Limitations*. The drug is administered in 3 applications of sugar syrup or 3 dustings at 4- to 5-day intervals. The drug should be fed early in the spring or fall and consumed by the bees

before main honey flow begins to avoid contamination of production honey. Remove at least 6 weeks prior to main honey flow.

[50 FR 32694, Aug. 14, 1985, as amended at 52 FR 48094, Dec. 18, 1987; 54 FR 18280, Apr. 28, 1989; 57 FR 42490, Sept. 15, 1992; 58 FR 42854, Aug. 12, 1993; 58 FR 43794, Aug. 18, 1993; 58 FR 3318, Jan. 21, 1994; 59 FR 41664, Aug. 15, 1994; 59 FR 56388, Nov. 14, 1994; 60 FR 40455, Aug. 9, 1995; 61 FR 2915, Jan. 30, 1996; 61 FR 52690, Oct. 8, 1996; 62 FR 23356, Apr. 30, 1997; 62 FR 35076, June 30, 1997; 63 FR 44384, Aug. 19, 1998; 64 FR 10389, Mar. 4, 1999; 64 FR 13341, Mar. 18, 1999; 64 FR 13678, Mar. 22, 1999]

§ 520.1696 Penicillin oral dosage forms.

§ 520.1696a Buffered penicillin powder, penicillin powder with buffered aqueous diluent.

(a) *Specifications*. When reconstituted, each milliliter contains penicillin G procaine equivalent to 20,000, 25,000, 40,000, 50,000, 80,000, or 100,000 units of penicillin G.

(b) *Sponsor*. [Reserved]

(c) *Related tolerances*. See § 556.510 of this chapter.

(d) *Conditions of use. Chickens*—It is used in drinking water as follows:

(1) *Amount*. 100,000 units per gallon.

(i) *Indications for use*. Treatment of chronic respiratory disease (air-sac infection) and bluecomb (nonspecific infectious enteritis).

(ii) *Limitations*. As penicillin G procaine; not for use in laying chickens; prepare fresh solution daily; withdraw 1 day before slaughter; as sole source of penicillin.

(2) *Amount*. 50,000 to 100,000 units per gallon.

(i) *Indications for use*. Prevention of chronic respiratory disease (air-sac infection) and bluecomb (nonspecific infectious enteritis).

(ii) *Limitations*. As penicillin G procaine; not for use in laying chickens; prepare fresh solution daily; withdraw 1 day before slaughter; as sole source of penicillin.

[57 FR 37326, Aug. 18, 1992]

§ 520.1696b Penicillin G potassium in drinking water.

(a) *Specifications*. When reconstituted, each milliliter contains penicillin G potassium equivalent to 20,000, 25,000,

§ 520.1696c

40,000, 50,000, 80,000, or 100,000 units of penicillin G.

(b) *Sponsors*. See Nos. 017144, 046573, 050604, and 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use. Turkeys*—(1) *Amount*. 1,500,000 units per gallon drinking water for 5 days.

(2) *Indications for use*. Treatment of erysipelas caused by *Erysipelothrix rhusiopathiae*.

(3) *Limitations*. Prepare concentrated stock solution for use with medication proportioners fresh every 24 hours. Prepare recommended use levels for gravity flow watering system fresh every 12 hours. For best results, treatment should be started at the first sign of infection. Discontinue treatment at least 1 day prior to slaughter. Not for use in turkeys producing eggs for human consumption.

[57 FR 37326, Aug. 18, 1992, as amended at 59 FR 42493, Aug. 18, 1994; 60 FR 26359, May 17, 1995; 62 FR 55160, Oct. 23, 1997]

§ 520.1696c Penicillin V potassium for oral solution.

(a) *Specifications*. When reconstituted, each milliliter contains 25 milligrams (40,000 units) of penicillin V.

(b) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(c) *National Academy of Sciences/National Research Council (NAS/NRC) status*. The conditions of use were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use. Dogs and cats*—(1) *Amount*. 10 to 15 milligrams per pound of body weight every 6 to 8 hours.

(2) *Indications for use*. Treatment of respiratory, urogenital, skin, and soft tissue infections and septicemia caused by pathogens susceptible to penicillin V potassium.

(3) *Limitations*. Administer orally 1 to 2 hours prior to feeding for maximum absorption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37326, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992]

21 CFR Ch. I (4–1–99 Edition)

§ 520.1696d Penicillin V potassium tablets.

(a) *Specifications*. Each tablet contains penicillin V potassium equivalent to 125 milligrams (200,000 units) or 250 milligrams (400,000 units) of penicillin V.

(b) *Sponsors*. See Nos. 017144, 050604, and 053501 in § 510.600(c) of this chapter.

(c) *National Academy of Sciences/National Research Council (NAS/NRC) status*. These conditions of use were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use. Dogs and Cats*—(1) *Amount*. 10 to 15 milligrams per pound of body weight every 6 to 8 hours.

(2) *Indications for use*. Treatment of respiratory, urogenital, skin and soft tissue infections and septicemia caused by pathogens susceptible to penicillin V potassium.

(3) *Limitations*. Administer orally 1 to 2 hours prior to feeding for maximum absorption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37327, Aug. 18, 1992, as amended at 59 FR 58775, Nov. 15, 1994]

§ 520.1720 Phenylbutazone oral dosage forms.

§ 520.1720a Phenylbutazone tablets and boluses.

(a) *Specifications*. Each tablet contains 100, 200, or 400 milligrams, or 1 gram of phenylbutazone. Each bolus contains 2 or 4 grams of phenylbutazone.

(b) *Sponsor*. See sponsor numbers in § 510.600(c) of this chapter, as follows:

(1) No. 000061 for use of 100- or 400-milligram or 1-gram tablets, or 2- or 4-gram boluses, in dogs and horses.

(2) No. 000010 for use of 100- or 200-milligrams or 1-gram tablets in dogs and horses.

(3) Nos. 000031, 000856, 000864, 015579, and 059130 for use of 100-milligram or 1-gram tablets in dogs and horses.

(4) No. 055246 for use of 100-milligram tablets in dogs.

Food and Drug Administration, HHS

§ 520.1720c

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount*. Twenty milligrams per pound of body weight daily.¹

(ii) *Indications for use*. The drug is used for the relief of inflammatory conditions associated with a musculoskeletal system.¹

(iii) *Limitations*. Administer in three divided doses daily. Do not exceed a total daily dose of 800 milligrams regardless of body weight. Administer at a relatively high dosage level for the first 48 hours and then reduce gradually to a maintenance dosage level with the lowest dosage maintained at a level capable of producing the desired clinical response. Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

(2) *Horses*—(i) *Amount*. One to two grams per 500 pounds weight daily.¹

(ii) *Indications for use*. This drug is used for the relief of inflammatory conditions associated with the musculoskeletal system.¹

(iii) *Limitations*. Do not exceed a daily dosage of 4 grams per day. Administer at a relatively high dosage level for the first 48 hours and then reduce gradually to a maintenance dosage level with the lowest dosage maintained at the level capable of producing the desired clinical response. Not for use in animals intended for food purposes. Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[42 FR 44227, Sept. 2, 1977, as amended at 45 FR 10333, Feb. 15, 1980; 45 FR 14023, Mar. 4, 1980; 46 FR 48642, Oct. 2, 1981; 47 FR 30968, July 16, 1982; 50 FR 49372, Dec. 2, 1985; 52 FR 36023, Sept. 25, 1987; 54 FR 22885, May 30, 1989; 55 FR 8462, Mar. 8, 1990; 59 FR 53585, Oct. 25, 1994; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997; 63 FR 36178, July 2, 1998]

§ 520.1720b Phenylbutazone granules.

(a) *Specifications*. The drug is in granular form. It is packaged to contain either 8 grams of phenylbutazone per package or 1 gram of phenylbutazone per package.

(b) *Sponsor*. See 000061 in § 510.600(c) for 8-gram package, see 050604 for 1-gram package.

(c) *NAS/NRC status*. The conditions of use have been NAS/NRC reviewed and

found effective. NADA's for approval of drugs for these conditions of use need not include effectiveness data specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use*—(1) *Horses*—(i) *Amount*. 1 to 2 grams per 500 pounds of body weight, not to exceed 4 grams, daily, as required.

(ii) *Indications*. For the treatment of inflammatory conditions associated with the musculoskeletal system.

(iii) *Limitations*. Administer orally by adding to a portion of the usual grain ration. Use a relatively high dose for the first 48 hours, then gradually reduce to a maintenance level at the lowest level capable of producing the desired clinical response. Treated animals should not be slaughtered for food use. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 18960, Mar. 27, 1981, as amended at 46 FR 48642, Oct. 2, 1981; 57 FR 2836, Jan. 24, 1992; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 520.1720c Phenylbutazone paste.

(a) *Specifications*. The paste contains 20 percent phenylbutazone.

(b) *Sponsor*. See 000061 and 010797 in § 510.600(c) of this chapter.

(c) *NAS/NRC status*. The conditions of use are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified in § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use in horses*—(1) *Amount*. 1 to 2 grams of phenylbutazone per 500 pounds of body weight, not to exceed 4 grams daily.

(2) *Indications for use*. For relief of inflammatory conditions associated with the musculoskeletal system.

(3) *Limitations*. Use a relatively high dose for the first 48 hours, then gradually reduce to a maintenance level of the lowest level capable of producing the desired clinical response. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 84762, Dec. 23, 1980, as amended at 58 FR 29777, May 24, 1993; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

¹ See footnote 1 to § 520.1660b.

§ 520.1720d

21 CFR Ch. I (4–1–99 Edition)

§ 520.1720d Phenylbutazone gel.

(a) *Specifications.* Each 30 grams of gel contains 4 grams of phenylbutazone.

(b) *Sponsor.* See 050604 in § 510.600(c) of this chapter.

(c) *NAS/NRC status.* The conditions of use are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified in § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use in horses—(1) Amount.* 1 to 2 grams of phenylbutazone per 500 pounds of body weight, not to exceed 4 grams daily.

(2) *Indications for use.* For relief of inflammatory conditions associated with the musculoskeletal system of horses.

(3) *Limitations.* Use a relatively high dose for the first 48 hours, then gradually reduce to a maintenance level at the lowest level capable of producing the desired clinical response. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 13561, Apr. 5, 1985, as amended at 50 FR 49372, Dec. 2, 1985; 55 FR 8462, Mar. 8, 1990]

§ 520.1802 Piperazine-carbon disulfide complex oral dosage forms.

§ 520.1802a Piperazine-carbon disulfide complex suspension.

(a) *Specifications.* Each fluid ounce of suspension contains 7.5 grams of piperazine-carbon disulfide complex. The piperazine-carbon disulfide complex contains equimolar parts of piperazine and carbon disulfide (1 gram contains 530 mgs of piperazine and 470 mgs of carbon disulfide).

(b) *Sponsor.* See 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use. Horses and ponies—(1) Amount.* One fluid ounce per 100 pounds of body weight.¹

(2) *Indications for use.* For removing ascarids (large roundworms, *Parascaris equorum*), bots (*Gastrophilus* spp.), small

strongyles, large strongyles (*Strongyles* spp.), and pinworms (*Oxyuris equi*).¹

(3) *Limitations.* Administer by stomach tube or dose syringe after withholding feed overnight or for 8 to 10 hours. Provide water as usual. Resume regular feeding 4 to 6 hours after treatment. Treatment of debilitated or anemic animals is contraindicated. Do not administer to animals that are or were recently affected with colic, diarrhea, or infected with a serious infectious disease. As with most anthelmintics, drastic cathartics and other gastrointestinal irritants should not be administered in conjunction with this drug. Animals in poor condition or heavily parasitized should be given one half the recommended dose and treated again in 2 or 3 weeks. Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[45 FR 52781, Aug. 8, 1980]

§ 520.1802b Piperazine-carbon disulfide complex boluses.

(a) *Specifications.* Each bolus contains 20 grams of piperazine-carbon disulfide complex.

(b) *Sponsor.* See 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use: Horses and ponies—(1) Amount.* For removal of ascarids and small strongyles, 1 bolus (20 grams) per 500 pounds body weight; removal of large strongyles, pinworms, and bots, 1 bolus per 250 pounds body weight.¹

(2) *Indications for use.* For removing ascarids (large roundworms, *Parascaris equorum*), large strongyles (*Strongylus* spp.) bots (*Gastrophilus* spp.), small strongyles, and pinworms (*Oxyuris equi*).¹

(3) *Limitations.* Withhold feed overnight or for 8 to 10 hours. Give water just before and/or after treatment. Resume regular feeding 4 to 6 hours after treatment. Treatment of debilitated or anemic animals is contraindicated. Do not administer to animals that are or were recently affected with colic, diarrhea, or infected with a serious infectious disease. As with most anthelmintics, drastic cathartics or other gastrointestinal irritants should not be administered in conjunction with this drug. Animals in poor condition or heavily parasitized should be

¹These conditions are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

Food and Drug Administration, HHS

§ 520.1805

given one half the recommended dose and treated again in 2 or 3 weeks. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.¹

[45 FR 52782, Aug. 8, 1980]

§ 520.1802c Piperazine-carbon disulfide complex with phenothiazine suspension.

(a) *Specifications.* Each fluid ounce contains 5 grams of piperazine-carbon disulfide complex and 0.83 gram of phenothiazine.

(b) *Sponsor.* See 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use: Horses and ponies—*(1) *Amount.* One fluid ounce per 100 pounds of body weight.

(2) *Indications for use.* For removing ascarids (large roundworms, *Parascaris equorum*), bots (*Gastrophilus* spp.), small strongyles, and large strongyles (*Strongylus* spp.).

(3) *Limitations.* See § 520.1802a(c)(3).

[45 FR 52782, Aug. 8, 1980]

§ 520.1803 Piperazine citrate capsules.

(a) *Specifications.* Piperazine citrate capsules contain piperazine citrate equivalent to 140 milligrams of piperazine base in each capsule.

(b) *Sponsor.* See No. 021091 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in dogs and cats for the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*).

(2) The contents of 1 capsule should be mixed with the food of the animal for each 5 pounds, or fraction thereof of body weight, except dogs weighing over 25 pounds should be given the contents of 6 capsules. The drug should be mixed in ½ of the regular feeding and when the animal has finished eating the dosed food, the remainder of the food may be given. Dogs and cats may be wormed at 6 to 8 weeks of age. The first treatment should be repeated 10 days later. Reinfection may occur. Repeat treatment if indicated.

(3) Severely debilitated animals should not be wormed except on the advice of a veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 54 FR 38515, Sept. 19, 1989]

§ 520.1804 Piperazine phosphate capsules.

(a) *Specifications.* Each capsule contains 120, 300, or 600 milligrams of piperazine phosphate monohydrate.

(b) *Sponsor.* See No. 050906 in § 510.600(c) of this chapter.

(c) *Conditions of use—*(1) *Amount.* 60 milligrams of piperazine phosphate monohydrate per pound of body weight.¹

(2) *Indications for use—*(i) *Dogs.* It is used for the removal of large roundworms (ascarids) *Toxocara canis* and *Toxascaris leonina*.¹

(ii) *Cats.* It is used for the removal of large roundworms (ascarids) *Toxocara mystax* and *Toxocaris leonina*.¹

(3) *Limitations.* Administer in animal's food or milk. For animals up to 1 year of age administer every 2 or 3 months; for animals over 1 year old, administer periodically as necessary. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.¹

[43 FR 6941, Feb. 17, 1978; 43 FR 9804, Mar. 10, 1978, as amended at 46 FR 20158, Apr. 3, 1981]

§ 520.1805 Piperazine phosphate with thenium closylate tablets.

(a) *Specifications.* Each scored tablet contains the equivalent of 250 milligrams piperazine hexahydrate (as piperazine phosphate) and 125 milligrams thenium (as thenium closylate) or 500 milligrams piperazine hexahydrate (as piperazine phosphate) and 250 milligrams thenium (as thenium closylate).

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use—*(1) *Amount.* Administer orally to dogs as follows:

NUMBER OF TABLETS AT EACH OF THE TWO DOSES

Animal weight (lb)	375 mg	750 mg
2 but less than 5	½	
5 but less than 10	1	½
10 or heavier	2	1

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

§ 520.1806

21 CFR Ch. I (4–1–99 Edition)

(2) *Indications for use.* For removal of immature (fourth stage larvae) and adult hookworms (*Ancylostoma caninum*, *A. braziliense*, and *Uncinaria stenocephala*) and ascarids (*Toxocara canis*) from weaned pups and adult dogs.

(3) *Limitations.* Do not use this product to treat dogs weighing less than 2 pounds, unweaned pups, or pups under 5 weeks of age. Maximum efficacy against hookworms necessitates two doses in 1 day of treatment. The interval between the doses should be not less than 4 hours or more than 24 hours. Administer the first dose in the morning before feeding. Do not permit dog to chew tablet. Feed the dog between doses. Do not feed milk or other fatty foods during treatment. Retreatment may be needed in 7 to 28 days as determined by laboratory fecal examinations or in animals kept in known contaminated quarters. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 32747, July 28, 1978, as amended at 47 FR 55476, Dec. 10, 1982; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 520.1806 Piperazine monohydrochloride liquid.

(a) *Specifications.* The product contains 4.77 percent piperazine monohydrochloride, equivalent to 3.35 percent piperazine base.

(b) *Sponsor.* See Nos. 017135 and 063765 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Indications for use.* For dogs for the removal of roundworms (*Toxocara canis* and *Toxascaris leonina*).

(2) *Dosage.* Administer 20 to 30 milligrams of piperazine base per pound of body weight as a single dose.

(3) *Limitations.* Administer by mixing into the animal's ration to be consumed at one feeding. For animals in heavily contaminated areas, reworm at monthly intervals. Not for use in unweaned pups or animals less than three weeks of age. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[47 FR 20758, May 14, 1982, as amended at 57 FR 48163, Oct. 22, 1992; 58 FR 44611, Aug. 24, 1993; 59 FR 28769, June 3, 1994; 63 FR 8348, Feb. 19, 1998]

§ 520.1840 Poloxalene.


(a) *Chemical name.* Polyoxypropylene-polyoxyethylene glycol nonionic block polymer.

(b) *Specifications.* (1) Molecular weight range: 2,850 to 3,150.

(2) Hydroxyl number: 35.7 to 39.4.

(3) Cloud point (10 percent solution): 42° C.–46° C.

(4) Structural formula:



(c) *Sponsor.* (1) See No. 000069 in § 510.600(c) of this chapter for the sponsor of the usage provided by paragraph (d)(1) of this section.

(2) See No. 000069 in § 510.600(c) of this chapter for sponsor of usage provided by paragraph (d)(3) of this section.

(3) See No. 036904 in § 510.600(c) of this chapter for sponsor of usage provided by paragraph (d)(2) of this section.

(4) See No. 017800 in § 510.600(c) of this chapter for sponsor of the usage provided by paragraph (d)(4) of this section.

(d) *Conditions of use.* (1) For treatment of legume (alfalfa, clover) bloat in cattle. Administer as a drench at the rate of 25 grams for animals up to 500 pounds and 50 grams for animals over 500 pounds of body weight.

(2) For control of legume (alfalfa, clover) bloat in cattle. Administer, in molasses block containing 6.6 percent poloxalene, at the rate of 0.8 oz. of block (1.5 grams poloxalene) per 100 lbs. of body weight per day.

(3) For prevention of legume (alfalfa, clover) and wheat pasture bloat in cattle. A 53-percent poloxalene top dressing on individual rations of ground feed. Dosage is 1 gram of poloxalene per 100 pounds of body weight daily. If bloating conditions are severe, the dose is doubled. Treatment should be started 2 to 3 days before exposure to bloat-producing conditions. Repeat use of the drug if animals are exposed to bloat-producing conditions for more than 12 hours after the last treatment. Do not exceed the double dose in any 24-hour period.

(4) For control of legume (alfalfa, clover) and wheat pasture bloat in cattle.

Food and Drug Administration, HHS

§ 520.1871

Administer in molasses block containing 6.6 percent poloxalene, at the rate of 0.8 ounce of block (1.5 grams of poloxalene) per 100 pounds of body weight per day. Provide access to blocks at least 7 days before exposure to bloat-producing conditions.

[40 FR 13838, Mar. 27, 1975, as amended at 40 FR 39857, Aug. 29, 1975; 42 FR 41854, Aug. 19, 1977; 50 FR 5385, Feb. 8, 1985; 54 FR 33501, Aug. 15, 1989; 56 FR 50653, Oct. 8, 1991; 58 FR 26523, May 4, 1993; 60 FR 55659, Nov. 2, 1995]

§ 520.1846 Polyoxyethylene (23) lauryl ether blocks.

(a) *Specifications.* Each molasses-based block contains 2.2 percent polyoxyethylene (23) lauryl ether.

(b) *Sponsor.* See No. 050112 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 2 grams of polyoxyethylene (23) lauryl ether per 100 kilograms of body weight per day (1 pound of block per 500 kilogram (1,100 pound) animal per day).

(2) *Indications for use.* For reduction of the incidence of bloat (alfalfa and clover) in pastured cattle.

(3) *Limitations.* Administer free-choice to beef cattle and nonlactating dairy cattle only. Initially, provide one block per five head of cattle. Start treatment 10 to 14 days before exposure to bloat-producing pastures. Do not allow cattle access to other sources of salt while being fed this product. Do not feed this product to animals without adequate forage/roughage consumption.

[50 FR 48189, Nov. 22, 1985, as amended at 56 FR 9841, Mar. 8, 1991]

§ 520.1870 Praziquantel tablets.

(a) *Specifications.* Each dog tablet contains 34 milligrams (mg) of praziquantel; each cat tablet contains 11.5 or 23 mg of praziquantel.

(b) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs*—(i) *Indications for use.* For removal of canine cestodes *Dipylidium caninum* and *Taenia pisiformis*. If labeled for use by or on the order of a licensed veterinarian, for removal of the canine cestode *Echinococcus granulosus*, and for removal and control of the canine cestode *Echinococcus multilocularis*.

(ii) *Dosage.* Dogs 5 pounds and under, ½ tablet (17 mg); 6 to 10 pounds, 1 tablet (34 mg); 11 to 15 pounds, 1½ tablets (51 mg); 16 to 30 pounds, 2 tablets (68 mg); 31 to 45 pounds, 3 tablets (102 mg); 46 to 60 pounds, 4 tablets (136 mg); over 60 pounds, 5 tablets maximum (170 mg).

(iii) *Limitations.* Administer directly by mouth or crumbled and in feed. Not intended for use in puppies less than 4 weeks of age. For over-the-counter (OTC) use: Consult your veterinarian before administering tablets to weak or debilitated animals, and for assistance in the diagnosis, treatment, and control of parasitism. For prescription use: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Indications for use.* For removal of feline cestodes *Dipylidium caninum* and *Taenia taeniaeformis*.

(ii) *Dosage.* Cats 4 pounds and under, 11.5 mg; 5 to 11 pounds, 23 mg; over 11 pounds, 34.5 mg.

(iii) *Limitations.* Administer directly by mouth or crumbled and in feed. Not intended for use in kittens less than 6 weeks of age. For OTC use: Consult your veterinarian before administering tablets to weak or debilitated animals, and for assistance in the diagnosis, treatment, and control of parasitism.

[46 FR 60570, Dec. 11, 1981, as amended at 47 FR 26377, June 18, 1982; 55 FR 2234, Jan. 23, 1990; 58 FR 7864, Feb. 10, 1993; 58 FR 42853, Aug. 12, 1993]

§ 520.1871 Praziquantel/pyrantel pamoate tablets.

(a) *Specifications.* Each cat tablet contains 18.2 milligrams (mg) praziquantel with 72.6 mg pyrantel (as pyrantel pamoate).

(b) *Sponsor.* See 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Cats*—(i) *Dosage.* 1.5 to 1.9 pounds, 1/4 tablet; 2 to 3 pounds, 1/2 tablet; 4 to 8 pounds, 1 tablet; 9 to 12 pounds, 1 1/2 tablets; 13 to 16 pounds, 2 tablets.

(ii) *Indications for use.* For removal of tapeworms (*Dipylidium caninum* and *Taenia taeniaeformis*), hookworms (*Ancylostoma tubaeforme*), and large roundworms (*Toxocara cati*) in cats and kittens.

§ 520.1872

(iii) *Limitations.* Not for use in kittens less than 1 month of age or weighing less than 1.5 pounds. May be given directly by mouth or in a small amount of food. Do not withhold food prior to or after treatment. If reinfection occurs, treatment may be repeated. Consult your veterinarian before giving to sick or pregnant animals. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) [Reserved]

[58 FR 58652, Nov. 3, 1993]

§ 520.1872 Praziquantel, pyrantel pamoate, and febantel tablets.

(a) *Specifications.* Each tablet contains either:

(1) Tablet No. 1: 22.7 milligrams praziquantel, 22.7 milligrams pyrantel base, and 113.4 milligrams febantel; or

(2) Tablet No. 2: 68 milligrams praziquantel, 68 milligrams pyrantel base, and 340.2 milligrams febantel.

(b) *Sponsor.* See 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Dogs—(i) Amount.* Administer as a single dose directly by mouth or in a small amount of food as follows:

Weight of animal		Number of tablets per dose	
Kilograms	Pounds	Tablet no. 1	Tablet no. 2
0.9 to 1.8	2 to 4	1/2	
2.3 to 3.2	5 to 7	1	
3.6 to 5.4	8 to 12	1 1/2	
5.9 to 8.2	13 to 18	2	
8.6 to 11.4	19 to 25	2 1/2	
11.8 to 13.6 ...	26 to 30		1
14.1 to 20.0 ...	31 to 44		1 1/2
20.4 to 27.2 ...	45 to 60		2
27.7 to 33.6 ...	61 to 74		2 1/2
34.0 to 40.9 ...	75 to 90		3
41.3 to 47.2 ...	91 to 104		3 1/2
47.7 to 54.5 ...	105 to 120		4

(ii) *Indications for use.* For the removal of tapeworms (*Dipylidium caninum*, *Taenia pisiformis*, *Echinococcus granulosus*); hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*); ascarids (*Toxocara canis*, *Toxascaris leonina*); and whipworms (*Trichuris vulpis*) and for the removal and control of tapeworm *Echinococcus multilocularis* in dogs.

(iii) *Limitations.* Do not use in pregnant animals. Do not use in dogs weighing less than 0.9 kilogram (2

21 CFR Ch. I (4–1–99 Edition)

pounds) or puppies less than 3 weeks of age. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[59 FR 33908, July 1, 1994, as amended at 61 FR 29651, June 12, 1996]

§ 520.1880 Prednisolone tablets.

(a) *Specifications.* Each tablet contains 5 or 20 milligrams prednisolone.

(b) *Sponsor.* See No. 061690 in § 510.600(c)(2) of this chapter.

(c) *Special considerations.* (1) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate parturition followed by dystocia, fetal death, retained placenta, and metritis.

(2) Do not use in viral infections. Systemic therapy with prednisolone is contraindicated in animals with peptic ulcer, corneal ulcer, and Cushingoid syndrome. The presence of diabetes, osteoporosis, predisposition to thrombophlebitis, hypertension, congestive heart failure, renal insufficiency, and active tuberculosis necessitates carefully controlled use. Some of the above conditions occur only rarely in dogs but should be kept in mind.

(3) Anti-inflammatory action of corticosteroids may mask signs of infection.

(d) *Conditions of use—(1) Amount.* Dogs: 2.5 milligrams per 4.5 kilograms (10 pounds) body weight per day. Administer total daily dose orally in equally divided doses 6 to 10 hours apart until response is noted or 7 days have elapsed. When response is attained, dosage should be gradually reduced until maintenance level is achieved.

(2) *Indications for use.* For use in dogs as an anti-inflammatory agent.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 4718, Feb. 7, 1992, as amended at 60 FR 57832, Nov. 22, 1995; 63 FR 148, Jan. 5, 1998]

§ 520.1900 Primidone tablets.

(a) *Specifications.* Each tablet contains 50 or 250 milligrams of primidone.

(b) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter for use of 250 milligram tablets; see No. 000856 in § 510.600(c) of this chapter for use of 50 and 250 milligram tablets.

(c) *Conditions of use in dogs—(1) Amount.* Twenty-five milligrams of primidone per pound of body weight (55 milligrams per kilogram of body weight) daily.¹

(2) *Indications for use.* For the control of convulsions associated with idiopathic epilepsy, epileptiform convulsions, viral encephalitis, distemper, and hardpad disease that occurs as a clinically recognizable lesion in certain entities in dogs.¹

(3) *Limitations.* The tablets may be administered whole or crushed and mixed with the food. When convulsions are frequent, the dosage should be divided and administered at intervals. Reduction in dosage should be made gradually and never be abruptly discontinued. Do not use in feline species, as primidone appears to have a specific neurotoxicity in cats. Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[42 FR 61594, Dec. 6, 1977, as amended at 43 FR 55386, Nov. 28, 1978; 46 FR 8467, Jan. 27, 1981; 46 FR 57477, Nov. 24, 1981; 53 FR 40727, Oct. 18, 1988; 56 FR 37473, Aug. 7, 1991; 62 FR 35076, June 30, 1997]

§ 520.1920 Prochlorperazine, isopropamide sustained release capsules.

(a) *Specifications.* Prochlorperazine, isopropamide sustained release capsules contain either:

(1) 3.33 milligrams of prochlorperazine (as the dimaleate) and 1.67 milligrams of isopropamide (as the iodide), or

(2) 10 milligrams of prochlorperazine (as the dimaleate) and 5 milligrams of isopropamide (as the iodide).

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used for the treatment of dogs in which gastrointestinal disturbances are associated with emotional stress.

(2)(i) Capsules described in paragraph (a)(1) of this section are administered orally to dogs weighing from 4 to 15 pounds at the rate of 1 capsule twice daily. These capsules are administered orally to dogs weighing from 16 to 30 pounds at the rate of 1 or 2 capsules twice daily. For dogs weighing less than 4 pounds, administer orally an appropriate fraction of the contents of one of these capsules.

(ii) Capsules described in paragraph (a)(2) of this section are given to dogs weighing 30 pounds and over at the rate of 1 capsule twice daily.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.1921 Prochlorperazine, isopropamide, with neomycin sustained-release capsules.

(a) *Specifications.* Each capsule contains either:

(1) Capsule No. 1: 3.33 milligrams of prochlorperazine (as the dimaleate), 1.67 milligrams of isopropamide (as the iodide), and 25 milligrams of neomycin base (as the sulfate); or

(2) Capsule No. 3: 10 milligrams of prochlorperazine (as the dimaleate), 5 milligrams of isopropamide (as the iodide), and 75 milligrams of neomycin base (as the sulfate).

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* Administer capsules orally twice daily to dogs as follows:

Animal weight (pounds)	Number of capsules per dose	
	Capsule No. 1	Capsule No. 3
10 to 20	1	
20 to 30	2	
Over 30	3	1
Over 60		2

(2) *Indications for use.* For treatment of dogs in which infectious bacterial

§ 520.1962

gastroenteritis is associated with emotional stress.

(3) *Limitations.* Do not continue medication longer than 5 days. Overdosage or prolonged administration may produce nephrotoxicity as manifested by albuminuria, presence of granular casts and depressed urinary output. If it is desirable to administer a vasoconstrictor, norepinephrine is the drug of choice. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[49 FR 14103, Apr. 10, 1984, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.1962 Promazine hydrochloride.

(a)(1) *Chemical name.* 10-[3-(Dimethylamino)propyl]phenothiazine monohydrochloride.

(2) *Specifications.* Conforms to N.F. XII.

(3) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(4) [Reserved]

(5) *Conditions of use.* (i) The drug is used for quieting excitable, unruly, or intractable horses. It is administered at a dosage level of 0.45 to 0.9 milligrams of promazine hydrochloride per pound of body weight mixed with an amount of feed that will be readily consumed.

(ii) Do not use in horses intended for food.

(iii) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b) [Reserved]

[40 FR 13838, Mar. 27, 1975, as amended at 43 FR 55386, Nov. 28, 1978; 59 FR 5705, Feb. 8, 1994]

§ 520.2002 Propiopromazine hydrochloride.

(a) *Chemical name.* 1-Propanone, 1-[10-[3-(dimethylamino) propyl] phenothiazine-2-yl]-, monohydrochloride.

(b) *Specifications.* The drug is administered in a chewable tablet containing 10 to 20 milligrams of propiopromazine hydrochloride.

(c) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

21 CFR Ch. I (4–1–99 Edition)

(d) *Conditions of use.* (1) The drug is intended for oral administration to dogs as a tranquilizer. It is used as an aid in handling difficult, excited, and unruly dogs, and in controlling excessive kennel barking, car sickness, and severe dermatitis. It is also indicated for use in minor surgery and prior to routine examinations, laboratory procedures, and diagnostic procedures.

(2) It is administered at the rate of 0.5 to 2 milligrams of propiopromazine hydrochloride per pound of body weight once or twice daily depending upon the degree of tranquilization desired.

NOTE: Not for use with organophosphates and/or procaine hydrochloride, as phenothiazine may potentiate the toxicity of organophosphates and the activity of procaine hydrochloride. Overdosage may produce significant depression.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 46 FR 60570, Dec. 11, 1981; 61 FR 5506, Feb. 13, 1996]

§ 520.2041 Pyrantel pamoate chewable tablets.

(a) *Specifications.* Each tablet contains pyrantel pamoate equivalent to 22.7 or 113.5 milligrams pyrantel base.

(b) *Sponsor.* See No. 017135 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* Provides at least 2.27 milligrams pyrantel base per pound body weight for dogs weighing more than 5 pounds, and at least 4.54 milligrams of pyrantel base per pound body weight for dogs weighing 5 pounds or less.

(2) *Indications for use—(i) In dogs and puppies.* For removal of ascarids (*Toxocara canis*; *Toxascaris leonina*) and hookworms (*Ancylostoma caninum*; *Uncinaria stenocephala*).

(ii) In puppies and adult dogs and in lactating bitches after whelping. To prevent reinfection of *Toxocara canis*.

(3) *Limitations.* Administer to puppies at 2, 3, 4, 6, 8, and 10 weeks of age. Administer to lactating bitches 2 to 3 weeks after whelping. Retreatment of adult dogs may be necessary at monthly intervals as determined by laboratory fecal examinations. Consult your

Food and Drug Administration, HHS

§ 520.2043

veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[52 FR 37937, Oct. 13, 1987, as amended at 57 FR 48163, Oct. 22, 1992; 58 FR 44611, Aug. 24, 1993]

§ 520.2042 Pyrantel pamoate tablets.

(a) *Specifications.* Each tablet contains pyrantel pamoate equivalent to 22.7, 45.4, or 113.5 milligrams of pyrantel base.

(b) *Sponsor.* See No. 017135 in § 510.600(c) of this chapter.

(c) *Conditions of use.* It is used for dogs as follows:

(1) *Amount.* For dogs weighing over 5 pounds, use at least 2.27 milligrams of pyrantel base per pound of body weight; for dogs weighing 5 pounds or less, use at least 4.54 milligrams of pyrantel base per pound of body weight.

(2) *Indications for use.* For removal and control of large roundworms (ascarids) (*Toxocara canis* and *Toxascaris leonina*), and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*).

(3) *Limitations.* Administer orally directly or in a small amount of food. To prevent reinfection of *T. canis* in puppies, lactating bitches after whelping, and adult dogs; treat puppies 2, 3, 4, 6, 8, and 10 weeks of age; treat lactating bitches 2 to 3 weeks after whelping; routinely treat adult dogs monthly. Do not withhold food prior to or after treatment. The presence of these parasites should be confirmed by laboratory fecal examination. A followup fecal examination should be conducted 2 to 4 weeks after first treatment regimen to determine the need for re-treatment. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[43 FR 52700, Nov. 14, 1978, as amended at 49 FR 22073, May 25, 1984; 57 FR 48163, Oct. 22, 1992; 58 FR 44611, Aug. 24, 1993]

§ 520.2043 Pyrantel pamoate suspension.

(a)(1) *Specifications.* Pyrantel pamoate suspension contains pyrantel pamoate equivalent to 50 milligrams of pyrantel base per milliliter.

(2) *Sponsors.* See Nos. 000069 and 059130 in § 510.600(c) of this chapter.

(3) *Conditions of use.* It is used in horses and ponies as follows:

(i) *Amount.* Equivalent of 3 milligrams pyrantel base per pound of body weight.

(ii) *Indications for use.* For the removal and control of infections from the following mature parasites: Large strongyles (*Strongylus vulgaris*, *Strongylus edentatus*, *Strongylus equinus*), small strongyles, pinworms (*Oxyuris*), and large roundworms (*Parascaris*).

(iii) *Limitations.* Administered as a single dose mixed with the usual grain ration, or by stomach tube, or by dose syringe. Not for use in horses and ponies to be slaughtered for food purposes. When the drug is for administration by stomach tube, it shall be labeled: "Federal law restricts this drug to use by or on the order of a licensed veterinarian." When the drug is not for administration by stomach tube, it shall be labeled: "Consult your veterinarian for assistance in the diagnosis, control, and treatment of parasitism."

(b)(1) *Specifications.* Pyrantel pamoate suspension contains pyrantel pamoate equivalent to 2.27 or 4.54 milligrams of pyrantel base per milliliter.

(2) *Sponsors.* See Nos. 000069, 011615, and 059130 for use of 2.27 and 4.54 milligrams per milliliter product. See No. 023851 for use of 4.54 milligrams per milliliter product.

(3) *Conditions of use.* It is used in puppies and dogs as follows:

(i) *Amount.* Equivalent of 2.27 milligrams of pyrantel base per pound of body weight.

(ii) *Indications for use.* For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*).

(iii) *Limitations.* Administer in the animal's feed bowl as a single dose by itself or mixed in a small quantity of food. Additional treatment may be required and should be confirmed by fecal examination within 2 to 4 weeks. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(4) *Conditions of use.* It is used in puppies and adult dogs and in lactating bitches after whelping as follows:

§ 520.2044

21 CFR Ch. I (4–1–99 Edition)

(i) *Amount.* Equivalent to 2.27 milligrams of pyrantel base per pound of body weight.

(ii) *Indications for use.* To prevent reinfections of *Toxocara canis*.

(iii) *Limitations.* Administer to puppies at 2, 3, 4, 6, 8, and 10 weeks of age. Administer to lactating bitches 2 to 3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

[42 FR 28534, June 3, 1977, as amended at 44 FR 36381, June 22 1979; 45 FR 10760, Feb. 19, 1980; 47 FR 16776, Apr. 20, 1982; 51 FR 11440, Apr. 3, 1986; 61 FR 29477, June 11, 1996; 61 FR 63712, Dec. 2, 1996; 62 FR 37712, July 15, 1997; 63 FR 39727, July 24, 1998; 63 FR 45945, Aug. 28, 1998]

§ 520.2044 Pyrantel pamoate paste.

(a) *Specifications.* Each milliliter of paste contains 180 milligrams of pyrantel base (as pyrantel pamoate).

(b) *Sponsor.* See 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* It is used in horses and ponies as follows:

(1) *Amount.* Equivalent of 3 milligrams pyrantel base per pound of body weight.

(2) *Indications for use.* For removal and control of infections from the following mature parasites: large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); small strongyles; pinworms (*Oxyuris equi*); and large roundworms (*Parascaris equorum*).

(3) *Limitations.* Administer as single dose by depositing paste on dorsum of the tongue using the dose syringe. Not for use in horses intended for food. It is recommended that severely debilitated animals not be treated with this preparation. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[47 FR 47377, Oct. 26, 1982; 48 FR 3367, Jan. 25, 1983]

§ 520.2045 Pyrantel tartrate powder; pyrantel tartrate pellets.

(a) *Specifications.* (1) Pyrantel tartrate powder horse wormer contains 11.3 percent and swine wormer 10.6 percent pyrantel tartrate.

(2) Pyrantel tartrate pellets colt and horse wormer contains 1.25 percent pyrantel tartrate.

(b) *Sponsor.* (1) See No. 000069 in § 510.600(c) of this chapter for conditions of use provided for in paragraphs (d) (1) and (2) of this section.

(2) See No. 060594 in § 510.600(c) of this chapter, for conditions of use provided for in paragraph (d)(3) of this section.

(c) *Related tolerances.* See § 556.560 of this chapter.

(d) *Conditions of use.* It is used in: (1) Horses and ponies:

(i) For the removal and control of infections from the following mature parasites: Large strongyles (*Strongylus vulgaris*, *Strongylus edentatus*, *Strongylus equinus*), small strongyles (*Trichonema spp.*, *Triodontophorus*), pinworms (*Oxyuris*), and large roundworms (*Parascaris*).

(ii) It is administered as a single dose at 0.57 gram of pyrantel tartrate per 100 pounds of body weight mixed with the usual grain ration.

(iii) It is recommended that severely debilitated animals not be treated with this drug. Do not administer by stomach tube or dose syringe. The drug should be used immediately after the package is opened.

(iv) Warning: Not for use in horses and ponies to be slaughtered for food purposes.

(2) Swine:

(i) For the removal and control of large roundworms (*Ascaris suum*) and nodular worm (*Oesophagostomum*) infections.

(ii) It is added to feed at 0.4 gram pyrantel tartrate per pound of nonpelleted ration. The ration is administered as a single treatment as the sole ration at the rate of 1 pound per 40 pounds of animal weight for animals up to 200 pounds. Animals 200 pounds and over are administered 5 pounds of ration per animal.

(iii) Fast pigs over night for optimum results. Water should be made available to animals during fasting and treatment periods. Consult veterinarian before using in severely debilitated animals. The drug should be used immediately after the package is opened.

(iv) Warning: Do not treat within 24 hours of slaughter.

(3) Horses and colts:

(i) For the removal and control of infections from the following mature parasites: Large strongyles (*Strongylus vulgaris*, *Strongylus edentatus*, *Strongylus equinus*), small strongyles (*Trichonema* spp., *Triodontophorus*), pinworms (*Oxyuris*), and large roundworms (*Parascaris*).

(ii) It is administered as a single dose at 12.5 milligrams of pyrantel tartrate per 2.2 pounds of body weight mixed with the usual grain ration.

(iii) It is recommended that severely debilitated animals not be treated with this drug.

(iv) Warning: Do not use in horses or colts intended for food.

[40 FR 13838, Mar. 27, 1975, as amended at 59 FR 28769, June 3, 1994]

§ 520.2087 Roxarsone soluble powder.

(a) *Specifications*. Each ounce (avoirdupois) of soluble powder contains 21.7 grams of roxarsone (monosodium 3-nitro-4-hydroxyphenylarsonate).

(b) *Sponsor*. See No. 046573 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.60 of this chapter.

(d) *NAS/NRC status*. These conditions of use are NAS/NRC reviewed and found effective. NADA's for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(e) *Conditions of use*—(1) *Growing chickens and growing turkeys*—(i) *Amount*. 0.002 percent roxarsone in drinking water (one packet per each 250 gallons of drinking water).

(ii) *Indications for use*. For increased rate of weight gain, improved feed efficiency, and improved pigmentation.

(iii) *Limitations*. Administer continuously throughout growing period. Withdraw 5 days before slaughter. Use as sole source of organic arsenic.

(2) *Swine*—(i) *Amount*. 0.01 percent roxarsone in drinking water (one packet per each 50 gallons of drinking water); or 30 milliliters of a 1.55 percent roxarsone solution (one packet per 3 pints of water) per 50 pounds of body weight as a drench.

(ii) *Indications for use*. As an aid in the treatment of swine dysentery

(hemorrhagic enteritis or bloody scours).

(iii) *Limitations*. Administer drinking water continuously for not more than 6 days. Administer drench once daily for 1 or 2 days. If no improvement is observed, consult a veterinarian. Treatment may be repeated after 5 days. Withdraw 5 days before slaughter. Use as sole source of organic arsenic.

[46 FR 41039, Aug. 14, 1981, as amended at 55 FR 8460, Mar. 8, 1990; 57 FR 8577, Mar. 11, 1992]

§ 520.2088 Roxarsone tablets.

(a)(1) *Specifications*. Each tablet contains 36 milligrams of roxarsone (3-nitro-4-hydroxyphenylarsonic acid).

(2) *Sponsor*. See No. 046573 in § 510.600(c) of this chapter.

(3) *Related tolerances*. See § 556.60 of this chapter.

(4) *NAS/NRC status*. The weight gain, feed efficiency, and pigmentation claims are NAS/NRC reviewed and found effective. NADA's for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(5) *Conditions of use*—(i) *Growing chickens and growing turkeys*—(a) *Amount*. Dissolve 2 tablets in each gallon of drinking water (0.002 percent roxarsone).

(b) *Indications for use*. For increased rate of weight gain, improved feed efficiency, and improved pigmentation.

(c) *Limitations*. Administer continuously throughout growing period. Withdraw 5 days before slaughter. Use as sole source of organic arsenic.

(ii) *Growing chickens*—(a) *Amount*. Dissolve 8 tablets in each gallon of drinking water (0.008 percent roxarsone).

(b) *Indications for use*. As an aid in the prevention of coccidiosis due to *Eimeria tenella*.

(c) *Limitations*. Administer for not more than 10 consecutive days. Treatment may be repeated after 5 days off medication. Withdraw 5 days before slaughter. Use as sole source of organic arsenic.

(b)(1) *Specifications*. Each tablet contains 400 milligrams of roxarsone (3-nitro-4-hydroxyphenylarsonic acid).

(2) *Sponsor*. See No. 046573 in § 510.600(c) of this chapter.

§ 520.2089

21 CFR Ch. I (4–1–99 Edition)

(3) *Related tolerances.* See § 556.60 of this chapter.

(4) *NAS/NRC status.* These conditions are NAS/NRC reviewed and found effective. NADA's for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(5) *Conditions of use*—(i) *Swine*—(a) *Amount.* 1 tablet (400 milligrams) per gallon of drinking water for no more than 6 days, or 1 tablet (400 milligrams) per 2 fluid ounces of warm water per 50 pounds of body weight as a drench once daily for 1 to 2 days.

(b) *Indications for use.* As an aid in the treatment of swine dysentery (hemorrhagic enteritis or bloody scours).

(c) *Limitations.* Treatment may be repeated after 5 days off medication. If no improvement is observed, consult a veterinarian. Treated animals must consume enough medicated water to provide a therapeutic dose. Withdraw 5 days before slaughter. Use as sole source of organic arsenic.

(ii) [Reserved]

(c)(1) *Specifications.* Each tablet contains 72 milligrams of roxarsone (3-nitro-4-hydroxyphenylarsonic acid).

(2) *Sponsor.* See No. 017144 in § 510.600(c) of this chapter.

(3) *Related tolerances.* See § 556.60 of this chapter.

(4) *Conditions of use in growing chickens and growing turkeys*—(i) *Amount.* 1 tablet in each gallon of drinking water (0.002 percent roxarsone).

(ii) *Indications for use.* For improved rate of weight gain, improved feed efficiency, and improved pigmentation.

(iii) *Limitations.* Administer continuously throughout growing period. Do not administer to chickens producing eggs for human consumption. Withdraw 5 days before slaughter. Use as sole source of organic arsenic. Overdosage or the lack of water intake may result in weakness or paralysis of legs.

[46 FR 41040, Aug. 14, 1981, as amended at 46 FR 42448, Aug. 21, 1981; 47 FR 15238, Apr. 9, 1982; 55 FR 8460, Mar. 8, 1990; 57 FR 8577, Mar. 11, 1992; 58 FR 65664, Dec. 16, 1993]

§ 520.2089 Roxarsone liquid.

(a) *Specifications.* Each teaspoon (5 milliliters) of solution contains 72 mil-

ligrams of roxarsone (3-nitro-4-hydroxyphenylarsonic acid).

(b) *Sponsor.* See No. 017144 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.60 of this chapter.

(d) *Conditions of use in growing chickens and growing turkeys*—(1) *Amount.* 1 teaspoon (5 milliliters) to each gallon of drinking water (0.002 percent roxarsone).

(2) *Indications for use.* For improved rate of weight gain, improved feed efficiency, and improved pigmentation.

(3) *Limitations.* Administer continuously throughout growing period. Do not administer to chickens producing eggs for human consumption. Withdraw 5 days before slaughter. Use as sole source of organic arsenic. Overdosage or the lack of water intake may result in weakness or paralysis of legs.

[58 FR 65665, Dec. 16, 1993]

§ 520.2095 Sarafloxacin soluble powder.

(a) *Specifications.* Each 145 grams (5.1 ounces) pouch contains sarafloxacin hydrochloride equivalent to 14.5 grams of sarafloxacin base.

(b) *Sponsor.* See No. 000074 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.594 of this chapter.

(d) *Conditions of use.* Used in drinking water as follows:

(1) *Amount.* Chickens—20 to 40 parts per million for 5 consecutive days as the only source of drinking water. Turkeys—30 to 50 parts per million for 5 consecutive days as the only source of drinking water.

(2) *Indications for use.* For control of mortality in growing turkeys and broiler chickens associated with *Escherichia coli* organisms susceptible to sarafloxacin.

(3) *Limitations.* No preslaughter drug withdrawal period is required when the product is used as directed. Use in a manner other than that indicated or with a dose in excess of that recommended may result in drug residues in edible tissues. Do not use in laying hens producing eggs for human consumption. The effects of sarafloxacin on the reproductive function of treated fowl have not been determined. Federal

Food and Drug Administration, HHS

§ 520.2122

law restricts this drug to use by or on the order of a licensed veterinarian.

[60 FR 50097, Sept. 28, 1995]

§ 520.2098 Selegiline hydrochloride tablets.

(a) *Specifications.* Each tablet contains either 2, 5, 10, 15, or 30 milligrams of selegiline hydrochloride.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use—Dogs—*(1) *Dosage.* 1 milligram per kilogram (0.45 milligram per pound) of body weight.

(i) *Indications for use.* For control of clinical signs associated with uncomplicated pituitary-dependent hyperadrenocorticism in dogs.

(ii) *Limitations.* Administer orally once daily. If no improvement in clinical signs or physical examination findings after 2 months of therapy, increase dose to a maximum of 2 milligrams per kilogram once daily. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Dosage.* 0.5 to 1.0 milligram per kilogram of body weight once daily.

(i) *Indications for use.* For the control of clinical signs associated with canine cognitive dysfunction syndrome.

(ii) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[62 FR 34632, June 27, 1997; 62 FR 55159, Oct. 23, 1997, as amended at 63 FR 29551, June 1, 1998; 64 FR 2122, Jan. 13, 1999]

§ 520.2100 Selenium, vitamin E capsules.

(a) *Specifications.* The capsules contain 2.19 milligrams of sodium selenite (equivalent to 1 milligram of selenium) and 56.2 milligrams of vitamin E (68 I.U.) (as d-alpha tocopheryl acid succinate) or 0.548 milligram of sodium selenite (equivalent to .25 milligram of selenium) and 14 milligrams of vitamin E (17 I.U.) (as d-alpha tocopheryl acid succinate.)

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is intended for use as an aid in alleviating and controlling inflammation, pain, and lameness associated with certain arthropathies in dogs.

(2) The capsules are administered orally with the larger capsules administered at a dosage level of 1 capsule per 20 pounds of body weight to a maximum of 5 capsules with the dosage repeated at 3 day intervals until a satisfactory therapeutic response is observed. A maintenance dosage is then administered consisting of 1 capsule per 40 pounds of body weight, with a minimum of 1 capsule per 40 pounds of body weight, with a minimum of 1 capsule, given every 3 days, or 7 days, or longer, as required to maintain improvement or an asymptomatic condition. For dogs under 20 pounds of body weight, the small capsules are administered orally at a dosage level of 1 per 5 pounds of body weight with a minimum of 1 capsule which dosage is repeated at 3 day intervals until a satisfactory response is observed then a maintenance regimen is initiated with 1 capsule per 10 pounds of body weight, minimum of 1 capsule, every 3 days, or 7 days, or longer as required to maintain continued improvement or an asymptomatic condition.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 52 FR 7832, Mar. 13, 1987; 52 FR 9756, Mar. 26, 1987]

§ 520.2122 Spectinomycin dihydrochloride oral solution.

(a) *Specifications.* The spectinomycin dihydrochloride pentahydrate used in manufacturing the drug is the antibiotic substance produced by growth of *Streptomyces flavopersicus* (var. Abbott) or the same antibiotic substance produced by any other means. The drug is packaged as an aqueous solution containing 50 milligrams of spectinomycin activity per milliliter.

(b) *Sponsors.* (1) See No. 050604 in § 510.600(c) of this chapter.

(2) See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used for the treatment and control of infectious bacterial enteritis (white scours) associated with *E. coli* in pigs under 4 weeks of age.

(2) It is administered orally at the rate of 50 milligrams per 10 pounds body weight twice daily for 3 to 5 days.

§ 520.2123

(3) Do not administer to pigs over 15 pounds body weight or over 4 weeks of age. Do not administer within 21 days of slaughter.

[40 FR 13838, Mar. 27, 1975, as amended at 45 FR 56798, Aug. 26, 1980; 46 FR 60570, Dec. 11, 1981; 61 FR 5506, Feb. 13, 1996]

§ 520.2123 Spectinomycin dihydrochloride pentahydrate oral dosage forms.

§ 520.2123a Spectinomycin dihydrochloride pentahydrate tablets.

(a) *Specifications.* The spectinomycin dihydrochloride pentahydrate used in manufacturing the drug is the antibiotic substance produced by growth of *Streptomyces flavopersicus* (var. Abbott) or the same antibiotic substance produced by any other means.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Special considerations.* The quantities of spectinomycin cited in this section refer to the equivalent weight of base activity for the drug.

(d) *Conditions of use.* (1) The tablets are administered orally to dogs in the treatment of infectious diarrhea and gastroenteritis caused by organisms susceptible to spectinomycin.

(2) The drug is administered orally to provide 10 milligrams per pound of body weight twice daily. The tablets may be placed in the animal's mouth or crushed and administered in milk or in the feed. Dosage may be continued for 4 consecutive days. Should no improvement be observed, discontinue drug and redetermine diagnosis.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 47 FR 14149, Apr. 2, 1982]

§ 520.2123b Spectinomycin dihydrochloride pentahydrate soluble powder.

(a) *Specifications.* The spectinomycin dihydrochloride pentahydrate used in manufacturing the drug is the antibiotic substance produced by growth of *Streptomyces flavopersicus* (var. Abbott) or the same antibiotic substance produced by any other means.

21 CFR Ch. I (4–1–99 Edition)

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Special considerations.* The quantities of spectinomycin cited in this section refer to the equivalent weight of base activity for the drug.

(d) *Related tolerances.* See § 556.600 of this chapter.

(e) *Conditions of use.* (1) It is administered in the drinking water of growing chickens at 2 grams of spectinomycin per gallon of water as the only source of drinking water for the first 3 days of life and for 1 day following each vaccination. It is administered as an aid in the prevention or control of losses due to CRD associated with *M. gallisepticum* (PPLO). Do not administer to laying chickens. Do not administer within 5 days of slaughter.

(2) It is administered in the drinking water of floor-raised broiler chickens at 0.5 gram of spectinomycin per gallon of water as the only source of drinking water for the first 3 days of life and for 1 day following each vaccination. It is administered for increased rate of weight gain and improved feed efficiency. Do not administer to laying chickens. Do not administer within 5 days of slaughter.

(3) It is administered in drinking water of broiler chickens at 1 gram of spectinomycin per gallon of water as the only source of drinking water for the first 3 to 5 days of life as an aid in controlling infectious synovitis due to *Mycoplasma synoviae*. Do not administer to laying chickens. Do not administer within 5 days of slaughter.

[40 FR 13838, Mar. 27, 1975, as amended at 45 FR 56798, Aug. 26, 1980]

§ 520.2150 Stanozolol oral dosage forms.

§ 520.2150a Stanozolol tablets.

(a) *Specifications.* Each tablet contains 2 milligrams of stanozolol.

(b) *Sponsor.* No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) Used as an anabolic steroid treatment in dogs and cats.

(2) Administered orally to cats and small breeds of dogs, ½ to 1 tablet twice daily for several weeks; to large breeds of dogs, 1 to 2 tablets twice daily for several weeks. The tablets

Food and Drug Administration, HHS

§ 520.2158c

may be crushed and administered in feed.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 46101, Oct. 6, 1975, as amended at 42 FR 36995, July 19, 1977. Redesignated at 50 FR 38114, Sept. 20, 1985, and amended at 55 FR 23076, June 6, 1990]

§ 520.2150b Stanozolol chewable tablets.

(a) *Specifications.* Each chewable tablet contains 2 milligrams of stanozolol.

(b) *Sponsor.* No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) Used as an anabolic steroid treatment in dogs.

(2) Administered orally to small breeds of dogs, ½ to 1 tablet twice daily for several weeks; to large breeds of dogs, 1 to 2 tablets twice daily for several weeks.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 38114, Sept. 20, 1985, as amended at 55 FR 23076, June 6, 1990]

§ 520.2158 Streptomycin/dihydrostreptomycin oral dosage forms.

§ 520.2158a Streptomycin sulfate oral solution.

(a) *Specifications.* Solution containing 25 percent streptomycin sulfate.

(b) *Sponsor.* See Nos. 033008 and 055462 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.610 of this chapter.

(d) *Conditions of use.* Use in drinking water as follows:

(1) *Calves and swine*—(i) *Amount.* 10 to 15 milligrams per pound (mg/pound) of body weight (1.0 to 1.5 grams per gallon).

(ii) *Indications for use.* Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella* spp. susceptible to streptomycin.

(iii) *Limitations.* Calves: Do not administer for more than 5 days. Swine: Do not administer for more than 4 days. Prepare fresh solution daily. Calves: Withdraw 2 days before slaughter. As sole source of streptomycin. Warning: Certain strains of bacteria may develop a tolerance for strepto-

mycin. Consult a veterinarian or animal pathologist for diagnosis.

(2) *Chickens*—(i) *Amount.* 10 to 15 mg/pound of body weight (0.6 to 0.9 grams per gallon).

(ii) *Indications for use.* Treatment of nonspecific infectious enteritis caused by organisms susceptible to streptomycin.

(iii) *Limitations.* Chickens: Do not administer for more than 5 days. Withdraw 4 days before slaughter. Do not administer to chickens producing eggs for human consumption. Prepare fresh solution daily. As sole source of streptomycin. Warning: Certain strains of bacteria may develop a tolerance for streptomycin. Consult a veterinarian or animal pathologist for diagnosis.

[57 FR 37327, Aug. 18, 1992, as amended at 58 FR 47211, Sept. 8, 1993; 63 FR 51821, Sept. 29, 1998]

§ 520.2158b Dihydrostreptomycin tablets.

(a) *Specifications.* Each tablet contains 37.5 milligrams dihydrostreptomycin (as the sulfate) with 375 milligrams chlorhexidine dihydrochloride.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See §§ 556.120 and 556.200 of this chapter.

(d) *Conditions of use.* *Calves*—(1) *Amount.* 150 milligrams of dihydrostreptomycin and 1.5 grams of chlorhexidine dihydrochloride per 100 pounds of body weight per day.

(2) *Indications for use.* Treatment of bacterial scours in calves.

(3) *Limitations.* Administer orally once a day for 5 days; withdraw 3 days before slaughter.

[57 FR 37327, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992]

§ 520.2158c Dihydrostreptomycin oral suspension.

(a) *Specifications.* Each milliliter contains 1.25 milligrams dihydrostreptomycin (as the sulfate) with 12.5 milligrams chlorhexidine dihydrochloride.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See §§ 556.120 and 556.200 of this chapter.

(d) *Conditions of use.* *Calves*—(1) *Amount.* 150 milligrams of dihydrostreptomycin and 1.5 grams of

§ 520.2160

chlorhexidine dihydrochloride per 100 pounds of body weight per day.

(2) *Indications for use.* Treatment of bacterial scours in calves.

(3) *Limitations.* Administer orally once a day for 5 days; withdraw 3 days before slaughter.

[57 FR 37327, Aug. 18, 1992]

§ 520.2160 Styrylpyridinium, diethylcarbazine oral dosage forms.

§ 520.2170 Sulfabromomethazine sodium boluses.

(a) *Specifications.* Each bolus contains 15 grams of sulfabromomethazine sodium.

(b) *Related tolerance.* See § 556.620 of this chapter.

(c) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(d) *NAS/NRC status.* These conditions of use are NAS/NRC reviewed and found effective. NADA's for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(e) *Conditions of use. Cattle—*(1) *Amount.* 90 milligrams per pound body weight.

(2) *Indications for use.* Treatment of necrotic pododermatitis (foot rot) and calf diphtheria caused by *Fusobacterium necrophorum*; colibacillosis (scours) caused by *Escherichia coli*; bacterial pneumonia and bovine respiratory disease complex (shipping fever complex) associated with *Pasteurella* spp.; acute metritis and acute mastitis caused by *Streptococcus* spp.

(3) *Limitations.* Administer orally; repeat in 48 hours if necessary; milk taken from animals within 96 hours (8 milkings) of latest treatment must not be used for food; do not administer within 18 days of slaughter; discontinue use if hematuria, crystalluria or severe depression are noticed; if signs persist after 2 or 3 days consult a veterinarian.

[47 FR 30243, July 13, 1982, as amended at 62 FR 63270, Nov. 28, 1997]

§ 520.2184 Sodium sulfachloropyrazine monohydrate.

(a) *Chemical name.* 2-Sulfamido-6-chloroxyrazine, sodium.

21 CFR Ch. I (4–1–99 Edition)

(b) *Sponsor.* See Nos. 010042 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.625 of this chapter.

(d) *Conditions of use.* It is used in the drinking water of broilers, breeder flocks, and replacement chickens as follows:

(1) *Amount.* 0.03 percent.

(2) *Indications for use.* Treatment of coccidiosis.

(3) *Limitations.* Administer in drinking water for 3 days as sole source of drinking water and sulfonamide medication; withdraw 4 days prior to slaughter; not to be administered to chickens producing eggs for human consumption.

[40 FR 13838, Mar. 27, 1975, as amended at 50 FR 41489, Oct. 11, 1985; 54 FR 12188, Mar. 24, 1989; 55 FR 8460, Mar. 8, 1990; 64 FR 15684, Apr. 1, 1999]

§ 520.2200 Sulfachlorpyridazine oral dosage forms.

§ 520.2200a Sulfachlorpyridazine bolus.

(a) *Chemical name.* N'-6-(Chloro-3-pyridazinyl) sulfanilamide.

(b) *Specifications.* Melting point range: 190° C to 191° C.

(c) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(d) *Related tolerances.* See § 556.630 of this chapter.

(e) *Conditions of use.* It is used in calves as follows:

(1) *Amount.* 30 to 45 milligrams per pound body weight per day.

(2) *Indications for use.* Treatment of diarrhea caused or complicated by *E. coli* (colibacillosis).

(3) *Limitations.* Administer in a bolus containing 2 grams of sulfachlorpyridazine for 1 to 5 days in divided doses twice daily; treated calves must not be slaughtered for food during treatment or for 7 days after the last treatment.

[40 FR 13838, Mar. 27, 1975, as amended at 50 FR 41489, Oct. 11, 1985]

§ 520.2200b Sulfachlorpyridazine medicated milk and drinking water.

(a) *Chemical name.* N'-(6-Chloro-3-pyridazinyl) sulfanilamide.

(b) *Specifications.* Melting point range: 190 °C to 191 °C.

Food and Drug Administration, HHS

§ 520.2220a

(c) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(d) *Related tolerances.* See § 556.630 of this chapter.

(e) *Conditions of use.* It is used as follows:

(1) *Calves*—(i) *Amount.* 30 to 45 milligrams per pound body weight per day.

(ii) *Indications for use.* Treatment of diarrhea caused or complicated by *E. coli* (colibacillosis).

(iii) *Limitations.* Administer as the sodium salt of sulfachlorpyridazine in milk or milk-replacer formulations for 1 to 5 days in divided doses twice daily; treated calves must not be slaughtered for food during treatment or for 7 days after the last treatment.

(2) *Swine*—(i) *Amount.* 20 to 35 milligrams per pound body weight per day.

(a) *Indications for use.* Treatment of diarrhea caused or complicated by *E. coli* (colibacillosis).

(b) *Limitations.* Administer as the sodium salt of sulfachlorpyridazine in drinking water for 1 to 5 days; for individual treatment, administer orally in divided doses twice daily; treated swine must not be slaughtered for food during treatment or for 4 days after the last treatment.

(ii) *Amount.* 20 to 35 milligrams per pound body weight per day.

(a) *Indications for use.* Treatment of diarrhea caused or complicated by *E. coli* (colibacillosis).

(b) *Limitations.* Administer individually in an oral suspension containing 50 milligrams of sulfachlorpyridazine per milliliter in divided doses twice daily for 1 to 5 days; treated swine must not be slaughtered for food during treatment or for 4 days after the last treatment.

[40 FR 13838, Mar. 27, 1975, as amended at 50 FR 41489, Oct. 11, 1985]

§ 520.2200c Sulfachlorpyridazine tablets.

(a) *Specifications.* Sulfachlorpyridazine tablets contain 250 milligrams of sulfachlorpyridazine per tablet.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs as a broad spectrum antibacterial agent to aid in the treatment of infectious tracheobronchitis and in-

fections caused by *E. coli*. It can also be used in the treatment of infections caused by other gram-positive and gram-negative organisms that are susceptible to sulfonamide therapy.

(2) It is administered orally at a dosage level of 500 milligrams per 10 to 15 pounds of body weight daily, in two or three divided doses.

(3) The administration of the drug should be discontinued if a response is not noted within 7 to 10 days.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 36622, Aug. 18, 1978, as amended at 50 FR 41489, Oct. 11, 1985]

§ 520.2220 Sulfadimethoxine oral dosage forms.

§ 520.2220a Sulfadimethoxine oral solution and soluble powder.

(a) *Approvals.* (1) For oral solution containing 12.5 percent (3.75 grams per ounce) sulfadimethoxine, see Nos. 000010, 000069, 051259, 057561, and 059130 in § 510.600(c).

(2) For soluble powder, each package containing the equivalent of 94.6 grams of sulfadimethoxine (as the sodium salt), see Nos. 000069, 051259, and 057561 in § 510.600(c).

(b) *Special considerations.* Chickens and turkeys that have survived fowl cholera outbreaks should not be kept for replacements or breeders.

(c) *Related tolerances.* See § 556.640 of this chapter.

(d) *Conditions of use.* The oral solution is administered as a cattle drench or diluted as directed to prepare drinking water. The powder is used to prepare a drench or drinking water. The concentrations and uses of the various solutions are as follows:

(1) *Broiler and replacement chickens only.* (i) *Amount.* 1.875 (0.05 percent) grams per gallon.

(ii) *Indications for use.* Treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

(iii) *Limitations.* Administer for 6 consecutive days; do not administer to chickens over 16 weeks of age; as sole source of drinking water and sulfonamide medication; as sulfadimethoxine solution or

§ 520.2220b

sulfadimethoxine soluble sodium salt; withdraw 5 days before slaughter.

(2) *Meat-producing turkeys only*—(i) *Amount.* 0.938 (0.025 percent) grams per gallon.

(ii) *Indications for use.* Treatment of disease outbreaks of coccidiosis and fowl cholera.

(iii) *Limitations.* Administer for 6 consecutive days; do not administer to turkeys over 24 weeks of age; as sole source of drinking water and sulfonamide medication; as sulfadimethoxine solution or sulfadimethoxine soluble sodium salt; withdraw 5 days before slaughter.

(3) *Dairy calves, dairy heifers, and beef cattle only*—(i) *Amount.* 1.18 to 2.36 (0.031 to 0.062 percent) grams per gallon.

(ii) *Indications for use.* Treatment of shipping fever complex, bacterial pneumonia, calf diphtheria, and foot rot.

(iii) Administer 2.5 grams per 100 pounds of body weight for first day, then 1.25 grams per 100 pounds of body weight per day for the next 4 consecutive days; in drinking water or drench; available as a sulfadimethoxine soluble powder or a 12.5 percent sulfadimethoxine sodium solution (3.75 grams sulfadimethoxine per fluid ounce); if no improvement within 2 to 3 days, reevaluate diagnosis; do not treat beyond 5 days; withdraw 7 days before slaughter.

[40 FR 13838, Mar. 27, 1975, as amended at 58 FR 6092, Jan. 26, 1993; 59 FR 56000, Nov. 10, 1994; 61 FR 4875, Feb. 9, 1996; 62 FR 8371, Feb. 25, 1997; 62 FR 23357, Apr. 30, 1997; 62 FR 35076, June 30, 1997; 62 FR 40932, July 31, 1997; 63 FR 59714, Nov. 5, 1998]

§ 520.2220b Sulfadimethoxine tablets and boluses.

(a) *Sponsors.* Approval to firms identified in § 510.600(c) of this chapter as follows:

(1) To 000069, approval for use as in paragraphs (d)(1), (d)(2), and (d)(3) of this section.

(2) To 000061, approval for use as in paragraph (d)(2).

(b) *Related tolerances.* See § 556.640 of this chapter.

(c) [Reserved]

(d) It is used as follows:

(1) *Cattle*—(i) *Amount.* 12.5 to 2.5 grams per 100 pounds body weight.

21 CFR Ch. I (4–1–99 Edition)

(ii) *Indications for use.* Treatment of foot rot, bacterial pneumonia, shipping fever, and calf diphtheria.

(iii) *Limitations.* Administer 2.5 grams per 100 pounds body weight for 1 day followed by 1.25 grams per 100 pounds body weight per day; treat from 4 to 5 days; do not administer within 7 days of slaughter; milk that has been taken from animals during treatment and 60 hours (5 milkings) after the latest treatment must not be used for food.

(2) *Dogs and cats.* (i) *Amount.* 12.5 to 25 milligrams per pound of body weight.

(ii) *Indications for use.* Treatment of sulfadimethoxine-susceptible bacterial infections.

(iii) *Limitations.* Administer 25 milligrams per pound of body weight on the first day followed by 12.5 milligrams per pound of body weight per day until the animal is free of symptoms for 48 hours. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(3) *Beef cattle and nonlactating dairy cattle*—(i) *Amount.* 12.5-gram-sustained-release bolus.

(ii) *Indications for use.* Treatment of shipping fever complex and bacterial pneumonia associated with organisms such as *Pasteurella* spp. sensitive to sulfadimethoxine; calf diphtheria and foot rot associated with *Sphaerophorus necrophorus* sensitive to sulfadimethoxine.

(iii) *Limitations.* Administer one bolus for the nearest 200 pounds of body weight, i.e., 62.5 milligrams per pound of body weight. Do not repeat treatment for 7 days. Do not use in lactating dairy cattle. Do not administer within 12 days of slaughter. During treatment make certain that animals maintain adequate water intake. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 40 FR 43488, Sept. 22, 1975; 49 FR 36830, Sept. 20, 1984; 59 FR 56000, Nov. 10, 1994; 61 FR 4875, Feb. 9, 1996; 62 FR 61625, Nov. 19, 1997; 64 FR 15684, Apr. 1, 1999]

§ 520.2220c Sulfadimethoxine oral suspension.

(a) *Chemical name.* *N*'-(2,6-Dimethoxy-4-pyrimidinyl) sulfanilamide.

Food and Drug Administration, HHS

§ 520.2240a

(b) *Specifications.* Each milliliter of the drug contains 50 milligrams of sulfadimethoxine.

(c) *Sponsor.* See Nos. 000061 and 000069 in § 510.600(c) of this chapter.

(1) It is intended for use in the treatment of sulfonamide susceptible bacterial infections in dogs and cats and enteritis associated with coccidiosis in dogs.

(2) On the first day of treatment administer an oral dose of 25 milligrams per pound of body weight, then follow with a daily dosage of 12.5 milligrams per pound of body weight. Length of treatment will depend upon clinical response. Continue treatment until patient is asymptomatic for 48 hours. Maintain adequate water intake during the treatment period.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 59 FR 56000, Nov. 10, 1994; 61 FR 4875, Feb. 9, 1996; 62 FR 61625, Nov. 19, 1997]

§ 520.2220d Sulfadimethoxine-ormetoprim tablets.

(a) *Specifications.* Each tablet contains 120 milligrams (100 milligrams of sulfadimethoxine and 20 milligrams of ormetoprim), 240 milligrams (200 milligrams of sulfadimethoxine and 40 milligrams of ormetoprim), 600 milligrams (500 milligrams of sulfadimethoxine and 100 milligrams of ormetoprim), or 1200 milligrams (1,000 milligrams of sulfadimethoxine and 200 milligrams of ormetoprim).

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* On the first day of treatment, administer 25 milligrams per pound (55 milligrams per kilogram) of body weight. Then follow with a daily dosage of 12.5 milligrams per pound (27.5 milligrams per kilogram) of body weight.

(2) *Indications of use.* Treatment of skin and soft tissue infections (wounds and abscesses) in dogs caused by strains of *Staphylococcus aureus* and *Escherichia coli* and urinary tract infections caused by *Escherichia coli*, *Staphylococcus* spp., and *Proteus mirabilis* susceptible to ormetoprim-potentiated sulfadimethoxine.

(3) *Limitations.* Continue treatment until patient is asymptomatic for 48

hours, but do not exceed a total of 21 consecutive days. Maintain adequate water intake during the treatment period. Safety in breeding animals has not been established. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[54 FR 48593, Nov. 24, 1989, as amended at 59 FR 56000, Nov. 10, 1994; 61 FR 4875, Feb. 9, 1996; 61 FR 46719, Sept. 5, 1996]

§ 520.2240 Sulfaethoxypyridazine.

§ 520.2240a Sulfaethoxypyridazine drinking water.

(a) *Chemical name.* *N'*-(6-Ethoxy-3-pyridazinyl) sulfanilamide.

(b) *Specifications.* Melting point range of 180° C. to 186° C.

(c) *Sponsor.* See No. 010042 in § 510.600(c) of this chapter.

(d) *Related tolerances.* See § 556.650 of this chapter.

(e) *Conditions of use.* It is used as follows:

(1) *Swine*—(i) *Amount.* 1.9 to 3.8 grams per gallon (0.05 percent to 0.1 percent).

(ii) *Indications for use.* Treatment of bacterial scours pneumonia enteritis, bronchitis, septicemia accompanying *Salmonella choleraesuis* infection.

(iii) *Limitations.* Administer 3.8 grams per gallon for first day followed by 1.9 grams per gallon for not less than 3 days nor more than 9 days as sodium sulfaethoxypyridazine; do not treat within 10 days of slaughter; as sole source of sulfonamide; for use by or on the order of a licensed veterinarian.

(2) *Cattle*—(i) *Amount.* 2.5 grams per gallon (0.066 percent).

(ii) *Indications for use.* Treatment of respiratory infections (pneumonia, shipping fever), foot rot, calf scours; as adjunctive therapy in septicemia accompanying mastitis and metritis.

(iii) *Limitations.* Administer at the rate of 1 gallon per 100 pounds of body weight per day for 4 days; as sodium sulfaethoxypyridazine; do not treat within 16 days of slaughter; as sole source of sulfonamide; for use by or on the order of a licensed veterinarian; milk that has been taken from animals during treatment and for 72 hours (6 milkings) after latest treatment must not be used for food.

§ 520.2240b

21 CFR Ch. I (4–1–99 Edition)

§ 520.2240b Sulfaethoxypyridazine tablets.

(a) *Chemical name.* N'-(6-Ethoxy-3-pyridazinyl) sulfanilamide.

(b) *Specifications.* Melting point range of 180° C to 186° C.

(c) *Sponsor.* See No. 010042 in § 510.600(c) of this chapter.

(d) *Related tolerances.* See § 556.650 of this chapter.

(e) *Conditions of use.* It is used for cattle as follows:

(1) *Amount.* 2.5 or 15 grams per tablet.

(i) *Indications for use.* Treatment of respiratory infections (pneumonia, shipping fever), foot rot, calf scours; as adjunctive therapy in septicemia accompanying mastitis and metritis.

(ii) *Limitations.* Administer 25 milligrams per pound of animal weight per day for 4 days; do not treat within 16 days of slaughter; as sole source of sulfonamide; milk that has been taken from animals during treatment and for 72 hours (6 milkings) after the latest treatment must not be used for food; for use only by or on the order of a licensed veterinarian.

(2) *Amount.* 15-gram controlled release tablets.

(i) *Indications for use.* Treatment of foot rot and respiratory infections (shipping fever and pneumonia) caused by sulfonamide-susceptible pathogens (*E. coli*, streptococci, staphylococci, *Sphaerophorus necrophorus* and Gram-negative rods including *Pasteurella*); for use prophylactically in cattle during periods of stress for reducing losses due to sulfonamide sensitive disease conditions.

(ii) *Limitations.* Administer 100 milligrams per pound of body weight; do not treat within 16 days of slaughter; as sole source of sulfonamide; not for use in lactating dairy cows; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

§ 520.2260 Sulfamethazine oral dosage forms.

§ 520.2260a Sulfamethazine oblets and boluses.

(a)(1) *Sponsor.* See No. 010042 in § 510.600(c) of this chapter for use of 2.5-, 5-, or 15-gram sulfamethazine oblet.

(2) *Related tolerance in edible products.* See § 556.670 of this chapter.

(3) *Conditions of use*—(i) *Amount.* Administer as a single dose 100 milligrams of sulfamethazine per pound of body weight the first day and 50 milligrams per pound of body weight on each following day.

(ii) *Indications for use.* For treatment of diseases caused by organisms susceptible to sulfamethazine.

(A) *Beef cattle and nonlactating dairy cattle.* Treatment of bacterial pneumonia and bovine respiratory disease complex (shipping fever complex) (*Pasteurella* spp.), colibacillosis (bacterial scours) (*Escherichia coli*), necrotic pododermatitis (foot rot) (*Fusobacterium necrophorum*), calf diphtheria (*Fusobacterium necrophorum*), acute mastitis (*Streptococcus* spp.), acute metritis (*Streptococcus* spp.), coccidiosis (*Eimeria bovis* and *Eimeria zurnii*).

(B) *Horses.* Treatment of bacterial pneumonia (secondary infections associated with *Pasteurella* spp.), strangles (*Streptococcus equi*), and bacterial enteritis (*Escherichia coli*).

(iii) *Limitations.* Administer daily until animal's temperature and appearance are normal. If symptoms persist after using for 2 or 3 days consult a veterinarian. Fluid intake must be adequate. Treatment should continue 24 to 48 hours beyond the remission of disease symptoms, but not to exceed 5 consecutive days. Follow dosages carefully. Not for use in lactating dairy animals. Do not treat cattle within 10 days of slaughter. Not to be used in horses intended for food.

(4) *NAS/NRC status.* The conditions of use specified in this section have been reviewed by NAS/NRC and are found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(b)(1) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter for use of 5-gram sulfamethazine bolus.

(2) *Related tolerances in edible products.* See § 556.670 of this chapter.

(3) *Conditions of use*—(i) *Amount.* Administer 10 grams (2 boluses) of sulfamethazine per 100 pounds of body weight the first day, then 5 grams (1 bolus) of sulfamethazine per 100 pounds

of body weight daily for up to 4 additional consecutive days.

(ii) *Indications for use.* Ruminating beef and dairy calves. For treatment of the following diseases caused by organisms susceptible to sulfamethazine: bacterial scours (colibacilloosis) caused by *E. coli*; necrotic pododermatitis (foot rot) and calf diphtheria caused by *F. necrophorum*; bacterial pneumonia associated with *Pasteurella* spp.; and coccidiosis caused by *E. bovis* and *E. zurnii*.

(iii) *Limitations.* Do not administer for more than 5 consecutive days. Do not treat calves within 11 days of slaughter. Do not use in calves to be slaughtered under 1 month of age or in calves being fed an all milk diet. Do not use in female dairy cattle 20 months of age or older; such use may cause drug residues in milk. Administer with adequate supervision. Follows recommended dosages carefully. Fluid intake must be adequate. If symptoms persist after 2 or 3 days, consult a veterinarian.

[54 FR 15751, Apr. 19, 1989; 54 FR 19283, May 4, 1989, as amended at 56 FR 50653, Oct. 8, 1991; 59 FR 22754, May 3, 1994; 61 FR 4875, Feb. 9, 1996]

§ 520.2260b Sulfamethazine sustained-release boluses.

(a)(1) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter for use of a 22.5-gram sulfamethazine prolonged-release bolus.

(2) *Conditions of use—(i) Amount.* Depending on the duration of therapeutic levels desired, administer boluses as a single dose as follows: 3½ days—1 bolus (22.5 grams) per 200 pounds of body weight; 5 days—1 bolus per 100 pounds of body weight.

(ii) *Indications for use.* Beef and nonlactating cattle for sustained treatment of shipping fever pneumonia caused or complicated by *Pasteurella multocida*; as an aid in the treatment of foot rot, mastitis, pneumonia, metritis, bacterial enteritis, calf diphtheria, and septicemia when caused or complicated by bacteria susceptible to sulfamethazine.

(iii) *Limitations.* Cattle that are acutely ill should be treated parenterally with a suitable antibacterial product to obtain immediate therapeutic blood levels; do not slaugh-

ter animals for food within 16 days of treatment; do not use in lactating dairy cattle; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter for use of a 27-gram sulfamethazine sustained-release bolus.

(2) *Conditions of use—(i) Amount.* 27 grams (1 bolus) for each 150 pounds of body weight as a single dose.

(ii) *Indications for use.* For nonlactating cattle for the treatment of infections caused by organisms sensitive to sulfamethazine such as hemorrhagic septicemia (shipping fever complex), bacterial pneumonia, foot rot, and calf diphtheria and as an aid in the control of bacterial diseases usually associated with shipping and handling of cattle.

(iii) *Limitations.* If no response within 2 to 3 days, reevaluate therapy; do not crush tablets; treated animals must not be slaughtered for food within 28 days after the latest treatment; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(c)(1) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter for use of a 32.1-gram sustained-release bolus.

(2) *Conditions of use—(i) Amount.* 32.1 grams (1 bolus) per 200 pounds of body weight.¹

(ii) *Indications for use.* For beef and nonlactating dairy cattle for the treatment of diseases caused by sulfamethazine-sensitive organisms as follows: bacterial pneumonia and bovine respiratory disease complex (shipping fever complex) caused by *Pasteurella* spp., colibacillosis (bacterial scours) caused by *E. coli*, necrotic pododermatitis (foot rot) and calf diphtheria caused by *Fusobacterium necrophorum*, and acute mastitis and acute metritis caused by *Streptococcus* spp.)¹

(iii) *Limitations.* After 72 hours, all animals should be reexamined for persistence of observable disease signs. If signs are present, consult a veterinarian. It is strongly recommended

¹These conditions are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

that a second dose be given to provide for an additional 72 hours of therapy, particularly in more severe cases. The dosage schedule should be used at each 72-hour interval. Animals should not receive more than 2 doses because of the possibility of incurring residue violations. This drug, like all sulfonamides, may cause toxic reactions and irreparable injury unless administered with adequate and continuous supervision; *follow dosages carefully*. Fluid intake must be adequate at all times throughout the 3-day therapy. Do not use in lactating dairy cattle. Do not treat animals within 12 days of slaughter.

(d)(1) *Sponsor*. See 000859 in § 510.600(c) of this chapter for use of a 22.5-gram sulfamethazine sustained release bolus.

(2) *Conditions of use*—(i) *Amount*. Administer 1 bolus (22.5 grams) per 200 pounds of body weight, as a single dose.

(ii) *Indications for use*. Beef and nonlactating dairy cattle for the prolonged treatment of the following diseases when caused by one or more of the listed pathogenic organisms sensitive to sulfamethazine: bovine respiratory disease complex (shipping fever complex) (*Pasteurella* spp.), bacterial pneumonia (*Pasteurella* spp.), necrotic pododermatitis (foot rot) (*Fusobacterium necrophorum*), colibacillosis (bacterial scours) (*Escherichia coli*), calf diphtheria (*Fusobacterium necrophorum*), acute mastitis (*Streptococcus* spp.) and acute metritis (*Streptococcus* spp.).

(iii) *Limitations*. Cattle that are acutely ill should be treated by injection with a suitable antibacterial product to obtain immediate therapeutic blood levels; do not slaughter animals for food within 16 days of treatment; do not use in lactating dairy cattle; if treated animals do not respond within 2 to 3 days, consult a veterinarian.

(e)(1) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter for use of an 8.02-gram sulfamethazine sustained-release bolus.

(2) *Conditions of use*—(i) *Amount*. Administer 2 boluses (8.02 grams per bolus) per 100 pounds of body weight, as a single dose.

(ii) *Indications for use*. Administer orally to ruminating calves for the prolonged treatment of the following dis-

eases when caused by one or more of the listed pathogenic organisms sensitive to sulfamethazine: bacterial pneumonia (*Pasteurella* spp.), colibacillosis (bacterial scours) (*E. coli*), and calf diphtheria (*Fusobacterium necrophorum*).

(iii) *Limitations*. For use in ruminating replacement calves only; 72 hours after dosing all animals should be reexamined for persistence of disease signs; if signs are present, consult a veterinarian; do not slaughter animals for food for at least 12 days after the last dose; this product has not been shown to be effective for nonruminating calves; exceeding two consecutive doses may cause violative tissue residue to remain beyond the withdrawal time; do not use in calves under 1 month of age or calves being fed an all milk diet.

(f)(1) *Sponsor*. See No. 000010 in § 510.600(c) of this chapter for use of a 30-gram sulfamethazine sustained-release bolus.

(2) *Conditions of use*—(i) *Amount*. Administer at the rate of 1 bolus (30 grams per bolus) per 200 pounds of body weight, as a single dose.

(ii) *Indications for use*. Administer orally to beef cattle and nonlactating dairy cattle for the treatment of the following diseases when caused by one or more of the listed pathogenic organisms sensitive to sulfamethazine: bovine respiratory disease complex (shipping fever complex) associated with *Pasteurella* spp.; bacterial pneumonia associated with *Pasteurella* spp.; necrotic pododermatitis (foot rot) and calf diphtheria caused by *Fusobacterium necrophorum*; colibacillosis (bacterial scours) caused by *Escherichia coli*; coccidiosis caused by *Eimeria bovis* and *E. zurnii*; acute mastitis and metritis caused by *Streptococcus* spp.

(iii) *Limitations*. For use in beef cattle and nonlactating dairy cattle only; if symptoms persist for 2 or 3 days after use, consult a veterinarian; do not slaughter animals for food for at least 8 days after the last dose; do not use in lactating dairy cattle; do not administer more than two consecutive doses.

(g) *Related tolerances*. See § 556.670 of this chapter.

(h)(1) *Sponsor*. See No. 000010 in § 510.600(c) of this chapter for use of an

Food and Drug Administration, HHS

§ 520.2261a

8.25-gram sulfamethazine sustained-release bolus.

(2) *Conditions of use*—(i) *Amount*. Administer at the rate of 1 bolus (8.25 grams per bolus) per 50 pounds of body weight, as a single dose. If signs of disease are significantly reduced, it is recommended that a second dose be given to provide an additional 72 hours of therapy.

(ii) *Indications for use*. Administer orally to ruminating beef and dairy calves for treatment of the following diseases when caused by one or more of the listed pathogenic organisms susceptible to sulfamethazine: bacterial pneumonia associated with *Pasteurella* spp.; colibacillosis (bacterial scours) caused by *Escherichia coli*; coccidiosis caused by *Eimeria bovis* and *E. zurnii*; and calf diphtheria caused by *Fusobacterium necrophorum*.

(iii) *Limitations*. Do not use in calves to be slaughtered under 1 month of age or calves being fed an all milk diet. Do not use in female dairy cattle 20 months of age or older. If symptoms persist after 3 days, consult a veterinarian. Do not administer more than 2 consecutive doses. Do not slaughter animals for food for at least 8 days after the last dose. Do not crush bolus.

[46 FR 36132, July 14, 1981, as amended at 48 FR 18803, Apr. 26, 1983; 48 FR 32760, July 19, 1983; 49 FR 29057, July 18, 1984; 50 FR 49372, Dec. 2, 1985; 51 FR 30212, Aug. 25, 1986; 53 FR 40727, Oct. 18, 1988; 54 FR 14341, Apr. 11, 1989; 55 FR 8462, Mar. 8, 1990; 56 FR 50653, Oct. 8, 1991; 59 FR 22754, May 3, 1994; 61 FR 4875, Feb. 9, 1996; 62 FR 35076, June 30, 1997]

§ 520.2260c Sulfamethazine sustained-release tablets.

(a) *Sponsor*. See No. 053501 in § 510.600(c) of this chapter for use of an 8-gram sulfamethazine sustained-release tablet.

(b) *Conditions of use*—(1) *Amount*. 8 grams (1 tablet) per 45 pounds of body weight as a single dose.

(2) *Indications for use*. In calves for sustained treatment of pneumonia caused by *Pasteurella* spp., colibacillosis (bacterial scours) caused by *Escherichia coli*; and calf diphtheria caused by *Fusobacterium necrophorum*.

(3) *Limitations*. If there is no response within 2 to 3 days, reevaluate therapy. Do not crush tablets. Treated animals must not be slaughtered for food with-

in 18 days after the latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 26763, June 10, 1983, as amended at 56 FR 50653, Oct. 8, 1991; 59 FR 22754, May 3, 1994; 61 FR 4875, Feb. 9, 1996]

§ 520.2261 Sulfamethazine sodium oral dosage forms.

§ 520.2261a Sulfamethazine sodium drinking water solution.

(a) *Sponsors*. See Nos. 017800 and 010042 in § 510.600(c) of this chapter for use of a 12.5-percent sulfamethazine sodium solution.

(b) *Related tolerances in edible products*. See § 556.670 of this chapter.

(c) *Conditions of use*—(1) *Amount*. Administer in drinking water to provide: Cattle and swine 112.5 milligrams of sulfamethazine sodium per pound of body weight per day on the first day and 56.25 milligrams per pound of body weight on subsequent days; Chickens, 61 to 89 milligrams of sulfamethazine sodium per pound of body weight per day, and turkeys 53 to 130 milligrams of sulfamethazine sodium per pound of body weight per day, depending upon the dosage, age, and class of chickens or turkeys, ambient temperature, and other factors.

(2) *Indications for use*. For treatment and control of diseases caused by organisms sensitive to sulfamethazine.

(i) *Beef and nonlactating dairy cattle*. Treatment of bacterial pneumonia and bovine respiratory disease complex (shipping fever complex) (*Pasteurella* spp.), colibacillosis (bacterial scours) (*Escherichia coli*), necrotic pododermatitis (foot rot) (*Fusobacterium necrophorum*), calf diphtheria (*Fusobacterium necrophorum*), acute mastitis (*Streptococcus* spp.), and acute metritis (*Streptococcus* spp.).

(ii) *Swine*. Treatment of porcine colibacillosis (bacterial scours) (*Escherichia coli*), and bacterial pneumonia (*Pasteurella* spp.).

(iii) *Chickens and turkeys*. In chickens for control of infectious coryza (*Haemophilus gallinarum*), coccidiosis (*Eimeria tenella*, *Eimeria necatrix*), acute fowl cholera (*Pasteurella multocida*), and pullorum disease (*Salmonella pullorum*). In turkeys for control of coccidiosis

(*Eimeria meleagritidis*, *Eimeria adenoides*). Medicate as follows: Infectious coryza in chickens, medicate for 2 consecutive days; acute fowl cholera and pullorum disease, in chickens, medicate for 6 consecutive days; coccidiosis, in chickens and turkeys, medicate as in paragraph (c) of this section, then reduce amount of medication to one-half for 4 additional days.

(3) *Limitations*. Add the required dose to that amount of water that will be consumed in 1 day. Consumption should be carefully checked. Have only medicated water available during treatment. Withdraw medication from cattle, chickens, and turkeys 10 days prior to slaughter for food. Withdraw medication from swine 15 days before slaughter for food. Not for use in lactating dairy cattle. Do not medicate chickens or turkeys producing eggs for human consumption. Treatment of all diseases should be instituted early. Treatment should continue 24 to 48 hours beyond the remission of disease symptoms, but not to exceed a total of 5 consecutive days in cattle or swine. Medicated cattle, swine, chickens, and turkeys must actually consume enough medicated water which provides the recommended dosages.

(d) *NAS/NRC status*. The conditions of use specified in this section have been reviewed by NAS/NRC and are found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

[47 FR 25322, June 11, 1982, as amended at 47 FR 25735, June 15, 1982]

§ 520.2261b Sulfamethazine sodium soluble powder.

(a) *Sponsor*. See No. 010042 in § 510.600(c) of this chapter for use of a soluble powder composed of 100 percent sulfamethazine sodium.

(b) *Related tolerances in edible products*. See § 556.670 of this chapter.

(c) *Conditions of use*—(1) *Amount*. Administer in drinking water to provide: Chickens 58 to 85 milligrams of sulfamethazine sodium per pound of body weight per day; turkeys 50 to 124 milligrams of sulfamethazine sodium per pound of body weight per day; depending upon the dosage, age, and class

of chickens or turkeys, ambient temperature, and other factors. Administer to cattle and swine in drinking water, or as a drench, to provide 108 milligrams of sulfamethazine sodium per pound of body weight on the first day and 54 milligrams of sulfamethazine sodium per pound of body weight per day on the second, third, and fourth days of administration.

(2) *Indications for use*. For treatment and control of disease caused by organisms sensitive to sulfamethazine.

(i) *Beef and nonlactating dairy cattle*. Treatment of bacterial pneumonia and bovine respiratory disease complex (shipping fever complex) (*Pasteurella* spp.), colibacillosis (bacterial scours) (*Escherichia coli*), necrotic pododermatitis (foot rot) (*Fusobacterium necrophorum*), calf diphtheria (*Fusobacterium necrophorum*), acute mastitis (*Streptococcus* spp.), and acute metritis (*Streptococcus* spp.).

(ii) *Swine*. Treatment of porcine colibacillosis (bacterial scours) (*Escherichia coli*), and bacterial pneumonia (*Pasteurella* spp.).

(iii) *Chickens and turkeys*. In chickens for control of infectious coryza (*Haemophilus gallinarum*), coccidiosis (*Eimeria tenella*, *Eimeria necatrix*), acute fowl cholera (*Pasteurella multocida*), and pullorum disease (*Salmonella pullorum*). In turkeys for control of coccidiosis (*Eimeria meleagritidis*, *Eimeria adenoides*). Medicate as follows: Infectious coryza in chickens, medicate for 2 consecutive days; acute fowl cholera and pullorum disease in chickens, medicate for 6 consecutive days; coccidiosis in chickens and turkeys, medicate as in paragraph (c) of this section for 2 days, then reduce drug concentration to one-half for 4 additional days.

(3) *Limitations*. Add the required dose to that amount of water that will be consumed in 1 day. Consumption should be carefully checked. Have only medicated water available during treatment. Withdraw medication from cattle, chickens, and turkeys 10 days prior to slaughter for food. Withdraw medication from swine 15 days prior to slaughter for food. Not for use in lactating dairy animals. Do not medicate chickens or turkeys producing eggs for human consumption. Treatment of all diseases should be instituted early.

Treatment should continue 24 to 48 hours beyond the remission of disease symptoms, but not to exceed a total of 5 consecutive days in cattle or swine. Medicated cattle, swine, chickens, and turkeys must actually consume enough medicated water which provides the recommended dosages.

(d) *NAS/NRC status.* The conditions of use specified in this section have been reviewed by NAS/NRC and are found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

[47 FR 25322, June 11, 1982]

§ 520.2280 Sulfamethizole and methenamine mandelate tablets.

(a) *Specifications.* Each tablet contains 250 milligrams of sulfamethizole and 250 milligrams of methenamine mandelate.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is indicated for the treatment of urinary tract infections in dogs and cats such as cystitis, nephritis, prostatitis, urethritis, and pyelonephritis. It is also used as an aid in the management of complications resulting from surgical manipulations of the urinary tract such as removal of calculi from the bladder, in ureterostomies, and in instrumentation of the urethra and bladder.

(2) It is administered at a dosage level of one tablet for each 20 pounds of body weight given three times per day. The drug should be given until all signs are alleviated. To reduce the possibility of a relapse, it is suggested that therapy be continued for a further period of a week to 10 days.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 50 FR 13561, Apr. 5, 1985]

§ 520.2320 Sulfanitrans and aklomide in combination.

(a) *Chemical names.* (1) Sulfanitrans: Acetyl-(p-nitrophenyl)-sulfanilamide.

(2) Aklomide: 2-Chloro-4-nitrobenzamide.

(b) *Specifications.* (1) Sulfanitrans conform to the following specifications:

(i) Melting point range: 260° C. to 261° C.

(ii) Assay (by sodium nitrite titration): 97 to 100.5 percent.

(iii) Moisture (Method No. 6.123, "Toluene Distillation Method—Official Final Action" in "Official Methods of Analysis of the Association of Official Analytical Chemists," 13th Ed., 1980, p. 83. Copies are available from the Association of Official Analytical Chemists, 2200 Wilson Blvd., Suite 400, Arlington, VA 22201-3301, or available for inspection at the Office of the Federal Register, 800 North Capitol Street, NW., suite 700, Washington, DC 20001): Not more than 2.0 percent.

(iv) Molecular weight: 335.34.

(v) Soluble in 0.1N sodium hydroxide, reprecipitating unchanged on acidification.

(2) Aklomide conforms to the following specifications:

(i) Minimum melting point: 170° C.

(ii) Moisture content: Not to exceed 1.0 percent.

(iii) Purity: Not less than 98 percent on an anhydrous basis.

(c) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(d) *Related tolerances.* See §§ 556.30 and 556.680 of this chapter.

(e) *Conditions of use.* It is used in the drinking water of chickens as follows:

(1) *Amount.* 374-747 milligrams of sulfanitrans with 477-954 milligrams of aklomide.

(2) *Indications for use.* As an aid in the treatment of coccidiosis caused by *E. tenella*, *E. necatrix*, and *E. acervulina*.

(3) *Limitations.* Administer for 2 days at 747 milligrams of sulfanitrans per gallon and 954 milligrams of aklomide per gallon, followed by 5 days at 374 milligrams of sulfanitrans per gallon and 477 milligrams of aklomide per gallon; do not treat birds over 6 weeks of age; do not administer within 5 days of slaughter; not for laying chickens.

[40 FR 13838, Mar. 27, 1975, as amended at 47 FR 9396, Mar. 5, 1982; 54 FR 18280, Apr. 28, 1989; 55 FR 8460, Mar. 8, 1990]

§ 520.2325

21 CFR Ch. I (4–1–99 Edition)

§ 520.2325 Sulfaquinoxaline oral dosage forms.

§ 520.2325a Sulfaquinoxaline drinking water.

(a) *Sponsor.* See § 510.600(c) of this chapter for identification of the sponsors.

(1) To No. 050749 for use of a 25-percent sulfaquinoxaline soluble powder and a 20-percent sulfaquinoxaline sodium solution as provided for in paragraph (c) of this section.

(2) To No. 060594 for use of 3.44- and 12.85-percent sulfaquinoxaline sodium solutions as provided for in paragraphs (c)(1), (c)(2), (c)(3), (c)(4)(i), and (c)(4)(ii) of this section.

(3) To No. 017144 for use of a 31.92-percent sulfaquinoxaline solution (sodium and potassium salts) as provided for in paragraphs (c)(1), (c)(2), (c)(3), (c)(4)(i), and (c)(4)(ii) of this section.

(4) No. 053501 for use of a 28.62-percent sulfaquinoxaline sodium solution as provided in paragraphs (c)(1), (c)(2), and (c)(3) of this section.

(b) *Related tolerances.* See § 556.685 of this chapter.

(c) *Conditions of use.* It is used in drinking water as follows:

(1) *Chickens.* (i) As an aid in the control of outbreaks of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, and *E. brunetti*.

(ii) Administer at the 0.04 percent level for 2 or 3 days, skip 3 days then administer at the 0.025 percent level for 2 more days. If bloody droppings appear, repeat treatment at the 0.025 percent level for 2 more days. Do not change litter unless absolutely necessary. Do not give flushing mash.

(2) *Turkeys.* (i) As an aid in the control of outbreaks of coccidiosis caused by *Eimeria meleagritidis* and *E. adenoeides*.

(ii) Administer at the 0.025 percent level for 2 days, skip 3 days, give for 2 days, skip 3 days and give for 2 more days. Repeat if necessary. Do not change litter unless absolutely necessary. Do not give flushing mash.

(3) *Chickens and turkeys.* (i) As an aid in the control of acute fowl cholera caused by *Pasteurella multocida* susceptible to sulfaquinoxaline and fowl typhoid caused by *Salmonella gallinarum* susceptible to sulfaquinoxaline.

(ii) Administer at the 0.04 percent level for 2 or 3 days. Move birds to clean ground. If disease recurs, repeat treatment. If cholera has become established as the respiratory or chronic form, use feed medicated with sulfaquinoxaline. Poultry which have survived typhoid outbreaks should not be kept for laying house replacements or breeders unless tests show they are not carriers.

(4) *Cattle and calves.* (i) For the control and treatment of outbreaks of coccidiosis caused by *Eimeria bovis* or *E. zurnii*.

(ii) Administer at the 0.015-percent level for 3 to 5 days in drinking water medicated with sulfaquinoxaline solution.

(iii) In lieu of treatment as provided in paragraph (e)(4)(ii) of this section, administer 1 teaspoon of 25-percent sulfaquinoxaline soluble powder per day for each 125 pounds of body weight for 3 to 5 days in drinking water.

(d) *Limitations.* Consult a veterinarian or poultry pathologist for diagnosis. May cause toxic reactions unless the drug is evenly mixed in water at dosages indicated and used according to directions. For control of outbreaks of disease, medication should be initiated as soon as the diagnosis is determined. Medicated chickens, turkeys, cattle, and calves must actually consume enough medicated water which provides a recommended dosage of approximately 10 to 45 milligrams per pound per day in chickens, 3.5 to 55 milligrams per pound per day in turkeys, and approximately 6 milligrams per pound per day in cattle and calves depending on the age, class of animal, ambient temperature, and other factors. A withdrawal period has not been established for sulfaquinoxaline in preruminating calves. Do not use in calves to be processed for veal. Not for use in lactating dairy cattle. Do not give to chickens, turkeys or cattle within 10 days of slaughter for food. Do not medicate chickens or turkeys producing eggs for human consumption. Make fresh drinking water daily.

[48 FR 3964, Jan. 28, 1983, as amended at 48 FR 26762, June 10, 1983; 55 FR 29843, July 23, 1990; 59 FR 28769, June 3, 1994; 59 FR 33197, June 28, 1994; 61 FR 24443, May 15, 1996; 61 FR 63711, Dec. 2, 1996; 62 FR 37712, July 15, 1997]

§ 520.2325b Sulfaminoxaline drench.

(a)–(b) [Reserved]

(c) *Sponsor.* See No. 050749 in § 510.600(c) of this chapter.

(d) *NAS/NRC status.* The conditions of use specified in this section have been reviewed by NAS/NRC and are found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency information. Applications must be accompanied by a written commitment to undertake the human safety studies required by FDA.

(e) *Conditions of uses.* As a 25-percent sulfaminoxaline soluble powder.

(1) For the control and treatment of outbreaks of coccidiosis in cattle and calves caused by *Eimeria bovis* or *E. zurnii*.

(2) Give one teaspoon of 25 percent sulfaminoxaline soluble powder for each 125 pounds of body weight for 3 to 5 days as a drench.

(f) *Limitations.* For control of outbreaks of disease, medication should be initiated as soon as the diagnosis is determined. Consult a veterinarian for diagnosis. Do not give to cattle within 10 days of slaughter for food. Not for use in lactating dairy cattle.

[48 FR 3964, Jan. 28, 1983, as amended at 55 FR 29843, July 23, 1990; 59 FR 33197, June 28, 1994]

§ 520.2330 Sulfisoxazole tablets.

(a) *Specifications.* Each tablet contains 260 milligrams (4 grains) of sulfisoxazole.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* Administer one tablet orally per 4 pounds of body weight.¹

(2) *Indications for use.* Use in dogs and cats as an aid in treatment of bacterial pneumonia and bacterial enteritis when caused by organisms sensitive to sulfisoxazole.¹

(3) *Limitations.* Repeat dosage at 24-hour intervals until 2 to 3 days after

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

disappearance of clinical symptoms. (Administration of one-half daily dosage at 12-hour intervals or one-third daily dosage at 8-hour intervals will provide a more constant blood level.) Provide adequate supply of drinking water. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian.¹

[43 FR 60895, Dec. 29, 1978]

§ 520.2345 Tetracycline oral dosage forms.**§ 520.2345a Tetracycline hydrochloride capsules.**

(a) *Specifications.* Each capsule contains 50, 100, 125, 250, or 500 milligrams of tetracycline hydrochloride.

(b) *Sponsor.* See § 510.600(c) of this chapter for identification of the sponsors:

(1) To No. 000009: 250 milligrams per capsule.

(2) To No. 000069: 125, 250, and 500 milligrams per capsule.

(3) To No. 000115: 50, 100, 250, and 500 milligrams per capsule.

(c) *Conditions of use. Dogs—(1) Amount.* 25 milligrams per pound of body weight per day in divided doses every 6 hours.

(2) *Indications for use.* Treatment of infections caused by organisms sensitive to tetracycline hydrochloride, such as bacterial gastroenteritis due to *E. coli* and urinary tract infections due to *Staphylococcus* spp. and *E. coli*.

(3) *Limitations.* Administer orally; continue treatment until symptoms of the disease have subsided and the temperature is normal for 48 hours; not for use in animals raised for food production; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37327, Aug. 18, 1992, as amended at 59 FR 59365, Nov. 17, 1994; 63 FR 5255, Feb. 2, 1998]

§ 520.2345b Tetracycline tablets.

(a) *Specifications.* Each tablet contains 100, 250, or 500 milligrams of tetracycline (as the hydrochloride).

(b) *Sponsor.* For 100, 250, or 500 milligrams per tablet, see No. 000069 in § 510.600(c) of this chapter. For 250 milligrams per tablet, see No. 000009 in § 510.600(c) of this chapter.

§ 520.2345c

(c) *Conditions of use. Dogs*—(1) *Amount.* 25 milligrams per pound of body weight per day in divided doses every 6 hours.

(2) *Indications for use.* Treatment of infections caused by organisms sensitive to tetracycline hydrochloride, such as bacterial gastroenteritis due to *E. coli* and urinary tract infections due to *Staphylococcus* spp. and *E. coli*.

(3) *Limitations.* Administer orally; continue treatment until symptoms of the disease have subsided and temperature is normal for 48 hours; not for use in animals raised for food production; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37327, Aug. 18, 1992]

§ 520.2345c Tetracycline boluses.

(a) *Specifications.* Each bolus contains 500 milligrams of tetracycline (as the hydrochloride).

(b) *Sponsors.* See No. 010042 in § 510.600(c) of this chapter for use as in paragraph (d)(1) of this section. See No. 000009 in § 510.600(c) of this chapter for use as in paragraph (d)(2) of this section.

(c) *Related tolerances.* See § 556.720 of this chapter.

(d) *Conditions of use. Calves*—(1) *Amount.* 10 milligrams per pound of body weight per day in divided doses.

(i) *Indications for use.* Control and treatment of bacterial enteritis (scours) caused by *E. coli* and bacterial pneumonia caused by *Pasteurella* spp., *Hemophilus* spp., and *Klebsiella* spp.

(ii) *Limitations.* Administer orally for 3 to 5 days; do not slaughter animals for food within 14 days of treatment; use as sole source of tetracycline.

(iii) *National Academy of Sciences/National Research Council (NAS/NRC) status.* The conditions of use specified in paragraph (d)(1)(i) of this section were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified in § 514.111 of this chapter, but may require bioequivalency and safety information.

(2) *Amount.* 10 milligrams per pound of body weight per day in two divided doses.

(i) *Indications for use.* Treatment of bacterial pneumonia caused by orga-

21 CFR Ch. I (4–1–99 Edition)

nisms susceptible to tetracycline, bacterial enteritis caused by *E. coli*, and salmonella organisms susceptible to tetracycline.

(ii) *Limitations.* Administer orally for not more than 5 days; do not slaughter animals for food within 12 days of treatment; use as sole source of tetracycline.

[57 FR 37328, Aug. 18, 1992]

§ 520.2345d Tetracycline hydrochloride soluble powder.

(a) *Sponsors.* The following sponsors listed in § 510.600(c) of this chapter hold approvals for the drug concentrations (i.e., grams of tetracycline hydrochloride per pound) and conditions of use indicated:

(1) 046573, 051259, 054273, 057561, and 059130 102.4 and 324 grams per pound as in paragraph (d) of this section.

(2) 000069, 25 grams per pound as in paragraphs (d)(3) and (d)(4) of this section.

(3) 010042, 102.4 and 324 grams per pound as in paragraphs (d)(1) and (d)(2) of this section.

(4) 046573, 102.4 and 324 grams per pound as in paragraph (d)(3) of this section.

(b) *Related tolerances.* See § 556.720 of this chapter.

(c) [Reserved]

(d) *Conditions of use in drinking water*—(1) *Calves*—(i) *Amount.* 10 milligrams per pound of body weight per day in divided doses.

(ii) *Indications for use.* Control and treatment of bacterial enteritis (scours) caused by *Escherichia coli* and bacterial pneumonia (shipping fever complex) associated with *Pasteurella* spp., *Actinobacillus pleuropneumoniae* (*Hemophilus* spp.), and *Klebsiella* spp., susceptible to tetracycline.

(iii) *Limitations.* Administer for 3 to 5 days; do not slaughter animals for food within 4 days of treatment for sponsor 010042 and within 5 days of treatment for sponsors 046573 and 000010; prepare a fresh solution daily; use as the sole source of tetracycline.

(2) *Swine*—(i) *Amount.* 10 milligrams per pound of body weight per day in divided doses.

(ii) *Indications for use.* Control and treatment of bacterial enteritis (scours) caused by *E. coli* and bacterial

Food and Drug Administration, HHS

§ 520.2345f

pneumonia associated with *Pasteurella* spp., *A. pleuropneumoniae* (*Hemophilus* spp.), and *Klebsiella* spp., susceptible to tetracycline.

(iii) *Limitations*. Administer for 3 to 5 days; do not slaughter animals for food within 7 days of treatment for sponsor 010042 and within 4 days of treatment for sponsors 046573 and 000010; prepare a fresh solution daily; use as the sole source of tetracycline.

(3) *Chickens*—(i) *Amount*. Chronic respiratory disease: 400 to 800 milligrams per gallon. Infectious synovitis: 200 to 400 milligrams per gallon.

(ii) *Indications for use*. Control of chronic respiratory disease (CRD or air-sac disease) caused by *Mycoplasma gallisepticum* and *E. coli*; control of infectious synovitis caused by *M. synoviae* susceptible to tetracycline.

(iii) *Limitations*. Administer for 7 to 14 days; do not slaughter for food within 4 days of treatment; not for use in chickens producing eggs for human consumption; prepare a fresh solution daily; use as the sole source of tetracycline.

(4) *Turkeys*—(i) *Amount*. For infectious synovitis: 400 milligrams per gallon. For complicating bacterial organisms associated with bluecomb (transmissible enteritis or coronaviral enteritis): 25 milligrams per pound of body weight per day.

(ii) *Indications for use*. Control of infectious synovitis caused by *M. synoviae*; control of bluecomb complicated by organisms sensitive to tetracycline.

(iii) *Limitations*. Administer for 7 to 14 days; do not slaughter for food within 4 days of treatment; not for use in turkeys producing eggs for human consumption; prepare a fresh solution daily; use as the sole source of tetracycline.

[59 FR 17693, Apr. 14, 1994, as amended at 59 FR 19133, Apr. 22, 1994; 62 FR 5319, Feb. 5, 1997; 62 FR 35076, June 30, 1997; 62 FR 46668, Sept. 4, 1997; 62 FR 55160, Oct. 23, 1997]

§ 520.2345e Tetracycline oral liquid.

(a) *Specifications*. Each milliliter contains the equivalent of either 25 or 100 milligrams of tetracycline hydrochloride.

(b) *Sponsor*. See No. 000069, in § 510.600(c) of this chapter for use of 25

or 100 milligrams per milliliter liquid in dogs as in paragraph (c)(1) of this section; see No. 000009 in § 510.600(c) of this chapter for use of 100 milligrams per milliliter liquid in dogs and cats as in paragraph (c)(2).

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount*. 25 milligrams per pound of body weight per day in divided doses every 6 hours.

(ii) *Indications for use*. Treatment of infections caused by organisms sensitive to tetracycline hydrochloride, such as bacterial gastroenteritis due to *Escherichia coli* and urinary tract infections due to *Staphylococcus* spp. and *E. coli*.

(iii) *Limitations*. Administer orally; continue treatment until symptoms have subsided and the temperature is normal for 48 hours; not for use in animals which are raised for food production; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(iv) *National Academy of Sciences/National Research Council (NAS/NRC) status*. These conditions were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(2) *Dogs and cats*—(i) *Amount*. 25 milligrams per pound of body weight per day in divided doses every 6 hours.

(ii) *Indications for use*. Treatment of infections caused by organisms susceptible to tetracycline hydrochloride, such as bacterial gastroenteritis due to *E. coli* and urinary tract infections due to *Staphylococcus* spp. and *E. coli*.

(iii) *Limitations*. Administer orally; continue treatment until the temperature has been normal for 48 hours; not for use in food-producing animals; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37329, Aug. 18, 1992]

§ 520.2345f Tetracycline phosphate complex and sodium novobiocin capsules.

(a) *Specifications*. Each capsule contains the equivalent of 60 milligrams of tetracycline hydrochloride and 60 milligrams of novobiocin.

(b) *Sponsor*. No. 000009 in § 510.600(c) of this chapter.

§ 520.2345g

(c) *Conditions of use. Dogs*—(1) *Amount.* 10 milligrams of each antibiotic per pound of body weight (1 capsule for each 6 pounds) every 12 hours.

(2) *Indications for use.* Treatment of acute or chronic canine respiratory infections such as tonsillitis, bronchitis, and tracheobronchitis when caused by pathogens susceptible to tetracycline and/or novobiocin, such as *Staphylococcus* spp. and *Escherichia coli*.

(3) *Limitations.* Continue treatment for at least 48 hours after the temperature has returned to normal and all evidence of infection has disappeared. As with all antibiotics, appropriate in vitro culturing and susceptibility tests of samples taken before treatment should be conducted. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37329, Aug. 18, 1992]

§ 520.2345g Tetracycline hydrochloride and sodium novobiocin tablets.

(a) *Specifications.* Each tablet contains the equivalent of 60 milligrams of tetracycline hydrochloride and 60 milligrams of novobiocin, or 180 milligrams of tetracycline hydrochloride and 180 milligrams of novobiocin.

(b) *Sponsor.* No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use. Dogs*—(1) *Amount.* 10 milligrams of each antibiotic per pound of body weight (one single-strength tablet for each 6 pounds or one triple-strength tablet for each 18 pounds).

(2) *Indications for use.* Treatment of acute or chronic canine respiratory infections such as tonsillitis, bronchitis, and tracheobronchitis when caused by pathogens susceptible to tetracycline and/or novobiocin, such as *Staphylococcus* spp. and *Escherichia coli*.

(3) *Limitations.* Continue treatment for at least 48 hours after the temperature has returned to normal and all evidence of infection has disappeared. As with all antibiotics, appropriate in vitro culturing and susceptibility tests of samples taken before treatment should be conducted. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37329, Aug. 18, 1992]

21 CFR Ch. I (4–1–99 Edition)

§ 520.2345h Tetracycline hydrochloride, sodium novobiocin, and prednisolone tablets.

(a) *Specifications.* Each tablet contains the equivalent of 60 milligrams of tetracycline hydrochloride, 60 milligrams of novobiocin, and 1.5 milligrams of prednisolone or 180 milligrams of tetracycline hydrochloride, 180 milligrams of novobiocin, and 4.5 milligrams of prednisolone.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use. Dogs*—(1) *Amount.* 10 milligrams of each antibiotic and 0.25 milligram of prednisolone per pound of body weight (one single-strength tablet for each 6 pounds or one triple-strength tablet for each 18 pounds) every 12 hours for 48 hours. Treatment is to be continued with novobiocin and tetracycline alone at the same dose schedule for an additional 3 days or longer as needed.

(2) *Indications for use.* Treatment of acute and chronic canine respiratory infections such as tonsillitis, bronchitis, and tracheobronchitis when caused by pathogens susceptible to tetracycline and/or novobiocin, such as *Staphylococcus* spp. and *Escherichia coli*, when it is necessary to initially reduce the severity of associated clinical signs.

(3) *Limitations.* As with all antibiotics, appropriate in vitro culturing and susceptibility tests of samples taken before treatment should be conducted. Administer for 48 hours only. Continue treatment if needed with tetracycline/novobiocin alone. The product is contraindicated in animals with tuberculosis, hyperadrenocorticalism, or peptic ulcers. Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37329, Aug. 18, 1992]

§ 520.2362 Thenium closylate tablets.

(a) *Chemical name.* (N,N-Dimethyl-N-2-phenoxyethyl-N-2'-thenylammonium)-p-chlorobenzene-sulfonate.

(b) *Specifications.* Thenium closylate tablets contain thenium closylate equivalent to 500 milligrams thenium as base in each tablet.

(c) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) The tablets are administered orally to dogs as a single day treatment of canine ancylostomiasis by the removal from the intestines of the adult forms of the species *Ancylostoma caninum* and *Uncinaria stenocephala* (hookworms). Dogs weighing 10 pounds and over are administered 1 tablet as a single dose. Dogs weighing 5 to 10 pounds are administered one-half tablet twice during a single day. All dosages are given for 1 day only. The treatment should be repeated after 2 or 3 weeks.

(2) Suckling puppies or recently weaned puppies weighing less than 5 pounds should not be treated with the drug. Animals that are severely infected, exhibiting evidence of intestinal hemorrhage, debilitation, and anemia, should be given supportive treatment.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 41 FR 53477, Dec. 7, 1976; 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 520.2380 Thiabendazole oral dosage forms.**§ 520.2380a Thiabendazole top dressing and mineral protein feed block.**

(a) *Chemical name.* 2-(4-Thiazolyl)-benzimidazole.

(b) *Specifications.* Conforms to N.F. XII.

(c) *Sponsor.* (1) See No. 017800 in § 510.600(c) of this chapter for the sponsor of the usage provided by paragraph (e)(1)(ii) of this section.

(2) See No. 050604 in § 510.600(c) of this chapter for the sponsor of the usages provided for by paragraph (e)(1)(ii) of this section.

(3) See No. 021930 in § 510.600(c) of this chapter for the sponsor of the usage

provided for by paragraph (e)(2) of this section.

(d) *Related tolerances.* See § 556.730 of this chapter.

(e) *Conditions of use.* It is used as follows:

(1) *Horses*—(i) *Route of administration.* In feed, as a top dressing.

(a) *Amount.* 2 grams per 100 pounds of body weight.

(b) *Indications for use.* For control of large strongyles, small strongyles, pinworms, and threadworms (including members of the genera *Strongylus*, *Cyathostomum*, *Cylicobrachytus*, and related genera, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*, *Oxyuris*, and *Strongyloides*).

(c) *Limitations.* Add to the usual feed of horses mixed into that amount of the feed normally consumed at one feeding. Warning: Not for use in horses intended for food.

(ii) *Route of administration.* In feed.

(a) *Amount.* 2 grams per 100 pounds of body weight.

(1) *Indications for use.* For control of large and small strongyles, *Strongyloides*, and pinworms of the genera *Strongylus*, *Cyathostomum*, *Cylicobrachytus* and related genera, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*, *Oxyuris*, and *Strongyloides*.

(2) *Limitations.* Administer in a single dosage mixed with the normal grain ration given at one feeding. Warning: Not for use in horses intended for food.

(b) *Amount.* 4 grams per 100 pounds of body weight.

(1) *Indications for use.* For control of ascarids of the genus *Parascaris*.

(2) *Limitations.* Administer in a single dosage mixed with the normal grain ration given at one feeding. Warning: Not for use in horses intended for food.

(2) *Cattle*—(i) *Route of administration.* In feed block.

(ii) *Amount.* 3.3 percent block consumed at the recommended level of 0.11 pound per 100 pounds of body weight per day.

(iii) *Indications for use.* For control of infections of gastrointestinal roundworms (*Trichostrongylus*, *Haemonchus*, *Ostertagia* and *Cooperia*).

(iv) *Limitations.* Administer to cattle on pasture or range accustomed to mineral protein block feeding for 3

§ 520.2380b

days. Milk taken from animals during treatment and within 96 hours (8 milkings) after the latest treatment must not be used for food. Do not treat cattle within 3 days of slaughter. For a satisfactory diagnosis, a microscopic fecal examination should be performed by a veterinarian or diagnostic laboratory prior to worming. Animals maintained under conditions of constant worm exposure may require re-treatment within 2 to 3 weeks. Animals that are severely parasitized, sick, or off feed should be isolated and a veterinarian consulted for advice concerning treatment.

[40 FR 13838, Mar. 27, 1975, as amended at 41 FR 9149, Mar. 3, 1976; 62 FR 63271, Nov. 28, 1997]

§ 520.2380b Thiabendazole drench or oral paste.

(a) *Chemical name.* 2-(4-Thiazolyl) benzimidazole.

(b) *Specifications.* Conforms to N.F. XII.

(c) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter for the sponsor of the usages provided for by paragraph (e) of this section.

(d) *Related tolerances.* See § 556.730 of this chapter.

(e) *Conditions of use.* It is used as follows:

(1) *Horses.* As a single liquid oral dose, administered as a drench or by stomach tube; or as an oral paste.

(i) *Amount.* 2 grams per 100 pounds of body weight.

(a) *Indications for use.* For the control of infections of large strongyles (*Strongylus vulgaris*, *Strongylus edentatus*), small strongyles (*Cyathostomum*, *Cylicobrachytus* and related genera, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*), pinworms (*Oxyuris*), and threadworms (*Strongyloides*).

(b) *Limitations.* Not for use in horses to be slaughtered for food purposes. When administered by stomach tube, for use only by or on the order of a licensed veterinarian. When for use as a liquid oral drench or an oral paste, consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(ii) *Amount.* 4 grams per 100 pounds of body weight.

21 CFR Ch. I (4–1–99 Edition)

(a) *Indications for use.* For control of infections of ascaridis (*Parascaris*).

(b) *Limitations.* Not for use in horses to be slaughtered for food purposes. When administered by stomach tube, use only by or on the order of a licensed veterinarian. When for use as a liquid oral drench or an oral paste, consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) *Pigs.* As an oral paste.

(i) *Amount.* 200 milligrams for each 5 to 7 pounds of body weight per dose.

(ii) *Indications for use.* For control of infections with *Strongyloides ransomi*. These infections are commonly found in Southeastern United States.

(iii) *Limitations.* Administer to baby pigs (1 to 8 weeks of age). Treatment may be repeated in 5 to 7 days if necessary. Before treatment, obtain an accurate diagnosis from a veterinarian or diagnostic laboratory. Do not treat within 30 days of slaughter.

(3) *Cattle.* Orally as a drench and in paste form using a dosing gun designed for the product.

(i) *Amount.* 3 grams per 100 pounds of body weight.

(a) *Indications for use.* Control of infections of gastrointestinal roundworms (*Trichostrongylus* spp., *Haemonchus* spp., *Nematodirus* spp., *Ostertagia* spp., and *Oesophagostomum radiatum*).

(b) *Limitations.* For most effective results, severely parasitized animals or those constantly exposed to helminth infection should be re-treated every 2 to 3 weeks. Milk taken from treated animals within 96 hours (8 milkings) after the latest treatment must not be used for food. Do not treat cattle within 3 days of slaughter. For a satisfactory diagnosis, a microscopic fecal examination should be performed prior to worming.

(ii) *Amount.* 5 grams per 100 pounds of body weight.

(a) *Indications for use.* Control of infections of *Cooperia* spp. or severe infections of other species in paragraph (e)(3)(i)(a) of this section.

(b) *Limitations.* For most effective results, severely parasitized animals or those constantly exposed to helminth infection should be re-treated every 2 to 3 weeks. Milk taken from treated

Food and Drug Administration, HHS

§ 520.2380c

animals within 96 hours (8 milkings) after the latest treatment must not be used for food. Do not treat cattle within 3 days of slaughter. For a satisfactory diagnosis, a microscopic fecal examination should be performed prior to worming.

(4) *Sheep and goats*. Orally, as a drench.

(i) *Amount*. 2 grams per 100 pounds of body weight.

(ii) *Indications for use*. Control of infections of gastrointestinal roundworms in sheep and goats. (*Trichostrongylus* spp., *Haemonchus* spp., *Ostertagia* spp., *Cooperia* spp., *Nematodirus* spp., *Bunostomum* spp., *Strongyloides* spp., *Chabertia* spp., and *Oesophagostomum* spp.); also active from 3 hours to 3 days following treatment against ova and larvae passed by sheep (good activity against *Trichostrongylus colubriformis* and *axei*, *Ostertagia* spp., *Bunostomum* spp., *Nematodirus* spp., and *Strongyloides* spp.; less effective against *Haemonchus contortus* and *Oesophagostomum* spp.).

(iii) *Limitations*. As a single oral dose; do not treat animals within 30 days of slaughter; milk taken from treated animals within 96 hours (8 milkings) after the latest treatment must not be used for food; in severe infections in sheep, treatment should be repeated in 2 to 3 weeks.

(5) *Goats*. Orally, as a drench.

(i) *Amount*. 3 grams per 100 pounds of body weight.

(ii) *Indications for use*. Control of severe infections of gastrointestinal roundworms (*Trichostrongylus* spp., *Haemonchus* spp., *Ostertagia* spp., *Cooperia* spp., *Nematodirus* spp., *Bunostomum* spp., *Strongyloides* spp., *Chabertia* spp., and *Oesophagostomum* spp.).

(iii) *Limitations*. As a single oral dose; do not treat animals within 30 days of slaughter; milk taken from treated animals within 96 hours (8 milkings) after the latest treatment must not be used for food; treatment should be repeated in 2 to 3 weeks.

[40 FR 13838, Mar. 27, 1975, as amended at 41 FR 9149, Mar. 3, 1976; 41 FR 47424, Oct. 29, 1976; 62 FR 63271, Nov. 28, 1997]

§ 520.2380c Thiabendazole bolus.

(a) *Chemical name*. 2-(4-Thiazolyl) benzimidazole.

(b) *Specifications*. Conforms to N.F. XII.

(c) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(d) *Related tolerances*. See § 556.730 of this chapter.

(e) *Conditions of use*. It is used as follows:

(1) *Cattle*. In a bolus.

(i) *Amount*. 3 grams per 100 pounds of body weight.

(a) *Indications for use*. Control of infections of gastrointestinal roundworms (general *Trichostrongylus* spp., *Haemonchus* spp., *Nematodirus* spp., *Ostertagia* spp., and *Oesophagostomum radiatum*).

(b) *Limitations*. As a single oral dose; may repeat once in 2 to 3 weeks; do not treat animals within 3 days of slaughter; milk taken from treated animals within 96 hours (8 milkings) after the latest treatment must not be used for food.

(ii) *Amount*. 5 grams per 100 pounds of body weight.

(a) *Indications for use*. Control of severe infections of gastrointestinal roundworms (genera *Trichostrongylus* spp., *Haemonchus* spp., *Nematodirus* spp., *Ostertagia* spp., and *Oesophagostomum radiatum*). Control of infections with *Cooperia* spp.

(b) *Limitations*. As a single oral dose; as a drench or bolus; may repeat once in 2 to 3 weeks; do not treat animals within 3 days of slaughter; milk taken from treated animals within 96 hours (8 milkings) after the latest treatment must not be used for food.

(2) *Sheep and goats*. In a bolus.

(i) *Amount*. 2 grams per 100 pounds of body weight.

(ii) *Indications for use*. Control of infections of gastrointestinal roundworms in sheep and goats (general *Trichostrongylus* spp., *Haemonchus* spp., *Ostertagia* spp., *Cooperia* spp., *Nematodirus* spp., *Bunostomum* spp., *Strongyloides* spp., *Chabertia* spp., and *Oesophagostomum* spp.); also active from 3 hours to 3 days following treatment against ova and larvae passed by sheep (good activity against *T. colubriformis* and *axei*, *Ostertagia* spp., *Bunostomum* spp., *Nematodirus* spp., and

§ 520.2380d

Strongyloides spp.; less effective against *Haemonchus contortus* and *Oesophagostomum* spp.).

(iii) *Limitations*. As a single oral dose; do not treat animals within 30 days of slaughter; milk taken from treated animals within 96 hours (8 milkings) after the latest treatment must not be used for food; in severe infections in sheep, treatment should be repeated in 2 to 3 weeks.

(3) *Goats*. In a bolus.

(i) *Amount*. 3 grams per 100 pounds of body weight.

(ii) *Indications for use*. Control of severe infections of gastrointestinal roundworms (genera *Trichostrongylus* spp., *Haemonchus* spp., *Ostertagia* spp., *Cooperia* spp., *Nematodirus* spp., *Bunostomum* spp., *Strongyloides* spp., *Chabertia* spp., and *Oesophagostomum* spp.).

(iii) *Limitations*. As a single oral dose; do not treat animals within 30 days of slaughter; milk taken from treated animals within 96 hours (8 milkings) after the latest treatment must not be used for food; treatment should be repeated in 2 to 3 weeks.

[40 FR 13838, Mar. 27, 1975, as amended at 41 FR 9149, Mar. 3, 1976; 62 FR 63271, Nov. 28, 1997]

§ 520.2380d Thiabendazole, piperazine citrate suspension.

(a) *Specifications*. Each fluid ounce of suspension contains 2 grams of thiabendazole and 2.5 grams of piperazine (from piperazine citrate).

(b) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) It is administered to horses by stomach tube or as a drench at the rate of 1 fluid ounce of suspension per 100 pounds of body weight for the control of large strongyles, small strongyles, pinworms, *Strongyloides* and ascarids (including members of the genera *Strongylus* spp., *Cyathostomum* spp., *Cylicobrachytus* spp. and related genera *Craterostomum* spp., *Oesophagodontus* spp., *Poteriostomum* spp., *Oxyuris* spp., *Strongyloides* spp., and *Parascaris* spp.).

(2) Do not use in horses intended to be used for food purposes.

21 CFR Ch. I (4–1–99 Edition)

(3) For use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 62 FR 63271, Nov. 28, 1997]

§ 520.2380e Thiabendazole with trichlorfon.

(a) *Specifications*. The drug contains 5 grams of thiabendazole with 4.5 grams of trichlorfon, or 20 grams of thiabendazole with 18 grams of trichlorfon.

(b) *Sponsor*. See No. 017135 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) Used for the treatment and control of bots (*Gasterophilus* spp.), large strongyles (*Strongylus* spp.), small strongyles (genera *Cyathostomum*, *Cylicobrachytus*, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*), pinworms (*Oxyuris* spp., *Strongyloides* spp.), and ascarids (*Parascaris* spp.) in horses.

(2) Administer 2 grams of thiabendazole with 1.8 grams of trichlorfon per 100 pounds of body weight sprinkled on the animals' usual daily ration of feed, or may be mixed in 5 to 10 fluid ounces of water and administered by stomach tube or drench.

(3) Do not re-treat more than once every 30 days, preferably every 6 to 8 weeks.

(4) Do not treat animals if sick or debilitated; less than 4 months of age; or mares in last month of pregnancy.

(5) Do not administer intravenous anesthetics, especially muscle relaxants, within 2 weeks of use.

(6) Not for animals intended for food use.

(7) Do not use within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(8) If the label bears directions for administration of the drug by stomach tube or drench it shall also bear the statement: Caution; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 23071, May 28, 1975, as amended at 48 FR 48229, Oct. 18, 1983]

§ 520.2380f Thiabendazole, piperazine phosphate powder.

(a) *Specifications*. Each ounce of water dispersible powder contains 6.67 grams

Food and Drug Administration, HHS

§ 520.2456

of thiabendazole and 8.33 grams of piperazine (as piperazine phosphate).

(b) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 2 grams of thiabendazole and 2.5 grams of piperazine (0.3 ounce of powder) per 100 pounds of body weight.

(2) *Indications for use*. Treatment of infections of large strongyles (genus *Strongylus*), small strongyles (genera *Cyathostomum*, *Cylicobrachytus*, and related genera *Craterostomum*, *Oesophagodontus*, *Poteriostomum*), pinworms (*Oxyuris*), threadworms (*Strongyloides*), and ascarids (*Parascaris*) in horses.

(3) *Limitations*. Use a single oral dose. Administer as a drench or by stomach tube suspended in 1 pint of warm water; by dose syringe suspended in ½ ounce of water for each 100 pounds of body weight; or sprinkled over a small amount of daily feed. Not for animals intended for food use. If the label bears directions for administration by stomach tube or drench, it shall also bear the statement “Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian;” if not labeled for use by stomach tube or drench, the label shall bear the statement, “Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.”

[46 FR 18963, Mar. 27, 1981, as amended at 46 FR 52330, Oct. 27, 1981; 62 FR 63271, Nov. 28, 1997]

§ 520.2455 Tiamulin soluble powder.

(a) *Specifications*. A water-soluble powder containing 45 percent tiamulin used to make a medicated drinking water containing 227 or 677 milligrams of tiamulin per gallon.

(b) *Sponsor*. See No. 000010 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.738 of this chapter.

(d) *Conditions of use in swine*—(1) *Amount*. 3.5 milligrams of tiamulin per pound of body weight for 5 days.

(i) *Indications for use*. For treatment of swine dysentery associated with *Treponema hyodysenteriae* susceptible to tiamulin.

(ii) *Limitations*. Use for 5 consecutive days. Withdraw 3 days before slaughter. Prepare fresh water daily. Not for

use in swine over 250 pounds body weight. Use as only source of drinking water.

(2) *Amount*. 10.5 milligrams of tiamulin per pound of body weight for 5 days.

(i) *Indications for use*. For treatment of swine pneumonia due to *Actinobacillus pleuropneumoniae* susceptible to tiamulin.

(ii) *Limitations*. Use for 5 consecutive days. Withdraw 7 days before slaughter. Prepare fresh water daily. Not for use in swine over 250 pounds body weight. Use as only source of drinking water. Do not allow consumption of feeds containing polyether ionophores (e.g., monensin, lasalocid, narasin or salinomycin) as adverse reactions may occur.

[52 FR 15718, Apr. 30, 1987, as amended at 58 FR 14313, Mar. 17, 1993; 62 FR 35076, June 30, 1997]

§ 520.2456 Tiamulin liquid concentrate.

(a) *Specifications*. A liquid concentrate containing 12.3 percent tiamulin used to make a medicated drinking water containing 227 milligrams or 681 milligrams of tiamulin per gallon.

(b) *Sponsor*. See 000010 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.738 of this chapter.

(d) *Conditions of use in swine*—(1) *Amount*. Dysentery: 3.5 milligrams of tiamulin per pound of body weight daily. Pneumonia: 10.5 milligrams of tiamulin per pound of body weight daily.

(2) *Indications for use*. For treatment of swine dysentery associated with *Treponema hyodysenteriae* and swine pneumonia due to *Actinobacillus pleuropneumoniae* susceptible to tiamulin.

(3) *Limitations*. Use for 5 consecutive days. When a dose is 3.5 milligrams per pound of body weight daily, withdraw medication 3 days before slaughter. When a dose is 10.5 milligrams per pound of body weight daily, withdraw 7 days before slaughter. Prepare fresh medicated water daily. Not for use in swine over 250 pounds body weight. Use as only source of drinking water. Do not allow consumption of feeds containing polyether ionophores (e.g.,

§ 520.2460

monensin, lasalocid, narasin, or salinomycin) as adverse reactions may occur.

[58 FR 14313, Mar. 17, 1993, as amended at 62 FR 35076, June 30, 1997]

§ 520.2460 Ticarbodine oral dosage forms.

§ 520.2460a Ticarbodine tablets.

(a) *Specifications.* Ticarbodine tablets, veterinary contain 90, 225, or 900 milligrams of ticarbodine per tablet.

(b) *Sponsor.* See No. 000986 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs for the removal of roundworms (*Toxocara canis*), hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*), and tapeworms (*Dipylidium caninum* and *Taenia pisiformis*).

(2) Dosage is administered at 45 milligrams of the drug per pound of body weight in a single dose. Dosage may be repeated in 21 days.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

§ 520.2460b Ticarbodine capsules.

(a) *Specifications.* Each capsule contains 90, 225, 450, or 900 milligrams of ticarbodine.

(b) *Sponsor.* See No. 000986 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs for removal of roundworms (*Toxocara canis*), hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*), and tapeworms (*Dipylidium caninum* and *Taenia pisiformis*).

(2) Dosage is administered orally as a single dose at 45 milligrams per lb. of body weight. Dosage may be repeated at 21-day intervals.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

§ 520.2473 Tioxidazole oral dosage forms.

§ 520.2473a Tioxidazole granules.

(a) *Specifications.* Each gram of granules contains 200 milligrams of tioxidazole.

21 CFR Ch. I (4–1–99 Edition)

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Horses*—(i) *Amount.* 5 milligrams per pound of body weight as a single dose.

(ii) *Indications for use.* Removal of mature large strongyles (*Strongylus edentatus*, *S. equinus*, and *S. vulgaris*), mature ascarids (*Parascaris equorum*), mature and immature (4th larval stage) pinworms (*Oxyuris equi*), and mature small strongyles (*Triodontophorus* spp.)

(iii) *Limitations.* For administration with feed: Sprinkle required amount of granules on a small amount of the usual grain ration and mix. Prepare for each horse individually. Withholding of feed or water not necessary. Not for use in horses intended for food. The reproductive safety of tioxidazole in breeding animals has not been determined. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. It is recommended that this drug be administered with caution to sick or debilitated horses.

(2) [Reserved]

[50 FR 52772, Dec. 26, 1985; 51 FR 2693, Jan. 21, 1986, as amended at 52 FR 7832, Mar. 13, 1987]

§ 520.2473b Tioxidazole paste.

(a) *Specifications.* Each plastic syringe contains 6.25 grams of tioxidazole.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Horses*—(i) *Amount.* 5 milligrams of tioxidazole per pound of body weight as a single dose.

(ii) *Indications for use.* Removal of mature large strongyles (*Strongylus edentatus*, *S. equinus*, and *S. vulgaris*), mature ascarids (*Parascaris equorum*), mature and immature (4th larval stage) pinworms (*Oxyuris equi*), and mature small strongyles (*Triodontophorus* spp.).

(iii) *Limitations.* Administer orally by inserting the nozzle of the syringe through the space between front and back teeth and deposit the required dose on the base of the tongue. Before dosing, make sure the horse's mouth contains no feed. Not for use in horses intended for food. The reproductive safety of tioxidazole in breeding animals has not been determined. Consult your veterinarian for assistance in the

Food and Drug Administration, HHS

§ 520.2520a

diagnosis, treatment, and control of parasitism. It is recommended that this drug be administered with caution to sick or debilitated horses.

(2) [Reserved]

[52 FR 43059, Nov. 9, 1987]

§ 520.2481 Triamcinolone acetonide tablets.

(a) *Specifications.* Each tablet contains either 0.5 milligram or 1.5 milligrams of the drug.

(b) *Sponsor.* See Nos. 000010 and 053501 in § 510.600(c) of this chapter.

(c) *NAS/NRC status.* The conditions of use specified in this section are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use.* (1) The drug is indicated for use in dogs and cats for its anti-inflammatory activity.

(2) An initial daily dosage of 0.05 milligram per pound of body weight is usually sufficient to control symptoms, although up to 0.1 milligram per pound of body weight may be given daily if response to the smaller dose is inadequate. As soon as feasible, and in any case within 2 weeks, dosage should be reduced gradually to maintenance levels of 0.0125 to 0.025 milligram per pound of body weight per day. Therapy should be discontinued by a gradual reduction in dosage after the condition has been controlled for several days. Therapy may be initiated with a single dose of sterile triamcinolone acetonide suspension veterinary in which case the tablet dosage should be administered beginning 5 to 7 days after the injection or when symptoms reappear.

(3) The labeling shall comply with the requirements of § 510.410 of this chapter.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 51 FR 26002, July 18, 1986; 53 FR 40727, Oct. 18, 1988; 62 FR 35076, June 30, 1997]

§ 520.2482 Triamcinolone acetonide oral powder.

(a) *Specifications.* Each 15 grams of triamcinolone acetonide oral powder

contains 10 milligrams of triamcinolone acetonide.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *NAS/NRC status.* The conditions of use specified in this section are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use.* (1) The drug is used as an anti-inflammatory agent for horses.

(2) It is administered at a dosage of 0.005 to 0.01 milligram triamcinolone acetonide per pound of body weight twice daily, sprinkled (top-dressed) on a small portion of feed. Treatment may be initiated with a single dose of sterile triamcinolone acetonide suspension USP followed after 3 or 4 days with the use of triamcinolone acetonide oral powder.

(3) The labeling shall comply with the requirements of § 510.410 of this chapter.

(4) Not for use in horses intended for food.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 24884, June 21, 1976, as amended at 50 FR 41489, Oct. 11, 1985; 51 FR 26002, July 18, 1986]

§ 520.2520 Trichlorfon oral dosage forms.

§ 520.2520a Trichlorfon oral.

(a) *Chemical name.* Dimethyl 2,2,2-trichloro-1-hydroxyethyl phosphonate.

(b) *Sponsor.* See No. 017800 in § 510.600(c) of this chapter.

(c) *Special considerations.* This drug is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(d) *Conditions of use.* (1) It is intended for use in horses for the removal of bots (*Gasterophilus spp.*), ascarids (*Parascaris equorum*), and pinworms (*Oxyuris equi*).

(2) Mix the drug, either dry or dissolved in water, in feed and administer

§ 520.2520b

at the rate of 4.5 grams of trichlorfon per 250 pounds of body weight. The drug is to be consumed at one feeding. Treatment should be repeated at 3-to 4-month intervals. Do not repeat treatment more frequently than every 30 days. Do not treat horses to be used for food. Do not treat sick or debilitated horses, colts under 4 months of age, mares in the last month of pregnancy, or animals other than horses. Do not administer intravenous anesthetics, especially muscle relaxants, for a period of 2 weeks after treatment.

[40 FR 13838, Mar. 27, 1975, as amended at 44 FR 63097, Nov. 2, 1979; 61 FR 34729, July 3, 1996]

§ 520.2520b Trichlorfon and atropine.

(a) *Chemical name.* (1) For trichlorfon: O,O-Dimethyl 2,2,2-trichloro-1-hydroxyethyl phosphonate.

(2) For atropine: Atropine N.F.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used for the treatment of *Syphacia obvelata* (pinworm) in laboratory mice.

(2) It is administered in distilled water as sole source of drinking water continuously for 7 to 14 days at 1.67 grams of trichlorfon and 7.7 milligrams of atropine per liter.

(3) Prepare fresh solution every 3 days. Do not use simultaneously with other drugs, insecticides, pesticides, or chemicals having cholinesterase activity, nor within 7 days before or after treatment with any other cholinesterase inhibitor.

(4) Restricted to use by or on the order of a licensed veterinarian.

§ 520.2520e Trichlorfon boluses.

(a) *Specifications.* Each bolus contains either 7.3, 10.9, 14.6, or 18.2 g of trichlorfon.

(b) *Sponsor.* See 000856 in § 510.600(c) of this chapter.

(c) *Special considerations.* Trichlorfon is a cholinesterase inhibitor. Do not use this product on animals simultaneously with, or within 2 weeks, before or after treatment with or exposure to, neuromuscular depolarizing agents (i.e., succinylcholine) or to cholinesterase-inhibiting drugs, pesticides, or chemicals.

21 CFR Ch. I (4-1-99 Edition)

(d) *NAS/NRC status.* Use of this drug has been NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter.

(e) *Conditions of use*—(1) *Amount.* 18.2 milligrams per pound of body weight, except for strongyles use 36.4 milligrams per pound of body weight.

(2) *Indications for use.* For horses for removal of bots (*Gastrophilus nasalis*, *Gastrophilus intestinalis*), large strongyles (*Strongylus vulgaris*), small strongyles, large roundworms (ascarids, *Parascaris equorum*), and pinworms (*Oxyuris equi*).

(3) *Limitations.* Do not fast horses before or after treatment. Treatment of mares in late pregnancy is not recommended. Surgery or any severe stress should be avoided for at least 2 weeks before or after treatment. Do not administer to sick, toxic, or debilitated horses. Not to be used in horses intended for use as food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 48127, July 18, 1980]

§ 520.2520f Trichlorfon granules.

(a) *Specifications.* Each package contains either 18.2 or 36.4 g of trichlorfon.

(b) *Sponsor.* See 000856 in § 510.600(c) of this chapter.

(c) *Special considerations.* Trichlorfon is a cholinesterase inhibitor. Do not use this product on animals simultaneously with, or within 2 weeks before or after treatment with or exposure to neuromuscular depolarizing agents (i.e., succinylcholine) or to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(d) *NAS/NRC status.* Use of this drug has been NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter.

(e) *Conditions of use*—(1) *Amount.* 18.2 milligrams per pound of body weight.

(2) *Indications for use.* For horses for removal of bots (*Gastrophilus nasalis*, *Gastrophilus intestinalis*), large roundworms (ascarids, *Parascaris equorum*), and pinworms (*Oxyuris equi*).

(3) *Limitations.* Do not fast horses before or after treatment. Treatment of mares in late pregnancy is not recommended. Surgery or any severe

Food and Drug Administration, HHS

§ 520.2604

stress should be avoided for at least 2 weeks before or after treatment. Do not administer to sick, toxic, or debilitated horses. Not to be used in horses intended for use as food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 48128, July 18, 1980]

§ 520.2520g Trichlorfon, phenothiazine, and piperazine dihydrochloride powder.

(a) *Specifications.* Each 54.10 grams (1.91 ounces) of water dispersible powder contains 9.10 grams of trichlorfon, 6.25 grams of phenothiazine, and the equivalent of 20.0 grams of piperazine base (as piperazine dihydrochloride).

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Special considerations.* Labeling shall bear the following statements: The drug is a cholinesterase inhibitor. Do not use this product in horses simultaneously with, or within 2 weeks before or after treatment with, or exposure to, neuromuscular depolarizing agents (e.g., succinylcholine) or to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(d) *Conditions of use—(1) Amount.* 18.2 milligrams of trichlorfon, 12.5 milligrams of phenothiazine, and 40.0 milligrams of piperazine base per pound of body weight.

(2) *Indications for use.* For horses for removal of bots (*Gastrophilus nasalis*, *Gastrophilus intestinalis*), large strongyles (*Strongylus vulgaris*), small strongyles, large roundworms (ascarids, *Parascaris equorum*), and pinworms (*Oxyuris equi*).

(3) *Limitations.* Mix powder and vial contents together in warm water to form suspension. Administer by stomach tube. Do not fast horses before or after treatment. Treatment of mares in late pregnancy is not recommended. Surgery or any severe stress should be avoided for at least 2 weeks before or after treatment. Do not administer to sick, toxic, or debilitated horses. Not to be used in horses intended for use as food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 2757, Jan. 21, 1983]

§ 520.2582 Triflupromazine hydrochloride tablets.

(a) *Specifications.* Each tablet contains either 10 milligrams or 25 milligrams of triflupromazine hydrochloride.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs and cats to relieve anxiety and to help control psychomotor overactivity as well as to increase the tolerance of animals to pain and pruritus. The drug is indicated in various office and clinical procedures which require the aid of a tranquilizer, antiemetic, or preanesthetic.¹

(2) The drug is administered orally to dogs and cats at a dosage level of 1 to 2 milligrams per pound of body weight daily; an initial dosage at the 2-milligrams level is suggested followed by daily doses at the 1-milligram level. Frequently, the drug may be withdrawn after 4 to 5 days, with drug effect continuing after withdrawal.¹

(3) Do not use in conjunction with organophosphates and/or procaine hydrochloride, because phenothiazines may potentiate the toxicity of organophosphates and the activity of procaine hydrochloride.¹

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13838, Mar. 27, 1975, as amended at 50 FR 41489, Oct. 11, 1985]

§ 520.2604 Trimeprazine tartrate and prednisolone tablets.

(a) *Specifications.* Each tablet contains: trimeprazine tartrate, 5 milligrams; and prednisolone, 2 milligrams.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is administered orally to dogs for the relief of itching regardless of cause; reduction of inflammation commonly associated with most skin disorders of dogs such as eczema, caused by internal disorders, otitis, and dermatitis, allergic, parasitic, pustular and nonspecific. It is also used in dogs as adjunctive therapy in various cough conditions including treatment of "kennel cough" or tracheobronchitis, bronchitis including allergic bronchitis, in

§ 520.2605

tonsillitis, acute upper respiratory infections and coughs of nonspecific origin. The product may also be administered to dogs suffering from acute or chronic bacterial infections, provided the infection is controlled by appropriate antibiotic or chemotherapeutic agents.¹

(2) The drug is administered orally at an initial dosage level of $\frac{1}{2}$ tablet twice daily to dogs weighing up to 10 pounds, one tablet twice daily to dogs weighing 11 to 20 pounds, two tablets twice daily to dogs weighing 21 to 40 pounds, and three tablets twice daily to dogs weighing over 40 pounds. After 4 days, the dosage is reduced to approximately $\frac{1}{2}$ the initial dosage or to an amount just sufficient to maintain remission of symptoms. Dosages in individual cases may vary and should be adjusted until proper response is obtained.¹

(3) Do not use the drug in cases of viral infections involving corneal ulceration or dendritic ulceration of the cornea.¹

(4) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.¹

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13838, Mar. 27, 1975, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.2605 Trimeprazine tartrate and prednisolone capsules.

(a) *Specifications.* Each capsule contains 3.75 milligrams of trimeprazine in sustained released form (as the tartrate) and 1 milligram of prednisolone (capsule no. 1) or 7.5 milligrams of trimeprazine in sustained release form

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

21 CFR Ch. I (4–1–99 Edition)

(as the tartrate) and 2 milligrams of prednisolone (capsule no. 2).

(b) *Sponsor.* See 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* Administer either capsule orally once daily to dogs as follows:

Animal weight (pounds)	Number of capsules per dose	
	Capsule No. 1	Capsule No. 2
Up to 10	1	
11 to 20	2	1
21 to 40	4	2
Over 40	6	3

(2) *Indications for use.* For the relief of itching regardless of cause, reduction of inflammation commonly associated with most skin disorders of dogs such as eczema caused by internal disorders, otitis, and dermatitis (allergic, parasitic, pustular, and nonspecific). It is also used in dogs as adjunctive therapy in various cough conditions including treatment of “kennel cough” or tracheobronchitis, bronchitis including allergic bronchitis, tonsillitis, acute upper respiratory infections, and coughs of nonspecific origin. The product may also be administered to dogs suffering from acute or chronic bacterial infections, provided the infection is controlled by appropriate antibiotic or chemotherapeutic agents.

(3) *Limitations.* After 4 days, reduce dosage to one-half the initial dose or to an amount sufficient to maintain remission of symptoms. Dosages in individual cases may vary and should be adjusted until proper response is obtained. Do not use the drug in cases of viral infections involving corneal ulceration or dendritic ulceration of the cornea. Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 19367, Apr. 29, 1983, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

Food and Drug Administration, HHS

§ 520.2611

§ 520.2610 Trimethoprim and sulfadiazine tablets.

(a) *Specifications.* Each tablet contains 30 milligrams (5 milligrams of trimethoprim and 25 milligrams of sulfadiazine), 120 milligrams (20 milligrams of trimethoprim and 100 milligrams of sulfadiazine), 480 milligrams (80 milligrams of trimethoprim and 400 milligrams of sulfadiazine) or 960 milligrams (160 milligrams of trimethoprim and 800 milligrams of sulfadiazine).

(b) *Sponsor.* See Nos. 000061 and 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs where systemic antibacterial action against sensitive organisms is required, either alone or as an adjunct to surgery or debridement with associated infection. The drug is indicated where control of bacterial infection is required during the treatment of acute urinary tract infections, acute bacterial complications of distemper, acute respiratory tract infections, acute alimentary tract infections, wound infections, and abscesses.

(2) The drug is given orally at 30 milligrams per kilogram of body weight per day (14 milligrams per pound per day), or as follows:

Animal body weight (pounds)	Number of tablets
30 MG TABLETS	
2.2	1
4.4	2
6.6	3
8.8	4
120 MG TABLETS	
Up to 9	1
10 to 19	2
20 to 29	3
30 to 40	4
480 MG TABLETS	
30 to 40	1
40 to 60	1½
60 to 80	2
80 to 110	3
Over 110	4

(3) The drug is given once daily. Alternatively, especially in severe infections, the initial dose may be followed by one-half the recommended daily dose every 12 hours. If no improvement is seen in 3 days, discontinue therapy and reevaluate diagnosis.

(4) Administer for 2 to 3 days after symptoms have subsided. Do not treat for more than 14 consecutive days.

(5) During long term treatment, periodic platelet counts and white and red blood cell counts are recommended.

(6) The drug should not be used in patients showing marked liver parenchymal damage or blood dyscrasia, nor in those with a history of sulfonamide sensitivity.

(7) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 3853, Jan. 27, 1976, as amended at 44 FR 32214, June 5, 1979; 46 FR 23231, Apr. 24, 1981; 47 FR 36814, Aug. 24, 1982; 50 FR 9800, Mar. 12, 1985; 50 FR 11852, Mar. 26, 1985; 61 FR 5506, Feb. 13, 1996; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 520.2611 Trimethoprim and sulfadiazine oral paste.

(a) *Specifications.* Each gram of oral paste contains 400 milligrams (67 milligrams of trimethoprim and 333 milligrams of sulfadiazine).

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter for product to be dosed at 5 grams per 150 pounds of body weight per day. See No. 000061 in § 510.600(c) of this chapter for product to be dosed at 3.75 grams per 110 pounds of body weight per day.

(c) *Conditions of use—(1) Dosage.* (i) 5 grams (335 milligrams of trimethoprim and 1,665 milligrams of sulfadiazine) per 150 pounds (68 kilograms) of body weight per day. (ii) 3.75 grams (250 milligrams of trimethoprim and 1,250 milligrams of sulfadiazine) per 110 pounds (50 kilograms) of body weight per day.

(2) *Indications for use.* For horses where systemic anti-bacterial action against sensitive organisms is required during treatment of acute strangles, respiratory infections, acute urogenital infections, and wound infections and abscesses.

(3) *Limitations.* Administer orally, once a day, as a single dose for 5 to 7 days; daily dose may also be halved and given morning and evening; for acute infection therapy continue treatment 2 to 3 days after clinical signs have subsided; if no improvement of acute infections is seen in 3 to 5 days, reevaluate diagnosis; a complete blood count

§ 520.2612

should be done periodically for prolonged use; not for use in horses intended for food; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 34469, July 29, 1983, as amended at 49 FR 26714, June 29, 1984; 53 FR 11063, Apr. 5, 1988; 61 FR 5506, Feb. 13, 1996; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 520.2612 Trimethoprim and sulfadiazine oral suspension.

(a) *Specifications.* Each milliliter of oral suspension contains 60 milligrams of drug (10 milligrams of trimethoprim and 50 milligrams of sulfadiazine).

(b) *Sponsor.* See No. 000061 in § 510.600 of this chapter.

(c) *Conditions of use. Dogs—(1) Dosage.* 1 milliliter (10 milligrams of trimethoprim and 50 milligrams of sulfadiazine) per 5 pounds of body weight.

(2) *Indications for use.* The drug is used in dogs where systemic antibacterial action against sensitive organisms is required, either alone or as an adjunct to surgery or debridement with associated infection. The drug is indicated where control of bacterial infection is required during the treatment of acute urinary tract infections, acute bacterial complications of distemper, acute respiratory tract infections, acute alimentary tract infections, wound infections, and abscesses.

(3) *Limitations.* For oral use only. Administer the recommended dose once daily or one-half the recommended daily dose every 12 hours. Administer for 2 to 3 days after symptoms have subsided. If no improvement is seen in 3 days, discontinue therapy and reevaluate diagnosis. Do not treat for more than 14 consecutive days. During long-term treatment, a complete blood count is recommended. The drug should not be used in patients showing marked liver parenchymal damage or blood dyscrasia, nor in those with a history of sulfonamide sensitivity. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 19168, May 7, 1985, as amended at 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

21 CFR Ch. I (4–1–99 Edition)

§ 520.2613 Trimethoprim and sulfadiazine powder.

(a) *Specifications.* Each gram of powder contains 67 milligrams of trimethoprim and 333 milligrams of sulfadiazine.

(b) *Sponsor.* See No. 058711 in § 510.600(c) of this chapter.

(c) *Conditions of use: Horses—(1) Dosage.* 3.75 grams of powder per 110 pounds (50 kilograms) of body weight per day.

(2) *Indications for use.* For control of bacterial infections of horses during treatment of acute strangles, respiratory tract infections, acute urogenital infections, wound infections, and abscesses.

(3) *Limitations.* Administer orally in a small amount of feed, as a single daily dose, for 5 to 7 days. Continue therapy for 2 to 3 days after clinical signs have subsided. If no improvement is seen in 3 to 5 days, reevaluate diagnosis. A complete blood count should be done periodically with prolonged use. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[58 FR 36135, July 6, 1993]

§ 520.2640 Tylosin.

(a) *Specifications.* Tylosin is the antibiotic substance produced by growth of *Streptomyces fradiae* or the same antibiotic substance produced by any other means. Tylosin, present as the tartrate salt, conforms to the appropriate antibiotic standard. Tylosin contains at least 95 percent tylosin as a combination of tylosin A, tylosin B, tylosin C, and tylosin D of which at least 80 percent is tylosin A as determined by a method entitled “Determination of Factor Content in Tylosin by High Performance Liquid Chromatography,” which is incorporated by reference. Copies are available from the Dockets Management Branch (HFA-305), Food and Drug Administration, rm. 1–23, 12420 Parklawn Dr., Rockville, MD 20857, or available for inspection at the Office of the Federal Register, 800 North Capitol Street, NW., suite 700, Washington, DC 20001.

(b) *Sponsor.* See No. 000986 in § 510.600(c) of this chapter.

(c) *Special considerations.* The quantities of antibiotic in paragraph (e) of

this section refer to the activity of the appropriate standard.

(d) *Related tolerances.* See §556.740 of this chapter.

(e) *Conditions of use.* It is used in drinking water of animals as follows:

(1) *Chickens*—(i) *Amount.* 2 grams per gallon.

(ii) *Indications for use.* Aid in the treatment of chronic respiratory disease (CRD) caused by *Mycoplasma gallisepticum* sensitive to tylosin in broiler and replacement chickens. For the control of chronic respiratory disease (CRD) caused by *Mycoplasma gallisepticum* sensitive to tylosin at time of vaccination or other stress in chickens. For the control of chronic respiratory disease (CRD) caused by *Mycoplasma synoviae* sensitive to tylosin in broiler chickens.

(iii) *Limitations.* Do not use in layers producing eggs for human consumption; administer from 1 to 5 days as sole source of drinking water; treated chickens should consume enough medicated drinking water to provide 50 milligrams of tylosin per pound of body weight per day; prepare a fresh solution every 3 days; do not administer within 24 hours of slaughter.

(2) *Turkeys*—(i) *Amount.* 2 grams per gallon.

(ii) *Indications for use.* Maintaining weight gains and feed efficiency in the presence of infectious sinusitis caused by *Mycoplasma gallisepticum* sensitive to tylosin.

(iii) *Limitations.* Do not use in layers producing eggs for human consumption; administer from 2 to 5 days as sole source of drinking water; treated turkeys should consume enough medicated drinking water to provide 60 milligrams of tylosin per pound of body weight per day; prepare a fresh solution every 3 days; when sinus swelling is present, inject the sinus with tylosin injectable simultaneously with the drinking water treatment; do not administer within 5 days of slaughter.

(3) *Swine*—(i) *Amount.* 0.25 gram per gallon.

(ii) *Indications for use.* For the control and treatment of swine dysentery (bloody scours) caused by pathogens sensitive to tylosin.

(iii) *Limitations.* As only source of drinking water for 3 to 10 days, depend-

ing on the severity of the condition being treated: mix fresh solution daily; medication must be withheld from animals 48 hours prior to slaughter.

[40 FR 13838, Mar. 27, 1975, as amended at 50 FR 49841, Dec. 5, 1985; 59 FR 14365, Mar. 28, 1994; 62 FR 39443, July 23, 1997]

PART 522—IMPLANTATION OR INJECTABLE DOSAGE FORM NEW ANIMAL DRUGS

Sec.

- 522.23 Acepromazine maleate injection.
- 522.44 Sterile sodium acetazolamide.
- 522.46 Alfaprostol.
- 522.56 Amikacin sulfate injection.
- 522.62 Aminopentamide hydrogen sulfate injection.
- 522.82 Aminopropazine fumarate sterile solution injection.
- 522.84 Beta-aminopropionitrile fumarate.
- 522.88 Sterile amoxicillin trihydrate for suspension.
- 522.90 Ampicillin implantation and injectable dosage forms.
- 522.90a Ampicillin trihydrate sterile suspension.
- 522.90b Ampicillin trihydrate for sterile suspension.
- 522.90c Ampicillin sodium for aqueous injection.
- 522.144 Arsenamide sodium aqueous injection.
- 522.147 Atipamezole hydrochloride.
- 522.150 Azaperone injection.
- 522.161 Betamethasone acetate and betamethasone disodium phosphate aqueous suspension.
- 522.163 Betamethasone dipropionate and betamethasone sodium phosphate aqueous suspension.
- 522.204 Boldenone undecylenate injection.
- 522.234 Butamisol hydrochloride.
- 522.246 Butorphanol tartrate injection.
- 522.311 Carfentanil citrate injection.
- 522.313 Ceftiofur sterile powder for injection.
- 522.314 Ceftiofur hydrochloride sterile suspension.
- 522.380 Chloral hydrate, pentobarbital, and magnesium sulfate sterile aqueous solution.
- 522.390 Chloramphenicol injection.
- 522.460 Cloprostenol sodium.
- 522.468 Colistimethate sodium powder for injection.
- 522.480 Repository corticotropin injection.
- 522.518 Cupric glycinate injection.
- 522.533 Deslorelin acetate.
- 522.535 Desoxycorticosterone pivalate.
- 522.536 Detomidine hydrochloride injection.
- 522.540 Dexamethasone injection.
- 522.542 Dexamethasone-21-isonicotinate suspension.

Pt. 522**21 CFR Ch. I (4–1–99 Edition)**

- 522.563 Diatrizoate meglumine and diatrizoate sodium injection.
- 522.575 Diazepam injection.
- 522.650 Dihydrostreptomycin sulfate injection.
- 522.690 Dinoprost tromethamine sterile solution.
- 522.723 Diprenorphine hydrochloride injection.
- 522.770 Doramectin.
- 522.775 Doxapram hydrochloride injection.
- 522.778 Doxycycline hyclate.
- 522.784 Doxylamine succinate injection.
- 522.800 Droperidol and fentanyl citrate injection.
- 522.812 Enrofloxacin solution.
- 522.820 Erythromycin injection.
- 522.840 Estradiol.
- 522.842 Estradiol benzoate and testosterone propionate in combination.
- 522.850 Estradiol valerate and norgestomet in combination.
- 522.863 Ethylisobutrazine hydrochloride injection.
- 522.883 Etorphine hydrochloride injection.
- 522.900 Euthanasia solution.
- 522.914 Fenprostalene solution.
- 522.940 Colloidal ferric oxide injection.
- 522.955 Florfenicol solution.
- 522.960 Flumethasone implantation or injectable dosage forms.
- 522.960a Flumethasone suspension.
- 522.960b Flumethasone acetate injection.
- 522.960c Flumethasone solution.
- 522.970 Flunixin meglumine solution.
- 522.995 Fluprostamol sodium injection.
- 522.1002 Follicle stimulating hormone.
- 522.1004 Fomepizole.
- 522.1010 Furosemide injection.
- 522.1020 Gelatin solution.
- 522.1044 Gentamicin sulfate injection.
- 522.1055 Gleptoferron injection.
- 522.1066 Glycopyrrolate injection.
- 522.1077 Gonadorelin injectable.
- 522.1078 Gonadorelin diacetate tetrahydrate injection.
- 522.1079 Serum gonadotropin and chorionic gonadotropin.
- 522.1081 Chorionic gonadotropin for injection; chorionic gonadotropin suspension.
- 522.1085 Guaifenesin sterile powder.
- 522.1086 Guaifenesin injection.
- 522.1125 Hemoglobin glutamer-200 (bovine).
- 522.1145 Hyaluronate sodium injection.
- 522.1150 Hydrochlorothiazide injection.
- 522.1155 Imidocarb dipropionate sterile powder.
- 522.1156 Imidocarb dipropionate solution.
- 522.1182 Iron dextran complex injection.
- 522.1183 Iron hydrogenated dextran injection.
- 522.1192 Ivermectin injection.
- 522.1193 Ivermectin and clorsulon injection.
- 522.1204 Kanamycin sulfate injection.
- 522.1222 Ketamine hydrochloride injectable dosage forms.
- 522.1222a Ketamine hydrochloride injection.
- 522.1222b Ketamine hydrochloride with promazine hydrochloride and aminopentamide hydrogen sulfate injection.
- 522.1225 Ketoprofen solution.
- 522.1228 [Reserved]
- 522.1244 Levamisole phosphate injection.
- 522.1258 Lidocaine injection with epinephrine.
- 522.1260 Lincomycin injection.
- 522.1289 Lufenuron suspension.
- 522.1290 Luprostiol sterile solution.
- 522.1335 Medetomidine hydrochloride injection.
- 522.1350 Melatonin implant.
- 522.1362 Melarsomine dihydrochloride for injection.
- 522.1372 Mepivacaine hydrochloride injection.
- 522.1380 Methocarbamol injection.
- 522.1410 Sterile methylprednisolone acetate suspension.
- 522.1452 Nalorphine hydrochloride injection.
- 522.1462 Naloxone hydrochloride injection.
- 522.1465 Naltrexone hydrochloride injection.
- 522.1468 Naproxen for injection.
- 522.1484 Neomycin sulfate sterile solution.
- 522.1503 Neostigmine methylsulfate injection.
- 522.1610 Oleate sodium solution.
- 522.1620 Orgotein for injection.
- 522.1642 Oxymorphone hydrochloride injection.
- 522.1660 Oxytetracycline injection.
- 522.1662 Oxytetracycline hydrochloride implantation or injectable dosage forms.
- 522.1662a Oxytetracycline hydrochloride injection.
- 522.1662b Oxytetracycline hydrochloride with lidocaine injection.
- 522.1680 Oxytocin injection.
- 522.1696 Penicillin G procaine implantation and injectable dosage forms.
- 522.1696a Penicillin G benzathine and penicillin G procaine sterile suspension.
- 522.1696b Penicillin G procaine aqueous suspension.
- 522.1696c Penicillin G procaine in oil.
- 522.1698 Pentazocine lactate injection.
- 522.1704 Sodium pentobarbital injection.
- 522.1720 Phenylbutazone injection.
- 522.1820 Pituitary luteinizing hormone for injection.
- 522.1850 Polysulfated glycosaminoglycan.
- 522.1862 Sterile pralidoxime chloride.
- 522.1870 Praziquantel injectable solution.
- 522.1881 Sterile prednisolone acetate aqueous suspension.
- 522.1883 Prednisolone sodium phosphate injection, sterile.
- 522.1884 Prednisolone sodium succinate injection.
- 522.1885 Prednisolone tertiary butylacetate suspension.
- 522.1890 Sterile prednisone suspension.
- 522.1920 Prochlorperazine, isopropamide for injection.

Food and Drug Administration, HHS

§ 522.44

- 522.1940 Progesterone and estradiol benzoate in combination.
- 522.1962 Promazine hydrochloride injection.
- 522.2002 Propiopromazine hydrochloride injection.
- 522.2005 Propofol injection.
- 522.2012 Prostalene solution.
- 522.2063 Pyrilamine maleate injection.
- 522.2095 Sarafloxacin solution for injection.
- 522.2100 Selenium, vitamin E injection.
- 522.2112 Sterile sometribove zinc suspension.
- 522.2120 Spectinomycin dihydrochloride injection.
- 522.2121 Spectinomycin sulfate solution.
- 522.2150 Stanazolol sterile suspension.
- 522.2200 Sulfachlorpyridazine.
- 522.2220 Sulfadimethoxine injection.
- 522.2240 Sulfaethoxypyridazine.
- 522.2260 Sulfamethazine injectable solution.
- 522.2340 Sulfomyxin.
- 522.2404 Thialbarbitone sodium for injection.
- 522.2424 Sodium thiamylal for injection.
- 522.2444 Sodium thiopental implantation or injectable dosage forms.
- 522.2444a Sodium thiopental for injection.
- 522.2444b Sodium thiopental, sodium pentobarbital for injection.
- 522.2470 Tiletamine hydrochloride and zolazepam hydrochloride for injection.
- 522.2471 Tilmicosin phosphate injection.
- 522.2474 Tolazoline hydrochloride injection.
- 522.2476 Trenbolone acetate.
- 522.2477 Trenbolone acetate and estradiol.
- 522.2478 Trenbolone acetate and estradiol benzoate.
- 522.2483 Sterile triamcinolone acetanide suspension.
- 522.2582 Triflupromazine hydrochloride injection.
- 522.2610 Trimethoprim and sulfadiazine sterile suspension.
- 522.2615 Tripelethamine hydrochloride injection.
- 522.2640 Tylosin injectable dosage forms.
- 522.2640a Tylosin injection.
- 522.2640b Tylosin tartrate for injection.
- 522.2662 Xylazine hydrochloride injection.
- 522.2670 Yohimbine injectable.
- 522.2680 Zeranol.

AUTHORITY: 21 U.S.C. 360b.

SOURCE: 40 FR 13858, Mar. 27, 1975, unless otherwise noted.

§ 522.23 Acepromazine maleate injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 10 milligrams of acepromazine maleate.

(b) *Conditions of use.* See No. 000856 in § 510.600(c) of this chapter for use in dogs, cats, and horses as follows:

(1) *Indications for use.* It is used in dogs, cats, and horses as a tranquilizer.

(2) *Amount.* Dogs: 0.25 to 0.5 milligram per pound of body weight; Cats: 0.5 to 1.0 milligram per pound of body weight; Horses: 2.0 to 4.0 milligrams per 100 pounds of body weight.

(c) *Conditions of use.* See No. 000010 in § 510.600(c) of this chapter for use in dogs as follows:

(1) *Indications for use.* It is used in dogs as an aid in tranquilization and as a preanesthetic agent.

(2) *Amount.* Dogs: 0.25 to 0.5 milligram per pound of body weight.

(3) *Limitations.* The drug is administered intravenously, intramuscularly or subcutaneously with the dosage individualized depending upon the degree of tranquilization required. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 43831, Sept. 1, 1981, as amended at 53 FR 40727, Oct. 18, 1988; 62 FR 35076, June 30, 1997]

§ 522.44 Sterile sodium acetazolamide.

(a) *Specifications.* Sterile sodium acetazolamide contains acetazolamide sodium complying with United States Pharmacopeia as a sterile powder with directions for reconstituting the product with sterile distilled water to furnish a product having a concentration of 100 milligrams acetazolamide activity per milliliter.

(b) *Sponsor.* See No. 010042 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used as an aid in the treatment of dogs with mild congestive heart failure and for rapid reduction of intraocular pressure.¹

(2) It is administered intramuscularly or intraperitoneally to dogs at a level of 5 to 15 milligrams per pound of body weight daily preferably administered in two or more divided doses.¹

(3) For use only by or on the order of a licensed veterinarian.¹

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

§ 522.46

§ 522.46 Alfaprostol.

(a) *Specifications.* Each milliliter of sterile solution contains 1 milligram of alfaprostol.

(b) *Sponsor.* No. 055882 in § 510.600(c) of this chapter.

(c) *Conditions of use.* It is used in horses as follows:

(1) *Amount.* For average mature mares, 6.0 micrograms per kilogram of body weight.

(2) *Indications for use.* To cause luteolysis in mares with active corpora lutea.

(3) *Limitations.* For intramuscular or subcutaneous use as a single injection. Not for horses intended for food. Alfaprostol is readily absorbed through the skin and can cause abortion and/or bronchial spasms. Women of child-bearing age, asthmatics, and persons with bronchial and other respiratory problems should exercise extreme caution when handling this product. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 43300, Sept. 23, 1983, as amended at 53 FR 40057, Oct. 13, 1988]

§ 522.56 Amikacin sulfate injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 50 milligrams of amikacin (as the sulfate).

(b) *Sponsor.* See Nos. 000856 and 059130 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* 5 milligrams per pound of body weight twice daily.

(2) *Indications for use.* The drug is used in dogs for treatment of genitourinary tract infections (cystitis) caused by susceptible strains of *Escherichia coli* and *Proteus* spp. and skin and soft tissue infections caused by susceptible strains of *Pseudomonas* spp. and *E. coli*.

(3) *Limitations.* The drug is administered intramuscularly or subcutaneously. Treat dogs with skin and soft tissue infections for a minimum of 7 days and those with genitourinary infections for 7 to 21 days or until culture is negative and asymptomatic. If no response is observed after 3 days of treatment, therapy should be discontinued and the case re-evaluated. Maximum duration of ther-

21 CFR Ch. I (4–1–99 Edition)

apy should not exceed 30 days. Systemic aminoglycoside therapy is contraindicated in dogs with seriously impaired renal function. Not for use in breeding dogs as reproductive studies have not been conducted. Use with extreme caution in dogs in which hearing acuity is required for functioning. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[52 FR 11816, Apr. 13, 1987; 52 FR 15412, Apr. 28, 1987, as amended at 53 FR 27851, July 25, 1988; 62 FR 23357, Apr. 30, 1997]

§ 522.62 Aminopentamide hydrogen sulfate injection.

(a) *Chemical name.* 4-(Dimethylamino)-2,2-diphenylvaleramide hydrogen sulfate.

(b) *Specifications.* It is sterile and each milliliter of aqueous solution contains 0.5 milligram of the drug.

(c) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) It is intended for use in dogs and cats only for the treatment of vomiting and/or diarrhea, nausea, acute abdominal visceral spasm, pylorospasm, or hypertrophic gastritis.

NOTE: Not for use in animals with glaucoma because of the occurrence of mydriasis.

(2) Dosage is administered by subcutaneous or intramuscular injection every 8 to 12 hours, as follows:

Weight of animal in pounds	Dosage in milligrams
Up to 10	0.1
11 to 20	0.2
21 to 50	0.3
51 to 100	0.4
Over 100	0.5

Dosage may be gradually increased up to a maximum of five times the suggested dosage. Following parenteral use dosage may be continued by oral administration of tablets.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 53 FR 27851, July 25, 1988]

§ 522.82 Aminopropazine fumarate sterile solution injection.

(a) *Specifications.* Each milliliter of aminopropazine fumarate sterile aqueous solution, veterinary, contains

Food and Drug Administration, HHS

§ 522.88

aminopropazine fumarate equivalent to 25 milligrams of aminopropazine base.

(b) *Sponsor*. See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) The drug is used for reducing excessive smooth muscle contractions, such as occur in urethral spasms associated with urolithiasis in cats and dogs and in colic spasms in horses.¹

(2) It is administered intramuscularly or intravenously to dogs and cats at a level of 1 to 2 milligrams per pound of body weight. It is administered intramuscularly or intravenously to horses at a level of 0.25 milligrams per pound of body weight. Dosage can be repeated every 12 hours, as indicated.¹

(3) Not for use in animals intended for food purposes.¹

(4) For use only by or on the order of a licensed veterinarian.¹

[40 FR 13858, Mar. 27, 1975, as amended at 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 522.84 Beta-aminopropionitrile fumarate.

(a) *Specifications*. Each vial contains 7.0 milligrams of beta-aminopropionitrile fumarate sterile lyophilized powder which is reconstituted for injection with 10 milliliters of sterile physiologic saline, USP.

(b) *Sponsor*. See No. 064146 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Horses*—(i) *Amount*. 7 milligrams (10 milliliters) intralesionally every other day for 5 treatments beginning about 30 days after initial injury.

(ii) *Indications for use*. For treatment of tendinitis of the superficial digital flexor tendon (SDFT) in the adult horse where there is sonographic evidence of fiber tearing.

(iii) *Limitations*. Single dose container for intralesional injection. Do not use in horses with dermal irritation or open skin lesions in the injection area. Do not administer

intraarticularly, into the tendon sheath, or in the presence of concurrent limb fractures. Do not use in breeding animals since the effects on fertility, pregnancy, or fetal health have not been determined. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[63 FR 44382, Aug. 19, 1998]

§ 522.88 Sterile amoxicillin trihydrate for suspension.

(a)(1) *Specifications*. Each vial contains 3 grams of amoxicillin as the trihydrate. The powder is reconstituted with sterile water for injection USP to a concentration of 100 or 250 milligrams per milliliter for use as in paragraph (d) of this section.

(2) Each vial contains 25 grams of amoxicillin as the trihydrate. The powder is reconstituted with sterile water for injection USP to a concentration of 250 milligrams per milliliter for use as in paragraph (e).

(b) *Sponsor*. See 000069 in § 510.600(c) of this chapter.

(c) *Related tolerance*. See § 556.38 of this chapter.

(d) *Conditions of use in dogs and cats*—(1) *Amount*. 5 milligrams per pound of body weight daily.

(2) *Indications for use*—(i) *Dogs*. Treatment of infections caused by susceptible strains of organisms as follows: Respiratory infections (tonsillitis, tracheobronchitis) due to *Staphylococcus aureus*, *Streptococcus* spp., *Escherichia coli*, and *Proteus mirabilis*; genitourinary infections (cystitis) due to *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*; gastrointestinal infections (bacterial gastroenteritis) due to *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*; bacterial dermatitis due to *S. aureus*, *Streptococcus* spp., and *P. mirabilis*; soft tissue infections (abscesses, lacerations, and wounds), due to *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*.

(ii) *Cats*. Treatment of infections caused by susceptible strains of organisms as follows: Upper respiratory infections due to *S. aureus*, *Staphylococcus* spp., *Streptococcus* spp., *Hemophilus* spp., *E. coli*, *Pasteurella* spp.,

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

§ 522.90

and *P. mirabilis*; genitourinary infections (cystitis) due to *S. aureus*, *Streptococcus* spp., *E. coli*, *P. mirabilis*, and *Corynebacterium* spp.; gastrointestinal infections due to *E. coli*, *Proteus* spp., *Staphylococcus* spp., and *Streptococcus* spp.; skin and soft tissue infections (abscesses, lacerations, and wounds) due to *S. aureus*, *Staphylococcus* spp., *Streptococcus* spp., *E. coli*, and *Pasteurella multocida*.

(3) *Limitations*. For use in dogs and cats only. Administer once daily for up to 5 days by intramuscular or subcutaneous injection. Continue treatment for 48 hours after the animal has become afebrile or asymptomatic. If no improvement is seen within 5 days, review the diagnosis and change therapy. As with all antibiotics, appropriate in vitro culturing susceptibility testing of samples taken before treatment should be conducted. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(e) *Condition of use*. Cattle—(1) *Amount*. 3 to 5 milligrams per pound of body weight once a day according to the animal being treated, the severity of infection, and the animal's response.

(2) *Indications for use*.— Treatment of diseases due to amoxicillin-susceptible organisms as follows: Respiratory tract infections (shipping fever, pneumonia) due to *P. multocida*, *P. hemolytica*, *Hemophilus* spp., *Staphylococcus* spp., and *Streptococcus* spp. and acute necrotic pododermatitis (foot rot) due to *Fusobacterium necrophorum*.

(3) *Limitations*. Administer once daily for up to 5 days by intramuscular or subcutaneous injection. Continue treatment for 48 to 72 hours after the animal has become afebrile or asymptomatic. Do not continue treatment beyond 5 days. Treated animals must not be slaughtered for food during treatment and for 25 days after the last treatment. As with all antibiotics, appropriate in vitro culturing and susceptibility testing of samples taken before treatment should be conducted. Milk from treated cows must not be used for human consumption during treatment or for 96 hours (8 milkings) after last treatment. Maximum volume per injection should not exceed 30 milliliters. Federal law restricts this drug to use

21 CFR Ch. I (4–1–99 Edition)

by or on the order of a licensed veterinarian.

[57 FR 37330, Aug. 18, 1992; 60 FR 55659, Nov. 2, 1995]

§ 522.90 Ampicillin implantation and injectible dosage forms.

§ 522.90a Ampicillin trihydrate sterile suspension.

(a) *Specifications*. Each milliliter contains ampicillin trihydrate equivalent to 200 milligrams of ampicillin.

(1) *Sponsor*. See No. 053501 in § 510.600(c) of this chapter.

(2) *Related tolerances*. See § 556.40 of this chapter.

(3) *Conditions of use*—(i) *Calves*.

(A) *Amount*. For enteritis: 3 milligrams per pound of body weight, intramuscularly, once or twice daily, for up to 3 days. For pneumonia: 3 milligrams per pound of body weight, intramuscularly, twice daily, for up to 3 days.

(B) *Indications for use*. Treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella* spp. susceptible to ampicillin.

(C) *Limitations*. Not for use in other animals raised for food production. Treated animals must not be slaughtered for food use during treatment or for 9 days after the last treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(ii) *Dogs*. (A) *Amount*. 3 to 6 milligrams per pound of body weight intramuscularly, once or twice daily.

(B) *Indications for use*. Treatment of respiratory tract infections due to *E. coli*, *Pseudomonas* spp., *Proteus* spp., *Staphylococcus* spp., and *Streptococcus* spp.; tonsillitis due to *E. coli*, *Pseudomonas* spp., *Streptococcus* spp., and *Staphylococcus* spp.; generalized infections (septicemia) associated with abscesses, lacerations, and wounds due to *Staphylococcus* spp. and *Streptococcus* spp.

(C) *Limitations*. Continue treatment at least 48 hours after the animal's temperature has returned to normal and other signs of infection have subsided. Usual treatment is 3 to 5 days. Federal law restricts this drug to use

Food and Drug Administration, HHS

§ 522.90b

by or on the order of a licensed veterinarian.

(iii) *Cats.* (A) *Amount.* 5 to 10 milligrams per pound of body weight intramuscularly or subcutaneously, once or twice daily.

(B) *Indications for use.* Treatment of generalized infections (septicemia) associated with abscesses, lacerations, and wounds due to *Staphylococcus* spp., *Streptococcus* spp., and *Pasteurella* spp.

(C) *Limitations.* Continue treatment at least 48 hours after the animal's temperature has returned to normal and other signs of infection have subsided. Usual treatment is 3 to 5 days. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(iv) *Swine.* (A) *Amount.* 3 milligrams per pound of body weight, intramuscularly, once or twice daily, for up to 3 days.

(B) *Indications for use.* Treatment of bacterial enteritis (colibacillosis) caused by *E. coli* and bacterial pneumonia caused by *Pasteurella* spp. susceptible to ampicillin.

(C) *Limitations.* Treated animals must not be slaughtered for food use during treatment or for 15 days after the last treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b) *Specifications.* Each milliliter contains ampicillin trihydrate equivalent to 150 milligrams of ampicillin.

(1) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(2) *Related tolerances.* See § 556.40 of this chapter.

(3) *Conditions of use. Dogs—(i) Amount.* 3 to 5 milligrams of ampicillin per pound of body weight, once a day for up to 4 days.

(ii) *Indications for use.* Treatment of bacterial infections of the upper respiratory tract (tonsillitis) due to *Streptococcus* spp., *Staphylococcus* spp., *E. coli*, *Proteus* spp., and *Pasteurella* spp., and soft tissue infections (abscesses, lacerations, and wounds) due to *Staphylococcus* spp., *Streptococcus* spp., and *E. coli*, when caused by susceptible organisms.

(iii) *Limitations.* Administer intramuscularly. If continued treatment is indicated, oral dosage is recommended. As with all antibiotics, ap-

propriate in vitro culturing and susceptibility tests of samples taken before treatment are recommended. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37330, Aug. 18, 1992, as amended at 60 FR 55659, Nov. 2, 1995]

§ 522.90b Ampicillin trihydrate for sterile suspension.

(a) *Specifications.* When reconstituted, each milliliter contains ampicillin trihydrate equivalent to 50, 100, or 250 milligrams of ampicillin.

(b) *Sponsor.* (1) See 000856 in § 510.600(c) of this chapter for use of 50, 100, and 250 milligrams per milliliter ampicillin suspension.

(2) See 010515 in § 510.600(c) of this chapter for use of 100 and 250 milligrams per milliliter ampicillin suspension.

(c) *Related tolerances.* See § 556.40 of this chapter.

(d) *Conditions of use. (1) Dogs—(i) Amount.* 3 milligrams per pound of body weight twice daily.

(ii) *Indications for use.* Treatment against strains of organisms susceptible to ampicillin and associated with respiratory tract infections, urinary tract infections, gastrointestinal infections, skin infections, soft tissue infections, and postsurgical infections.

(iii) *Limitations.* Administer by subcutaneous or intramuscular injection. Treatment should be continued for 48 to 72 hours after the animal has become afebrile or asymptomatic. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats—(i) Amount.* 3 milligrams per pound of body weight twice daily.

(ii) *Indications for use.* Treatment against strains of organisms susceptible to ampicillin and associated with respiratory tract infections, urinary tract infections, gastrointestinal infections, skin infections, soft tissue infections, and postsurgical infections.

(iii) *Limitations.* Administer by subcutaneous or intramuscular injection. Treatment should be continued for 48 to 72 hours after the animal has become afebrile or asymptomatic. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

§ 522.90c

(3) *Cattle*—(i) *Amount*. 2 to 5 milligrams per pound of body weight once daily by intramuscular injection.

(ii) *Indications for use*. Treatment of respiratory tract infections caused by organisms susceptible to ampicillin, bacterial pneumonia (shipping fever, calf pneumonia, and bovine pneumonia) caused by *Aerobacter* spp., *Klebsiella* spp., *Staphylococcus* spp., *Streptococcus* spp., *Pasteurella multocida*, and *Escherichia coli*.

(iii) *Limitations*. Do not treat cattle for more than 7 days. Milk from treated cows must not be used for food during treatment and for 48 hours (4 milkings) after the last treatment. Cattle must not be slaughtered for food during treatment and for 144 hours (6 days) after the last treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37331, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992; 58 FR 18304, Apr. 8, 1993; 63 FR 41420, Aug. 4, 1998]

§ 522.90c Ampicillin sodium for aqueous injection.

(a) *Specifications*. When reconstituted, each milliliter contains ampicillin sodium equivalent to 300 milligrams of ampicillin.

(b) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use*. *Horses*—(1) *Amount*: 3 milligrams per pound of body weight twice daily.

(2) *Indications for use*. Treatment of respiratory tract infections (pneumonia and strangles) due to *Staphylococcus* spp., *Escherichia coli*, and *Proteus mirabilis*, and skin and soft tissue infections (abscesses and wounds) due to *Staphylococcus* spp., *Streptococcus* spp., *E. coli*, and *P. mirabilis*, when caused by susceptible organisms.

(3) *Limitations*. Administer either intravenously or intramuscularly. Treatment should be continued 48 hours after all symptoms have subsided. If no response is seen in 4 to 5 days, reevaluate diagnosis. Not for use in horses or other animals which are raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37331, Aug. 18, 1992, as amended at 60 FR 55659, Nov. 2, 1995]

21 CFR Ch. I (4–1–99 Edition)

§ 522.144 Arsenamide sodium aqueous injection.

(a) *Chemical name*. [[(p-Carbamoylphenyl) arsylene]dithio diacetic acid, sodium salt.

(b) *Specifications*. The drug is a sterile aqueous solution and each milliliter contains 10.0 milligrams of arsenamide sodium.

(c) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(d) *Conditions of use*. (1) For the treatment and prevention of canine heartworm disease caused by *Dirofilaria immitis*.

(2) It is administered intravenously at 0.1 milliliter per pound of body weight (1.0 milliliter for every 10 pounds) twice a day for 2 days. For dogs in poor condition, particularly those with evidence of reduced liver function, a more conservative dosage schedule of 0.1 milliliter per pound of body weight daily for 15 days is recommended.

(3) Restricted to use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 43 FR 27785, June 27, 1978; 45 FR 56798, Aug. 26, 1980; 55 FR 26683, June 29, 1990]

§ 522.147 Atipamezole hydrochloride.

(a) *Specifications*. Each milliliter of sterile injectable solution contains 5.0 milligrams of atipamezole hydrochloride.

(b) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount*. Inject intramuscularly the same volume as that of medetomidine used.

(2) *Indications for use*. To reverse clinical effects of the sedative and analgesic agent medetomidine hydrochloride.

(3) *Limitations*. For intramuscular use only. Not recommended for use in pregnant or lactating animals, or animals intended for breeding. Atipamezole has not been evaluated in breeding animals. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[61 FR 48830, Sept. 17, 1996]

Food and Drug Administration, HHS

§ 522.163

§ 522.150 Azaperone injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 40 milligrams of azaperone.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Indications for use.* Control of aggressiveness when mixing or regrouping weanling or feeder pigs weighing up to 80 pounds.

(2) *Dosage.* 2.2 milligrams per kilogram (1 milligram per pound).

(3) *Limitations.* Inject by deep intramuscular injection. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 48229, Oct. 18, 1983, as amended at 62 FR 61625, Nov. 19, 1997]

§ 522.161 Betamethasone acetate and betamethasone disodium phosphate aqueous suspension.

(a) *Chemical names.* Betamethasone acetate: 9- α -Fluoro-16- β -methylprednisolone - 21 - acetate (C₂₄H₃₁FO₆). Betamethasone disodium phosphate: 9- α -Fluoro-16- β -methylprednisolone-21-disodium phosphate (C₂₂H₂₈FN₂O₈P).

(b) *Specifications.* The drug is a sterile aqueous suspension and each cubic centimeter contains: 12 milligrams of betamethasone acetate (equivalent to 10.8 milligrams of betamethasone), 3.9 milligrams of betamethasone disodium phosphate (equivalent to 3 milligrams of betamethasone), 2 milligrams of dibasic sodium phosphate, 5 milligrams of sodium chloride, 0.1 milligram of disodium EDTA, 0.5 milligram of polysorbate 80, 9 milligrams of benzyl alcohol, 5 milligrams of sodium carboxymethylcellulose, 1.8 milligrams of methylparaben, 0.2 milligram of propylparaben, hydrochloric acid and/or sodium hydroxide to adjust pH, and water for injection q.s.

(c) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(d) *Conditions of use.* It is used or intended for use by intra-articular injection of horses for the treatment of various inflammatory joint conditions; for example, acute and traumatic lameness involving the carpal and fetlock joints. Administer from 2.5 to 5 cubic centimeters per dose. Dose may be repeated when necessary depending upon the duration of relief obtained. Not for

use in horses intended for food. For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 52 FR 7832, Mar. 13, 1987]

§ 522.163 Betamethasone dipropionate and betamethasone sodium phosphate aqueous suspension.

(a) *Specifications.* Betamethasone dipropionate and betamethasone sodium phosphate aqueous suspension is a sterile aqueous suspension. Each milliliter of the suspension contains the equivalent of 5 milligrams of betamethasone as betamethasone dipropionate and 2 milligrams of betamethasone as betamethasone sodium phosphate.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs.* (i) It is used as an aid in the control of pruritus associated with dermatoses.

(ii) It is administered by intramuscular injection at a dosage of 0.25 to 0.5 milliliter per 20 pounds of body weight, depending on the severity of the condition. Frequency of dosage depends on recurrence of pruritic symptoms. Dosage may be repeated every 3 weeks or when symptoms recur, not to exceed a total of 4 injections.

(2) *Horses.* (i) It is used as an aid in the control of inflammation associated with various arthropathies.

(ii) It is administered aseptically by intraarticular injection at a dosage of 2.5 to 5 milliliters per joint, depending on the severity of the condition and the joint size. Dosage may be repeated upon recurrence of clinical signs. Injection into the joint cavity should be preceded by withdrawal of synovial fluid.

(iii) Not for use in horses intended for food.

(3) *Clinical and experimental data.* It has been demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

§ 522.204

(4) *Restrictions.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 41 FR 27316, July 2, 1976; 52 FR 7832, Mar. 13, 1987]

§ 522.204 Boldenone undecylenate injection.

(a) *Specifications.* Each milliliter contains 25 or 50 milligrams of boldenone undecylenate in a sesame oil base.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is intended for use as an aid in treating debilitated horses following disease or overwork and overexertion when an improvement in weight, hair coat, or general physical condition is desired. The drug is given only as adjunctive therapy to other specific and supportive therapy for diseases, surgical cases, and traumatic injuries. Optimal results can be expected only when good management and feeding practices are followed.

(2) It is administered intramuscularly at a dosage level of 0.5 milligram per pound of body weight. Treatment may be repeated at 3-week intervals.

(3) For use in horses only. Do not administer to horses intended for use as food. The effectiveness of the drug in stallions and pregnant mares has not been established, nor has the drug been shown not to be teratogenic in pregnant mares; therefore, this drug should not be used in stallions and pregnant mares.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 50 FR 41489, Oct. 11, 1985]

§ 522.234 Butamisol hydrochloride.

(a) *Specifications.* The drug contains 11 milligrams of butamisol per milliliter in a solution consisting of 70 percent propylene glycol, 4 percent benzyl alcohol and distilled water.

(b) *Sponsor.* See Nos. 000859 and 043781 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is administered by subcutaneous injection to dogs for the treatment of infections with whipworms (*Trichuris*

21 CFR Ch. I (4–1–99 Edition)

vulpis), and the hookworm (*Ancylostoma caninum*).

(2) The drug is administered subcutaneously at the rate of 0.1 milliliter per pound of body weight. In problem cases, retreatment for whipworms may be necessary in approximately 3 months. For hookworms, a second injection should be given 21 days after the initial treatment.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 15625, Apr. 14, 1978. Redesignated at 43 FR 60883, Dec. 29, 1978, and amended at 45 FR 29789, May 6, 1980; 51 FR 19329, May 29, 1986]

§ 522.246 Butorphanol tartrate injection.

(a) *Specifications.* Each milliliter of aqueous solution contains either 0.5, 2 or 10 milligrams of butorphanol (as butorphanol tartrate).

(b) *Sponsors.* Approval to firms identified in § 510.600(c) of this chapter for use as indicated:

(1) See No. 057926 for use as in paragraph (c)(2) of this section.

(2) See No. 000856 for use as in paragraphs (c)(1), (c)(2), and (c)(3) of this section.

(c) *Conditions of use*—(i) *Dogs*—(i) *Amount.* 0.025 milligram of butorphanol base activity per pound of body weight (equivalent to 0.5 milliliter per 10 pounds), using 0.5 milligram per milliliter solution.

(ii) *Indications for use.* For the relief of chronic nonproductive cough associated with tracheo-bronchitis, tracheitis, tonsillitis, laryngitis, and pharyngitis associated with inflammatory conditions of the upper respiratory tract.

(iii) *Limitations.* For subcutaneous injection in dogs only. Repeat at intervals of 6 to 12 hours as required. If necessary, increase dose to maximum of 0.05 milligram per pound of body weight. Treatment should not normally be required for longer than 7 days. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Horses*—(i) *Amount.* 0.05 milligram of butorphanol base activity per pound of body weight (0.1 milligram/kilogram) using 10 milligrams per milliliter solution.

(ii) *Indications for use.* For the relief of pain associated with colic and postpartum pain in adult horses and yearlings.

(iii) *Limitations.* For intravenous use in horses only. Dose may be repeated within 3 to 4 hours. Treatment should not exceed 48 hours. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(3) *Cats*—(i) *Amount.* 0.2 milligram of butorphanol base activity per pound of body weight (0.4 milligram/kilogram), using 2 milligrams per milliliter solution.

(ii) *Indications for use.* For the relief of pain in cats caused by major or minor trauma, or pain associated with surgical procedures.

(iii) *Limitations.* For subcutaneous injection in cats only. Dose may be repeated up to 4 times per day. Do not treat for more than 2 days. Safety for use in pregnant female cats, breeding male cats or kittens less than 4 months of age has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 29276, May 2, 1980, as amended at 50 FR 24508, June 11, 1985; 53 FR 27851, July 25, 1988; 59 FR 41665, Aug. 15, 1994; 63 FR 66432, Dec. 2, 1998]

§ 522.311 Carfentanil citrate injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 3 milligrams of carfentanil citrate.

(b) *Sponsor.* See No. 053923 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 5 to 20 micrograms per kilogram (.005 to .020 milligram per kilogram) of body weight.

(2) *Indications for use.* For immobilizing free ranging and confined members of the family Cervidae (deer, elk, and moose).

(3) *Limitations.* Inject into large muscle of neck, shoulder, back, or hindquarter. Avoid intrathoracic, intra-abdominal, or subcutaneous injection. To reverse effect, use 7 milligrams of diprenorphine for each milligram of carfentanil citrate, given intravenously or one-half intravenously and one-half intramuscularly or subcutaneously. Do not use in domestic animals intended for food. Do not use

30 days before or during hunting season. Do not use in animals that display clinical signs of severe cardiovascular or respiratory disease. Available data are inadequate to recommend use in pregnant animals. Avoid use during breeding season. Federal law restricts this drug to use by or on the order of a licensed veterinarian. The licensed veterinarian shall be a veterinarian engaged in zoo and exotic animal practice, wildlife management programs, or research.

[53 FR 40057, Oct. 13, 1988]

§ 522.313 Ceftiofur sterile powder for injection.

(a) *Specifications.* Ceftiofur sodium sterile powder for injection is reconstituted to form an aqueous solution containing the equivalent of 50 milligrams ceftiofur per milliliter.

(b) *Sponsor.* See 000009 in § 510.600 of this chapter.

(c) *Related tolerances.* See § 556.113 of this chapter.

(d) *Conditions of use*—(1) *Cattle*—(i) *Amount.* 0.5 to 1.0 milligram of ceftiofur per pound of body weight intramuscularly.

(ii) *Indications for use.* Treatment of bovine respiratory disease (shipping fever, pneumonia) associated with *Pasteurella hemolytica*, *P. multocida*, and *Haemophilus somnus* in beef and dairy cattle. Also, for the treatment of acute bovine interdigital necrobacillosis (foot rot, pododermatitis) associated with *Fusobacterium necrophorum* and *Bacteroides melaninogenicus*.

(iii) *Limitations.* Treatment should be repeated once every 24 hours for 3 days. Treat for an additional 2 days if animals do not show a satisfactory response. Do not use in animals previously found to be hypersensitive to the drug. Use of doses in excess of those indicated may result in illegal residues in tissues. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Swine*—(i) *Amount.* 3 to 5 milligrams per kilogram (1.36 to 2.27 milligrams per pound) of body weight.

(ii) *Indications for use.* For treatment and control of swine bacterial respiratory disease (swine bacterial pneumonia) associated with *Actinobacillus* (*Haemophilus*) *pleuropneumoniae*,

§ 522.314

21 CFR Ch. I (4–1–99 Edition)

Pasteurella multocida, *Salmonella choleraesuis*, and *Streptococcus suis* Type 2.

(iii) *Limitations*. For intramuscular use only. Treatment should be repeated at 24 hour intervals for a total of 3 consecutive days. Do not use in animals previously found to be hypersensitive to the drug. Use of dosages in excess of those indicated or route of administration other than that recommended may result in illegal residues in tissues. Safety of ceftiofur has not been determined in breeding swine. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(3) *Day-old chickens*—(i) *Amount*. 0.08 to 0.20 milligram per chick.

(ii) *Indications for use*. For control of early mortality associated with *Escherichia coli* organisms susceptible to ceftiofur.

(iii) *Limitations*. For subcutaneous use in the neck of day-old chicks only. As a single dose only. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(4) *Day-old turkey poults*—(i) *Amount*. 0.17 to 0.5 milligram per poult.

(ii) *Indications for use*. For control of early mortality associated with *E. coli* organisms susceptible to ceftiofur.

(iii) *Limitations*. For subcutaneous use in the neck of day-old poults only. As a single dose only. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(5) *Horses*—(i) *Amount*. 2.2 to 4.4 milligrams per kilogram (1.0 to 2.0 milligrams per pound) of body weight.

(ii) *Indications for use*. For treatment of respiratory infections in horses associated with *Streptococcus zooepidemicus*.

(iii) *Limitations*. For intramuscular use only. Treatment should be repeated every 24 hours, continued for 48 hours after clinical signs have disappeared, and should not exceed 10 days. A maximum of 10 milliliters should be administered per injection site. Not for use in horses intended for food. Do not use in animals previously found to be hypersensitive to the drug. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(6) *Dogs*—(i) *Amount*. 1.0 milligrams per pound (2.2 milligrams per kilogram) of body weight.

(ii) *Indications for use*. Treatment of canine urinary tract infections associated with *Escherichia coli* and *Proteus mirabilis*.

(iii) *Limitations*. For subcutaneous use only. Treatment should be repeated at 24-hour intervals, continued for 48 hours after clinical signs have disappeared, for 5 to 14 days. Do not use in animals found to be hypersensitive to the drug. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(7) *Sheep*—(i) *Amount*. 0.5 to 1.0 milligram per pound (1.1 to 2.2 milligrams per kilogram) of body weight.

(ii) *Indications for use*. For treatment of sheep respiratory disease (pneumonia) associated with *Pasteurella haemolytica* and/or *P. multocida*.

(iii) *Limitations*. For intramuscular use only. Treatment should be repeated at 24 hour intervals for a total of 3 consecutive days. Additional treatments may be given on days 4 and 5 for animals which do not show satisfactory response. Use of dosages in excess of those indicated or by unapproved routes of administration may result in illegal residues in tissues. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[53 5369, Feb. 24, 1988, as amended at 55 FR 13768, Apr. 12, 1990; 56 FR 12119, Mar. 22, 1991; 57 FR 41862, Sept. 14, 1992; 59 FR 41666, Aug. 15, 1994; 59 FR 54518, Nov. 1, 1994; 60 FR 51719, Oct. 3, 1995; 61 FR 35130, July 5, 1996; 61 FR 66583, Dec. 18, 1996]

§ 522.314 Ceftiofur hydrochloride sterile suspension.

(a) *Specifications*. Each milliliter contains ceftiofur hydrochloride equivalent to 50 milligrams of ceftiofur.

(b) *Sponsor*. See No. 000009 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.113 of this chapter.

(d) *Conditions of use*. (1) *Swine*—(i) *Amount*. 3 to 5 milligrams per kilogram (1.36 to 2.27 milligrams per pound) of body weight.

(ii) *Indications for use*. For treatment and control of swine bacterial respiratory disease (swine bacterial pneumonia) associated with *Actinobacillus (Haemophilus) pleuropneumoniae*, *Pasteurella multocida*, *Salmonella*

Food and Drug Administration, HHS

§ 522.460

choleraesuis, and *Streptococcus suis* Type 2.

(iii) *Limitations*. For intramuscular use only. Treatment should be repeated at 24-hour intervals for a total of 3 consecutive days. Do not use in animals previously found to be hypersensitive to the drug. Use of dosages in excess of those indicated or route of administration other than that recommended may result in illegal residues in tissues. Safety of ceftiofur has not been determined in breeding swine. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cattle*—(i) *Dosage*. 1.1 to 2.2 milligrams per kilogram (0.5 to 1.0 milligrams per pound) of body weight, at 24-hour intervals for 3 to 5 consecutive days. In addition, for bovine respiratory disease, administer 2.2 milligrams per kilogram (1.0 milligram per pound) of body weight every other day on days 1 and 3 (48-hour interval).

(ii) *Indications for use*. For treatment of bovine respiratory disease (BRD, shipping fever, pneumonia) associated with *Pasteurella haemolytica*, *P. multocida*, and *Haemophilus somnus* and acute bovine interdigital necrobacillosis (foot rot, pododermatitis) associated with *Fusobacterium necrophorum* and *Bacteroides melaninogenicus*.

(iii) *Limitations*. For intramuscular or subcutaneous use only. Do not inject more than 15 milliliters at each intramuscular injection site. Do not slaughter treated cattle for 48 hours (2 days) after last treatment. A withdrawal period has not been established in preruminating calves. Do not use in calves to be processed for veal. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[61 FR 29479, June 11, 1996, as amended at 63 FR 53578, Oct. 6, 1998]

§ 522.380 Chloral hydrate, pentobarbital, and magnesium sulfate sterile aqueous solution.

(a) [Reserved]

(b)(1) *Specifications*. Chloral hydrate, pentobarbital, and magnesium sulfate sterile aqueous solution contains 42.5 milligrams of chloral hydrate, 8.86 milligrams of pentobarbital, and 21.2 milligrams of magnesium sulfate in each milliliter of sterile aqueous solution

containing water, 33.8 percent propylene glycol, and 14.25 percent ethyl alcohol.

(2) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(3) *Conditions of use*. (i) It is used for general anesthesia and as a sedative-relaxant in cattle and horses.

(ii) For intravenous use only. The drug is administered at a dosage level of 20 to 50 milliliters per 100 pounds of body weight for general anesthesia until the desired effect is produced. Cattle usually require a lower dosage on the basis of body weight. When used as a sedative-relaxant, it is administered at a level of one-fourth to one-half of the anesthetic dosage level.

(iii) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 45 FR 16482, Mar. 14, 1980]

§ 522.390 Chloramphenicol injection.

(a) *Specifications*. Each milliliter contains 100 milligrams of chloramphenicol.

(b) *Sponsor*. See Nos. 000069 and 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use*. *Dogs*—(1) *Amount*. 5 to 15 milligrams per pound of body weight, intramuscularly or intravenously, every 6 hours. In severe infections, use 4 to 6 hour treatment intervals the first day. If no response is obtained in 3 to 5 days, discontinue use and reevaluate diagnosis.

(2) *Indications for use*. Treatment of infections of the respiratory tract, the urinary tract, and enteritis and tonsillitis caused by organisms susceptible to chloramphenicol.

(3) *Limitations*. Not for use in animals raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37331, Aug. 18, 1992]

§ 522.460 Cloprostenol sodium.

(a)(1) *Specifications*. Each milliliter of the aqueous solution contains 263 micrograms of cloprostenol sodium (equivalent to 250 micrograms of cloprostenol) in a sodium citrate, anhydrous citric acid and sodium chloride buffer containing 0.1 percent w/v chlorocresol B.P. as a bactericide.

§ 522.468

21 CFR Ch. I (4–1–99 Edition)

(2) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(3) *Conditions of use.* For intramuscular use in beef and dairy cattle to induce luteolysis.

(i) *Amount.* 2 milliliters (equivalent to 500 micrograms of cloprostenol).

(ii) *Indications.* (a) For scheduling estrus and ovulation to control the time at which cycling cows or heifers can be bred.

(1) *Single cloprostenol injection.* Treat only animals with a mature corpus luteum. Estrus should occur in 2 to 5 days, and cattle should be inseminated at the usual time relative to the detection of estrus. If estrus is not observed, treated animals may be inseminated either once at 72 hours post injection or twice at 72 and 96 hours post injection.

(2) *Double cloprostenol injection.* Give cattle a second injection 11 days after the first injection. Estrus should occur 2 to 5 days after the second injection, and cattle should be inseminated at the usual time relative to the detection of estrus. If estrus is not observed, treated animals may be inseminated either once at about 72 hours post injection or twice at 72 and 96 hours following the second injection.

(b) *Single cloprostenol injection for terminating unwanted pregnancies from mismatings from 1 week after mating until 5 months after conception, or for treating unobserved (non-detected) estrus, mummified fetus, and luteal cysts.*

(c) *Single cloprostenol injection for the treatment of pyometra.*

(iii) Do not administer to pregnant animals where the calf is not to be aborted.

(iv) Women of childbearing age, asthmatics, and persons with bronchial and other respiratory problems should exercise extreme caution when handling this product. Cloprostenol is readily absorbed through the skin and may cause abortion and/or bronchospasms. Accidental spillage on the skin should be washed off immediately with soap and water.

(v) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Specifications.* Each milliliter of sterile aqueous solution contains 131.5

micrograms of cloprostenol sodium (equivalent to 125 micrograms of cloprostenol).

(2) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(3) *Special considerations.* Labeling shall bear the statements prescribed in paragraphs (a)(3) (iii) and (iv) of this section.

(4) *Conditions of use—(i) Amount.* 3 milliliters (equivalent to 375 micrograms of cloprostenol) intramuscularly per animal as a single dose.

(ii) *Indications for use.* To induce abortion in pregnant feedlot heifers from 1 week after mating until 4½ months of gestation.

(iii) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[47 FR 4678, Feb. 2, 1982, as amended at 48 FR 15619, Apr. 12, 1983; 49 FR 5100, Feb. 10, 1984; 49 FR 29957, July 25, 1984]

§ 522.468 Colistimethate sodium powder for injection.

(a) *Specifications.* Each vial contains colistimethate sodium equivalent to 10 grams colistin activity and mannitol to be reconstituted with 62.5 milliliters sterile saline or sterile water for injection. The resulting solution contains colistimethate sodium equivalent to 133 milligrams per milliliter colistin activity.

(b) *Sponsor.* See 046573 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use.* (1) 1- to 3-day-old chickens.

(i) *Dosage.* 0.2 milligram colistin activity per chicken.

(ii) *Indications for use.* Control of early mortality associated with *Escherichia coli* organisms susceptible to colistin.

(iii) *Limitations.* For subcutaneous injection in the neck of 1- to 3-day-old chickens. Not for use in laying hens producing eggs for human consumption. Do not use in turkeys. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[63 FR 13123, Mar. 18, 1998]

§ 522.480 Repository corticotropin injection.

(a)(1) *Specifications.* The drug conforms to repository corticotropin injection U.S.P. It contains 40 or 80 U.S.C. (I.U.) units per cubic centimeter.

(2) *Sponsor.* See No. 037990 in § 510.600(c) of this chapter.

(3) *Special considerations.* The drug should be refrigerated. With prolonged use supplement daily diet with potassium chloride at one gram for small animals and from 5 to 10 grams for large animals.

(4) *Conditions of use.* (i) It is used as an intramuscular or subcutaneous injection in cattle and small animals for stimulation of the adrenal cortex where there is a general deficiency of corticotropin (ACTH). It is also a therapeutic agent for primary bovine ketosis.

(ii) It is administered to cattle initially at 200 to 600 units followed by a dose daily or every other day of 200 to 300 units and to small animals at one unit per pound of body weight to be repeated as indicated.

(iii) For use only by or on the order of a licensed veterinarian.

(b)(1) *Specifications.* The drug conforms to repository corticotropin injection U.S.P. It contains 40 or 80 U.S.P. units per milliliter.

(2) *Sponsor.* See No. 000864 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) For intramuscular injection in dogs as a diagnostic aid to test for adrenal dysfunction. For intramuscular or subcutaneous injection in dogs and cats for stimulation of the adrenal cortex where there is a general deficiency of ACTH.

(ii) For diagnostic use: Administer at one unit per pound of body weight intramuscularly. For therapeutic use: Administer at one unit per pound of body weight intramuscularly or subcutaneously, initially, to be repeated as indicated.

(iii) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(c) *National Academy of Sciences/National Research Council (NAS/NRC) status.* The therapeutic indication for use has been reviewed by NAS/NRC and found to be effective. Applications for

this use need not include effectiveness data as specified in § 514.111 of this chapter, but may require bioequivalency and safety information.

[40 FR 13858, Mar. 27, 1985, as amended at 50 FR 40966, Oct. 8, 1985; 53 FR 45760, Nov. 14, 1988]

§ 522.518 Cupric glycinate injection.

(a) *Specifications.* Each milliliter (mL) of sterile aqueous suspension contains 200 milligrams of cupric glycinate (equivalent to 60 milligrams of copper).

(b) *Sponsor.* See No. 049185 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* 200 milligrams (1 mL) for calves 300 pounds and under; 400 milligrams (2 mL) for calves over 300 pounds and adult cattle.

(2) *Indications for use.* For beef calves and beef cattle for the prevention of copper deficiency, or when labeled for veterinary prescription use, for the prevention and/or treatment of copper deficiency alone or in association with molybdenum toxicity.

(3) *Limitations.* For subcutaneous use only; repeat dose in 3 months in young calves, in 6 months in cattle; discontinue use 30 days before treated animals are slaughtered for food use; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 20159, Apr. 3, 1981, as amended at 52 FR 7832, Mar. 13, 1987; 62 FR 28630, May 27, 1997]

§ 522.533 Deslorelin acetate.

(a) *Specifications.* Each implant contains 2.1 milligrams deslorelin acetate.

(b) *Sponsor.* See 064288 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use—(1) Horses and ponies—(i) Amount.* One implant per mare.

(ii) *Indications for use.* For inducing ovulation within 48 hours in estrous mares with an ovarian follicle greater than 30 millimeters in diameter. Follicular size should be determined by rectal palpation and/or ultrasonography prior to treatment.

(iii) *Limitations.* Administer subcutaneously in the neck. Not for use in horses or ponies intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

§ 522.535

(2) [Reserved]

[63 FR 44383, Aug. 19, 1998]

§ 522.535 Desoxycorticosterone pivalate.

(a) *Specifications.* Each milliliter of sterile aqueous suspension contains 25 milligrams of desoxycorticosterone pivalate.

(b) *Sponsor.* See No. 058198 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use—(1) Dogs—(i) Amount.* Dosage requirements are variable and must be individualized on the basis of the response of the patient to therapy. Initial dose of 1 milligram per pound (0.45 kilogram) of body weight every 25 days, intramuscularly. Usual dose is 0.75 to 1.0 milligram per pound of body weight every 21 to 30 days.

(ii) *Indications for use.* For use as replacement therapy for the mineralocorticoid deficit in dogs with primary adrenocortical insufficiency.

(iii) *Limitations.* For intramuscular use only. Do not use in pregnant dogs, dogs suffering from congestive heart disease, severe renal disease, or edema. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[63 FR 13122, Mar. 18, 1998]

§ 522.536 Detomidine hydrochloride injection.

(a) *Specification.* Each milliliter of sterile aqueous solution contains 10 milligrams of detomidine hydrochloride.

(b) *Sponsor.* See 052483 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* For sedation, analgesia, or sedation and analgesia: 20 or 40 micrograms per kilogram (0.2 or 0.4 milliliter per 100 kilogram or 220 pounds) by body weight, depending on depth and duration required.

(2) *Indication for use.* As a sedative and analgesic to facilitate minor surgical and diagnostic procedures in mature horses and yearlings.

(3) *Limitations.* For sedation administer intravenously (IV) or intramuscularly (IM); for analgesia by IV; for both sedation and analgesia by

21 CFR Ch. I (4–1–99 Edition)

IV. Do not use in horses with pre-existing atrioventricular or sinoauricular block, with severe coronary insufficiency, cerebrovascular disease, respiratory disease, or chronic renal failure. Do not use in breeding animals. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[54 FR 50365, Dec. 6, 1989; 54 FR 51551, Dec. 15, 1989]

§ 522.540 Dexamethasone injection.

(a)(1) *Specifications.* The drug is a sterile aqueous solution. Each milliliter contains 2 mg of dexamethasone.

(2) *Sponsor.* See Nos. 000061 and 059130 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) The drug is indicated for the treatment of primary bovine ketosis and as an anti-inflammatory agent in dogs, cats, cattle, and horses.¹

(ii) The drug is administered intravenously or intramuscularly and dosage may be repeated if necessary, as follows:¹

(a) Canine—0.25 to 1 mg.

(b) Feline—0.125 to 0.5 mg.

(c) Equine—2.5 to 5 mg.

(d) Bovine—5 to 20 mg depending on the severity of the condition.

(iii) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(iv) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Specifications.* The drug is a sterile aqueous solution. Each milliliter contains either 2.0 milligrams of dexamethasone or 4.0 milligrams of dexamethasone sodium phosphate (equivalent to 3.0 milligrams dexamethasone).

(2) *Sponsor.* See number in § 510.600(c) of this chapter as follows:

(i) No. 000864 for use of 2.0 milligrams dexamethasone or 4.0 milligrams dexamethasone sodium phosphate injections.

(ii) No. 000402 for use of 2.0 milligrams dexamethasone or 4.0 milligrams dexamethasone sodium phosphate injections.

(3) *Conditions of use.* (i) The drug is used in dogs for the treatment of inflammatory conditions, as supportive therapy in canine posterior paresis, as supportive therapy before or after surgery to enhance recovery of poor surgical risks, and as supportive therapy in nonspecific dermatosis.¹

(ii) The drug is administered intravenously at 0.25 to 1 milligram initially. The dose may be repeated for 3 to 5 days or until a response is noted. If continued treatment is required, oral therapy may be substituted. When therapy is withdrawn after prolonged use, the daily dose should be reduced gradually over several days.¹

(iii) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(iv) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(c)(1) *Specifications.* The drug is a sterile aqueous solution. Each milliliter contains 2.0 milligrams of dexamethasone or 4.0 milligrams of dexamethasone sodium phosphate (equivalent to 3.0 milligrams of dexamethasone).

(2) *Sponsor.* See Nos. 000402 and 000864 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) The drug is used as a rapid adrenal glucocorticoid and/or anti-inflammatory agent in horses.¹

(ii) The drug is administered intravenously at a dosage of 2.5 to 5.0 milligrams. If permanent corticosteroid effect is required, oral therapy may be substituted. When therapy is withdrawn after prolonged use, the daily

dose should be reduced gradually over several days.¹

(iii) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(iv) Not for use in horses intended for food.

(v) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d)(1) *Specifications.* The drug is a sterile aqueous solution. Each milliliter contains 2.0 milligrams of dexamethasone or 4.0 milligrams of dexamethasone sodium phosphate (equivalent to 3.0 milligrams of dexamethasone).

(2) *Sponsors.* See the following numbers in § 510.600(c) of this chapter:

(i) Nos. 000069 and 050604 for intravenous or intramuscular use of 2.0 milligrams dexamethasone injection.

(ii) No. 000069 for intravenous use of 2.0 milligrams dexamethasone injection.

(3) *Conditions of use.* (i) The drug is used as an anti-inflammatory agent in dogs, cats, and horses.

(ii) It is administered intravenously or intramuscularly as follows: dogs—0.25 to 1 milligram; cats—0.125 to 0.5 milligram; horses—2.5 to 5 milligrams.

(iii) Clinical and experimental data have demonstrated that corticosteroids administered orally or by injection to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(iv) Not for use in horses intended for food.

(v) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(e)(1) *Specifications.* The drug is a sterile aqueous solution. Each milliliter contains 4.0 milligrams of dexamethasone sodium phosphate (equivalent to 3 milligrams of dexamethasone).

¹These conditions are NAS/NRC-reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

§ 522.542

(2) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) The drug is given for glucocorticoid and anti-inflammatory effect in dogs and horses.

(ii) Administer intravenously as follows: Dogs—0.25 to 1 milligram initially; may be repeated for 3 to 5 days or until response is noted. Horses—2.5 to 5 milligrams. If permanent glucocorticoid effect is required, oral therapy may be substituted. When therapy is to be withdrawn after prolonged use, the daily dose should be reduced gradually over several days.

(iii) Clinical and experimental data have demonstrated that corticosteroids administered orally or by injection may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(iv) Do not use in viral infections. Anti-inflammatory action of corticosteroids may mask signs of infections. Except when used for emergency therapy, the product is contraindicated in animals with tuberculosis, chronic nephritis, cushingoid syndrome, or peptic ulcers.

(v) Not for use in horses intended for food.

(vi) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 28265, July 9, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 522.540, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 522.542 Dexamethasone-21-isonicotinate suspension.

(a) *Specifications.* Each milliliter of sterile suspension contains 1 milligram of dexamethasone-21-isonicotinate.

(b) *Sponsor.* No. 000010 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in the treatment of various inflammatory conditions associated with the musculoskeletal system in dogs, cats, and horses.

(2) It is recommended for intramuscular administration as follows: Dogs—0.25 to 1 milligram; cats—

21 CFR Ch. I (4–1–99 Edition)

0.125 to 0.5 milligram; horses—5 to 20 milligrams. Dosage may be repeated.

(3) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition following by dystocia, fetal death, retained placenta, and metritis.

(4) Not for use in horses intended for food.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[42 FR 37543, July 22, 1977, as amended at 47 FR 14703, Apr. 6, 1982]

§ 522.563 Diatrizoate meglumine and diatrizoate sodium injection.

(a) *Specifications.* Diatrizoate meglumine and diatrizoate sodium injection contains 34.3 percent diatrizoate meglumine and 35 percent diatrizoate sodium, or 66 percent diatrizoate meglumine and 10 percent diatrizoate sodium, in sterile aqueous solution.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is indicated for use in dogs and cats for visualization in excretion urography, including renal angiography, uretography, cystography, and urethrography; aortography; angiocardiology, peripheral arteriography, and venography; selective coronary arteriography; cerebral angiography; lymphography; arthrography; discography; and sialography. It is also useful as an aid in delineating peritoneal hernias and fistulous tracts.

(2) For excretion urography administer 0.5 to 1.0 milliliter per pound of body weight to a maximum of 30 milliliters intravenously. For cystography remove urine, administer 5 to 25 milliliters directly into the bladder via catheter. For urethrography administer 1.0 to 5 milliliters via catheter into the urethra to provide desired contrasts delineation. For angiocardiology (including aortography) rapidly inject 5 to 10 milliliters directly into the heart via catheter or intraventricular puncture. For cerebral angiography rapid injection of 3 to 10

milliliters via carotid artery. For peripheral arteriography and/or venography and selective coronary arteriography rapidly inject 3 to 10 milliliters intravascularly into the vascular bed to be delineated. For lymphography slowly inject 1.0 to 10 milliliters directly into the lymph vessel to be delineated. For arthrography slowly inject 1.0 to 5 milliliters directly into the joint to be delineated. For discography slowly inject 0.5 to 1.0 milliliter directly into the disc to be delineated. For sialography slowly inject 0.5 to 1.0 milliliter into the duct to be delineated. For delineation of fistulous tracts slowly inject quantity necessary to fill the tract. For delineation of peritoneal hernias inject 0.5 to 1.0 milliliter per pound of body weight directly into the peritoneal cavity.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 12993, Mar. 9, 1979, as amended at 50 FR 41489, Oct. 11, 1985]

§ 522.575 Diazepam injection.

(a) *Specification.* Each milliliter of sterile solution contains 5 milligrams of diazepam.

(b) *Sponsor.* See 000004 in § 510.600(c) of this chapter.

(c) *Conditions of use. Dogs—(1) Indications for use.* As a preanesthetic agent to reduce the amount of barbiturate required for short duration anesthesia.

(2) *Dosage.* Intravenously, 0.2 milligram per kilogram of body weight 3–5 minutes before anesthesia is to be induced using a short acting barbiturate.

(3) *Limitations.* Not for use in dogs with known sensitivity to benzodiazepines. Safety in animals intended for breeding and pregnant animals has not been established. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[58 FR 500, Jan. 6, 1993]

§ 522.650 Dihydrostreptomycin sulfate injection.

(a) *Specifications.* Each milliliter contains dihydrostreptomycin sulfate equivalent to 500 milligrams of dihydrostreptomycin.

(b) *Sponsor.* See Nos. 000069 and 055529 in § 510.600(c) of this chapter.

(c) *National Academy of Sciences/National Research Council (NAS/NRC) status.* The conditions of use were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter but may require bioequivalency and safety information.

(d) *Conditions of use—(1) Amount.* 5 milligrams per pound of body weight every 12 hours.

(2) *Indications for use.* Treatment of leptospirosis in dogs and horses due to *Leptospira canicola*, *L. icterohemorrhagiae*, and *L. pomona*; in cattle due to *L. pomona*; and in swine due to *L. pomona*; and *L. grippotyphosa*.

(3) *Limitations.* Administer by deep intramuscular injection only. Treatment should be continued for 3 to 5 days or until the urine is free of leptospira for at least 72 hours as measured by darkfield microscopic examination. Treatment with subtherapeutic dosages, excessive duration of therapy, or inappropriate use of this antibiotic may lead to the emergence of streptomycin or dihydrostreptomycin resistant organisms. Discontinue use 30 days before slaughter for food. Not for use in animals producing milk because use of the drug will contaminate the milk. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37331, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992]

§ 522.690 Dinoprost tromethamine sterile solution.

(a) *Specifications.* Each milliliter of sterile solution contains the equivalent of 5 milligrams of dinoprost.

(b) *Sponsor.* See Nos. 000009 and 059130 in § 510.600(c) of this chapter.

(c) *Special considerations.* Women of child-bearing age, asthmatics, and persons with bronchial and other respiratory problems should exercise extreme caution when handling this product. Dinoprost tromethamine is readily absorbed through the skin and can cause abortion and bronchospasms. Accidental spillage on the skin should be washed off immediately with soap and water. Use of this product in excess of the approved dose may result in drug residues. Do not administer to pregnant cattle unless abortion is desired.

§522.723

Do not administer intravenously; this may potentiate adverse reactions.

(d) *Conditions of use*—(1) *Mares*—(i) *Amount*. Equivalent of 1 milligram of dinoprost per 100 pounds of body weight.

(ii) *Indications*. For its luteolytic effect to control timing of estrus in estrus cycling mares and in clinically anestrus mares that have a corpus luteum.

(iii) *Limitations*. For use once as a single intramuscular injection. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cattle*—(i) *Amount*. 5 milliliters (equivalent to 25 milligrams of dinoprost).

(ii)(a) *Indications*. For its luteolytic effect to control timing of estrus and ovulation in estrus cycling cattle that have a corpus luteum.

(b) *Limitations*. For use in beef cattle and nonlactating dairy heifers, as follows: Inject a dose of 5 milliliters intramuscularly either once or twice at a 10- to 12-day interval. With a single injection, cattle should be bred at the usual time relative to estrus. With the two injections, cattle can be bred after the second injection either at the usual time relative to detected estrus or at about 80 hours after the second injection. Estrus is expected to occur 1 to 5 days after injection if a corpus luteum was present. Cattle that do not become pregnant to breeding at estrus on days 1 to 5 after injection will be expected to return to estrus in about 18 to 24 days. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(iii)(a) *Indications*. For treatment of pyometra (chronic endometritis).

(b) *Limitations*. For intramuscular use as a single injection. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(3) *Nonlactating cattle*—(i) *Amount*. Five milliliters intramuscularly as a single injection.

(ii) *Indications*. For its abortifacient effect in nonlactating cattle.

(iii) *Limitations*. For intramuscular use only, during first 100 days of gestation. Cattle that abort will abort within 35 days after injection. Federal law

21 CFR Ch. I (4–1–99 Edition)

restricts this drug to use by or on the order of a licensed veterinarian.

(4) *Lactating dairy cattle*—(i) *Amount*. Five milliliters intramuscularly as a single injection.

(ii) *Indications*. For treatment of unobserved (silent) estrus in lactating dairy cattle that have a corpus luteum.

(iii) *Limitations*. Breed cattle as they are detected in estrus. If estrus has not been observed by 80 hours after injection, breed at 80 hours. If cattle return to estrus breed at the usual time relative to estrus. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(5) *Swine*—(i) *Amount*. 2 milliliters (equivalent to 10 milligrams of dinoprost).

(ii) *Indications*. For parturition induction in swine when injected within 3 days of normal predicted farrowing.

(iii) *Limitations*. For use in swine as follows: Inject a dose of 2 milliliters intramuscularly within 3 days of predicted farrowing. The response to treatment varies by individual animals with a mean interval from administration to parturition of approximately 30 hours. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 4818, Feb. 2, 1976, as amended at 46 FR 13214, Feb. 20, 1981; 46 FR 39127, July 31, 1981; 48 FR 6331, Feb. 11, 1983; 48 FR 46023, Oct. 11, 1983; 49 FR 4373, Feb. 6, 1984; 64 FR 15685, Apr. 1, 1999]

§522.723 Diprenorphine hydrochloride injection.

(a) *Chemical name*. N-(Cyclopropylmethyl)-6,7,8,14-tetrahydro-7- α -(1-hydroxy - 1 - methylethyl) - 6,14 - endoethanonoripavine hydrochloride.

(b) *Specifications*. Each milliliter of diprenorphine hydrochloride injection, veterinary, contains 2 mg of diprenorphine hydrochloride in sterile aqueous solution.

(c) *Sponsors*. See No. 053923 in §510.600(c) of this chapter.

(d) *Conditions of use*. (1) The drug is used for reversing the effects of etorphine hydrochloride injection, veterinary, the use of which is provided for in §522.883, in wild and exotic animals.

(2) It is administered intramuscularly or intravenously at a

Food and Drug Administration, HHS

§ 522.778

suitable dosage level depending upon the species.

(3) For use in wild or exotic animals only. Do not use in domestic food-producing animals. Do not use 30 days before, or during, the hunting season in free-ranging wild animals that might be used for food.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian. Distribution is restricted to veterinarians engaged in zoo and exotic animal practice, wildlife management programs and researchers.

[40 FR 13858, Mar. 27, 1975, as amended at 48 FR 16241, Apr. 15, 1983; 60 FR 39847, Aug. 4, 1995; 64 FR 15684, Apr. 1, 1999]

§ 522.770 Doramectin.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 10 milligrams of doramectin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.225 of this chapter.

(d) *Conditions of use*—(1) *Cattle*—(i) *Amount.* 200 micrograms per kilogram (10 milligrams per 110 pounds).

(ii) *Indications for use.* For treatment and control of gastrointestinal roundworms, lungworms, eyeworms, grubs, sucking lice, and mange mites. To control infections and to protect from reinfection with *Cooperia oncophora* and *Haemonchus placei* for 14 days, *Ostertagia ostertagi* for 21 days, and *C. punctata*, *Oesophagostomum radiatum*, and *Dictyocaulus viviparus* for 28 days after treatment.

(iii) *Limitations.* Administer as a single subcutaneous or intramuscular injection. Do not slaughter cattle within 35 days of treatment. Not for use in female dairy cattle 20 months of age or older. Do not use in calves to be processed for veal.

(2) *Swine*—(i) *Amount.* 300 micrograms per kilogram (10 milligrams per 75 pounds).

(ii) *Indications for use.* For treatment and control of gastrointestinal roundworms, lungworms, kidney worms, sucking lice, and mange mites.

(iii) *Limitations.* Administer as a single intramuscular injection. Do not slaughter swine within 24 days of treatment. Consult your veterinarian for as-

sistance in the diagnosis, treatment, and control of parasitism.

[61 FR 53321, Oct. 11, 1996, as amended at 62 FR 44410, Aug. 21, 1997; 62 FR 62242, Nov. 21, 1997; 63 FR 68183, Dec. 10, 1998; 64 FR 13509, Mar. 19, 1999]

§ 522.775 Doxapram hydrochloride injection.

(a) *Specifications.* The drug is a sterile aqueous solution containing 20 milligrams doxapram hydrochloride per milliliter.

(b) *Sponsor.* See No. 000031 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) Administer to dogs, cats, and horses to stimulate respiration during and after general anesthesia; to speed awakening and return of reflexes after anesthesia. Administer to neonate dogs and cats to initiate respiration following dystocia or caesarean section; to stimulate respiration following dystocia or caesarean section.

(2) For intravenous use in dogs and cats at a dose of 2½ to 5 milligrams of doxapram hydrochloride per pound of body weight in barbiturate anesthesia, 0.5 mg per lb. in gas anesthesia; for intravenous use in horses at 0.25 mg per lb. of body weight in barbiturate anesthesia, 0.2 mg per lb. in inhalation anesthesia, 0.25 mg per lb. with chloral hydrate with or without magnesium sulfate; for subcutaneous, sublingual, or umbilical vein administration in neonate puppies at a dose rate of 1 to 5 mg; for subcutaneous or sublingual use in neonate kittens at 1 to 2 mg. Dosage may be repeated in 15 to 20 minutes if necessary.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 17838, Apr. 23, 1975]

§ 522.778 Doxycycline hyclate.

(a) *Specifications.* Doxycycline hyclate solution contains 8.5 percent doxycycline activity. A syringe of *N*-methyl-2-pyrrolidone and poly (DL-lactide) mixed with a syringe of doxycycline produces 0.5 milliliter of solution.

(b) *Sponsor.* See 063604 in § 510.600(c) of this chapter.

(c) [Reserved]

§ 522.784

21 CFR Ch. I (4–1–99 Edition)

(d) *Conditions of use*—(1) *Dogs*—(i) *Amount*. Apply subgingivally to periodontal pocket(s) of affected teeth.

(ii) *Indications for use*. For treatment and control of periodontal disease.

(iii) *Limitations*. Do not use in dogs less than 1-year old. Use of tetracyclines during tooth development has been associated with permanent discoloration of teeth. Do not use in pregnant bitches. Use in breeding dogs has not been evaluated. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[63 FR 8349, Feb. 19, 1998]

§ 522.784 Doxylamine succinate injection.

(a) *Specifications*. Each milliliter of the drug contains 11.36 mg of doxylamine succinate.

(b) *Sponsor*. See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) The drug is used in conditions in which antihistaminic therapy may be expected to alleviate some signs of disease in horses, dogs, and cats.¹

(2) It is administered to horses at a dosage level of 25 mg per hundred pounds of body weight. It is administered to dogs and cats at a dosage level of 0.5 to 1 mg per pound of body weight. Doses may be repeated at 8 to 12 hours, if necessary, to produce desired effect. Intravenous route is not recommended for dogs and cats and should be injected slowly in horses. Intramuscular and subcutaneous administration should be by divided injection sites.¹

(3) Not for use in horses intended for food.¹

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13858, Mar. 27, 1975, as amended at 42 FR 60140, Nov. 25, 1977; 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 522.800 Droperidol and fentanyl citrate injection.

(a) *Specifications*. Droperidol and fentanyl citrate injection is a sterile

solution containing 20 milligrams of droperidol and 0.4 milligram of fentanyl citrate per cubic centimeter.

(b) *Sponsor*. See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) It is used in dogs as an analgesic and tranquilizer and for general anesthesia.

(2) It is administered as follows:

(i) For analgesia and tranquilization administer according to response desired, as follows:

(a) Intramuscularly at the rate of 1 cubic centimeter per 15 to 20 pounds of body weight in conjunction with atropine sulfate administered at the rate of 0.02 milligram per pound of body weight, or

(b) Intravenously at the rate of 1 cubic centimeter per 25 to 60 pounds of body weight in conjunction with atropine sulfate administered at the rate of 0.02 milligram per pound of body weight.

(ii) For general anesthesia administer according to response desired, as follows:

(a) Intramuscularly at the rate of 1 cubic centimeter per 40 pounds of body weight in conjunction with atropine sulfate administered at the rate of 0.02 milligram per pound of body weight and followed in 10 minutes by an intravenous administration of sodium pentobarbital at the rate of 3 milligrams per pound of body weight, or

(b) Intravenously at the rate of 1 cubic centimeter per 25 to 60 pounds of body weight in conjunction with atropine sulfate administered at the rate of 0.02 milligram per pound of body weight and followed within 15 seconds by an intravenous administration of sodium pentobarbital at the rate of 3 milligrams per pound of body weight.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 64 FR 15684, Apr. 1, 1999]

§ 522.812 Enrofloxacin solution.

(a) *Specifications*. Each milliliter of sterile solution contains either 22.7 milligrams of enrofloxacin when intended for use in dogs or 100 milligrams of enrofloxacin when intended for use in cattle.

(b) *Sponsor*. See No. 000859 in § 510.600(c) of this chapter.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter.

(c) *Related tolerance.* See § 556.228 of this chapter.

(d) *Conditions of use*—(1) *Dogs*—(i) *Amount.* 2.5 milligrams per kilogram (1.13 milligrams per pound) of body weight as an initial dose only.

(ii) *Indications for use.* Dogs for management of diseases associated with bacteria susceptible to enrofloxacin.

(iii) *Limitations.* As a single, intramuscular, initial dose followed by use of tablets twice daily for 2 to 3 days beyond cessation of clinical signs to a maximum of 10 days. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cattle*—(i) *Amount.* Single-dose therapy: 7.5 to 12.5 milligrams enrofloxacin per kilogram of body weight (3.4 to 5.7 milliliters per 100 pounds). Multiple-day therapy: 2.5 to 5.0 milligrams per kilogram of body weight (1.1 to 2.3 milliliters per 100 pounds) administered once daily for 3 to 5 days.

(ii) *Indications for use.* For the treatment of bovine respiratory disease (BRD) associated with *Pasteurella haemolytica*, *P. multocida*, and *Haemophilus somnus*.

(iii) *Limitations.* For subcutaneous use in cattle only. Do not inject more than 20 milliliters at each site. Do not slaughter within 28 days of last treatment. Do not use in cattle intended for dairy production. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. The effect of enrofloxacin on bovine reproductive performance, pregnancy, and lactation have not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian. Federal law prohibits the extra-label use of this drug in food-producing animals.

[55 FR 26683, June 29, 1990, as amended at 62 FR 38907, July 21, 1997; 63 FR 49003, Sept. 14, 1998]

§ 522.820 Erythromycin injection.

(a) *Sponsor.* See 050604 in § 510.600(c) of this chapter.

(b) *NAS/NRC status.* The conditions of use have been reviewed by NAS/NRC and found effective.

(c) *Dogs and cats*—(1) *Specifications.* Each milliliter of polyethylene glycol

vehicle contains 100 milligrams of erythromycin base with 2 percent butyl aminobenzoate.

(2) *Conditions of use*—(i) *Amount.* 3 to 5 milligrams per pound of body weight, intramuscularly, two to three times daily, for up to 5 days.

(ii) *Indications for use*—(A) *Dogs.* For the treatment of bacterial pneumonia, upper respiratory infections (tonsillitis, bronchitis, tracheitis, pharyngitis, pleurisy), endometritis and metritis, and bacterial wound infections caused by *Staphylococcus* spp., *Streptococcus* spp., and *Corynebacterium* spp., sensitive to erythromycin.

(B) *Cats.* For the treatment of bacterial pneumonia, upper respiratory infections (rhinitis, bronchitis), secondary infections associated with panleukopenia, and bacterial wound infections caused by *Staphylococcus* spp. and *Streptococcus* spp., susceptible to erythromycin.

(iii) *Limitations.* Administer by deep intramuscular injection into the heavy muscles of the neck and limbs. Do not administer intravenously or intraperitoneally. Avoid subcutaneous use. Do not administer from moist or wet syringe. As with all antibiotics, appropriate in vitro culturing and susceptibility testing of samples taken before treatment should be conducted. Do not administer in conjunction with penicillin. As with all antibiotics, excessive continuous use may result in an overgrowth of nonsusceptible organisms. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d) *Cattle*—(1) *Specifications.* Each milliliter of nonaqueous, buffered, alcohol base sterile solution contains 200 milligrams of erythromycin base.

(2) *Related tolerances.* See § 556.230 of this chapter.

(3) *Conditions of use*—(i) *Amount.* 4 milligrams of erythromycin base per pound of body weight once daily for up to 5 days.

(ii) *Indications for use.* For the treatment of bovine respiratory disease (shipping fever complex and bacterial pneumonia) associated with *Pasteurella multocida* susceptible to erythromycin.

(iii) *Limitations.* For intramuscular use only. Do not use in female dairy cattle over 20 months of age. Do not

§ 522.840

slaughter treated animals within 6 days of last treatment. To avoid excess trim, do not slaughter within 21 days of last injection.

[58 FR 43795, Aug. 18, 1993]

§ 522.840 Estradiol.

(a) *Specifications.* Each silicone rubber implant contains 25.7 or 43.9 milligrams of estradiol.

(b) *Sponsor.* See 000986 in § 510.600(c) of this chapter.

(c) *Conditions of use.* It is used for implantation in steers and heifers as follows:

(1) *Amount.* Insert one 25.7-milligram implant every 200 days; insert one 43.9-milligram implant every 400 days.

(2) *Indications for use.* For increased rate of weight gain in suckling and pastured growing steers; for improved feed efficiency and increased rate of weight gain in confined steers and heifers.

(3) *Limitations.* For subcutaneous ear implantation in steers and heifers only. A second implant may be used if desired. No additional effectiveness may be expected from reimplanting in less than 200 days for the 25.7-milligram implant or 400 days for the 43.9 milligram implant. Increased sexual activity (bulling, riding, and excitability) has been reported in implanted animals.

[51 FR 22276, June 19, 1986, as amended at 57 FR 41861, Sept. 14, 1992]

§ 522.842 Estradiol benzoate and testosterone propionate in combination.

(a) *Chemical names.* (1) Estradiol benzoate: 1,3,5(10)-Estratriene-3,17 beta-diol 3-benzoate.

(2) Testosterone propionate: 17beta-Hydroxyandrost-4-en-3-one propionate.

(b) *Sponsor.* See Nos. 000856 and 021641 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See §§ 556.240 and 556.710 of this chapter.

(d) *Conditions of use.* It is used for implantation in heifers as follows:

(1) *Amount.* 20 milligrams of estradiol benzoate and 200 milligrams of testosterone propionate per dose.

(2) *Indications for use.* Growth promotion and improved feed efficiency.

(3) *Limitations.* For heifers weighing 400 pounds or more; for subcutaneous ear implantation, one dose per animal;

21 CFR Ch. I (4–1–99 Edition)

not for use in dairy or beef replacement heifers.

(e) *NAS/NRC status.* These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety data.

[40 FR 13858, Mar. 27, 1975, as amended at 49 FR 29778, July 24, 1984; 61 FR 5506, Feb. 13, 1996]

§ 522.850 Estradiol valerate and norgestomet in combination.

(a) *Specifications.* The product is a two-component drug consisting of the following:

(1) An implant containing 6.0 milligrams of norgestomet.

(2) An injectable solution (sesame oil) containing 3.0 milligrams of norgestomet and 5.0 milligrams of estradiol valerate per 2 milliliters.

(b) *Sponsor.* See 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use—*(1) *Amount.* One implant and 2 milliliters of injection at time of implantation.

(2) *Indications for use.* For synchronization of estrus/ovulation in cycling beef cattle and non-lactating dairy heifers.

(3) *Limitations.* Insert implant subcutaneously in the ear only; then immediately inject solution intramuscularly only. Counting the day of implantation as day 1, remove the implant on day 10. Collect all implants as they are removed and burn them. While animals are restrained for artificial insemination, avoid other treatments such as vaccinations, dipping, pour-on grub and louse prevention, spraying, etc. When inseminating without estrus detection, the entire treated group should be started at 48 hours after the last implant has been removed and should be completed within 6 hours. Where estrus detection is preferred, insemination should be approximately 12 hours after first detection of estrus. Those that do not conceive can be re-bred when they return to estrus approximately 17 to 25 days after implant removal. Do not use in

Food and Drug Administration, HHS

§ 522.900

cows producing milk for human consumption.

[47 FR 55477, Dec. 10, 1982, as amended at 48 FR 49656, Oct. 27, 1983; 51 FR 33592, Sept. 22, 1986; 54 FR 1165, Jan. 12, 1989]

§ 522.863 Ethylisobutrazine hydrochloride injection.

(a) *Specifications.* The drug is a sterile aqueous solution. Each milliliter contains 50 milligrams of ethylisobutrazine hydrochloride.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in dogs as a tranquilizer.¹

(2) It is administered intramuscularly at a dosage level of 2 to 5 milligrams of ethylisobutrazine hydrochloride per pound of body weight for profound tranquilization. It is administered intravenously at a dosage level of 1 to 2 milligrams of ethylisobutrazine hydrochloride per pound of body weight to effect.¹

(3) It is not to be used in conjunction with organophosphates and/or procaine hydrochloride because phenothiazines may potentiate the toxicity of organophosphates and the activity of procaine hydrochloride.¹

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13858, Mar. 27, 1975, as amended at 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 522.883 Etorphine hydrochloride injection.

(a) *Chemical name.* 6,7,8,14 - tetrahydro - alpha - methyl - alpha - propyl - 6,14 - endo-ethenooripavine-alpha-methanol hydrochloride.

(b) *Specifications.* Each milliliter of etorphine hydrochloride injection, veterinary, contains 1 mg of etorphine hydrochloride in sterile aqueous solution.

(c) *Sponsors.* See No. 053923 in § 510.600(c) of this chapter.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use.* (1) The drug is used for the immobilization of wild and exotic animals.

(2) It is administered intramuscularly by hand syringe or syringe dart at a suitable dosage level depending upon the species.

(3) Do not use the drug unless diprenorphine hydrochloride injection, veterinary, as provided for in § 522.723, is available for use in reversing the effects of etorphine hydrochloride injection, veterinary.

(4) For use in wild or exotic animals only. Do not use in domestic food-producing animals. Do not use 30 days before, or during, the hunting season in free-ranging wild animals that might be used for food.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian. Distribution is restricted to veterinarians engaged in zoo and exotic animal practice, wildlife management programs, and researchers.

[40 FR 13858, Mar. 27, 1975, as amended at 48 FR 16241, Apr. 15, 1983; 61 FR 260, Jan. 4, 1996]

§ 522.900 Euthanasia solution.

(a) [Reserved]

(b)(1) *Specifications.* Each milliliter of nonsterile solution contains 390 milligrams of pentobarbital sodium and 50 milligrams of phenytoin sodium.

(2) *Sponsor.* Nos. 000061 and 059079 in § 510.600(c) of this chapter.

(3) *Conditions of use—(i) Indications for use.* For the humane, painless, and rapid euthanasia of dogs.

(ii) *Amount.* One milliliter (390 milligrams of pentobarbital sodium and 50 milligrams of phenytoin sodium) for each 10 pounds of body weight.

(iii) *Limitations.* For intravenous injection or intracardiac injection when intravenous use is impractical. Do not use for therapeutic purposes. Do not use in animals intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(c)(1) *Specifications.* Each milliliter of nonsterile, aqueous solution contains 400 milligrams of secobarbital sodium and 25 milligrams of dibucaine hydrochloride.

(2) *Sponsor.* See 000033 in § 510.600(c) of this chapter.

§ 522.914

(3) *Conditions of use*—(i) *Indications for use.* To induce rapid, humane, painless euthanasia of dogs.

(ii) *Amount.* For dogs, 1 milliliter for each 10 pounds of body weight.

(iii) *Limitations.* For intravenous injection. Do not use for therapeutic purposes. Do not use in animals intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 23232, Apr. 24, 1981, as amended at 47 FR 15327, Apr. 9, 1982; 48 FR 16241, Apr. 15, 1983; 52 FR 7832, Mar. 13, 1987; 56 FR 9623, Mar. 7, 1991; 59 FR 14367, Mar. 28, 1994]

§ 522.914 Fenprostalene solution.

(a) *Specifications*—(1) *Cattle.* Each milliliter of sterile solution contains 0.5 milligram of fenprostalene.

(2) *Swine.* Each milliliter of sterile solution contains 0.25 milligram of fenprostalene.

(b) *Sponsor.* See 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.277 of this chapter.

(d) *Special considerations.* Labeling shall bear the following statements: Women of childbearing age, asthmatics, and persons with bronchial and other respiratory problems should exercise extreme caution when handling this product. It is readily absorbed through the skin and may cause abortion and/or bronchospasms. Accidental spillage on the skin should be washed off immediately with soap and water.

(e) *Conditions of use*—(1) *Cattle*—(i) *Amount.* 1 milligram (2 milliliters) subcutaneously per animal.

(ii) *Indications for use.* For feedlot heifers to induce abortion when pregnant 150 days or less. For beef or nonlactating dairy cattle for estrus synchronization.

(iii) *Limitations.* Subcutaneous use in cattle only. Feedlot heifers to induce abortion, single dose. Beef or nonlactating dairy cattle for estrus synchronization, a single dose or two doses 11 to 13 days apart. Do not use in pregnant animals unless abortion is desired. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

21 CFR Ch. I (4–1–99 Edition)

(2) *Swine*—(i) *Amount.* 0.25 milligram (1 milliliter) subcutaneously once per animal.

(ii) *Indications for use.* For sows and gilts pregnant at least 112 days for the induction of parturition.

(iii) *Limitations.* Subcutaneous use in swine only. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 7164, Feb. 18, 1983, as amended at 49 FR 26715, June 29, 1984; 54 FR 400, Jan. 6, 1989; 61 FR 5506, Feb. 13, 1996]

§ 522.940 Colloidal ferric oxide injection.

(a) *Specifications.* Each milliliter of the drug contains colloidal ferric oxide equivalent to 100 milligrams of iron stabilized with a low-viscosity dextrin and contains 0.5 percent phenol as a preservative.

(b) *NAS/NRC status.* Use of this drug has been NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(c)(1) *Sponsor.* See Nos. 010042 and 017800 in § 510.600(c) of this chapter.

(2) *Conditions of use.* It is used in baby pigs as follows:

(i) For the prevention of anemia due to iron deficiency, administer an initial intramuscular injection of 1 milliliter of the drug to each animal at any time between 2 to 5 days of age. Dosage may be repeated at 2 weeks of age.

(ii) For the treatment of anemia due to iron deficiency, administer an intramuscular injection of from 1 to 2 milliliters of the drug to each animal at any time between 5 to 28 days of age.

[40 FR 13858, Mar. 27, 1975, as amended at 49 FR 38938, Oct. 2, 1984; 50 FR 23298, June 3, 1985; 50 FR 25216, June 18, 1985; 51 FR 14989, Apr. 22, 1986; 51 FR 18314, May 19, 1986]

§ 522.955 Florfenicol solution.

(a) *Specifications.* Each milliliter of sterile solution contains 300 milligrams of florfenicol.

(b) *Sponsor.* See 000061 in § 510.600(c) of this chapter.

(c) *Related tolerance.* See § 556.283 of this chapter.

(d) *Conditions of use*—(1) *Cattle*—(i) *Treatment of disease*—(A) *Amount.* 20

milligrams per kilogram of body weight (3 milliliters per 100 pounds) as an intramuscular injection. A second dose should be given 48 hours later. Alternatively, 40 milligrams per kilogram of body weight (6 milliliters per 100 pounds) as a single subcutaneous injection may be used.

(B) *Indications for use.* For treatment of bovine respiratory disease (BRD) associated with *Pasteurella haemolytica*, *P. multocida*, and *Haemophilus somnus*. For treatment of bovine interdigital phlegmon (foot rot, acute interdigital necrobacillosis, infectious pododermatitis) associated with *Fusobacterium necrophorum* and *Bacteroides melaninogenicus*.

(ii) *Control of disease—(A) Amount.* 40 milligrams per kilogram of body weight (6 milliliters per 100 pounds) as a single subcutaneous injection.

(B) *Indications for use.* For control of respiratory disease in cattle at high risk of developing bovine respiratory disease (BRD) associated with *Pasteurella haemolytica*, *P. multocida*, and *Haemophilus somnus*.

(iii) *Limitations.* For intramuscular or subcutaneous use only. Do not inject more than 10 milliliters at each site. Injection should be given in the neck only. Do not slaughter within 28 days of last intramuscular treatment or within 38 days of subcutaneous treatment. Do not use in female dairy cattle 20 months of age or older. Use may cause milk residues. A withdrawal period has not been established in preruminating calves. Do not use in calves to be processed for veal. Not for use in cattle of breeding age. The effect of florfenicol on bovine reproductive performance, pregnancy, and lactation have not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[61 FR 42383, Aug. 15, 1996, as amended at 63 FR 26981, May 15, 1998; 63 FR 41191, Aug. 3, 1998; 64 FR 5596, Feb. 4, 1999; 64 FR 9435, Feb. 26, 1999]

§ 522.960 Flumethasone implantation or injectable dosage forms.

§ 522.960a Flumethasone suspension.

(a) *Chemical name.* 6 α ,9 α -Difluoro-11 β ,17,21 - trihydroxy - 16 α - methylpregna - 1,4 - diene - 3,20 - dione.

(b) *Specifications.* Flumethasone suspension is sterile and each milliliter of the drug contains: 2 milligrams of flumethasone, 20 milligrams of propylene glycol, 9 milligrams of benzyl alcohol (as preservative), 8 milligrams of sodium chloride, 0.02 milligram of polysorbate-80, 0.1 milligram of citric acid, and water for injection q.s.

(c) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) It is recommended in the various disease states involving synovial structures (joints) of horses where excessive synovial fluid of inflammatory origin is present and where permanent structural changes do not exist. Such conditions include arthritis, carpalitis, and osselets.

(2) The drug is administered intraarticularly at a dosage level of 6 to 10 milligrams per injection. The dosage level is dependent upon the size of the involved synovial structure and the degree of severity of the condition under treatment. The dosage is limited to a single injection per week in any one synovial structure.

(3) Clinical and experimental data have demonstrated that corticosteroids administered orally and parenterally to animals during the last trimester of pregnancy may induce the first stage of parturition and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis. The drug is not to be used in horses intended for slaughter for food purposes.

(4) For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975. Redesignated at 44 FR 16011, Mar. 16, 1979, as amended at 61 FR 5506, Feb. 13, 1996]

§ 522.960b

21 CFR Ch. I (4–1–99 Edition)

§ 522.960b Flumethasone acetate injection.

(a) *Chemical name.* 6- α ,9- α -difluoro - 16 - α - methylprednisolone 21-acetate.

(b) *Specifications.* Flumethasone injection is sterile and contains per cubic centimeter: 2 milligrams of flumethasone acetate; 20 milligrams of propylene glycol; 9 milligrams of benzyl alcohol (as preservative); 8 milligrams of sodium chloride; 1 milligram of polysorbate 80; 0.1 milligram of citric acid; water for injection q.s.

(c) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) It is recommended in certain acute and chronic canine dermatoses of varying etiology to help control the pruritus, irritation, and inflammation associated with these conditions.

(2) The drug is administered intramuscularly at the following recommended daily dosage:

Weight of animal in pounds	Dosage in milligrams
Up to 10	1.0
10 to 25	2.0
25 and over	4.0

Dosage should be adjusted according to the weight of the animal, the severity of the symptoms, and the response noted. Dosage by injection should not exceed 3 days of therapy. With chronic conditions intramuscular therapy may be followed by oral administration of flumethasone tablets at a daily dose of from 0.0625 to 0.25 milligram per animal.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975. Redesignated at 44 FR 16011, Mar. 16, 1979, as amended at 61 FR 5507, Feb. 13, 1996]

§ 522.960c Flumethasone solution.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 0.5 milligram flumethasone.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* It is used as follows:

(1) *Horses*—(i) *Amount.* 1.25 to 2.5 milligrams daily, intravenously, intramuscularly, or intra-articularly.

(ii) *Indications for use.* It is used for the treatment of musculoskeletal conditions due to inflammation, where permanent structural changes do not exist, e.g., bursitis, carpalis, osselets, and myositis; and allergic states, e.g., hives, urticaria, and insect bites.

(iii) *Limitations*—(a) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(b) When a long-term therapy is used, the dose should be individually adjusted to the minimum maintenance dose. A protein-rich diet is useful in dogs and cats on long-term therapy to counteract nitrogen loss if it should occur. A small amount of potassium chloride daily in the diet will counteract excessive potassium loss if this is present.

(c) It has been demonstrated that corticosteroids, especially at high dose levels, may result in delayed wound and fracture healing.

(d) Flumethasone may be administered to animals with bacterial diseases provided appropriate antibacterial therapy is administered simultaneously.

(e) The drug is not to be used in horses intended for slaughter for food purposes.

(f) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Dogs*—(i) *Amount.* 0.0625 to 0.25 milligram daily, intravenously, intramuscularly, or subcutaneously; 0.125 to 1.0 milligram daily, intralesionally, depending on the size and location of the lesion; 0.166 to 1.0 milligram daily, intra-articularly, depending on the severity of the condition and the size of the involved joint.

(ii) *Indications for use.* It is used for the treatment of musculoskeletal conditions due to inflammation of muscles or joints and accessory structures where permanent structural changes do not exist, e.g., arthritis, osteoarthritis, disc syndrome, and myositis (in septic arthritis, appropriate antibacterial

therapy should be concurrently administered); certain acute and chronic dermatoses of varying etiology to help control associated pruritus, irritation, and inflammation; otitis externa in conjunction with topical medication; allergic states, e.g., hives, urticaria, and insect bites; and shock and shock-like states by intravenous administration.

(iii) *Limitations.* See paragraph (c)(1)(iii) of this section.

(3) *Cats*—(i) *Amount.* 0.03125 to 0.125 milligram daily intravenously, intramuscularly, or subcutaneously.

(ii) *Indications for use.* It is used for the treatment of certain acute and chronic dermatoses of varying etiology to help control associated pruritus, irritation, and inflammation.

(iii) *Limitations.* See paragraph (c)(1)(iii) of this section.

[44 FR 16011, Mar. 16, 1978, as amended at 61 FR 5507, Feb. 13, 1996]

§ 522.970 Flunixin meglumine solution.

(a) *Specifications.* The drug contains 50 milligrams of flunixin per milliliter of aqueous solution.

(b) *Sponsors.* See 000061 in § 510.600(c) of this chapter for use as in paragraph (d) of this section. See 000856 and 059130 for use as in paragraph (d)(1) of this section only.

(c) *Related tolerances.* See § 556.286 of this chapter.

(d) *Conditions of use*—(1) *Horses*—(i) *Amount.* 0.5 milligram of flunixin per pound of body weight (1 milliliter per 100 pounds) per day.

(ii) *Indications for use.* For alleviation of inflammation and pain associated with musculoskeletal disorders, and alleviation of visceral pain associated with colic.

(iii) *Limitations.* For musculoskeletal disorders, administer intravenously or intramuscularly for up to 5 days. For colic, administer a single dose intravenously—treatment may be repeated when signs of colic recur. Caution: The effect of this drug on pregnancy has not been determined. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Beef cattle and nonlactating dairy cattle*—(i) *Amount.* 1.1 to 2.2 milligrams per kilogram of body weight (0.5 to 1

milligram per pound, 1 to 2 milliliters per 100 pounds), once a day as a single dose or divided into 2 doses administered at 12-hour intervals for up to 3 days.

(ii) *Indications for use.* For control of pyrexia associated with bovine respiratory disease and endotoxemia. Also indicated for control of inflammation in endotoxemia.

(iii) *Limitations.* Do not slaughter for food use within 4 days of last treatment. Not for use in lactating or dry dairy cows. A withdrawal period has not been established for use in preruminating calves. Do not use in calves to be processed for veal. Do not use in bulls intended for breeding as reproductive effects in this class of cattle have not been studied. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[42 FR 39103, Aug. 2, 1977, as amended at 52 FR 7832, Mar. 13, 1987; 60 FR 54942, Oct. 27, 1995; 62 FR 22888, Apr. 28, 1997; 63 FR 38749, July 20, 1998]

§ 522.995 Fluprostenol sodium injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains fluprostenol sodium equivalent to 50 micrograms of fluprostenol.

(b) *Sponsor.* See 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 0.55 microgram fluprostenol per kilogram of body weight.

(2) *Indications for use.* The drug is used in mares for its luteolytic effect to control the timing of estrus in estrous cycling and in clinically anestrous mares that have a corpus luteum.

(3) *Limitations.* Administer by intramuscular injection only. *Warning:* Not for use in horses intended for food. For veterinary use only. Federal law restricts this drug to use by or on the order of a licensed veterinarian. Women of childbearing age, asthmatics, and persons with bronchial and other respiratory problems should exercise extreme caution when handling this product. In the early stages, women may be unaware of their pregnancies. Fluprostenol is readily absorbed through the skin and can cause abortion and/or bronchospasms. Direct

§ 522.1002

contact with the skin should therefore be avoided. Accidental spillage on the skin should be washed off immediately with soap and water.

[44 FR 52191, Sept. 7, 1979, as amended at 47 FR 22092, May 21, 1982]

§ 522.1002 Follicle stimulating hormone.

(a)(1) *Specifications.* Each package contains 2 vials. One vial contains dry, powdered, porcine pituitary gland equivalent to 75 units (NIH-FSH-S1) of follicle stimulating hormone. The other vial contains 10 milliliters of aqueous diluent.

(2) *Sponsor.* See 059521 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) *Dosage.* 12.5 units of follicle stimulating hormone twice a day for 3 days (a total of 75 units). To effect regression of the corpus luteum, prostaglandin should be given with the 5th dose.

(ii) *Indications for use.* For induction of superovulation in cows for procedures requiring the production of multiple ova at a single estrus.

(iii) *Limitations.* For intramuscular use in cows that are not pregnant and have a normal corpus luteum. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Specifications.* The drug is a lyophilized pituitary extract material. Each 10-milliliter vial contains an amount equivalent to 50 milligrams of standard porcine follicle stimulating hormone and is reconstituted for use by addition of 10 milliliters of 0.9 percent aqueous sodium chloride solution.

(2) *Sponsor.* See 063112 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) *Dosage.* Cattle and horses, 10–50 milligrams; sheep and swine, 5–25 milligrams; dogs, 5–15 milligrams.

(ii) *Indications for use.* The drug is used as a supplemental source of follicle stimulating hormone where there is a general deficiency in cattle, horses, sheep, swine, and dogs.

(iii) *Limitations.* Administer intramuscularly, subcutaneously, or intravenously. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[58 FR 47377, Sept. 9, 1993, as amended at 62 FR 62242, Nov. 21, 1997]

21 CFR Ch. I (4–1–99 Edition)

§ 522.1004 Fomepizole.

(a) *Specifications.* Two vials, one containing 1.5 grams fomepizole (1.5 milliliter of 1.0 gram fomepizole per milliliter sterile aqueous solution), and one vial containing 30 milliliters of 0.9 percent sodium chloride injection USP (as a diluent).

(b) *Sponsor.* See 062161 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs—*(1) *Amount.* 20 milligrams per kilogram initially, 15 milligrams per kilogram at 12 and 24 hours, and 5 milligrams per kilogram at 36 hours.

(2) *Indications for use.* As an antidote for ethylene glycol (antifreeze) poisoning in dogs who have ingested or are suspected of having ingested ethylene glycol.

(3) *Limitations.* Administer intravenously. For use by or on the order of a licensed veterinarian.

[61 FR 68147, Dec. 27, 1996]

§ 522.1010 Furosemide injection.

(a) *Specifications.* Each milliliter of sterile solution contains 50 milligrams of furosemide as the diethanolamine salt.

(b) *Sponsor.* See No. 012799 in § 510.600(c) of this chapter for use in dogs and cats as in paragraph (c)(1) of this section, horses as in paragraph (c)(2)(i) of this section, and cattle as in paragraph (c)(3) of this section. See Nos. 000010 and 000864 in § 510.600(c) for use in horses as in paragraph (c)(2)(ii) of this section. See No. 000010 in § 510.600(c) of this chapter for use in dogs as in paragraph (c)(1) of this section.

(c) *Conditions of use—*(1) *Dogs and cats.* (i) It is used for the treatment of edema (pulmonary congestion, ascites) associated with cardiac insufficiency and acute noninflammatory tissue edema.

(ii) The drug is administered intramuscularly or intravenously at a dosage of 12.5 to 25 milligrams per 10 pounds of body weight; once or twice daily after a 6- to 8-hour interval. The lower dosage is suggested for cats. The dosage should be adjusted to the individual animal's response. In refractory or severe edematous cases, the dosage

may be doubled or increased by increments of 1 milligram per pound of body weight to establish the effective dose. The established effective dose should be administered once or twice daily on an intermittent daily schedule. Diuretic therapy should be discontinued after reduction of edema, or when necessary, maintained after determining a programmed dosage schedule to prevent recurrence.

(2) *Horses.* (i) It is used for the treatment of edema (pulmonary congestion, ascites) associated with cardiac insufficiency and acute noninflammatory tissue edema.

(a) Administer intramuscularly or intravenously at 250 to 500 milligrams per animal once or twice daily at 6- to 8-hours intervals until desired results are achieved.

(b) Do not use in horses intended for food.

(ii) It is used for treatment of acute noninflammatory tissue edema.

(a) Administer intramuscularly or intravenously at 0.5 milligram per pound of body weight (1.0 milligram per kilogram); once or twice daily at 6- to 8-hour intervals.

(b) The dosage should be adjusted to the individual's response. In refractory or severe edematous cases, the dosage may be doubled or increased by increments of 1 milligram per pound of body weight to establish the effective dose. The established effective dose should be administered once or twice daily on an intermittent daily schedule, i.e., every other day or 2 to 4 consecutive days weekly. Concurrent therapy for treatment of systemic conditions causing edema (pulmonary congestion, ascites, cardiac insufficiency) should be instituted.

(3) *Cattle.* (i) It is used for the treatment of physiological parturient edema of the mammary gland and associated structures.

(ii) The drug is administered intramuscularly or intravenously at a dosage of 500 milligrams per animal once daily or 250 milligrams per animal twice daily at 12-hour intervals, treatment not to exceed 48 hours postparturition.

(iii) Milk taken during treatment and for 48 hours (four milkings) after

the last treatment must not be used for food.

(iv) Cattle must not be slaughtered for food within 48 hours following last treatment.

(4) The drug if given in excessive amounts may result in dehydration and electrolyte imbalance.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 60051, Dec. 31, 1975, as amended at 41 FR 10426, Mar. 11, 1976; 48 FR 36572, Aug. 12, 1983; 49 FR 26715, June 29, 1984; 53 FR 40727, Oct. 18, 1988; 62 FR 35076, June 30, 1997]

§ 522.1020 Gelatin solution.

(a) *Specifications.* It is sterile and each 100 cubic centimeters contains 8 grams of gelatin in an 0.85 percent sodium chloride solution.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used to restore circulatory volume and maintain blood pressure in animals being treated for shock.

(2) The exact dosage to be administered must be determined after evaluating the animal's condition and will vary according to the size of the animal and the degree of shock. A suggested dosage range for small animals such as dogs is 4 to 8 cubic centimeters per pound body weight. The suggested dosage range for large animals such as sheep, calves, cows, or horses is 2 to 4 cubic centimeters per pound of body weight. It is administered intravenously at a rate of 10 cubic centimeters per minute in small animals and 20 to 30 cubic centimeters per minute in large animals. The solution is administered aseptically and must be between 50 to 70 °F. when injected.

(3) A few animals will exhibit signs of allergic reaction. This solution can cause transient reversible nephrosis. This product is not intended to replace whole blood in cases of anemia and should not be used in the presence of renal dysfunction. Unused portions remaining in bottles should be discarded.

(4) For use only by or on the order of a licensed veterinarian.

§ 522.1044 Gentamicin sulfate injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains gentamicin sulfate equivalent to either 5, 50, or 100 milligrams of gentamicin.

(b) *Sponsors.* (1) See No. 000061 in § 510.600(c) of this chapter for use of: 5-milligrams-per-milliliter solution in swine as in paragraph (d)(4) of this section, 50-milligrams-per-milliliter solution in dogs and cats as in paragraph (d)(1) of this section, 50- and 100-milligrams-per-milliliter solution in chickens and turkeys as in paragraphs (d)(2) and (d)(3) of this section.

(2) [Reserved]

(3) See No. 000010 for use of 50 milligrams-per-milliliter solution in dogs as in paragraph (d)(5) of this section.

(4) See No. 050604 for use of 100 milligram-per-milliliter solution in turkeys as in paragraph (d)(2) of this section and in chickens as in paragraph (d)(3) of this section.

(c) *Related tolerances.* See § 556.300 of this chapter.

(d) *Conditions of use—(1) Dogs and cats—(i) Amount.* Two milligrams of gentamicin per pound of body weight, twice daily on the first day, once daily thereafter, using a 50 milligram-per-milliliter solution.

(ii) *Indications for use—(a) Dogs.* For the treatment of infections of urinary tract (cystitis, nephritis), respiratory tract (tonsillitis, pneumonia, tracheobronchitis), skin and soft tissue (pyodermatitis, wounds, lacerations, peritonitis).

(b) *Cats.* For the treatment of infections of urinary tract (cystitis, nephritis), respiratory tract (pneumonitis, pneumonia, upper respiratory tract infections), skin and soft tissue (wounds, lacerations, peritonitis), and as supportive therapy for secondary bacterial infections associated with panleucopenia.

(iii) *Limitations.* Administer intramuscularly or subcutaneously. If response is not noted after 7 days, the antibiotic sensitivity of the infecting organism should be retested. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Turkeys—(i) Amount.* One milligram of gentamicin per 0.2 milliliter dose, using the 50- or 100-milligrams-

per-milliliter product diluted with sterile saline to a concentration of 5 milligrams-per-milliliter.

(ii) *Indications for use.* As an aid in the prevention of early mortality due to Arizona paracolon infections susceptible to gentamicin.

(iii) *Limitations.* For 1- to 3-day old turkey poults. Administer subcutaneously in the neck. Injected poults must not be slaughtered for food for at least 9 weeks after treatment.

(3) *Chickens—(i) Amount.* 0.2 milligram of gentamicin per 0.2 milliliter dose, using the 50- or 100-milligrams-per-milliliter product diluted with sterile saline to a concentration of 1.0 milligram-per-milliliter.

(ii) *Indications for use.* In day-old chickens, for prevention of early mortality caused by *Escherichia coli*, *Salmonella typhimurium*, and *Pseudomonas aeruginosa* that are susceptible to gentamicin.

(iii) *Limitations.* For use in day-old chickens only. Administer aseptically, injecting the diluted product subcutaneously in the neck. Do not slaughter treated animals for food for at least 5 weeks after treatment.

(4) *Swine—(i) Amount.* 5 milligrams of gentamicin as a single intramuscular dose using 5 milligram-per-milliliter solution.

(ii) *Indications for use.* In piglets up to 3 days old for treatment of porcine colibacillosis caused by strains of *E. coli* sensitive to gentamicin.

(iii) *Limitations.* For single intramuscular dose in pigs up to 3 days of age only. Do not slaughter treated animals for food for at least 40 days following treatment.

(5) *Dogs—(i) Amount.* 2 milligrams of gentamicin per pound of body weight, twice daily on the first day, then once daily.

(ii) *Indications for use.* For use in the treatment of urinary tract infections (cystitis) caused by *Proteus mirabilis*, *Escherichia coli*, and *Staphylococcus aureus*.

(iii) *Limitations.* Administer intramuscularly or subcutaneously. If no improvement is seen after 3 days, treatment should be discontinued and the diagnosis reevaluated. Treatment

Food and Drug Administration, HHS

§522.1079

not to exceed 7 days. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 1942, Jan. 13, 1978, as amended at 48 FR 791, Jan. 7, 1983; 51 FR 15606, Apr. 25, 1986; 52 FR 7832, Mar. 13, 1987; 53 FR 40727, Oct. 18, 1988; 60 FR 29985, June 7, 1995; 61 FR 24441, May 15, 1996; 62 FR 45157, Aug. 26, 1997; 63 FR 59714, Nov. 5, 1998; 63 FR 68182, Dec. 10, 1998]

EDITORIAL NOTE: At 62 FR 35076, June 30, 1997, §520.1044 was amended in paragraph (b)(3) by removing "054273" and adding in its place "000010". The amendment could not be incorporated because paragraph (b)(3) was not added until Aug. 26, 1997, at 62 FR 45157.

§522.1055 Gleptoferron injection.

(a) *Specifications.* Each milliliter contains the equivalent of 200 milligrams of elemental iron as gleptoferron (complex of ferric hydroxide and dextran glucoheptonic acid), and 0.5 percent phenol as a preservative.

(b) *Sponsor.* See 062408 in §510.600(c) of this chapter.

(c) *Conditions of use.* It is used in baby pigs as follows:

(1) For prevention of iron deficiency anemia, administer 200 milligrams of elemental iron intramuscularly on or before 3 days of age.

(2) For treatment of iron deficiency anemia, administer 200 milligrams of elemental iron intramuscularly.

[45 FR 61288, Sept. 16, 1980, as amended at 61 FR 18672, Apr. 29, 1996]

§522.1066 Glycopyrrolate injection.

(a) *Specifications.* Each milliliter of aqueous solution contains 0.2 milligram of glycopyrrolate.

(b) *Sponsor.* See No. 000031 in §510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is indicated as a preanesthetic agent in dogs and cats.

(2) It is administered intravenously, intramuscularly, or subcutaneously in dogs and intramuscularly in cats at a dosage level of 5 micrograms per pound of body weight (0.25 milliliter per 10 pounds of body weight).

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 21567, May 13, 1983]

§522.1077 Gonadorelin injectable.

(a) *Specifications.* Each milliliter sterile aqueous solution contains 50 micrograms of gonadorelin (as hydrochloride).

(b) *Sponsor.* See No. 000856 in §510.600(c) of this chapter.

(c) *Conditions of use in cattle.*—(1) *Amount.* 100 micrograms per cow intramuscularly.

(2) *Indications for use.* For the treatment of cystic ovaries (ovarian follicular cysts) in cattle to reduce the time to first estrus.

(3) *Limitations.* For intramuscular use only. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[54 FR 50235, Dec. 5, 1989]

§522.1078 Gonadorelin diacetate tetrahydrate injection

(a) *Specifications.* The drug contains 50 micrograms of gonadorelin diacetate tetrahydrate in each milliliter of sterile solution.

(b) *Sponsor.* See Nos. 050604 and 057926 in §510.600(c) of this chapter.

(c) *Conditions of use.* It is used in dairy cows as follows:

(1) *Amount.* 100 micrograms per cow.

(2) *Indications for use.* The drug is used for the treatment of ovarian cysts.

(3) *Limitations.* Administer as a single intramuscular or intravenous injection. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 9804, Mar. 10, 1978, as amended at 45 FR 56798, Aug. 26, 1980; 61 FR 37682, July 19, 1996]

§522.1079 Serum gonadotropin and chorionic gonadotropin.

(a) *Specifications.* Each dose consists of 400 international units (I.U.) serum gonadotropin and 200 I.U. chorionic gonadotropin as a freeze-dried powder to be reconstituted with 5 milliliters of sterile aqueous diluent.

(b) *Sponsor.* See No. 057926 in §510.600(c) of this chapter.

(c) *Conditions of use in swine.* (1) *Amount.* 400 I.U. serum gonadotropin with 200 I.U. chorionic gonadotropin per 5 milliliters dose per animal.

§ 522.1081

21 CFR Ch. I (4–1–99 Edition)

(2) *Indications for use.* (i) *Gilts.* For induction of fertile estrus (heat) in healthy prepuberal (noncycling) gilts.

(ii) *Sows.* For induction of estrus in healthy weaned sows experiencing delayed return to estrus.

(3) *Limitations.* For subcutaneous use only.

(i) *Gilts.* For use only in gilts over 5 1/2 months of age and weighing at least 85 kilograms (187 pounds).

(ii) *Sows.* Delayed return to estrus is most prevalent after the first litter. The effectiveness has not been established after later litters. Delayed return to estrus often occurs during periods of adverse environmental conditions, and sows mated under such conditions may farrow smaller than normal litters.

[55 FR 1405, Jan. 16, 1990, as amended at 58 FR 52222, Oct. 7, 1993]

§ 522.1081 Chorionic gonadotropin for injection; chorionic gonadotropin suspension.

(a)(1) *Specifications.* Chorionic gonadotropin for injection is supplied in vials containing 5,000, 10,000 or 20,000 U.S.P. units of lyophilized powder for reconstitution with the accompanying sterile diluent to a 10 milliliter solution.

(2) *Sponsor.* See sponsor numbers in § 510.600(c) of this chapter, as follows:

(i) Nos. 000402 and 053501 for use of 10,000 U.S.P. units intramuscularly, 2,500 to 5,000 U.S.P. units intravenously, and 500 to 2,500 U.S.P. units intrafollicularly.

(ii) Nos. 058639 and 063323 for use of 10,000 U.S.P. units intramuscularly and 500 to 2,500 U.S.P. units intrafollicularly.

(iii) No. 057926 for use of 10,000 U.S.P. units intramuscularly.

(3) *Conditions of use*—(i) *Amount.* (a) 10,000 U.S.P. units as a single, deep intramuscular injection.¹

(b) 500 to 2,500 U.S.P. units for intrafollicular injection.¹

(c) 2,500 to 5,000 U.S.P. units intravenously.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified as § 514.111 of this chapter.

(ii) *Indications for use.* For parenteral use in cows for treatment of nymphomania (frequent or constant heat) due to cystic ovaries.¹

(iii) *Limitations.* Dosage may be repeated in 14 days if the animal's behavior or rectal examination of the ovaries indicates the necessity for retreatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

(b)(1) *Specifications.* Chorionic gonadotropin suspension, veterinary contains in each milliliter, 750 I.U. of chorionic gonadotropin suspended in white wax and sesame oil.

(2) *Sponsor.* See No. 000986 in § 510.600(c) of this chapter.

(3) *Conditions of use*—(i) *Amount.* 2 milliliters (1,500 I.U.) subcutaneously, at the time of insemination, in the neck or shoulder region.

(ii) *Indications for use.* The drug is used as an aid in increasing pregnancy rate of estrus-synchronized and normalcycling heifers.

(iii) *Limitations.* The drug is not to be used to induce multiple ovulations. Doses higher than recommended may reduce pregnancy rate. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[42 FR 58167, Nov. 8, 1977, as amended at 45 FR 81038, Dec. 9, 1980; 50 FR 41489, Oct. 11, 1985; 50 FR 45603, Nov. 1, 1985; 52 FR 25212, July 6, 1987; 56 FR 67175, Dec. 30, 1991; 56 FR 14642, Apr. 11, 1991; 63 FR 51822, Sept. 29, 1998]

§ 522.1085 Guaifenesin sterile powder.

(a) *Specifications.* It is a sterile powder containing guaifenesin.

(b) *Sponsor.* See No. 000031 and 037990 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is indicated for intravenous use as a muscle relaxant in horses.

(2) A solution is prepared by dissolving the drug in sterile water for injection to make a solution containing 50 milligrams of guaifenesin per milliliter of solution. It is administered by rapid intravenous infusion at a fixed dosage of 1 milliliter of prepared solution per pound of body weight.

(3) Not to be used in horses intended for food.

Food and Drug Administration, HHS

§522.1145

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[49 FR 48039, Dec. 10, 1984, as amended at 60 FR 27223, May 23, 1995]

§522.1086 Guaifenesin injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 50 milligrams of guaifenesin and 50 milligrams of dextrose.

(b) *Sponsor.* See Nos. 037990 and 059130 in §510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use.* (1) The drug is used intravenously in horses as a skeletal muscle relaxant.

(2) Administer rapidly at a dosage of 1 milliliter per pound of body weight.

(3) No to be used in horses intended for food.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[60 FR 27223, May 23, 1995, as amended at 63 FR 29352, May 29, 1998]

§522.1125 Hemoglobin glutamer-200 (bovine).

(a) *Specifications.* Each 125 milliliter bag contains 13 grams per deciliter of polymerized hemoglobin of bovine origin in modified Lactated Ringer's Solution. It is a sterile, clear, dark purple solution.

(b) *Sponsor.* See No. 063075 in §510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use—*(1) *Amount.* One-time dose of 30 milliliters per kilogram of body weight administered intravenously at a rate of up to 10 milliliters per kilogram per hour.

(2) *Indications for use.* For the treatment of anemia in dogs by increasing systemic oxygen content (plasma hemoglobin concentration) and improving the clinical signs associated with anemia for at least 24 hours, regardless of the cause of anemia (hemolysis, blood loss, or ineffective erythropoiesis).

(3) *Limitations.* For intravenous use only. Overdosage or an excessive rate of administration (greater than 10 milliliters per kilogram per hour) may result in circulatory overload. Federal

law restricts this drug to use by or on the order of a licensed veterinarian.

[63 FR 11598, Mar. 10, 1998]

§522.1145 Hyaluronate sodium injection.

(a)(1) *Specifications.* Each milliliter of sterile aqueous solution contains 10 milligrams of hyaluronate sodium.

(2) *Sponsor.* See 000009 in §510.600(c).

(3) *Conditions of use—*(i) *Amount.* Small and medium-size joints (carpal, fetlock)—20 milligrams; larger joint (hock)—40 milligrams.

(ii) *Indications for use.* Treatment of joint dysfunction in horses due to non-infectious synovitis associated with equine osteoarthritis.

(iii) *Limitations.* For intra-articular injection in horses only. Treatment may be repeated at weekly intervals for a total of three treatments. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Specifications.* Each milliliter of sterile aqueous solution contains 5 milligrams of hyaluronate sodium.

(2) *Sponsor.* See 053501 in §510.600(c) of this chapter.

(3) *Conditions of use—*(i) *Amount.* Small and medium-size joints (carpal, fetlock)—10 milligrams; larger joint (hock)—20 milligrams.

(ii) *Indications for use.* Treatment of joint dysfunction in horses due to non-infectious synovitis associated with equine osteoarthritis.

(iii) *Limitations.* For intraarticular injection in horses only. Treatment may be repeated at weekly intervals for a total of four treatments. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(c)(1) *Specifications.* Each milliliter of sterile aqueous solution contains 10 milligrams of hyaluronate sodium.

(2) *Sponsor.* See 000856 in §510.600(c) of this chapter.

(3) *Conditions of use—*(i) *Amount.* Small and medium-size joints (carpal, fetlock)—20 milligrams.

(ii) *Indications for use.* Treatment of carpal or fetlock joint dysfunction in horses due to acute or chronic non-infectious synovitis associated with equine osteoarthritis.

§ 522.1150

(iii) *Limitations.* For intraarticular injection in horses only. Treatment may be repeated after 1 or more weeks but not to exceed 2 injections per week for a total of 4 weeks. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d)(1) *Specifications.* Each milliliter of sterile aqueous solution contains 10 milligrams of hyaluronate sodium.

(2) *Sponsor.* See 000061 in § 510.600(c) of this chapter.

(3) *Conditions of use*—(i) *Amount.* 50 milligrams in carpal and fetlock joints.

(ii) *Indications for use.* For treatment of equine carpal and fetlock joint dysfunction caused by traumatic and/or degenerative joint disease of mild to moderate severity.

(iii) *Limitations.* For intraarticular injection in horses only. Not for use in horses intended for food. Not intended for use in breeding animals. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(e)(1) *Specifications.* Each milliliter of sterile aqueous solution contains 10 milligrams of hyaluronate sodium.

(2) *Sponsor.* See 000859 in § 510.600(c)(2) of this chapter.

(3) *Conditions of use*—(i) *Amount.* Intraarticular: 20 milligrams in the carpus or fetlock. Intravenous: 40 milligrams slowly into the jugular vein.

(ii) *Indications for use.* Treatment of carpal or fetlock joint dysfunction in horses due to noninfectious synovitis associated with equine osteoarthritis.

(iii) *Limitations.* For intraarticular or intravenous use in horses only. Treatment may be repeated at weekly intervals for a total of three treatments. Not for use in horses intended for food. The safety of use of this drug in breeding animals has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(f)(1) *Specifications.* Each milliliter of sterile aqueous solution contains 11 milligrams of hyaluronate sodium.

(2) *Sponsor.* See 060865 in § 510.600(c).

(3) *Conditions of use*—(i) *Amount.* Small and medium-size joints (carpal, fetlock)—22 milligrams; larger joint (hock)—44 milligrams.

(ii) *Indications for use.* Treatment of joint dysfunction in horses due to non-

21 CFR Ch. I (4–1–99 Edition)

infectious synovitis associated with equine osteoarthritis.

(iii) *Limitations.* For intra-articular injection in horses only. Treatment may be repeated at weekly intervals for a total of three treatments. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[49 FR 45124, Nov. 15, 1984, as amended at 51 FR 11438, Apr. 3, 1986; 51 FR 25032, July 10, 1986; 53 FR 19773, May 31, 1988; 53 FR 22297, June 15, 1988; 56 FR 50814, Oct. 9, 1991; 57 FR 2837, Jan. 24, 1992; 59 FR 33198, June 28, 1994; 61 FR 59003, Nov. 20, 1996; 63 FR 59216, Nov. 3, 1998]

§ 522.1150 Hydrochlorothiazide injection.

(a) *Specifications.* Each milliliter contains 25 milligrams of hydrochlorothiazide.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 5 to 10 milliliters (125 to 250 milligrams) intravenously or intramuscularly once or twice a day. After onset of diuresis, treatment may be continued with an orally administered maintenance dose.

(2) *Indications for use.* For use in cattle as an aid in the treatment of postparturient udder edema.¹

(3) *Limitations.* Animals should be regularly and carefully observed for early signs of fluid and electrolyte imbalance. Take appropriate countermeasures if this should occur. Milk taken from dairy animals during treatment and for 72 hours (6 milkings) after the latest treatment must not be used for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[43 FR 59058, Dec. 19, 1978, as amended at 62 FR 63271, Nov. 28, 1997]

§ 522.1155 Imidocarb dipropionate sterile powder.

(a) *Specifications.* Imidocarb dipropionate powder is reconstituted with sterile water. Each milliliter of

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

Food and Drug Administration, HHS

§522.1182

solution contains 100 milligrams of imidocarb base.

(b) *Sponsor*. No. 000061 in §510.600(c) of this chapter.

(c) *Conditions of use*. The drug is used in horses and zebras as follows:

(1) *Amount*. For *Babesia caballi* infections, use intramuscularly 2 milligrams of imidocarb base per kilogram of body weight, repeating dosage once after 24 hours. For *Babesia equi* infections, use 4 milligrams of imidocarb base per kilogram of body weight, repeating dosage four times at 72-hour intervals.

(2) *Indications for use*. For the treatment of babesiosis (piroplasmosis) caused by *Babesia caballi* and *Babesia equi*.

(3) *Limitations*. Administer intramuscularly in the neck region. Do not inject intravenously. Do not use for other equidae or for animals of other species. Do not use in horses less than 1 year old. Do not use for animals in near-term pregnancies. Imidocarb dipropionate is a cholinesterase inhibitor. Do not use this product simultaneously with or a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals. Do not use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian. Imidocarb dipropionate is sold only under permit issued by the Director of the National Program Planning Staff, Veterinary Services, APHIS, USDA, to licensed or full-time State, Federal, or military veterinarians.

[43 FR 40455, Sept. 12, 1978, as amended at 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§522.1156 Imidocarb dipropionate solution.

(a) *Specifications*. Each milliliter of injectable solution contains 120 milligrams of imidocarb.

(b) *Sponsor*. See No. 000061 in §510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Dogs*—(i) *Amount*. 6.6 milligrams imidocarb per kilogram (3 milligrams per pound) of body weight.

(ii) *Indications for use*. Treatment of clinical signs of babesiosis and/or dem-

onstrated *Babesia* organisms in the blood.

(iii) *Limitations*. Use subcutaneously or intramuscularly. Not for intravenous use. Repeat the dose after 2 weeks for a total of two treatments. Imidocarb is a cholinesterase inhibitor. Do not use simultaneously with or a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[62 FR 66984, Dec. 23, 1997]

§522.1182 Iron dextran complex injection.

(a)(1) *Specifications*. Each milliliter of sterile solution contains ferric hydroxide dextran complex equivalent to 100 milligrams of elemental iron. It contains 0.5 percent phenol as a preservative.

(2) [Reserved]

(3)(i) *Sponsor*. See No. 000856 in §510.600(c) of this chapter.

(ii) *Conditions of use*. It is used in baby pigs as follows:

(a) For the prevention of anemia due to iron deficiency, administer an initial intramuscular injection of 100 milligrams of elemental iron to each animal at 2 to 4 days of age. Dosage may be repeated in 14 to 21 days.

(b) For the treatment of anemia due to iron deficiency, administer and intramuscular injection of 200 milligrams of elemental iron.

(4)(i) *Sponsor*. See Nos. 000061 and 062408 in §510.600(c) of this chapter.

(ii) *Conditions of use*. It is used in baby pigs as follows:

(a) For the prevention of iron deficiency anemia, administer intramuscularly an amount of drug containing 100 to 150 milligrams of elemental iron to animals from 1 to 3 days of age.

(b) For the treatment of iron deficiency anemia, administer intramuscularly an amount of drug containing 100 to 200 milligrams of elemental iron per animal. Dosage may be repeated in 10 days to 2 weeks.

(b)(1) *Specifications*. Each milliliter of sterile solution contains ferric hydroxide in complex with dextran equivalent

§ 522.1183

to 200 milligrams of elemental iron. It contains 0.5 percent phenol as a preservative.

(2)(i) *Sponsor*. See Nos. 000010 and 059130 in § 510.600(c) of this chapter.

(ii) *Conditions of use*. It is used in baby pigs as follows:

(a) For prevention of baby pig anemia due to iron deficiency, intramuscularly inject 200 milligrams of elemental iron (1 milliliter) at 1 to 3 days of age.

(b) For treatment of baby pig anemia due to iron deficiency, intramuscularly inject 200 milligrams of elemental iron at the first sign of anemia.

[49 FR 38938, Oct. 2, 1984, as amended 52 FR 7832, Mar. 13, 1987; 53 FR 40727, Oct. 18, 1988; 54 FR 41442, Oct. 10, 1989; 61 FR 18672, Apr. 29, 1996; 62 FR 35076, June 30, 1997; 63 FR 53578, Oct. 6, 1998]

§ 522.1183 Iron hydrogenated dextran injection.

(a) *Specifications*. Each milliliter contains 100 milligrams of elemental iron stabilized with a low molecular weight hydrogenated dextran and 0.5 percent phenol as a preservative.

(b) [Reserved]

(c)(1) *Sponsor*. See No. 000986 in § 510.600(c) of this chapter.

(2) *Conditions of use*. It is used in baby pigs as follows:

(i) For the prevention of anemia due to iron deficiency, administer an initial intramuscular injection of 100 milligrams of elemental iron to each animal at 2 to 5 days of age. Dosage may be repeated at 2 weeks of age.

(ii) For the treatment of anemia due to iron deficiency, administer an intramuscular injection of 100 milligrams of elemental iron to each animal when indicated between 5 and 28 days of age.

(d)(1) *Sponsor*. See No. 000003 in § 510.600(c) of this chapter.

(2) *Conditions of use*. It is used in baby pigs as follows:

(i) For the prevention of anemia due to iron deficiency, administer by intramuscular or subcutaneous injection of 100 milligrams of elemental iron to each animal at 2 to 4 days of age.

(ii) For the treatment of anemia due to iron deficiency, administer by intramuscular or subcutaneous injection of 100 milligrams of elemental iron in baby pigs up to 4 weeks of age.

21 CFR Ch. I (4–1–99 Edition)

(e)(1) *Sponsors*. See Nos. 000010, 017287, 050604, and 059130 in § 510.600(c) of this chapter.

(2) *Conditions of use*. It is used in baby pigs as follows:

(i) For the prevention of iron deficiency anemia, administer intramuscularly 100 milligrams at 2 to 4 days of age.

(ii) For the treatment of iron deficiency anemia, administer intramuscularly 100 milligrams. Treatment may be repeated in 10 days.

[42 FR 53955, Oct. 4, 1977, as amended at 46 FR 39128, July 31, 1981; 50 FR 23298, June 3, 1985; 52 FR 18691, May 19, 1987; 52 FR 36023, Sept. 25, 1987; 53 FR 40728 and 40729, Oct. 18, 1988; 55 FR 8462, Mar. 8, 1990; 55 FR 33670, Aug. 17, 1990; 62 FR 35076, June 30, 1997; 63 FR 44384, Aug. 19, 1998]

§ 522.1192 Ivermectin injection.

(a) *Specifications*—(1) *Horses*. Each milliliter of sterile aqueous solution contains 20 milligrams of ivermectin (2 percent).

(2) *Cattle, reindeer, swine, and American bison*. Each milliliter of sterile aqueous solution contains 10 milligrams of ivermectin (1 percent).

(3) *Piglets 70 pounds or less and ranch-raised foxes*. Each milliliter of sterile aqueous solution contains 2.7 milligrams of ivermectin (0.27 percent).

(b) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.344 of this chapter.

(d) *Conditions of use*—(1) *Horses*—(i) *Amount*. 20 milligrams per 100 kilograms (220 pounds) of body weight.

(ii) *Indications for use*. It is used in horses for the treatment and control of large strongyles (adult) (*Strongylus vulgaris*, *Strongylus edentatus*, *Triodontophorus* spp.), small strongyles (adult and fourth stage larvae) (*Cyathostomum* spp., *Cylicocyclus* spp., *Cylicostephanus* spp.), pinworms (adult and fourth-stage larvae) (*Oxyuris equi*), large roundworms (adult) (*Parascaris equorum*), hairworms (adult) (*Trichostrongylus axei*), large mouth stomach worms (adult) (*Habronema muscae*), neck threadworms (*microfilariae*) (*Onchocerca* spp.), and stomach bots (*Gastrophilus* spp.).

(iii) *Limitations*. For intramuscular use only. Do not use intravenously. Not

for use in horses intended for food. Effects of this drug on pregnant mares have not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cattle*—(i) *Amount*. 10 milligrams per 50 kilograms (110 pounds) body weight (200 micrograms per kilogram).

(ii) *Indications for use*. It is used in cattle for the treatment and control of gastrointestinal nematodes (adults and fourth-stage larvae) (*Haemonchus placei*, *Ostertagia ostertagi* (including inhibited larvae), *O. lyrata*, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia oncophora*, *C. punctata*, *C. pectinata*, *Oesophagostomum radiatum*, *Nematodirus helvetianus* (adults only), *N. spathiger* (adults only), *Bunostomum phlebotomum*); lungworms (adults and fourth-stage larvae) (*Dictyocaulus viviparus*); grubs (first, second, and third instars) (*Hypoderma bovis*, *H. lineatum*); lice (*Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*); mites (*Psoroptes ovis* (syn. *P. communis* var. *bovis*), *Sarcoptes scabiei* var. *bovis*). It is also used to control infections of *D. viviparus* and *O. ostertagi* for 21 days after treatment, and *H. placei*, *T. axei*, *C. punctata*, *C. oncophora*, and *Oesophagostomum radiatum* for 14 days after treatment.

(iii) *Limitations*. For subcutaneous use only. Not for intramuscular use. Do not treat cattle within 35 days of slaughter. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. Do not use in other animal species because severe adverse reactions, including fatalities in dogs, may result. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(3) *Reindeer*—(i) *Amount*. 10 milligrams per 50 kilograms (110 pounds) body weight.

(ii) *Indications for use*. It is used in reindeer for treatment and control of warbles (*Oedemagena tarandi*).

(iii) *Limitations*. For subcutaneous use only. Not for intramuscular use. Do not treat reindeer within 56 days of slaughter. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(4) *Swine*—(i) *Amount*. 300 micrograms per kilogram (2.2 pounds).

(ii) *Indications for use*. It is used in swine for treatment and control of gastrointestinal roundworms (adults and fourth-stage larvae) (large roundworm, *Ascaris suum*; red stomach worm, *Hyoststrongylus rubidus*; nodular worm, *Oesophagostomum* spp.; threadworm, *Strongyloides ransomi* (adults only)); somatic roundworm larvae (threadworm, *Strongyloides ransomi* (somatic larvae)); lungworms (*Metastrongylus* spp. (adults only)); lice (*Haematopinus suis*); and mites (*Sarcoptes scabiei* var. *suis*).

(iii) *Limitations*. For subcutaneous injection in the neck of swine only. Do not treat swine within 18 days of slaughter. Do not use in other animal species as severe adverse reactions, including fatalities in dogs, may result. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(5) *Ranch-raised foxes*—(i) *Amount*. 200 micrograms per kilogram body weight. Repeat in 3 weeks.

(ii) *Indications for use*. For treatment and control of ear mites (*Otodectes cynotis*).

(iii) *Limitations*. For subcutaneous use only. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(6) *American bison*—(i) *Amount*. 200 micrograms per kilogram (10 milligrams per 110 pounds) of body weight.

(ii) *Indications for use*. It is used in American bison for the treatment and control of grubs (*Hypoderma bovis*).

(iii) *Limitations*. For subcutaneous use. Do not slaughter within 56 days of last treatment. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[49 FR 5344, Feb. 13, 1984, as amended at 50 FR 30268, July 25, 1985; 51 FR 25686, July 16, 1986; 51 FR 27021, July 29, 1986; 51 FR 29463, Aug. 18, 1986; 53 FR 11064, Apr. 5, 1988; 56 FR 14020, Apr. 5, 1991; 60 FR 45041, Aug. 30, 1995; 62 FR 14634, Mar. 27, 1997; 62 FR 63271, Nov. 28, 1997; 63 FR 7702, Feb. 17, 1998]

§522.1193 Ivermectin and clorsulon injection.

(a) *Specifications*. Each milliliter of sterile aqueous solution contains 10 milligrams (1 percent) of ivermectin and 100 milligrams (10 percent) of clorsulon.

§ 522.1204

(b) *Sponsor*. See 050604 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See §§ 556.163 and 556.344 of this chapter.

(d) *Conditions of use*—(1) *Amount*. 1 milliliter (10 milligrams of ivermectin and 100 milligrams of clorsulon) per 50 kilograms (110 pounds).

(2) *Indications for use*. It is used in cattle for the treatment and control of gastrointestinal nematodes (adults and fourth-stage larvae) (*Haemonchus placei*, *Ostertagia ostertagi* (including inhibited larvae), *O. lyrata*, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia oncophora*, *C. punctata*, *C. pectinata*, *Oesophagostomum radiatum*, *Nematodirus helvetianus* (adults only), *N. spathiger* (adults only), *Bunostomum phlebotomum*; lungworms (adults and fourth-stage larvae) (*Dictyocaulus viviparus*); liver flukes (adults only) (*Fasciola hepatica*); grubs (parasitic stages) (*Hypoderma bovis*, *H. lineatum*); lice (*Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*); mites (*Psoroptes ovis* (syn. *P. communis* var. *bovis*), *Sarcoptes scabiei* var. *bovis*). It is also used to control infections of *D. viviparus* and *O. Ostertagi* for 21 days after treatment, and *H. placei*, *T. axei*, *C. punctata*, *C. oncophora*, and *O. radiatum* for 14 days after treatment.

(3) *Limitations*. For subcutaneous use only. Not for intravenous or intramuscular use. Do not treat cattle within 49 days of slaughter. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. Do not use in other animal species because severe adverse reactions, including fatalities in dogs, may result. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[55 FR 38984, Sept. 24, 1990, as amended at 62 FR 14302, Mar. 26, 1997; 62 FR 63271, Nov. 28, 1997]

§ 522.1204 Kanamycin sulfate injection.

(a) *Specifications*. Each milliliter of kanamycin sulfate injection veterinary contains either 50 or 200 milligrams of kanamycin.

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) It is used in the treatment of bacterial infections

21 CFR Ch. I (4–1–99 Edition)

due to kanamycin sensitive organisms in dogs and cats.

(2) It is administered subcutaneously or intramuscularly at 5 milligrams per pound of body weight per day in equally divided doses at 12-hour intervals.

(3) Its label shall bear an appropriate expiration date.

(4) Restricted to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 53 FR 27851, July 25, 1988; 64 FR 403, Jan. 5, 1999]

EFFECTIVE DATE NOTE: At 64 FR 403, Jan. 5, 1999, § 522.1204 was amended by revising paragraph (a), effective May 20, 1999. For the convenience of the user, the superseded text is set forth as follows:

§ 522.1204 Kanamycin sulfate injection.

(a) *Specifications*. Kanamycin sulfate injection veterinary conforms to the standards of identify, strength, quality, and purity prescribed by § 444.230(a) of this chapter, except that each milliliter contains either 50 or 200 milligrams of kanamycin.

* * * * *

§ 522.1222 Ketamine hydrochloride injectable dosage forms.

§ 522.1222a Ketamine hydrochloride injection.

(a) [Reserved]

(b) *Specifications*. The drug is a sterile aqueous solution and each milliliter contains: Ketamine hydrochloride equivalent to 100 milligrams ketamine base activity and 1:10,000 benzethonium chloride.

(c) *Sponsors*. See Nos. 000010, 000856, 059130, 061690, and 064408 in § 510.600(c) of this chapter.

(d) [Reserved]

(e) *Conditions of use*. (1) In cats:

(i) It is used for restraint or as the sole anesthetic agent in diagnostic or minor, brief surgical procedures that do not require skeletal muscle relaxation.

(ii) It is administered intramuscularly at a recommended dose that ranges from 5 to 15 milligrams per pound of body weight depending on the effect desired.

(2) In subhuman primates:

(i) It is used for restraint.

(ii) It is administered intramuscularly at a recommended

Food and Drug Administration, HHS

§ 522.1244

dose that ranges from 3 to 15 milligrams per kilogram of body weight depending upon the species, general condition, and age of the subject.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 59342, Dec. 23, 1975, as amended at 53 FR 27851, July 25, 1988; 59 FR 41976, Aug. 16, 1994; 59 FR 49291, Sept. 27, 1994; 60 FR 49339, Sept. 25, 1995; 62 FR 22888, Apr. 28, 1997; 62 FR 35076, June 30, 1997; 63 FR 51822, Sept. 29, 1998; 63 FR 65553, Nov. 27, 1998]

§ 522.1222b Ketamine hydrochloride with promazine hydrochloride and aminopentamide hydrogen sulfate injection.

(a) *Chemical name.* Ketamine hydrochloride, (±),-2-(*o*-chlorophenyl)-2-(methylamino) cyclohexanone hydrochloride, with promazine hydrochloride, 10-[3-(dimethylamino) propyl] phenothiazine monohydrochloride, and aminopentamide hydrogen sulfate.

(b) *Specifications.* The drug is a sterile aqueous solution and each milliliter contains: Ketamine hydrochloride equivalent to 100 milligrams ketamine base activity, 7.5 milligrams of promazine hydrochloride, and 0.0625 milligram of aminopentamide hydrogen sulfate, with 1:10,000 benzethonium chloride.

(c) *Sponsor.* See Code No. 000856 in § 510.600(c) of this chapter.

(d) *Special considerations.* Store in a cool place. Protect from light. Do not use if precipitate appears.

(e) *Conditions of use.* (1) It is used in cats as the sole anesthetic agent for ovariohysterectomy and general surgery.

(2) It is administered intramuscularly at a recommended dose from 15 to 20 milligrams ketamine base per pound of body weight, depending on the effect desired.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 59342, Dec. 23, 1975, as amended at 42 FR 3838, Jan. 21, 1977; 53 FR 27851, July 25, 1988]

§ 522.1225 Ketoprofen solution.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 100 milligrams of ketoprofen.

(b) *Sponsor.* See 000856 in 21 CFR 510.600(c) of this chapter.

(c) *Conditions of use in horses*—(1) *Amount.* 1.0 milligram per pound of body weight once daily for up to 5 days.

(2) *Indications for use.* For alleviation of inflammation and pain associated with musculoskeletal disorders in horses.

(3) *Limitations.* For intravenous use only. Do not use in breeding animals. Effects on fertility, pregnancy, or fetal health have not been determined. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[55 FR 40653, Oct. 4, 1990]

§ 522.1228 [Reserved]

§ 522.1244 Levamisole phosphate injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains levamisole phosphate equivalent to 136.5 or 182 milligrams of levamisole hydrochloride (13.65 or 18.2 percent).

(b) *Sponsor.* See No. 000061 in § 510.600 of this chapter for use of 13.65 percent injection, and see No. 043781 for use of 13.65 and 18.2 percent injection.

(c) *Conditions of use*—(1) *Amount.* 2 milliliters per 100 pounds of body weight, subcutaneously in the neck.

(2) *Indications for use.* (i) The 13.65 percent injection is used as an anthelmintic in cattle for treatment of the following parasites: stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*, *Chabertia*), and lungworms (*Dictyocaulus*).

(ii) The 18.2 percent injection is used as an anthelmintic in cattle for treatment of the following parasites: stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*) and lungworms (*Dictyocaulus*).

(3) *Limitations.* Do not administer more than 10 milliliters per site. Cattle that are severely parasitized or maintained under conditions of constant helminth exposure may require retreatment within 2 to 4 weeks after

§ 522.1258

first treatment. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Consult your veterinarian before using in severely debilitated animals or animals under severe stress. Do not administer to cattle within 7 days of slaughter. Do not administer to dairy animals of breeding age.

[43 FR 20489, May 12, 1978, as amended at 43 FR 29289, July 7, 1978; 43 FR 60895, Dec. 29, 1978; 47 FR 10807, Mar. 12, 1982; 62 FR 61625, Nov. 19, 1997]

§ 522.1258 Lidocaine injection with epinephrine.

(a) *Specifications.* Each milliliter of the drug contains 20 milligrams (2 percent) of lidocaine hydrochloride, 0.01 milligram of epinephrine, with sodium chloride, and with methylparaben as a preservative, in water for injection.

(b) *Sponsor.* See No. 000402 in § 510.600(c) of this chapter.

(c) *Conditions of use*¹—(1) *Amount.* The drug is administered by injection as a 2 percent solution or diluted with bacteriostatic water for injection to a 0.5 percent solution for local anesthesia of large and small animals, as follows:

(i) *Cats:* Administer approximately 2 milliliters of 2 percent solution with epinephrine by caudal injection.

(ii) *Cattle:* Administer 5 milliliters of 2 percent solution with epinephrine by epidural injection (standing animal). Administer 10 to 20 milliliters of 2 percent solution with epinephrine by cornual nerve block injection. For teat operations and infiltration, inject 0.5 percent solution with epinephrine to effect.

(iii) *Dogs:* Administer 2 to 10 milliliters of 2 percent solution with epinephrine by caudal injection. Do not give intravascularly. For infiltration, administer 0.5 percent solution with epinephrine to effect.

(iv) *Horses:* Administer 5 to 10 milliliters of 2 percent solution with epinephrine by volar nerve block. Administer 10 to 15 milliliters of 2 percent so-

¹ These conditions are National Academy of Science/National Research Council reviewed and deemed effective for this drug. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

21 CFR Ch. I (4–1–99 Edition)

lution with epinephrine by epidural injection. For standing animal, apply slowly and observe individual sensitivity. For infiltration, administer 0.5 percent solution with epinephrine to effect.

(2) *Limitations.* (i) The drug is contraindicated in the presence of sepsis in the region of proposed injection, shock and heart block, neurologic disease, spinal deformities, septicemia, and hypertension.

(ii) Do not give intravascularly.

(iii) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 46300, Oct. 6, 1978; 43 FR 59059, Dec. 19, 1978, as amended at 52 FR 25212, July 6, 1987]

§ 522.1260 Lincomycin injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains lincomycin hydrochloride equivalent to 25, 50, 100, or 300 milligrams of lincomycin.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Special considerations.* When common labeling for use of the drug in dogs, cats, and swine is included with the drug, all such uses are subject to the labeling requirements of § 201.105 of this chapter.

(d) *Related tolerances.* See § 556.360 of this chapter.

(e) *Conditions of use.* It is used for animals as follows:

(1) *Dogs and cats*—(i) *Amount.* 5 to 10 milligrams per pound of body weight per day.

(ii) *Indications for use.* Infections caused by Gram-positive organisms, particularly streptococci and staphylococci.

(iii) *Limitations.* Administer intramuscularly 10 milligrams per pound of body weight once a day or 5 milligrams per pound of body weight twice daily or intravenously 5 to 10 milligrams per pound of body weight one or two times daily by slow injection. May be diluted with 5 percent glucose in water or normal saline and given as an infusion; as lincomycin hydrochloride monohydrate; for use by or on the order of a licensed veterinarian.

(2) *Swine*—(i) *Amount.* 5 milligrams per pound of body weight per day.

Food and Drug Administration, HHS

§ 522.1335

(ii) *Indications for use.* Treatment of infectious arthritis and mycoplasma pneumonia.

(iii) *Limitations.* Administer intramuscularly as a single daily dose for 3 to 7 days; as lincomycin hydrochloride monohydrate: do not treat within 48 hours of slaughter.

[40 FR 13858, Mar. 27, 1975, as amended at 50 FR 31351, Aug. 2, 1985]

§ 522.1289 Lufenuron suspension.

(a) *Specifications.* Each milliliter of sterile aqueous suspension contains 10 milligrams of lufenuron.

(b) *Sponsor.* See No. 058198 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Cats*—(i) *Amount.* 10 milligrams per kilogram (4.5 milligrams per pound) of body weight every 6 months, subcutaneously.

(ii) *Indications for use.* For use in cats 6 weeks of age and older, for control of flea populations. Lufenuron controls flea populations by preventing the development of flea eggs and does not kill adult fleas. Concurrent use of insecticides may be necessary for adequate control of adult fleas.

(iii) *Limitations.* For subcutaneous use in cats only. The safety of this product in reproducing animals has not been established. Do not use in dogs. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[63 FR 29552, June 1, 1998]

§ 522.1290 Luprostiol sterile solution.

(a) *Specifications.* Each milliliter of sterile solution contains 7.5 milligrams of luprostiol.

(b) *Sponsor.* See No. 057926 in § 510.600(c) of this chapter.

(c) *Special considerations.* Labeling shall bear the following statements: *Warning:* Women of childbearing age, asthmatics, and persons with bronchial and other respiratory problems should exercise extreme caution when handling this product. In the early stages, women may be unaware of their pregnancies. Luprostiol is readily absorbed through the skin and can cause abortion and/or bronchospasms. Direct

contact with the skin should therefore be avoided. Accidental spillage on the skin should be washed off immediately with soap and water.

(d) *Conditions of use*—(1) *Amount.* 7.5 milligrams per mare.

(2) *Indications for use.* The drug is used in mares for estrus control and termination of pregnancy.

(3) *Limitations.* Administer by intramuscular injection only. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[55 FR 1185, Jan. 12, 1990, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995; 61 FR 66582, Dec. 18, 1996]

§ 522.1335 Medetomidine hydrochloride injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 1.0 milligram of medetomidine hydrochloride.

(b) *Sponsor.* See 052483 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 750 micrograms intravenously (IV) or 1,000 micrograms intramuscularly per square meter of body surface. The IV route is more efficacious for dental care.

(2) *Indications for use.* As a sedative and analgesic in dogs over 12 weeks of age to facilitate clinical examinations, clinical procedures, minor surgical procedures not requiring muscle relaxation, and minor dental procedures not requiring intubation. The intravenous route of administration is more efficacious for dental care.

(3) *Limitations.* Do not use in dogs with cardiac disease, respiratory disorders, liver or kidney diseases, dogs in shock, dogs which are severely debilitated, or dogs which are stressed due to extreme heat, cold, or fatigue. Allow agitated dogs to rest quietly before administration. Do not repeat dosing in dogs not responding satisfactorily to treatment. Do not use in breeding or pregnant animals. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[61 FR 21075, May 9, 1996]

§ 522.1350

§ 522.1350 Melatonin implant.

(a) *Specifications.* The drug is a silicone rubber elastomer implant containing 2.7 milligrams of melatonin.

(b) *Sponsor.* See No. 053923 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* One implant per mink.

(2) *Indications for use.* For use in healthy male and female kit and adult female mink (*Mustela vison*) to accelerate the fur priming cycle.

(3) *Limitations.* For subcutaneous implantation in mink only. Do not implant potential breeding stock. Do not use in food-producing animals.

[59 FR 37422, July 22, 1994]

§ 522.1362 Melarsomine dihydrochloride for injection.

(a) *Specifications.* The drug consists of a vial of lyophilized powder containing 50 milligrams of melarsomine dihydrochloride which is reconstituted with the provided 2 milliliters of sterile water for injection.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* For asymptomatic to moderate (class 1 to class 2) heartworm disease: 2.5 milligrams per kilogram of body weight (1.1 milligram per pound) twice, 24 hours apart. The series can be repeated in 4 months depending on the response to the first treatment and the condition, age, and use of the dog. For severe (class 3) heartworm disease: Single injection of 2.5 milligrams per kilogram followed, approximately 1 month later, by 2.5 milligrams per kilogram administered twice, 24 hours apart.

(2) *Indications.* Treatment of stabilized, class 1, 2, and 3 heartworm disease (asymptomatic to mild, moderate, and severe, respectively) caused by immature (4 month-old, stage L₅) to mature adult infections of *Dirofilaria immitis* in dogs.

(3) *Limitations.* Administer only by deep intramuscular injection in the lumbar muscles (L₃-L₅). Use a 23 gauge 1 inch needle for dogs less than or equal to 10 kilograms (22 pounds) and a 22 gauge 1 1/2 inch needle for dogs greater than 10 kilograms (22 pounds). Use alternate sides with each administration. The drug is contraindicated in

21 CFR Ch. I (4-1-99 Edition)

dogs with class 4 (very severe) heartworm disease (Caval Syndrome). Not for use in breeding animals and lactating or pregnant bitches. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[60 FR 49340, Sept. 25, 1995]

§ 522.1372 Mepivacaine hydrochloride injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 20 milligrams of mepivacaine hydrochloride.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is intended for use in horses as a local anesthetic for infiltration, nerve block, intra-articular and epidural anesthesia and topical and/or infiltration anesthesia of the laryngeal mucosa prior to ventriculectomy.

(2) It is administered as follows: for nerve block, 3 to 15 milliliters; for epidural anesthesia, 5 to 20 milliliters; for intra-articular anesthesia, 10 to 15 milliliters; for infiltration, as required; for anesthesia of the laryngeal mucosa prior to ventriculectomy, by topical spray, 25 to 40 milliliters, by infiltration, 20 to 50 milliliters.

(3) Not for use in horses intended for food.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[42 FR 5349, Jan. 28, 1977, as amended at 42 FR 36995, July 19, 1977; 55 FR 23076, June 6, 1990]

§ 522.1380 Methocarbamol injection.

(a) *Specifications.* The product is a sterile, pyrogen-free solution, each milliliter containing 100 milligrams of methocarbamol, 0.5 milliliter of polyethylene glycol 300, and water for injection q.s. Its pH is 3.5 to 6.0.

(b) *Sponsor.* See 000031 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*—(i) *Dogs and cats.* 20 milligrams per pound of body weight for moderate conditions, 25 to 100 milligrams per pound of body weight for severe conditions (tetanus and strychnine poisoning), total cumulative dose not to exceed 150 milligrams per pound of body weight.

(ii) *Horses.* 2 to 10 milligrams per pound of body weight for moderate conditions, 10 to 25 milligrams per pound of body weight for severe conditions (tetanus), additional amounts may be needed to relieve residual effects and to prevent recurrence of symptoms.

(2) *Indications for use.* As an adjunct for treating acute inflammatory and traumatic conditions of the skeletal muscles and to reduce muscular spasms.

(3) *Limitations.* For intravenous use only. For dogs, administer rapidly half the estimated dose, pause until the animal starts to relax, then continue administration to effect. For horses, administer rapidly to effect. Not for horses intended for food use. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 79758, Dec. 2, 1980, as amended at 46 FR 18964, Mar. 27, 1981]

§ 522.1410 Sterile methylprednisolone acetate suspension.

(a) *Specifications.* Each milliliter of aqueous suspension contains 20 or 40 milligrams of methylprednisolone acetate.¹

(b) *Sponsors.* See Nos. 000009 and 000010 in § 510.600(c) of this chapter.

(c) *Special considerations.* (1) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(2) Systemic therapy with methylprednisolone acetate, as with other corticoids, is contraindicated in animals with arrested tuberculosis, peptic ulcer, and Cushing's syndrome. The presence of active tuberculosis, diabetes mellitus, osteoporosis, renal insufficiency, predisposition to thrombophlebitis, hypertension, or congestive heart failure necessitates

carefully controlled use of corticosteroids.

Intrasynovial, intratendinous, or other injections of corticosteroids for local effect are contraindicated in the presence of acute infectious conditions. Exacerbation of pain, further loss of joint motion, with fever and malaise following injection may indicate that the condition has become septic. Appropriate antibacterial therapy should be instituted immediately.

(d) *Conditions of use—(1) Amount—(i) Intramuscular.* Dosage may be repeated when necessary, as follows: dogs—2 to 40 milligrams (up to 120 milligrams in extremely large breeds or dogs with severe involvement); cats—10 to 20 milligrams; horses—200 milligrams.¹

(ii) *Intrasynovial.* Dosage may be repeated when necessary, as follows: horses—40 to 240 milligrams; dogs—up to 20 milligrams.¹

(2) *Indications for use.* Treatment of inflammation and related disorders in dogs, cats, and horses;¹ treatment of allergic and dermatologic disorders in dogs and cats; and as supportive therapy to antibacterial treatment of severe infections in dogs and cats.

(3) *Limitations.* Not for use in horses intended for food. Not for human use. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 59058, Dec. 19, 1978, as amended at 51 FR 741, Jan. 8, 1986; 53 FR 40728, Oct. 18, 1988; 62 FR 35076, June 30, 1997]

§ 522.1452 Nalorphine hydrochloride injection.

(a) *Specifications.* Each milliliter of aqueous solution contains 5 milligrams of nalorphine hydrochloride.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* One milligram per 5 pounds; intravenously, intramuscularly, or subcutaneously.

(2) *Indications for use.* Respiratory and circulatory depression in dogs resulting from overdosage of, or unusual sensitivity to, morphine and certain other narcotics. Not for depression due to any other cause.

(3) *Limitations.* Successive doses of the drug gradually lose their analeptic effect and eventually induce respiratory depression equal to that of

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

§ 522.1462

opiates. Therefore, do not exceed therapeutic dosage. Do not mix drug with meperidine solutions because the buffer will cause precipitation. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 6707, Feb. 2, 1979, as amended at 47 FR 36418, Aug. 20, 1982; 62 FR 63271, Nov. 28, 1997]

§ 522.1462 Naloxone hydrochloride injection.

(a) *Specifications.* Naloxone hydrochloride injection is an aqueous sterile solution containing 0.4 milligram of naloxone hydrochloride per milliliter.

(b) *Sponsor.* See No. 060951 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used as a narcotic antagonist in dogs.

(2) It is administered by intravenous, intramuscular, or subcutaneous injection at an initial dose of 0.04 milligram per kilogram of body weight. When given intravenously, the dosage may be repeated at 2- to 3-minute intervals as necessary. Onset of action by intramuscular or subcutaneous injection is slightly longer than it is by intravenous injection, and repeated dosages must be administered accordingly.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 47 FR 20757, May 14, 1982; 54 FR 32632, Aug. 9, 1989; 63 FR 7701, Feb. 17, 1998]

§ 522.1465 Naltrexone hydrochloride injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 50 milligrams of naltrexone hydrochloride.

(b) *Sponsor.* See 053923 in § 510.600(c) of this chapter.

(c) *Conditions of use in elk and moose—*

(1) *Amount.* 100 milligrams of naltrexone hydrochloride for each milligram of carfentanil citrate administered. One-quarter of the dose should be administered intravenously and three-quarters of the dose should be administered subcutaneously.

(2) *Indications for use.* As an antagonist to carfentanil citrate immobilization in free-ranging or confined elk and moose (*Cervidae*).

(3) *Limitations.* Available data are inadequate to recommend use in pregnant animals. Avoid using during

21 CFR Ch. I (4–1–99 Edition)

breeding season. Do not use in domestic food-producing animals. Do not use in free-ranging animals for 45 days before or during hunting season. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[62 FR 5320, Feb. 5, 1997]

§ 522.1468 Naproxen for injection.

(a) *Specifications.* The drug is a lyophilized powder which is reconstituted with sterile water for injection to form a 10 percent sterile aqueous solution (100 milligrams per milliliter).

(b) *Sponsor.* See 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use in horses.* (1) *Dosage.* Five milligrams per kilogram of body weight intravenously followed by maintenance oral therapy of 10 milligrams per kilogram of body weight twice daily for up to 14 consecutive days.

(2) *Indications for use.* For the relief of inflammation and associated pain and lameness exhibited with arthritis, as well as myositis and other soft tissue diseases of the musculoskeletal system of the horse.

(3) *Limitations.* Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 26763, May 15, 1981. Redesignated and amended at 51 FR 24525, July 7, 1986; 61 FR 5507, Feb. 13, 1996]

§ 522.1484 Neomycin sulfate sterile solution.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 50 milligrams of neomycin sulfate (equivalent to 35 milligrams of neomycin base).¹

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use—*(1) *Amount.* 5 milligrams per pound of body weight daily divided into portions administered every 6 to 8 hours for 3 to 5 days.¹

(2) *Indications for use.* Administer to dogs and cats for the treatment of acute and chronic bacterial infections

¹These claims are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter.

Food and Drug Administration, HHS

§ 522.1620

due to organisms susceptible to neomycin.¹

(3) *Limitations.* For intramuscular or intravenous use only. Neomycin is not for use parenterally in food-producing animals because of prolonged residues in edible tissues. Labeling shall bear an appropriate expiration date. For use by or on the order of a licensed veterinarian.¹

[43 FR 48996, Oct. 20, 1978, as amended at 64 FR 403, Jan. 5, 1999]

EFFECTIVE DATE NOTE: At 64 FR 403, Jan. 5, 1999, § 522.1484 was amended by removing the second sentence of paragraph (a), effective May 20, 1999. For the convenience of the user, the superseded text is set forth as follows:

§ 522.1484 Neomycin sulfate sterile solution.

(a) * * * The neomycin sulfate used in manufacturing the drug conforms to the standards of identity, strength, quality, and purity prescribed by § 444.42a(a)(1) of this chapter.

* * * * *

§ 522.1503 Neostigmine methylsulfate injection.

(a) *Specifications.* Neostigmine methylsulfate injection contains two milligrams of neostigmine methylsulfate in each milliliter of sterile aqueous solution.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is intended for use for treating rumen atony; initiating peristalsis which causes evacuation of the bowel; emptying the urinary bladder; and stimulating skeletal muscle contractions. It is a curare antagonist.

(2) It is administered to cattle and horses at a dosage level of 1 milligram per 100 pounds of body weight subcutaneously. It is administered to sheep at a dosage level of 1 to 1½ milligrams per 100 pounds body weight subcutaneously. It is administered to swine at a dosage level of 2 to 3 milligrams per 100 pounds body weight intramuscularly. These doses may be repeated as indicated.

(3) The drug is contraindicated in mechanical, intestinal or urinary obstruction, late pregnancy, and in animals treated with other cholinesterase inhibitors.

(4) Not for use in animals producing milk, since this use will result in contamination of the milk.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 62 FR 61625, Nov. 19, 1997]

§ 522.1610 Oleate sodium solution.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 50 milligrams of sodium oleate.

(b) *Sponsor.* See No. 037990 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) It is used in horses to stimulate infiltration of cellular blood components that subsequently differentiate into fibrous and/or fibrocartilagenous tissue.

(2) The drug is administered by parenteral injection dependent upon the area of response desired. An injection of 1 milliliter will produce a response of approximately 15 square centimeters. Do not inject more than 2 milliliters per injection site. Regardless of the number of injection sites, the total volume used should not exceed 10 milliliters.

(3) Not for use in horses intended for food.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 27034, July 1, 1976, as amended at 50 FR 40966, Oct. 8, 1985]

§ 522.1620 Orgotein for injection.

(a) *Specifications.* Orgotein for injection is packaged in a vial containing 5 milligrams of orgotein and 10 milligrams of sucrose as lyophilized sterile nonpyrogenic powder with directions for dissolving the contents of the vial in 2 milliliters of diluent which is sodium chloride injection, U.S.P.

(b) *Sponsor.* See No. 024991 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Horses.* (i) It is used in the treatment of soft tissue inflammation associated with the musculoskeletal system.

(ii) It is administered by deep intramuscular injection at a dosage level of 5 milligrams every other day for 2 weeks and twice weekly for 2 to 3 more weeks. Severe cases, both acute

§ 522.1642

and chronic, may benefit more from daily therapy initially. Dosage may be continued beyond 5 weeks if satisfactory improvement has not been achieved.

(iii) Not for use in horses intended for food.

(2) *Dogs.* (i) It is used for the relief of inflammation associated with ankylosing spondylitis, spondylosis, and disc disease. When severe nerve damage is present, response will occur much more slowly, if at all.

(ii) It is administered by subcutaneous injection at a dosage level of 5 milligrams every day for 6 days, and thereafter, every other day for 8 days. In less severe conditions, shorter courses of therapy may be indicated.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 41 FR 32583, Aug. 4, 1976]

§ 522.1642 Oxymorphone hydrochloride injection.

(a) *Specifications.* The drug contains 1 or 1.5 milligrams of oxymorphone hydrochloride per milliliter of aqueous solution containing 0.8 percent sodium chloride.

(b) *Sponsor.* See No. 060951 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is a narcotic analgesic, preanesthetic, anesthetic, and substitute anesthetic adjuvant for intramuscular, subcutaneous or intravenous administration to cats and dogs as follows:

Animal	Body weight (pounds)	Dosage (milligram)
Dogs	2 to 5	0.75
	5 to 15	0.75–1.5
	15 to 30	1.5–2.5
	30 to 60	2.5–4.0
	Over 60	4.0
Cats	Small	0.4–0.75
	Large	0.75–1.5

(2) Do not mix with a barbiturate in the same syringe to preclude precipitation.

(3) It tends to depress respiration. Naloxone hydrochloride and other narcotic antagonists are used to counter over-dosing.

21 CFR Ch. I (4–1–99 Edition)

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 63 FR 7701, Feb. 17, 1998]

§ 522.1660 Oxytetracycline injection.

(a) *Specifications.* Each milliliter of sterile solution contains 200 milligrams of oxytetracycline base.

(b) *Sponsor.* See 000010, 000069, 053389, 059130, and 061623 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.500 of this chapter.

(d) *Conditions of use—*(1) *Beef cattle, dairy cattle, and calves including preruminating (veal) calves.*—(i) *Amount.* 3 to 5 milligrams of oxytetracycline per pound of body weight per day; 5 milligrams per pound of body weight per day for treatment of anaplasmosis, severe foot-rot, and advanced cases of other indicated diseases; 9 milligrams per pound of body weight as a single dosage where retreatment for anaplasmosis is impractical; 9 milligrams per pound of body weight as single dosage where retreatment of calves and yearlings for bacterial pneumonia is impractical; 9 milligrams per pound of body weight as a single dosage for treatment of infectious bovine keratoconjunctivitis.

(ii) *Indications for use.* Treatment of diseases due to oxytetracycline-susceptible organisms as follows: Pneumonia and shipping fever complex associated with *Pasteurella* spp. and *Haemophilus* spp., foot-rot and diphtheria caused by *Fusobacterium necrophorum*, bacterial enteritis (scours) caused by *Escherichia coli*, wooden tongue caused by *Actinobacillus lignieresii*, leptospirosis caused by *Leptospira pomona*, wound infections and acute metritis caused by *Staphylococcus* spp. and *Streptococcus* spp., and infectious bovine keratoconjunctivitis (pinkeye) caused by *Moraxella bovis*. If labeled for use by or on the order of a licensed veterinarian, it may also be used for treatment of anaplasmosis caused by *Anaplasma marginale* and anthrax caused by *Bacillus anthracis*.

(iii) *Limitations.* Administer intramuscularly or intravenously at the 3 to 5 milligrams level, intramuscularly at the 9 milligrams

Food and Drug Administration, HHS

§ 522.1662a

level. Sponsor 000010, may also administer subcutaneously at the 3 to 5 milligrams and 9 milligrams levels. Treatment of all diseases should be instituted early and continued for 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 4 consecutive days. Consult your veterinarian if no improvement is noted within 48 hours. Do not inject more than 10 milliliters per site in adult cattle, reducing the volume according to age and body size to 1 to 2 milliliters in small calves. Exceeding the highest recommended dose, administering at recommended levels for more than 4 consecutive days, and/or exceeding 10 milliliters intramuscularly per injection site may result in antibiotic residues beyond the withdrawal time. Discontinue treatment at least 28 days prior to slaughter. For sponsors 000010, 053389, 059130, and 061623: Not for use in lactating dairy cattle. For sponsor 000069: Milk taken from animals during treatment and for 96 hours after the last treatment must not be used for food; use subcutaneously with a maximum of 10 milliliters per injection site in adult cattle as well as intramuscularly and intravenously.

(2) *Swine*—(i) *Amount*. 3 to 5 milligrams of oxytetracycline per pound of body weight per day; 9 milligrams per pound of body weight as a single dosage where re-treatment for pneumonia is impractical. Sows: Administer once 3 milligrams of oxytetracycline per pound of body weight, approximately 8 hours before farrowing or immediately after completion of farrowing.

(ii) *Indications for use*. Treatment of bacterial enteritis (scours, colibacillosis) caused by *Escherichia coli*, pneumonia caused by *Pasteurella multocida*, and leptospirosis caused by *Leptospira pomona*. Sows: as an aid in control of infectious enteritis (baby pig scours, colibacillosis) in suckling pigs caused by *Escherichia coli*.

(iii) *Limitations*. Administer intramuscularly. Do not inject more than 5 milliliters per site in adult swine. Discontinue treatment at least 42 days prior to slaughter when provided by 000010 and 28 days prior to

slaughter when provided by 000069, 053389, 059130, or 061623.

[45 FR 16479, Mar. 14, 1980, as amended at 46 FR 20160, Apr. 3, 1981; 46 FR 27913, May 22, 1981; 52 FR 19502, May 26, 1987; 60 FR 14218, Mar. 16, 1995; 60 FR 29755, June 6, 1995; 61 FR 31028, June 19, 1996; 61 FR 36291, July 10, 1996; 62 FR 13825, Mar. 24, 1997; 62 FR 27692, May 21, 1997; 63 FR 52158, Sept. 30, 1998]

§ 522.1662 Oxytetracycline hydrochloride implantation or injectable dosage forms.

§ 522.1662a Oxytetracycline hydrochloride injection.

(a)(1) *Specifications*. The drug contains 50 milligrams of oxytetracycline hydrochloride in each milliliter of sterile solution.

(2) *Sponsor*. See No. 000010 in § 510.600(c) of this chapter.

(3) *Conditions of use*. (i) The drug is intended for use in beef cattle, beef calves, nonlactating dairy cattle, and dairy calves for treatment of disease conditions caused by one or more of the following oxytetracycline sensitive pathogens listed as follows: pneumonia and shipping fever complex (*Pasteurella* spp.; *Hemophilis* spp.; *Klebsiella* spp.), bacterial enteritis (scours) (*E. coli*), foot-rot (*Spherophorus necrophorus*), diphtheria (*Spherophorus necrophorus*), wooden tongue (*Actinobacillus lignieresii*), leptospirosis (*Leptospira pomona*), and wound infections; acute metritis; traumatic injury (caused by a variety of bacterial organisms (such as streptococcal and staphylococcal organisms).)

(ii) It is administered by intramuscular injection of 3 to 5 milligrams of oxytetracycline hydrochloride per pound of body weight per day. Leptospirosis, severe foot-rot and severe forms of the indicated diseases should be treated with 5 milligrams per pound of body weight per day. Treatment should be continued for 24 to 48 hours following remission of disease symptoms; however, not to exceed a total of 4 consecutive days. Only 2 milliliters of the drug should be injected per site in case of calves weighing 100 pounds or less and not more than 10 milliliters should be injected per site in adult cattle.

(iii) Discontinue treatment with the drug at least 20 days prior to slaughter

of the animal. When administered to animals within 30 days of slaughter, muscle discoloration may necessitate trimming of injection site and surrounding tissues.

(iv) For use only in beef cattle, beef calves, nonlactating dairy cattle, and dairy calves.

(b)(1) *Specifications*. Each milliliter of sterile solution contains 50 or 100 milligrams of oxytetracycline (as oxytetracycline hydrochloride).

(2) *Sponsor*. See 000010 in § 510.600(c) of this chapter.

(3) *Conditions of use*—(i) *Beef cattle and nonlactating dairy cattle*—(a) *Amount*. Three to 5 milligrams of oxytetracycline per pound of body weight per day; 5 milligrams per pound of body weight per day for the treatment of anaplasmosis, severe foot-rot, and severe cases of other indicated diseases.

(b) *Indications for use*. Treatment of diseases due to oxytetracycline-susceptible organisms as follows: Pneumonia and shipping fever complex associated with *Pasteurella spp.*, *Hemophilus spp.*, and *Klebsiella spp.*, foot-rot and diphtheria caused by *Spherophorus necrophorus*, bacterial enteritis (scours) caused by *Escherichia coli*, wooden tongue caused by *Actinobacillus lignieresii*, leptospirosis caused by *Leptospira pomona*, and wound infections and acute metritis caused by *Staphylococcus spp.* and *Streptococcus spp.* If labeled for use by or on the order of a licensed veterinarian, it may be used for the treatment of anaplasmosis caused by *Anaplasma marginale*.

(c) *Limitations*. For 50-milligram-per-milliliter solution, administer intramuscularly or intravenously; for 100-milligram-per-milliliter solution, administer intramuscularly only. Treatment of all diseases should be instituted early and continue for 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 4 consecutive days. Consult your veterinarian if no improvement is noted within 48 hours. Do not inject more than 10 milliliters per site in adult cattle, reducing the volume according to age and body size to 0.5 to 2 milliliters in small calves. Exceeding the highest recommended dose of 5 milligrams per pound of body weight, administering at

recommended levels for more than 4 consecutive days, and/or exceeding 10 milliliters intramuscularly per injection site may result in antibiotic residues beyond the withdrawal time. Discontinue treatment at least 18 days prior to slaughter. Not for use in lactating dairy cattle.

(ii) *Swine*—(a) *Amount*. Three to 5 milligrams of oxytetracycline per pound of body weight per day. Sows: 3 milligrams of oxytetracycline per pound of body weight, approximately 8 hours before farrowing or immediately after completion of farrowing.

(b) *Indications for use*. For treatment of bacterial enteritis (scours, colibacillosis) caused by *Escherichia coli*, pneumonia caused by *Pasteurella multocida*, and leptospirosis caused by *Leptospira pomona*. Sows: as an aid in control of infectious enteritis (baby pig scours, colibacillosis) in suckling pigs caused by *Escherichia coli*.

(c) *Limitations*. Administer intramuscularly. Do not inject more than 5 milliliters per site. Do not use for more than 4 consecutive days. Discontinue treatment at least 26 days before slaughter.

(c)(1) *Specifications*. The drug contains 50 or 100 milligrams of oxytetracycline hydrochloride in each milliliter of sterile solution.

(2) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(3) *Conditions of use*. (i) The drug is intended for use in the treatment of disease due to oxytetracycline-susceptible organisms in beef cattle and nonlactating dairy cattle. It is indicated in the treatment of pneumonia and shipping fever complex associated with *Pasteurella spp.*, *Hemophilus spp.*, *Klebsiella spp.*, foot-rot and diphtheria caused by *Spherophorus necrophorus*, bacterial enteritis (scours) caused by *Escherichia coli*, wooden tongue caused by *Actinobacillus lignieresii*, acute metritis, and wound infections caused by staphylococcal and streptococcal organisms.

(ii) It is administered to cattle at a dosage level of 3 to 5 milligrams per pound of body weight per day. It may be administered intramuscularly or intravenously from a 50 milligram per milliliter solution. It is administered intravenously from a 100 milligram per

milliliter solution. Severe foot-rot and the severe forms of the indicated diseases should be treated with 5 milligrams per pound of body weight. Treatment should be continued 24 to 48 hours following remission of disease symptoms, however, not to exceed a total of 4 consecutive days. If no improvement is noted within 24 hours, consult a veterinarian. When injecting the drug intramuscularly, do not inject more than 10 milliliters per site in adult cattle. Reduce the amount injected at each site according to the size of the animal. For very small calves do not use more than 2 milliliters per injection site.

(iii) Not for use in lactating dairy cattle. Discontinue treatment at least 19 days prior to slaughter. When administered intramuscularly within 30 days of slaughter, muscle discoloration may necessitate trimming of the injection site and surrounding tissues.

(d)(1) *Specifications.* The drug contains 50 milligrams of oxytetracycline hydrochloride in each milliliter of sterile solution.

(2) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) In beef cattle and nonlactating dairy cattle as follows:

(a) It is used for the treatment of pneumonia and shipping fever complex associated with *Pasteurella spp.* and *Hemophilus spp.*; foot-rot and diphtheria caused by *Spherophorus necrophorus*; bacterial enteritis (scours) caused by *Escherichia coli*; wooden tongue caused by *Actinobacillus lignieresii*; leptospirosis caused by *Leptospira pomona*; wound infections and acute metritis caused by staphylococcal and streptococcal organisms.

(b) Administer by intravenous or intramuscular injection at 3 to 5 milligrams of oxytetracycline per pound of body weight per day. In the treatment of severe foot-rot and severe forms of the indicated diseases, a dosage level of 5 milligrams per pound of body weight per day is recommended.

(c) If the labeling of the drug bears the statement "Federal law restricts this drug to use by or on the order of a licensed veterinarian," it may include additional directions for use in beef cattle and nonlactating dairy cattle for

the treatment of anaplasmosis caused by *Anaplasma marginale*, and anthrax caused by *Bacillus anthracis* in which case the drug is given at 3 to 5 milligrams of oxytetracycline per pound of body weight per day for anthrax, and at 5 milligrams per pound of body weight per day for anaplasmosis.

(ii) In swine as follows:

(a) It is used for the treatment of bacterial enteritis (scours, colibacillosis) caused by *Escherichia coli*; pneumonia caused by *Pasteurella multocida*; and leptospirosis caused by *Leptospira pomona*. Administered to sows as an aid in the control of infectious enteritis (baby pig scours, colibacillosis) in suckling pigs caused by *Escherichia coli*.

(b) Administer by intramuscular injection at 3 to 5 milligrams of oxytetracycline per pound of body weight per day to swine. Administered to sows at 3 milligrams of oxytetracycline per pound of body weight approximately 8 hours before farrowing or immediately after farrowing.

(iii) In poultry (broilers, turkeys, and breeding chickens) as follows:

(a) It is used for the treatment of air sacculitis (air-sac disease, chronic respiratory disease) caused by *Mycoplasma gallisepticum* and *Escherichia coli*; fowl cholera caused by *Pasteurella multocida*; infectious sinusitis caused by *Mycoplasma gallisepticum*; and infectious synovitis caused by *Mycoplasma synoviae*.

(b) Administered subcutaneously to chickens 1 day to 2 weeks of age at 6.25 milligrams of oxytetracycline per bird per day diluted with 1 part of the drug to 3 parts of sterile water; to chickens 2 to 4 weeks of age using the same diluted product at 12.5 milligrams of oxytetracycline per bird; to chickens 4 to 8 weeks of age without dilution at 25 milligrams of oxytetracycline per bird; to chickens 8 weeks of age (broilers and light pullets) at 50 milligrams of oxytetracycline per bird; to adult chickens at 100 milligrams of oxytetracycline per bird.

(c) Administered subcutaneously to turkeys 1 day to 2 weeks of age and 2 to 4 weeks of age at the same dosage as chickens; to turkeys 4 to 6 weeks of age at 50 milligrams of oxytetracycline as

the undiluted product per bird; to turkeys 6 to 9 weeks of age at 100 milligrams of oxytetracycline per bird; to turkeys 9 to 12 weeks of age at 150 milligrams of oxytetracycline per bird; to turkeys 12 weeks of age and older at 200 milligrams of oxytetracycline per bird. In light turkey breeds, no more than 25 milligrams per pound of body weight is administered. For the treatment of infectious sinusitis in turkeys, $\frac{1}{4}$ to $\frac{1}{2}$ milliliter of the drug is injected directly into each swollen sinus depending upon the age of the bird and the severity of the condition. At the time that the sinuses are treated, the drug should also be administered subcutaneously to the birds according to the dosage schedule given in paragraph (d)(3)(iii)(c) of this section. If refilling of the sinuses occurs, the treatment may be repeated in 5 to 7 days.

(iv) Treatment of all diseases should be instituted early. Treatment should continue for 24 to 48 hours beyond the remission of disease symptoms, but not exceed a total of 4 consecutive days. If no improvement is noted within 24 to 48 hours, diagnosis and therapy should be reevaluated.

(v) When injecting intramuscularly in adult livestock, do not inject more than 10 milliliters at any one site. The volume administered per injection site should be reduced according to age and body size so that 1 or 2 milliliters are injected in smaller animals such as small calves and young pigs. Intravenous administration is recommended in cattle when daily dosage exceeds 50 milliliters.

(vi) Treatment must be discontinued at least 5 days prior to slaughter for chickens and turkeys and at least 22 days prior to slaughter for cattle and swine. When administered intramuscularly to animals within 30 days of slaughter, muscle discoloration may necessitate trimming of the injection site(s) and surrounding tissues during the dressing procedure.

(vii) Not for use in lactating dairy animals. Do not administer to laying hens unless the eggs are used for hatching only.

(e)(1) *Specifications.* Each milliliter of sterile solution contains 100 milligrams of oxytetracycline hydrochloride.

(2) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(3) *Conditions of use—(i) Beef cattle and nonlactating dairy cattle—(a) Amount.* 3 to 5 milligrams of oxytetracycline per pound of body weight per day; 5 milligrams per pound of body weight per day for treatment of anaplasmosis, severe foot-rot, and severe cases of other indicated diseases.

(b) *Indications for use.* Treatment of diseases due to oxytetracycline-susceptible organisms as follows: Pneumonia and shipping fever complex associated with *Pasteurella* spp. and *Hemophilus* spp., foot-rot and diphtheria caused by *Fusobacterium necrophorum*, bacterial enteritis (scours) caused by *Escherichia coli*, wooden tongue caused by *Actinobacillus lignieresii*, leptospirosis caused by *Leptospira pomona*, and wound infections and acute metritis caused by *Staphylococcus* spp. and *Streptococcus* spp. If labeled for use by or on the order of a licensed veterinarian, it may be used for the treatment of anaplasmosis caused by *Anaplasma marginale* and anthrax caused by *Bacillus anthracis*.

(c) *Limitations.* Administer intramuscularly. Treatment of all diseases should be instituted early and continue for 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 4 consecutive days. Consult your veterinarian if no improvement is noted within 48 hours. Do not inject more than 10 milliliters per site in adult cattle, reducing the volume according to age and body size to 1 to 2 milliliters in small calves. Exceeding the highest recommended dose of 5 milligrams per pound of body weight, administering at recommended levels for more than 4 consecutive days, and/or exceeding 10 milliliters intramuscularly per injection site may result in antibiotic residues beyond the withdrawal time. Discontinue treatment at least 15 days prior to slaughter. Not for use in lactating dairy cattle.

(ii) *Swine—(a) Amount.* 3 to 5 milligrams of oxytetracycline per pound of body weight per day. Sows: 3 milligrams of oxytetracycline per pound of body weight, administered once, approximately 8 hours before farrowing

or immediately after completion of farrowing.

(b) *Indications for use.* For treatment of bacterial enteritis (scours, colibacillosis) caused by *Escherichia coli*, pneumonia caused by *Pasteurella multocida*, and leptospirosis caused by *Leptospira pomona*. Sows: as an aid in control of infectious enteritis (baby pig scours, colibacillosis) in suckling pigs caused by *Escherichia coli*.

(c) *Limitations.* Administer intramuscularly. Do not inject more than 5 milliliters per site in adult swine, reducing the volume according to age and body size to 1 to 2 milliliters in young pigs. Discontinue treatment at least 22 days prior to slaughter.

(f) [Reserved]

(g)(1) *Specifications.* Each milliliter of sterile solution contains 100 milligrams of oxytetracycline as oxytetracycline hydrochloride.

(2) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(3) *Conditions of use.* The drug is used for the treatment of diseases due to oxytetracycline-susceptible organisms as follows:

(i) *Beef cattle, beef calves, nonlactating dairy cattle, and dairy calves—(a) Amount.* 3 to 5 milligrams of oxytetracycline per pound of body weight per day.

(b) *Indications for use.* For the treatment of pneumonia and shipping fever complex associated with *Pasteurella spp.*, *Hemophilus spp.*, or *Klebsiella spp.*

(c) *Limitations.* Administer by intramuscular, intravenous, or subcutaneous injection. In severe forms of the indicated diseases, administer 5 milligrams of oxytetracycline per pound of body weight per day. Continue treatment 24 to 48 hours following remission of disease symptoms, not to exceed a total of 4 consecutive days. If no improvement is noted within 48 hours, consult a veterinarian. Do not inject more than 10 milliliters per injection site intramuscularly in adult cattle; no more than 1 milliliter per site in calves weighing 100 pounds or less. Do not slaughter cattle for 13 days after intramuscular or intravenous treatment, or 2 days after subcutaneous treatment. Exceeding the highest recommended dosage or duration of treatment (not more than 4 consecu-

tive days) may result in residues beyond the withdrawal period. A withdrawal period has not been established for use of this product in preruminating calves. Do not use in calves to be processed for veal.

(ii) *Swine—(a) Amount.* 3 to 5 milligrams of oxytetracycline per pound of body weight per day. Sows: Administer once 3 milligrams of oxytetracycline per pound of body weight, approximately 8 hours before farrowing or immediately after completion of farrowing.

(b) *Indications for use.* For treatment of bacterial enteritis (scours, colibacillosis) caused by *Escherichia coli*, pneumonia caused by *Pasteurella multocida*, and leptospirosis caused by *Leptospira pomona*. Sows: As an aid in control of infectious enteritis (baby pig scours, colibacillosis) in suckling pigs caused by *Escherichia coli*.

(c) *Limitations.* Administer intramuscularly. If no improvement is noted within 24 hours, consult a veterinarian. Do not inject more than 5 milliliters per site. Discontinue treatment at least 20 days prior to slaughter.

(h)(1) *Specifications.* Each milliliter of sterile solution contains 50 or 100 milligrams of oxytetracycline hydrochloride.

(2) *Sponsors.* See 000010 in § 510.600(c) of this chapter for use of 50 and 100 milligrams/milliliter solution, and see No. 059130 in § 510.600(c) for use of 100 milligrams/milliliter solution.

(3) *Conditions of use—(i) Amount.* The drug is used in beef cattle, beef calves, nonlactating dairy cattle, and dairy calves as follows: 3 to 5 milligrams of oxytetracycline hydrochloride per pound of body weight per day; 5 milligrams per pound of body weight per day for treatment of severe forms of the indicated diseases.

(ii) *Indications for use.* The drug is used for treatment of bacterial pneumonia and shipping fever complex associated with *Pasteurella spp.*; foot-rot and calf diphtheria caused by *Spherophorus necrophorus*; bacterial enterities (scours) caused by *Escherichia coli*; wooden tongue caused by *Actinobacillus lignieresii*; wound infections, acute metritis, and traumatic injury caused by staphylococcal and streptococcal organisms.

(iii) *Limitations.* Administer 50-milligram-per-milliliter solution intramuscularly; administer 100-milligram-per-milliliter solution intravenously. Continue treatment 24 to 48 hours following remission of disease symptoms, not to exceed a total of 4 consecutive days. If no improvement is noted within 24 to 48 hours, consult a veterinarian for diagnosis and therapy. When injecting the drug intramuscularly, do not inject more than 10 milliliters per site in adult cattle. Reduce the volume administered per injection site according to age and body size. In calves weighing 100 pounds or less, do not inject more than 2 milliliters intramuscularly per site. Discontinue treatment at least 22 days before slaughter. Not for use in lactating dairy animals.

(i)(1) *Specifications.* Each milliliter of sterile solution contains 50 milligrams of oxytetracycline hydrochloride.

(2) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(3) *Conditions of use—(i) Amount.* The drug is used in beef cattle, beef calves, nonlactating dairy cattle, and dairy calves as follows: Administer 3 to 5 milligrams of the oxytetracycline hydrochloride intramuscularly per pound of body weight per day.

(ii) *Indications for use.* The drug is used for treatment of bacterial pneumonia and shipping fever complex associated with *Pasteurella* spp.; foot-rot and diphtheria caused by *Spherophorus necrophorus*; bacterial enteritis (scours) caused by *Escherichia coli*; wooden tongue caused by *Actinobacillus lignieresii*; wound infections and acute metritis caused by staphylococcal and streptococcal organisms susceptible to oxytetracycline.

(iii) *Limitations.* In severe forms of the indicated diseases, administer the equivalent of 5 milligrams of oxytetracycline hydrochloride per pound of body weight per day. Continue treatment 24 to 48 hours following remission of disease symptoms, not to exceed a total of 4 consecutive days. If no improvement is noted within 24 to 48 hours, consult a veterinarian for diagnosis and therapy. In adult livestock, do not inject more than 10 milliliters at any one site. Reduce the volume administered per injection site according

to age and body size. In calves weighing 100 pounds or less inject only 2 milliliters per site. Discontinue treatment at least 18 days before slaughter. Not for use in lactating dairy cattle.

(j) [Reserved]

(k)(1) *Specifications.* Each milliliter of sterile solution contains either 50 or 100 milligrams of oxytetracycline hydrochloride.

(2) *Sponsor.* See No. 000864 in § 510.600(c) of this chapter.

(3) *Conditions of use in beef cattle and nonlactating dairy cattle.* (i) *Amount.* 3 to 5 milligrams per pound of body weight daily, 5 milligrams per pound for anaplasmosis, severe foot rot, and severe forms of other diseases.

(ii) *Indications for use.* Treatment of diseases due to oxytetracycline-susceptible organisms as follows: pneumonia and shipping fever complex associated with *Pasteurella* spp. and *Hemophilus* spp.; foot rot and diphtheria caused by *Fusobacterium necrophorum*; bacterial enteritis (scours) caused by *Escherichia coli*; wooden tongue caused by *Actinobacillus lignieresii*; leptospirosis caused by *Leptospira pomona*; acute metritis and wound infections caused by staphylococcal and streptococcal organisms; if labeled for use by or on the order of a licensed veterinarian, it may be used for treatment of anaplasmosis caused by *Anaplasma marginale* and anthrax caused by *Bacillus anthracis*.

(iii) *Limitations.* Administer by intravenous injection. Treatment should be continued 24 to 48 hours following remission of disease symptoms, but not to exceed a total of 4 consecutive days. If no improvement occurs within 24 to 48 hours, reevaluate diagnosis and therapy. Discontinue use at least 19 days prior to slaughter. Not for use in lactating dairy cattle.

[40 FR 13858, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 522.1662a, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 522.1662b Oxytetracycline hydrochloride with lidocaine injection.

(a) *Specifications.* The drug contains 50 or 100 milligrams of oxytetracycline hydrochloride and 2 percent lidocaine in each milliliter of sterile aqueous solution.

Food and Drug Administration, HHS

§ 522.1696a

(b) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) The drug is indicated for use in the treatment of diseases of dogs caused by pathogens sensitive to oxytetracycline hydrochloride including treatment for the following conditions in dogs caused by susceptible microorganisms: Bacterial infections of the urinary tract caused by *Hemolytic staphylococcus*, *Streptococcus spp.*, Bacterial pulmonary infections caused by *Brucella bronchiseptica*, *Streptococcus pyogenes*, *Staphylococcus aureus*, secondary bacterial infections caused by *Micrococcus pyogenes* var. *albus*, *Brucella bronchiseptica*, *Streptococcus spp.*

(2) The drug is administered intramuscularly at a recommended daily dosage to dogs at 5 milligrams per pound of body weight administered in divided doses at 6 to 12 hour intervals. Therapy should be continued for at least 24 hours after all symptoms have subsided.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 48 FR 30615, July 5, 1983]

§ 522.1680 Oxytocin injection.

(a) *Specifications*. Each milliliter of oxytocin injection contains 20 U.S.P. units of oxytocin.

(b) *Sponsors*. See Nos. 000010, 000856, 000857, 000864, 050604, 058639, and 059130 in § 510.600(c) of this chapter.

(c) *Conditions of use*¹—(1) *Amount*—(i) *Obstetrical*. Administer drug intravenously, intramuscularly, or subcutaneously under aseptic conditions as indicated. The following dosages are recommended and may be repeated as conditions require:

	ml	U.S.P. units
Cats	0.25 to 0.5	5 to 10.
Dogs	0.25 to 1.5	5 to 30.
Ewes, Sows	1.5 to 2.5	30 to 50.
Cows, Horses	5.0	100.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bio-equivalency and safety information.

(ii) *Milk letdown*. Intravenous administration is desirable. The following dosage is recommended and may be repeated as conditions require:

	ml	U.S.P. units
Cows	0.5 to 1.0	10 to 20.
Sows	0.25 to 1.0	5 to 20.

(2) *Indications for use*. Oxytocin may be used as a uterine contractor to precipitate and accelerate normal parturition and postpartum evacuation of uterine debris. In surgery it may be used postoperatively following cesarean section to facilitate involution and resistance to the large inflow of blood. It will contract smooth muscle cells of the mammary gland for milk letdown if the udder is in proper physiological state.

(3) *Limitations*. Do not use in dystocia due to abnormal presentation of fetus until correction is accomplished. For prepartum usage, full relaxation of the cervix should be accomplished either naturally or by administration of estrogen prior to oxytocin therapy. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 63097, Nov. 2, 1979; 45 FR 1019, Jan. 4, 1980, as amended at 52 FR 18691, May 19, 1987; 52 FR 25212, July 6, 1987; 52 FR 36023, Sept. 25, 1987; 53 FR 32610, Aug. 26, 1988; 53 FR 40728, Oct. 18, 1988; 54 FR 41442, Oct. 10, 1989; 55 FR 8462, Mar. 8, 1990; 56 FR 14642, Apr. 11, 1991, 56 FR 16002, Apr. 19, 1991; 59 FR 31139, June 17, 1994; 62 FR 35076, June 30, 1997; 62 FR 38906, July 21, 1997]

§ 522.1696 Penicillin G procaine implantation and injectable dosage forms.

§ 522.1696a Penicillin G benzathine and penicillin G procaine sterile suspension.

(a) *Specifications*. Each milliliter of aqueous suspension contains penicillin G benzathine and penicillin G procaine, each equivalent to 150,000 units of penicillin G.

(b) *Sponsors*. See sponsors in § 510.600(c) of this chapter for the conditions of use in paragraph (d) of this section as follows:

(1) See Nos. 000008, 049185, 000856, 000864, and 010515 for use as in paragraph (d)(1) of this section.

§ 522.1696b

(2) See Nos. 049185 and 000856 for use as in paragraph (d)(2) of this section.

(3) See Nos. 000069, 000864, and 010515 for use as in paragraph (d)(3) of this section.

(c) *Related tolerances.* See § 556.510 of this chapter.

(d) *Conditions of use*—(1) *Horses, dogs, and beef cattle.* Treatment of bacterial infections susceptible to penicillin G. Repeat dosage in 48 hours. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(i) *Horses.* 2 milliliters per 150 pounds of body weight intramuscularly. Do not use in horses intended for food purposes.

(ii) *Dogs.* 1 milliliter per 10 to 25 pounds of body weight intramuscularly or subcutaneously.

(iii) *Beef cattle.* 2 milliliters per 150 pounds of body weight intramuscularly or subcutaneously. Treatment should be limited to two doses. Not to be used in beef cattle within 30 days of slaughter.

(iv) *National Academy of Sciences/National Research Council (NAS/NRC) status.* The conditions of use were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter but may require bioequivalency and safety information.

(2) *Beef cattle.* Treatment of bacterial pneumonia (*Streptococcus* spp., *Corynebacterium pyogenes*, *Staphylococcus aureus*); upper respiratory infections such as rhinitis or pharyngitis (*C. pyogenes*); blackleg (*Clostridium chauvoei*); and prophylaxis of bovine shipping fever in 300 to 500 pound beef calves.

(i) *Amount.* 2 milliliters per 150 pounds of body weight subcutaneously. Repeat dosage in 48 hours. Limit treatment to two doses. Not for use within 30 days of slaughter.

(ii) *NAS/NRC status.* The conditions of use were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter but may require bioequivalency and safety information.

(3) *Beef cattle.* Treatment of bacterial pneumonia (shipping fever) (*Streptococcus* spp., *C. pyogenes*, *S. aureus*); upper respiratory infections

21 CFR Ch. I (4–1–99 Edition)

such as rhinitis or pharyngitis (*C. pyogenes*); and blackleg (*C. chauvoei*).

(i) 2 milliliters per 150 pounds of body weight subcutaneously. Repeat dosage in 48 hours. Limit treatment to two doses. Not for use within 30 days of slaughter.

(ii) [Reserved]

[57 FR 37332, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992, as amended at 58 FR 65285, Dec. 14, 1993; 59 FR 38119, July 27, 1994]

§ 522.1696b Penicillin G procaine aqueous suspension.

(a) *Specifications.* Each milliliter contains penicillin G procaine equivalent to 300,000 units of penicillin G.

(b) *Related tolerances.* See § 556.510 of this chapter.

(c) *National Academy of Sciences/National Research Council (NAS/NRC) status.* The conditions of use were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter for use as in paragraph (d) of this section, see No. 000864 for use as in paragraph (d)(3) of this section.

(1) *Dogs*—(i) *Amount.* 10,000 units per pound of body weight daily at 24-hour intervals.

(ii) *Indications for use.* Treatment of infections caused by penicillin-sensitive organisms.

(iii) *Limitations.* For intramuscular use only. Continue treatment at least 48 hours after symptoms disappear. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Amount.* 10,000 units per pound of body weight daily at 24-hour intervals.

(ii) *Indications for use.* Treatment of infections caused by penicillin-sensitive organisms.

(iii) *Limitations.* For intramuscular use only. Continue treatment at least 48 hours after symptoms disappear. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(3) *Cattle, sheep, swine, and horses*—(i) *Amount.* 3,000 units per pound of body

weight (1 milliliter per 100 pounds body weight) daily.

(ii) *Indications for use.* Treatment of cattle and sheep for bacterial pneumonia (shipping fever) caused by *Pasteurella multocida*; swine for erysipelas caused by *Erysipelothrix insidiosa*; and horses for strangles caused by *Streptococcus equi*.

(iii) *Limitations.* Administer by deep intramuscular injection. Continue treatment at least 48 hours after symptoms disappear but do not exceed 7 days of treatment in nonlactating dairy and beef cattle, sheep, and swine, or 5 days in lactating cattle. Milk that has been taken during treatment and for 48 hours (four milkings) after the last treatment must not be used for food. Discontinue treatment for the following number of days before slaughter: Nonruminating cattle (calves)—7; all other cattle—4; sheep—8; and swine—6. Not for use in horses intended for food.

(e) *Sponsor.* See No. 055529 in § 510.600(c) of this chapter.

(1) *Cattle, sheep, swine, and horses—(i) Amount.* 3,000 units per pound of body weight (1 milliliter per 100 pounds body weight) daily.

(ii) *Indications for use.* Treatment of cattle and sheep for bacterial pneumonia (shipping fever) caused by *Pasteurella multocida*; swine for erysipelas caused by *Erysipelothrix insidiosa*; and horses for strangles caused by *Streptococcus equi*.

(iii) *Limitations.* For intramuscular use only. Continue treatment at least 1 day after symptoms disappear (usually 2 or 3 days). Treatment should not exceed 4 consecutive days. Do not exceed 10 milliliters per injection site. Milk that has been taken during treatment and for 72 hours (six milkings) after the latest treatment must not be used for food. Discontinue treatment for the following number of days before slaughter: Cattle—10, sheep—9, and swine—7. Not for use in horses intended for food.

(2) [Reserved]

(f) *Sponsor.* See Nos. 000069 and 010515 in § 510.600(c) of this chapter. See paragraph (d) of this section for conditions of use, except that milk taken during treatment and for 48 hours (four

milkings) after the latest treatment shall not be used for food.

[57 FR 37332, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992, as amended at 58 FR 11964, Mar. 2, 1993]

§ 522.1696c Penicillin G procaine in oil.

(a) *Specifications.* Each milliliter contains penicillin G procaine equivalent to 300,000 units of penicillin G.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *National Academy of Sciences/National Research Council (NAS/NRC) status.* The conditions of use were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use.* (1) *Amount.* Dogs and cats—10,000 units per pound of body weight once daily. Horses—3,000 units per pound of body weight once daily.

(2) *Indications for use.* Treatment of infections of dogs, cats, and horses caused by penicillin-susceptible organisms such as Streptococci, Staphylococci, and Corynebacteria.

(3) *Limitations.* Not for use in food-producing animals. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37333, Aug. 18, 1992]

§ 522.1698 Pentazocine lactate injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains pentazocine lactate equivalent to 30 milligrams of pentazocine base.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Horses—(i) Amount.* 0.15 milligram of pentazocine base per pound of body weight per day.

(ii) *Indications for use.* For symptomatic relief of pain due to colic.

(iii) *Limitations.* Administer intravenously or intramuscularly. Intravenous injections are given slowly in the jugular vein. In cases of severe pain, a second dose is recommended intramuscularly 10 to 15 minutes after the initial dose at the same level. Not

§ 522.1704

for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Dogs*—(i) *Amount*. 0.75 to 1.50 milligrams of pentazocine base per pound of body weight.

(ii) *Indications for use*. For amelioration of pain accompanying post-operative recovery, fracture, trauma, and spinal disorders.

(iii) *Limitations*. Administer intramuscularly only. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[42 FR 31450, June 21, 1977, as amended at 42 FR 36995, July 19, 1977; 47 FR 5409, Feb. 5, 1982; 55 FR 23076, June 6, 1990]

§ 522.1704 Sodium pentobarbital injection.

(a)(1) *Specifications*. Sodium pentobarbital injection is sterile and contains in each milliliter 64.8 milligrams of sodium pentobarbital.

(2) *Sponsor*. See No. 000061 in § 510.600(c) of this chapter.

(3) *Conditions of use*. (i) The drug is indicated for use as a general anesthetic in dogs and cats. Although it may be used as a general surgical anesthetic for horses, it is usually given at a lower dose to cause sedation and hypnosis and may be supplemented with a local anesthetic. It may also be used in dogs for the symptomatic treatment of strychnine poisoning.

(ii) The drug is administered intravenously “to effect”. For general surgical anesthesia, the usual dose is 11 to 13 milligrams per pound of body weight. For sedation, the usual dose is approximately 2 milligrams per pound of body weight. For relieving convulsive seizures in dogs, when caused by strychnine, the injection should be administered intravenously “to effect”. The drug may be given intraperitoneally if desired. However, the results of such injections are less uniform. When given intraperitoneally, it is administered at the same dosage level as for intravenous administration. The dose must be reduced for animals showing under-nourishment, toxemia, shock and similar conditions.

(iii) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

21 CFR Ch. I (4–1–99 Edition)

(b)(1) *Specifications*. Sodium pentobarbital injection is sterile and contains in each milliliter 65 milligrams of sodium pentobarbital.

(2) *Sponsor*. See 000402 in § 510.600(c) of this chapter.

(3) *Conditions of use*. (i) The drug is indicated for use as a general anesthetic in dogs and cats.

(ii) The drug is administered intravenously “to effect.” For general anesthesia, the usual dose is 13 milligrams per pound of body weight.

(iii) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 45 FR 83483, Dec. 19, 1980; 52 FR 25212, July 6, 1987; 62 FR 61625, Nov. 19, 1997]

§ 522.1720 Phenylbutazone injection.

(a) *Specifications*. The drug contains 100 or 200 milligrams of phenylbutazone in each milliliter of sterile aqueous solution.

(b) *Sponsors*. (1) Approval for use of the 200 milligrams per milliliter drug in dogs and horses: See sponsor Nos. 000031, 000061, 015579, and 059130 in § 510.600(c) of this chapter.

(2) Approval for use of the 200 milligrams per milliliter drug for use in horses: See sponsor Nos. 000010 and 000864 in § 510.600(c) of this chapter.

(3) Approval for use of the 100 milligrams per milliliter drug in dogs and horses: See sponsor No. 000856 in § 510.600(c) of this chapter.

(4) Approval for use of the 200 milligrams per milliliter drug in dogs: See sponsor No. 000864 in § 510.600(c) of this chapter.

(c) *Conditions of use for dogs*. (1) It is used for the relief of inflammatory conditions associated with the musculoskeletal system.

(2) It is administered intravenously at a dosage level of 10 milligrams per pound of body weight daily in 3 divided doses, not to exceed 800 milligrams daily regardless of weight. Limit intravenous administration to 2 successive days. Oral medication may follow.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d) *Conditions of use for horses*. (1) It is used for the relief of inflammatory

Food and Drug Administration, HHS

§ 522.1862

conditions associated with the musculoskeletal system.

(2) It is administered intravenously at a dosage level of 1 to 2 grams per 1,000 pounds of body weight daily in 3 divided doses, not to exceed 4 grams daily. Limit intravenous administration to not more than 5 successive days.

(3) Not for use in animals intended for food.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 522.1720, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 522.1820 Pituitary luteinizing hormone for injection.

(a) *Specifications.* The drug is a lyophilized pituitary extract. Each 6-milliliter vial contains an amount equivalent to 25 milligrams of standard pituitary luteinizing hormone and is reconstituted for use by addition of 5 milliliters of 0.9 percent aqueous sodium chloride solution.

(b) *Sponsor.* No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is an aid in the treatment of breeding disorders related to pituitary hypofunction in cattle, horses, swine, sheep, and dogs.

(2) Preferably given by intravenous injection, it may be administered subcutaneously; dosage is as follows: Cattle and horses, 25 mg; swine, 5 mg; sheep, 2.5 mg, and dogs, 1.0 mg. Treatment may be repeated in 1 to 4 weeks, or as indicated.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 52 FR 7832, Mar. 13, 1987]

§ 522.1850 Polysulfated glycosaminoglycan.

(a) *Specifications.* Each 1-milliliter ampule of sterile aqueous solution contains 250 milligrams of polysulfated glycosaminoglycan; each 5-milliliter ampule or vial contains 500 milligrams.

(b) *Sponsor.* See No. 010797 in § 510.600(c) of this chapter.

(c) *Conditions of use—horses.* (1) *Indications for use.* Polysulfated glycosaminoglycan is for the treatment of noninfectious degenerative and/or traumatic joint dysfunction and associated lameness of the carpal and hock joints in horses.—

(2) *Amount—(i) Intra-articular use (carpal):* 250 milligrams once a week for 5 weeks. The joint area must be shaved, cleaned, and sterilized as in a surgical procedure prior to injection. If the joint reacts with excessive inflammation, after intra-articular treatment, cease therapy.

(ii) *Intramuscular use (carpal and hock):* 500 milligrams every 4 days for 28 days. Injection site must be thoroughly cleansed prior to injection.

(3) *Limitations.* Not for use in horses intended for food. Safe use in breeding animals has not been established. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d) *Conditions of use—dogs—(1) Indications for use.* For control of signs associated with noninfectious degenerative and/or traumatic arthritis of canine synovial joints.

(2) *Dosage.* 2 milligrams per pound of body weight by intramuscular injection.

(3) *Limitations.* Administer intramuscularly twice weekly for up to 4 weeks (maximum of 8 injections). Do not exceed recommended dose or regimen. Do not mix with other drugs or solvents. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[54 FR 53053, Dec. 27, 1989, as amended at 61 FR 54333, Oct. 18, 1996; 62 FR 45158, Aug. 26, 1997]

§ 522.1862 Sterile pralidoxime chloride.

(a) *Chemical name.* 2-Formyl-1-methylpyridinium chloride oxime.

(b) *Specifications.* Sterile pralidoxime chloride is packaged in vials. Each vial contains 1 gram of sterile pralidoxime chloride powder and includes directions for mixing this gram with 20 cubic centimeters of sterile water for injection prior to use.

(c) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) It is used in horses, dogs, and cats as an antidote in

§522.1870

the treatment of poisoning due to those pesticides and chemicals of the organophosphate class which have anticholinesterase activity in horses, dogs, and cats.

(2) It is administered as soon as possible after exposure to the poison. Before administration of the sterile pralidoxime chloride, atropine is administered intravenously at a dosage rate of 0.05 milligram per pound of body weight, followed by administration of an additional 0.15 milligram of atropine per pound of body weight administered intramuscularly. Then the appropriate dosage of sterile pralidoxime chloride is administered slowly intravenously. The dosage rate for sterile pralidoxime chloride when administered to horses is 2 grams per horse. When administered to dogs and cats, it is 25 milligrams per pound of body weight. For small dogs and cats, sterile pralidoxime chloride may be administered either intraperitoneally or intramuscularly. A mild degree of atropinization should be maintained for at least 48 hours. Following severe poisoning, a second dose of sterile pralidoxime chloride may be given after 1 hour if muscle weakness has not been relieved.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 49 FR 32061, Aug. 10, 1984]

§522.1870 Praziquantel injectable solution.

(a) *Specification.* Each milliliter contains 56.8 milligrams of praziquantel.

(b) *Sponsor.* See 000859 in §510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount.* For dogs 5 pounds and under, 0.3 milliliter (17.0 milligrams); for 6 to 10 pounds, 0.5 milliliter (28.4 milligrams); for 11 to 25 pounds, 1.0 milliliter (56.8 milligrams); if over 25 Pounds, 0.2 milliliter (11.4 milligrams) per 5 pounds body weight to a maximum of 3 milliliters (170.4 milligrams).

(ii) *Indications for use.* For removal of canine cestodes *Dipylidium caninum*, *Taenia pisiformis*, and *Echinococcus granulosus*, and removal and control of canine cestode *Echinococcus multilocularis*.

21 CFR Ch. I (4–1–99 Edition)

(iii) *Limitations.* For subcutaneous or intramuscular use; not intended for use in puppies less than 4 weeks of age; Federal law restricts the drug to use by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Amount.* For cats under 5 pounds, 0.2 milliliter (11.4 milligrams); 5 to 10 pounds, 0.4 milliliter (22.7 milligrams); 11 pounds and over, 0.6 milliliter (34.1 milligrams) maximum.

(ii) *Indications for use.* For removal of feline cestodes *Dipylidium caninum* and *Taenia taeniaeformis*.

(iii) *Limitations.* For subcutaneous or intramuscular injection only. Not intended for use in kittens less than 6 weeks of age. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 10464, Feb. 3, 1981, as amended at 47 FR 6617, Feb. 16, 1982; 58 FR 42853, Aug. 12, 1993]

§522.1881 Sterile prednisolone acetate aqueous suspension.

(a) *Specifications.* Each milliliter of sterile aqueous suspension contains 25 milligrams of prednisolone acetate.

(b) *Sponsor.* See No. 000061 in §510.600(c) of this chapter.

(c) *NAS/NRC status.* The conditions of use are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified in §514.111 of this chapter but may require bioequivalency and safety information.

(d) *Conditions of use.* (1) The drug is indicated in the treatment of dogs, cats, and horses for conditions requiring an anti-inflammatory agent. The drug is indicated for the treatment of acute musculoskeletal inflammations such as bursitis, carpalitis, and spondylitis. The drug is indicated as supportive therapy in nonspecific dermatosis such as summer eczema and atopy. The drug may be used as supportive therapy pre- and post-operatively and for various stress conditions when corticosteroids are required while the animal is being treated for a specific condition.

(2) The drug is administered to horses intra-articularly at a dosage level of 50 to 100 milligrams. The dose may be repeated when necessary. If no response

is noted after 3 or 4 days, the possibility must be considered that the condition is unresponsive to prednisolone therapy. The drug is administered to dogs and cats intramuscularly at a dosage level of 10 to 50 milligrams. The dosage may be repeated when necessary. If the condition is of a chronic nature, an oral corticosteroid may be given as a maintenance dosage. The drug may be given intra-articularly to dogs and cats at a dosage level of 5 to 25 milligrams. The dose may be repeated when necessary after 7 days for two or three doses.

(3) The labeling shall comply with the requirements of § 510.410 of this chapter for corticosteroids.

(4) Not for use in horses intended for food.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[52 FR 23032, June 17, 1987]

§ 522.1883 Prednisolone sodium phosphate injection, sterile.

(a)(1) *Specifications.* Each milliliter contains 20 milligrams of prednisolone sodium phosphate (equivalent to 14.88 milligrams of prednisolone) in sterile aqueous solution.

(2) *Sponsor.* See No. 000864 in § 510.600(c) of this chapter.

(3) *Conditions of use*—(i) It is used in treatment of dogs when a rapid adrenal glucocorticoid and/or anti-inflammatory effect is necessary.¹

(ii) It is administered intravenously in a dosage of 2½ to 5 milligrams of prednisolone sodium phosphate per pound of body weight, initially for shock and shock-like states, followed by equal maintenance doses at 1-, 3-, 6-, or 10-hour intervals as determined by the condition of the animal. If permanent use is required, oral therapy (tablets) may be substituted. If therapy is to be withdrawn after prolonged use, reduce daily dose gradually over a number of days.

(iii) Do not use in viral infections. Except in emergency therapy, do not use with tuberculosis, chronic nephritis, Cushing's disease, or peptic ulcers. With infections, use appropriate antibacterial therapy with, and for at least 3 days after, discontinuance of use and disappearance of all signs of infection.¹

(iv) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(v) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

(b) [Reserved]

[43 FR 29769, July 11, 1978, as amended at 49 FR 23834, June 8, 1984]

§ 522.1884 Prednisolone sodium succinate injection.

(a) *Chemical name.* 11 beta, 17, 21-Trihydroxypregna-1, 4-diene-3, 20-dione 21-succinate sodium salt.

(b) *Specifications.* Each milliliter of prednisolone sodium succinate injection contains: Prednisolone sodium succinate equivalent in activity to 10, 20, or 50 milligrams of prednisolone.

(c) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter for products containing 10, 20, and 50 milligrams equivalent prednisolone activity per milliliter for use in horses, dogs, and cats as provided in paragraphs (d)(1), (2) (i), (ii), and (iii) of this section. See No. 000402 in § 510.600(c) of this chapter for a product containing 10 milligrams equivalent prednisolone activity per milliliter for use in horses as provided in paragraph (d)(2)(iv) of this section.

(d) *Conditions of use.* (1) The drug is intended for the treatment of horses, dogs, and cats.¹

(2)(i) The dosage for horses is 50 to 100 milligrams as an initial dose given intravenously over a period of one-half to 1 minute, or intramuscularly, and may be repeated in inflammatory, allergic, or other stress conditions at intervals of 12, 24, or 48 hours, depending upon the size of the animal, the severity of the condition and the response to treatment.¹

¹These conditions are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(ii) In dogs, the drug is administered intravenously at a range of 2.5 to 5 milligrams per pound of body weight as an initial dose followed by maintenance doses at 1, 3, 6, or 10 hour intervals, as determined by the condition of the animal, for treatment of shock.

(iii) In dogs and cats, the drug may be given intramuscularly for treatment of inflammatory, allergic and less severe stress conditions, where immediate effect is not required, at 1 to 5 milligrams ranging upward to 30 to 50 milligrams in large breeds of dogs. Dosage may be repeated in 12 to 24 hours and continued for 3 to 5 days if necessary. If permanent corticosteroid effect is required oral therapy with prednisolone tablets may be substituted.

(iv) In horses, 50 to 100 milligrams as an initial dose given intravenously over a period of ½ to 1 minute. May be repeated in inflammatory, allergic, or other stress conditions at intervals of 12, 24, or 48 hours, depending upon the size of the animal, the severity of the condition and the response to treatment. Not for use in horses intended for food. Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered late in pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13858, Mar. 27, 1975, as amended at 46 FR 13215, Feb. 20, 1981; 46 FR 33513, June 30, 1981; 52 FR 25212, July 6, 1987]

§ 522.1885 Prednisolone tertiary butylacetate suspension.

(a) *Specifications.* Prednisolone tertiary butylacetate (Pregna-1,4-diene-3, 20-dione-11B, 17α 21-triol 21-(3,3, dimethyl butyrate) suspension contains 20 milligrams of prednisolone tertiary butylacetate per milliliter. It is sterile.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used as an anti-inflammatory agent in horses, dogs, and cats.¹

(2) It is administered to horses intramuscularly at a dosage level of 100

to 300 milligrams and intrasynovially at a dosage level of 50 to 100 milligrams. It is administered intramuscularly to dogs and cats at a dosage level of 1 milligram per 5 pounds of body weight and intrasynovially at a dosage level of 10 to 20 milligrams. Intramuscular retreatment of horses in 24 to 48 hours may be necessary, depending on the general condition of the animal and the severity and duration of the disease.¹

(3) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered late in pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.¹

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13858, Mar. 27, 1975, as amended at 62 FR 63271, Nov. 28, 1997]

§ 522.1890 Sterile prednisone suspension.

(a) [Reserved]

(b)(1) *Specifications.* Each milliliter of sterile aqueous suspension contains 10 to 40 milligrams of prednisone.

(2) *Sponsor.* See 000061 in § 510.600(c) of this chapter.

(3) *Conditions of use—(i) Amount.* Administer intramuscularly as follows:

(a) *Horses.* 100 to 400 milligrams, repeating if necessary. If no response is observed after 3 to 4 days of therapy, reevaluate diagnosis.¹

(b) *Dogs and cats.* 0.25 to 1.0 milligram per pound of body weight for 3 to 5 days or until a response is noted. Treatment may be continued with an orally administered dose.¹

(ii) *Indications for use.* It is used for conditions requiring an anti-inflammatory agent.¹

(iii) *Limitations.*¹ Do not use in viral infections. Except in emergency therapy, do not use in animals with tuberculosis, chronic nephritis, or

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may

Food and Drug Administration, HHS

§ 522.1940

Cushings's disease. With infections, use appropriate antibacterial therapy with and for at least 3 days after discontinuance of use and disappearance of all signs of infection. Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 13446, Feb. 29, 1980, as amended at 50 FR 6160, Feb. 14, 1985; 52 FR 7832, Mar. 13, 1987]

§ 522.1920 Prochlorperazine, isopropamide for injection.

(a) *Specifications.* Prochlorperazine, isopropamide for injection, veterinary, contains in each milliliter, 6 milligrams of prochlorperazine edisylate (equivalent to 4 milligrams prochlorperazine), and 0.38 milligrams of isopropamide iodide (equivalent to 0.28 milligrams of isopropamide) in buffered aqueous solution.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs and cats in which gastrointestinal disturbances are associated with emotional stress.

(2) Dosage is administered by subcutaneous injection twice daily as follows:

Weight of animal in pounds	Dosage in Milliliters
Up to 4	0.25
5 to 14	0.5-1
15 to 30	2-3
30 to 45	3-4
45 to 60	4-5
Over 60	6

Following the last injection, administer prochlorperazine and isopropamide sustained release capsules as indicated.

require bioequivalency and safety information.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 522.1940 Progesterone and estradiol benzoate in combination.

(a) [Reserved]

(b) *Sponsor.* See 000856 for use as provided in paragraphs (d)(1) and (d)(2) of this section; see 021641 for use as provided in paragraphs (d)(1) and (d)(2)(i) through (d)(2)(iii)(A) of this section.

(c) *Related tolerances.* See §§ 556.240 and 556.540 of this chapter.

(d) *Conditions of use.* It is used for implantation in animals as follows:

(1) *Suckling beef calves—(i) Amount.* 100 milligrams of progesterone and 10 milligrams of estradiol benzoate per dose.

(ii) *Indications for use.* Increased rate of weight gain.

(iii) *Limitations.* For use in suckling beef calves (at least 45 days of age) up to 400 pounds of body weight. For subcutaneous ear implantation, one dose per animal. Do not use in bull calves intended for reproduction.

(2) *Steers—(i) Amount.* 200 milligrams of progesterone and 20 milligrams of estradiol benzoate per dose.

(ii) *Indications for use.* For increased rate of weight gain and improved feed efficiency.

(iii) *Limitations.* (A) For animals weighing 400 pounds or more; for subcutaneous ear implantation, one dose per animal.

(B) For additional improvement in rate of weight gain in steers fed in confinement for slaughter, reimplant at approximately day 70.

(3) *Steers fed in confinement for slaughter—(i) Amount.* Reimplant 200 milligrams of progesterone and 20 milligrams of estradiol benzoate on approximately day 70 following an initial implant of 100 milligrams of progesterone and 10 milligrams of estradiol benzoate or 200 milligrams of progesterone and 20 milligrams of estradiol benzoate.

(ii) *Indications for use.* For additional improvement in rate of weight gain.

§ 522.1962

(iii) *Limitations.* For subcutaneous ear implantation.

[40 FR 13858, Mar. 27, 1975, as amended at 48 FR 48659, Oct. 20, 1983; 49 FR 13873, Apr. 9, 1984; 51 FR 21746, June 16, 1986; 52 FR 45312, Nov. 27, 1987; 53 FR 7406, Feb. 21, 1989; 55 FR 13769, Apr. 12, 1990; 59 FR 49808, Sept. 30, 1994; 61 FR 5507, Feb. 13, 1996; 62 FR 8372, Feb. 25, 1997; 63 FR 45945, Aug. 28, 1998]

§ 522.1962 Promazine hydrochloride injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 50 milligrams of promazine hydrochloride.

(b) *Sponsor.* In § 510.600(c) of this chapter, see No. 000008 for conditions of use as in paragraph (c)(1)(i) of this section; see No. 000856 for conditions of use as in paragraph (c)(1)(ii) of this section; see No. 000864 for conditions of use as in paragraph (c)(1)(iii) of this section.

(c) *Conditions of use.* (1)(i) To horses either intramuscularly or intravenously at a dosage of 0.2 to 0.5 milligram per pound of body weight, and to dogs and cats 1 to 3 milligrams per pound of body weight, every 4 to 6 hours as a tranquilizer or preanesthetic.¹

(ii) To horses either intramuscularly or intravenously at a dosage of 0.2 to 0.5 milligram per pound of body weight, and to dogs and cats at 1 to 2 milligrams per pound of body weight, every 4 to 6 hours as a tranquilizer, preanesthetic, for minor operative procedures in conjunction with local anesthesia, as adjunctive therapy for tetanus, and as an antiemetic in dogs and cats prior to worming, or to prevent motion sickness in dogs.¹

(iii) To horses intravenously at a dosage of 0.2 to 0.5 milligram per pound of body weight, as a tranquilizer and preanesthetic, as required.¹

(2) Not for use in conjunction with organophosphates because their toxicity may be potentiated, nor with procaine hydrochloride as its activity may be increased.¹

¹These conditions are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

21 CFR Ch. I (4–1–99 Edition)

(3) Not for use in horses intended for food.¹

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[46 FR 18962, Mar. 27, 1981]

§ 522.2002 Propiopromazine hydrochloride injection.

(a) *Chemical name.* 1-Propanone, 1-[10-[3-(dimethylamino) propyl] phenothiazine-2-yl]-, monohydrochloride.

(b) *Specifications.* Propiopromazine hydrochloride injection contains 5 or 10 milligrams of the drug in each milliliter of sterile aqueous solution.

(c) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) It is administered either intravenously or intramuscularly to dogs and cats for tranquilization at a dosage level of 0.05–0.5 milligram per pound of body weight and is also administered intravenously to dogs and cats as a preanesthetic at a dosage level of 0.25 milligram per pound of body weight.

(2) It is not to be used in conjunction with organophosphates and/or procaine hydrochloride since phenothiazines may potentiate the toxicity of organophosphates and the activity of procaine hydrochloride.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 46 FR 60570, Dec. 11, 1981; 61 FR 5507, Feb. 13, 1996]

§ 522.2005 Propofol injection.

(a) *Specifications.* The drug is a sterile, nonpyrogenic, oil-in-water emulsion containing 10 milligrams of propofol per milliliter.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter for use as in paragraphs (c)(1) and (c)(2) of this section. See No. 000074 in § 510.600(c) of this chapter for use as in paragraph (c)(1) of this section.

(c) *Conditions of use—(1) Dogs.* (i) The drug is indicated for use as an anesthetic as follows: As a single injection

to provide general anesthesia for procedures lasting up to 5 minutes; for induction and maintenance of general anesthesia using incremental doses to effect; for induction of general anesthesia where maintenance is provided by inhalant anesthetics.

(ii) The drug is administered by intravenous injection as follows: For induction of general anesthesia without the use of preanesthetics the dosage is 5.5 to 7.0 milligrams per kilogram (2.5 to 3.2 milligrams per pound); for the maintenance of general anesthesia without the use of preanesthetics the dosage is 1.1 to 3.3 milligrams per kilogram (0.5 to 1.5 milligrams per pound). The use of preanesthetic medication reduces propofol dose requirements.

(iii) Adequate data concerning safe use of propofol in pregnant and breeding dogs have not been obtained. Doses may need adjustment for geriatric or debilitated patients. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats.* (i) The drug is indicated for use as an anesthetic as follows: As a single injection to provide general anesthesia for short procedures, for induction and maintenance of general anesthesia using incremental doses to effect, and for induction of general anesthesia where maintenance is provided by inhalant anesthetics.

(ii) The drug is administered by intravenous injection as follows: For induction of general anesthesia without the use of preanesthetics the dosage is 8.0 to 13.2 milligrams per kilogram (3.6 to 6.0 milligrams per pound). For the maintenance of general anesthesia without the use of preanesthetics the dosage is 1.1 to 4.4 milligrams per kilogram (0.5 to 2.0 milligrams per pound). The use of preanesthetic medication reduces propofol dose requirements.

(iii) Adequate data concerning safe use of propofol in pregnant and breeding cats have not been obtained. Doses may need adjustment for geriatric or debilitated patients. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[61 FR 66582, Dec. 18, 1996, as amended at 62 FR 61625, Nov. 19, 1997; 63 FR 24420, May 4, 1998; 64 FR 13510, Mar. 19, 1999]

§ 522.2012 Prostalene solution.

(a) *Specifications.* Each milliliter of sterile solution contains 1 milligram of prostalene.

(b) *Sponsor.* No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use—Horses.* (1) It is used in mares for the control of estrus.

(2) It is administered at a dose of 5 micrograms per kilogram of body weight as a single subcutaneous injection.

(3) Not for use in horses intended for food.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 26854, June 30, 1976, as amended at 61 FR 5507, Feb. 13, 1996]

§ 522.2063 Pyrilamine maleate injection.

(a) *Specifications.* The drug is a sterile aqueous solution with each milliliter containing 20 milligrams of pyrilamine maleate.

(b) *Sponsors.* See No. 000061 in § 510.600(c) of this chapter for uses in paragraph (c)(2)(i) of this section; see No. 000864 in § 510.600(c) of this chapter for uses in paragraph (c)(2)(ii) of this section.

(c) *Conditions of use.* (1) It is intended for treating horses in conditions in which antihistaminic therapy may be expected to lead to alleviation of some signs of disease.¹

(2)(i) It is administered intramuscularly, subcutaneously, or intravenously. Local injection at the site of insect bites may be indicated in severe cases. Intravenous injections must be given slowly to avoid symptoms of overdosage. Dosage may be repeated every 6 to 12 hours whenever necessary. Horses, 40 to 60 milligrams per 100 pounds body weight; foals, 20 milligrams per 100 pounds body weight.¹

(ii) It is administered intravenously. Intravenous injections must be given

¹These conditions are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

§ 522.2095

slowly to avoid symptoms of overdosage. Dosage may be repeated every 6 to 12 hours if necessary. Horses, 40 to 60 milligrams per 100 pounds body weight; foals, 20 milligrams per 100 pounds body weight.¹

(3) Do not use in horses intended for food purposes.¹

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13858, Mar. 27, 1975; 41 FR 9150, Mar. 3, 1976, as amended at 42 FR 13549, Mar. 11, 1977; 42 FR 61256, Dec. 2, 1977; 51 FR 41477, Nov. 17, 1986; 52 FR 7832, Mar. 13, 1987; 54 FR 1164, Jan. 12, 1989]

§ 522.2095 Sarafloxacin solution for injection.

(a) *Specifications.* Each milliliter contains sarafloxacin hydrochloride equivalent to 50 milligrams of sarafloxacin in a 20 percent propylene glycol solution.

(b) *Sponsor.* See No. 000074 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.594 of this chapter.

(d) *Conditions of use.* 18-day embryonated broiler eggs and day-old broiler chickens:

(1) *Amount*—(i) 18-day embryonated broiler eggs: 0.05 milligram sarafloxacin in 0.1 milliliter dose in single in ovo injection.

(ii) Day-old broiler chickens: 0.1 milligrams sarafloxacin per 0.2 milliliter dose in single subcutaneous injection in the neck.

(2) *Indications for use.* For control of early chick mortality associated with *Escherichia coli* organisms susceptible to sarafloxacin.

(3) *Limitations.* Dilute 1 milliliter with 99 milliliters of sterile water or physiologic saline for use. Use entire contents of diluted solution within 24 hours. No preslaughter drug withdrawal period is required when the product is used as directed. Use in a manner other than that indicated or with dosages in excess of that recommended may result in illegal drug residues in edible tissues. Do not use in laying hens producing eggs for human consumption. Do not use in eggs intended for human consumption. The effects of sarafloxacin on the reproductive function of treated fowl have not

21 CFR Ch. I (4–1–99 Edition)

been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[60 FR 57833, Nov. 22, 1995, as amended at 62 FR 10220, Mar. 6, 1997]

§ 522.2100 Selenium, vitamin E injection.

(a)(1) *Specifications.* The drug is an emulsion containing in each milliliter, 5.48 milligrams sodium selenite (equivalent to 2.5 milligrams selenium), 50 milligrams of vitamin E (68 I.U.) (as d-alpha tocopheryl acetate), 250 milligrams polyoxyethylated vegetable oil, and 2.0 percent benzyl alcohol, and water for injection.

(2) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) The drug is intended for use for the prevention and treatment of selenium-tocopherol deficiency syndrome in horses.

(ii) The drug is administered by intravenous or deep intramuscular injection in divided doses in 2 or more sites in the gluteal or cervical muscles at a dosage level of 1 milliliter per 100 pounds of body weight and may be repeated at 5 to 10 day intervals.

(iii) Do not use in horses intended for food.

(iv) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Specifications.* The drug contains in each milliliter 2.19 milligrams of sodium selenite (equivalent to 1 milligram of selenium), 50 milligrams of vitamin E (68 I.U.) (as d-alpha tocopheryl acetate), 100 milligrams of polyoxyethylated vegetable oil, 1:10,000 thimerosal, and water for injection.

(2) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) The drug is intended for use as an aid in alleviating and controlling inflammation, pain and lameness associated with certain arthropathies in dogs.

(ii) The drug is administered subcutaneously or intramuscularly in divided doses in 2 or more sites at a dosage level of 1 milliliter per 20 pounds of body weight with a minimum dosage of ¼ milliliter and a maximum dosage of 5 milliliters. The dosage is

Food and Drug Administration, HHS

§522.2112

repeated at 3 day intervals until a satisfactory therapeutic response is observed. A maintenance regimen is then initiated which consists of 1 milliliter per 40 pounds of body weight with a minimum dosage of ¼ milliliter which is repeated every 3 days or 7 days, or longer, as required to maintain continued improvement or an asymptomatic condition; or the drug may be used in capsule form for oral maintenance therapy.

(iii) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(c)(1) *Specifications.* Each milliliter contains 2.19 milligrams of selenite sodium (equivalent to 1 milligram selenium), 50 milligrams vitamin E (68 U.S.P. units), 250 milligrams polysorbate 80, 2 percent benzyl alcohol, water for injection q. s.

(2) *Sponsor.* See No. 000061 in §510.600(c) of this chapter.

(3) *Conditions of use—(i) Dosage.* Calves: 2.5 to 3.75 milliliters per 100 pounds of body weight. Lambs 2 weeks of age or older: 1 milliliter per 40 pounds, minimum 1 milliliter. Ewes: 2.5 milliliters per 100 pounds. Sows: 1 milliliter per 40 pounds. Weanling pigs: 1 milliliter per 40 pounds, minimum 1 milliliter.

(ii) *Indications for use.* Calves, lambs, and ewes: prevention and treatment of white muscle disease (selenium-tocopherol deficiency syndrome). Sows and weanling pigs: an aid in the prevention and treatment of selenium-tocopherol deficiency.

(iii) *Limitations.* For subcutaneous or intramuscular use. Not for use in newborn pigs. Do not use in pregnant ewes. Calves: Discontinue use 30 days before treated calves are slaughtered for human consumption. Lambs, ewes, sows, or pigs: Discontinue use 14 days before treated lambs, ewes, sows, or pigs are slaughtered for human consumption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d)(1) *Specifications.* Each milliliter contains 10.95 milligrams selenite sodium (equivalent to 5 milligrams selenium), 50 milligrams vitamin E (68 U.S.P. units), 250 milligrams polysorbate 80, 2 percent benzyl alcohol, water for injection q. s.

(2) *Sponsor.* See Nos. 000061 and 000856 in §510.600(c) of this chapter.

(3) *Conditions of use—(i) Dosage.* Breeding beef cows: 1 milliliter per 200 pounds of body weight during the middle third of gestation, and 30 days before calving. Weanling calves: 1 milliliter per 200 pounds of body weight.

(ii) *Indications for use.* Weanling calves and breeding beef cows: For the prevention and treatment of selenium-tocopherol deficiency syndrome.

(iii) *Limitations.* For subcutaneous or intramuscular use. Discontinue use 30 days before treated cattle are slaughtered for human consumption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(e)(1) *Specifications.* Each milliliter contains 0.55 milligram selenite sodium (equivalent to 0.25 milligram selenium), 50 milligrams (68 U.S.P. units) vitamin E, 250 milligrams polysorbate 80, 2 percent benzyl alcohol, water for injection q. s.

(2) *Sponsor.* See No. 000061 in §510.600(c) of this chapter.

(3) *Conditions of use—(i) Dosage.* New-born lambs: 1 milliliter. Lambs 2 weeks of age or older: 4 milliliters. Baby pigs: 1 milliliter (or treat the sow during the last week of pregnancy).

(ii) *Indications for use.* Lambs: for prevention and treatment of white muscle disease (selenium-tocopherol deficiency syndrome). Baby pigs: an aid in the prevention and treatment of selenium-tocopherol deficiency.

(iii) *Limitations.* For subcutaneous or intramuscular use only. Discontinue use 14 days before treated animals are slaughtered for human consumption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 52 FR 7832, Mar. 13, 1987; 57 FR 21209, May 19, 1992; 58 FR 57556, Oct. 26, 1993; 60 FR 57833, Nov. 22, 1995]

§522.2112 Sterile sometribove zinc suspension.

(a) *Specifications.* The drug product consists of a single-dose syringe containing 500 milligrams of sometribove zinc in a sterile, prolonged-release suspension.

(b) *Sponsor.* See No. 059945 in §510.600(c) of this chapter.

(c) *Special considerations.* Use may result in reduced pregnancy rates and, in first calf heifers, an increase in days open. Use of the product has also been associated with increases in cystic ovaries and disorders of the uterus during the treatment period. Also, the incidence of retained placenta may be higher following subsequent calving. Treated cows are at an increased risk for clinical mastitis and subclinical mastitis. In some herds, use has been associated with increases in somatic cell counts in milk. Care should be taken to differentiate increased body temperature due to use of this product from an increased body temperature that may occur due to illness. Use may result in an increase in digestive disorders such as indigestion, bloat, and diarrhea. There may be an increase in the number of cows experiencing periods of “off-feed” (reduced feed intake) during treatment. Cows treated with this product may have increased numbers of enlarged hocks and lesions of the knee (carpal region), and second lactation or older cows may have more disorders of the foot region. Use has been associated with reductions in hemoglobin and hematocrit values during treatment. Human warning: Avoid prolonged or repeated contact with eyes and skin.

(d) *Conditions of use*—(1) *Amount.* 500 milligrams of sometribove zinc every 14 days beginning during the ninth week after calving and continuing until the end of lactation.

(2) *Indications for use.* For use in healthy lactating dairy cows to increase the production of marketable milk.

(3) *Limitations.* For use in lactating dairy cows only. Administer subcutaneously. Safety to replacement bulls born to treated dairy cows has not been established. To minimize injection site blemishes on carcass at time of slaughter, avoid injections within 2 weeks of expected slaughter. No milk discard or preslaughter withdrawal period is required.

[58 FR 59947, Nov. 12, 1993]

§ 522.2120 Spectinomycin dihydrochloride injection.

(a) *Specifications.* The spectinomycin dihydrochloride pentahydrate used in

manufacturing the drug is the antibiotic substance produced by the growth of *Streptomyces flavopersicus* (var. Abbott) or the same antibiotic substance produced by any other means. Each milliliter of the drug contains the following amount of spectinomycin activity from spectinomycin dihydrochloride pentahydrate:

(1) 5 milligrams when used as provided in paragraph (d)(1) of this section.

(2) [Reserved]

(3) 100 milligrams when used as provided in paragraphs (d) (2), (3), and (4) of this section.

(b) *Sponsor.* In § 510.600 of this chapter, see Nos. 000033 and 050604 for conditions of use as in paragraph (d) of this section, and see No. 000009 for conditions of use as in paragraph (d)(2) and (d)(4) of this section.

(c) *Special considerations.* The quantity of spectinomycin referred to in this section refers to the equivalent weight of base activity for the drug.

(d) *Conditions of use.* It is administered as spectinomycin dihydrochloride pentahydrate as follows:

(1) Subcutaneously in the treatment of 1-to-3-day-old turkey poults at the rate of 1 to 2 milligrams per poult as an aid in the prevention of mortality associated with Arizona group infection.

(2) Subcutaneously in the treatment of 1-to-3-day old:

(i) Turkey poults at the rate of 5 milligrams per poult as an aid in the control of chronic respiratory disease (CRD) associated with *E. coli*.

(ii) Baby chicks at the rate of 2.5 to 5 milligrams per chick as an aid in the control of mortality and to lessen severity of infections caused by *M. synoviae*, *S. typhimurium*, *S. infantis*, and *E. coli*.

(3) Intramuscularly in the treatment of dogs:

(i) At a dosage level of 2.5 milligrams to 5.0 milligrams per pound of body weight twice daily. Treatment may be continued for 4 days. For treatment of infections caused by gram-negative and gram-positive organisms susceptible to spectinomycin.

(ii) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Food and Drug Administration, HHS

§ 522.2220

(4) Administer single injection of 0.1 milliliter (10 milligrams) subcutaneously in nape of neck of 1- to 3-day-old turkey poults as an aid in control of airsacculitis associated with *M. meleagridis* sensitive to spectinomycin.

[40 FR 13858, Mar. 27, 1975, as amended at 43 FR 9273, Mar. 7, 1978; 46 FR 18964, Mar. 27, 1981; 47 FR 14149, Apr. 2, 1982; 61 FR 5507, Feb. 13, 1996; 61 FR 31028, June 19, 1996]

§ 522.2121 Spectinomycin sulfate solution.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains spectinomycin sulfate tetrahydrate equivalent to 100 milligrams of spectinomycin.

(b) *Sponsor.* See 000009 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.600 of this chapter.

(d) *Conditions of use*—(1) *Cattle*—(i) *Dose.* 10 to 15 milligrams per kilogram of body weight, at 24-hour intervals for 3 to 5 consecutive days.

(ii) *Indications for use.* For the treatment of bovine respiratory disease (pneumonia) associated with *Pasteurella haemolytica*, *P. multocida*, and *Haemophilus somnus*.

(iii) *Limitations.* For subcutaneous injection in the neck. Do not inject more than 50 milliliters at each site. Do not slaughter within 11 days of last treatment. Do not use in female dairy cattle 20 months of age or older. Use in this class of cattle may cause residues in milk. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[63 FR 24107, May 1, 1998]

§ 522.2150 Stanozolol sterile suspension.

(a) *Specifications.* Each milliliter of sterile suspension contains 50 milligrams of stanozolol.

(b) *Sponsor.* No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) Used as an anabolic steroid treatment in dogs, cats, and horses.

(2) Administered to dogs and cats by deep intramuscular injection in the thigh at weekly intervals, for several weeks. For cats and small breeds of dogs, 25 milligrams. For larger dogs, 50 milligrams.

(3) Administered to horses by deep intramuscular injection in the gluteal region at weekly intervals, for not more than 4 weeks; 25 milligrams per 100 pounds of body weight.

(4) Not for use in horses intended for food.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 46101, Oct. 6, 1975, as amended at 42 FR 36995, July 19, 1977; 55 FR 23076, June 6, 1990]

§ 522.2200 Sulfachlorpyridazine.

(a) *Chemical name.* *N*¹-(6-Chloro-3-pyridazinyl) sulfanilamide.

(b) *Specifications.* Melting point range 190° C to 191° C.

(c) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(d) *Related tolerances.* See § 556.630 of this chapter.

(e) *Conditions of use.* It is used for injection into calves as follows:

(1) *Amount.* 30 to 45 milligrams per pound of body weight per day.

(2) *Indications for use.* Treatment of diarrhea caused or complicated by *E. coli* (colibacillosis).

(3) *Limitations.* Administer as the sodium salt of sulfachlorpyridazine intravenously in aqueous solution for 1 to 5 days in divided doses twice daily; treated calves must not be slaughtered for food during treatment or for 5 days after the last treatment.

[40 FR 13858, Mar. 27, 1975, as amended at 50 FR 41489, Oct. 11, 1985]

§ 522.2220 Sulfadimethoxine injection.

(a)(1) *Specifications.* Sulfadimethoxine injection containing 400 milligrams per milliliter.

(2) *Sponsor.* (i) See No. 000069 in § 510.600(c) of this chapter for conditions of use as in paragraphs (a)(3)(i) through (a)(3)(iii) of this section.

(ii) See No. 057561 for conditions of use as in paragraph (a)(3) of this section.

(iii) See No. 059130 for use as in paragraph (a)(3)(iii) of this section.

§522.2240

21 CFR Ch. I (4–1–99 Edition)

(3) *Conditions of use.* (i) It is used or intended for use in dogs and cats as follows:

(a) For the treatment of respiratory, genitourinary tract, enteric, and soft tissue infections when caused by Streptococci, Staphylococci, Escherichia, Salmonella, Klebsiella, Proteus, or Shigella organisms sensitive to sulfadimethoxine, and in the treatment of canine bacterial enteritis associated with coccidiosis and canine Salmonellosis.

(b) It is administered by intravenous or subcutaneous injection at an initial dose of 55 milligrams per kilogram of body weight followed by 27.5 milligrams per kilogram of body weight every 24 hours.

(c) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(ii) It is used or intended for use in horses as follows:

(a) For the treatment of respiratory disease caused by Streptococcus equi (strangles).

(b) It is administered by intravenous injection at an initial dose of 55 milligrams per kilogram of body weight followed by 27.5 milligrams per kilogram of body weight every 24 hours until the patient is asymptomatic for 48 hours.

(c) Not for use in horses intended for food.

(d) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(iii) It is used or intended for use in cattle as follows:

(a) For the treatment of shipping fever complex, bacterial pneumonia, calf diphtheria, and foot-rot.

(b) It is administered by intravenous injection at an initial dose of 25 milligrams per pound of body weight followed by 12.5 milligrams per pound of body weight every 24 hours until the animal is asymptomatic for 48 hours.

(c) Milk taken from animals during treatment and for 60 hours (5 milkings) after the latest treatment must not be used for food. Do not administer within 5 days of slaughter.

(d) Tissue damage may result from perivascular infiltration.

(b) [Reserved]

(c)(1) *Specifications.* Sulfadimethoxine containing 100 milligrams per milliliter.

(2) *Sponsor.* See No. 000010 in §510.600(c) of this chapter.

(3) *Conditions of use.* (i) It is used or intended for use in the treatment of sulfadimethoxine-susceptible bacterial infections in dogs.

(ii) It is administered by subcutaneous, intramuscular, or intravenous injection at an initial dose of 25 milligrams per pound of body weight followed by 12.5 milligrams per pound of body weight every 24 hours thereafter. Continue treatment until the animal is free from symptoms for 48 hours.

(iii) For use by or on the order of a licensed veterinarian.

(d) *Related tolerances.* See §556.640 of this chapter.

[40 FR 13858, Mar. 27, 1975, as amended at 40 FR 34112, Aug. 14, 1975; 40 FR 42007, Sept. 10, 1975; 50 FR 254, Jan. 3, 1985; 53 FR 40728, Oct. 18, 1988; 54 FR 30205, July 19, 1989; 58 FR 38972, July 21, 1993; 59 FR 56000, Nov. 10, 1994; 61 FR 4875, Feb. 9, 1996; 62 FR 23128, Apr. 29, 1997; 62 FR 35076, June 30, 1997]

§522.2240 Sulfathiazopyridazine.

(a) *Chemical name.* N¹-(6-Ethoxy-3-pyridazinyl) sulfanilamide.

(b) *Specifications.* Melting point range of 180° C to 186° C.

(c) *Sponsor.* See No. 010042 in §510.600(c) of this chapter.

(d) *Related tolerances.* See §556.650 of this chapter.

(e) *Conditions of use.* It is used for injection into cattle as follows:

(1) *Amount.* 2.5 grams per 100 pounds of body weight per day.

(2) *Indications for use.* Treatment of respiratory infection (pneumonia, shipping fever), foot rot, calf scours; as adjunctive therapy in septicemia accompanying mastitis and metritis.

(3) *Limitations.* Administer intravenously for not more than 4 days; or first treatment may be followed by 3 days of treatment with sulfathiazopyridazine in drinking water, feed, or tablet in accordance with §558.579(e) or §§520.2240a(e) and 520.2240b(e) of this chapter; as sodium sulfathiazopyridazine; do not treat within 16 days of slaughter; as sole source of sulfonamide; milk that has

Food and Drug Administration, HHS

§ 522.2404

been taken from animals during treatment and for 72 hours (6 milkings) after the latest treatment must not be used for food; for use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 41 FR 11011, Mar. 15, 1976]

§ 522.2260 Sulfamethazine injectable solution.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 250 milligrams of sulfamethazine sodium.

(b) *Sponsor.* See No. 010042 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.670 of this chapter.

(d) *NAS/NRC status.* The conditions of use are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(e) *Conditions of use—*(1) *Amount.* 20 milliliters for each 50 pounds of body weight (100 milligrams per pound) initially, 20 milliliters per 100 pounds of body weight (50 milligrams per pound) daily thereafter for cattle.

(2) *Indications for use.* For cattle for treatment of bacterial pneumonia and bovine respiratory disease complex (shipping fever complex) (*Pasteurella* spp.), colibacillosis (bacterial scours) (*Escherichia coli*), necrotic pododermatitis (foot rot) (*Fusobacterium necrophorum*), calf diphtheria (*Fusobacterium necrophorum*), acute mastitis and acute metritis (*Streptococcus* spp.) when caused by one or more pathogenic organisms sensitive to sulfamethazine.

(3) *Limitations.* For intravenous use only. Not for use in lactating dairy animals. Withdraw medication from cattle 10 days prior to slaughter for food. If symptoms persist for 2 or 3 days, consult a veterinarian. Adequate water intake is important for animals treated with sulfonamides. Treatment should continue 24 to 48 hours beyond the remission of disease symptoms, but not to exceed a total of 5 consecutive days.

[46 FR 62055, Dec. 22, 1981]

§ 522.2340 Sulfomyxin.

(a) *Specifications.* Sulfomyxin for injection is sterile. It is derived from the antibiotic substance produced by the growth of *Bacillus polymyxa* or is the same substance produced by any other means.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Special considerations.* The quantities of antibiotic in paragraph (e) of this section refer to the activity of the appropriate standard.

(d) *Related tolerances.* See § 556.700 of this chapter.

(e) *Conditions of use.* (1) It is used or intended for use in chickens and turkeys as an aid in the treatment of disease caused or complicated by *E. coli*, such as colibacillosis and complicated chronic respiratory disease.

(2) It is administered by subcutaneous injection as follows:

Age of birds in days	Antibiotic activity	
	Chickens (units)	Turkeys (units)
1 to 14	12,500	12,500
15 to 28	25,000	25,000
29 to 63	50,000	50,000
Over 63	50,000	100,000

(3) A second injection may be given 3 days later if symptoms persist.

(4) Not for use in laying hens; do not treat chickens within 5 days of slaughter; do not treat turkeys within 7 days of slaughter.

§ 522.2404 Thialbarbitone sodium for injection.

(a) *Specifications.* Thialbarbitone sodium for injection when reconstituted with sterile distilled water provides 94 milligrams of thialbarbitone sodium per milliliter of solution.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is administered as a general anesthetic in surgical procedures on dogs, cats, swine, sheep, cattle, and horses. The drug is used for procedures of relatively short duration. However, the period of anesthesia can be lengthened by slower initial injection and supplemental administration during surgery.

(2) It is administered intravenously. The drug is injected slowly to dogs, cats, cattle, sheep, and swine. For

§ 522.2424

horses, it is recommended that a pre-anesthetic sedation be administered to the horse 30 minutes before the drug is administered. The drug is then injected rapidly and completely. The drug is used at the following dosage levels:

Species	Weight of animal in pounds	Dosage in milligrams per pound
Dog	Over 50	14.1
Do	30-50	18.8
Do	10-30	23.5
Do	Under 10	28.2
Cat		31.3-37.6
Horse		6.3-7.8
Cattle and swine		6.7-9.4
Calves and sheep		9.4-11.8

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

§ 522.2424 Sodium thiamylal for injection.

(a) *Specifications.* The drug is a sterile dry powder. It is reconstituted aseptically with sterile distilled water, water for injection, or sodium chloride injection, to a desired concentration of 0.5 to 4 percent sodium thiamylal.

(b) *Sponsors.* See code Nos. 000010 and 000856 in § 510.500(c) of this chapter.

(c) *Conditions of use.* (1) It is used as an ultra-short-acting anesthetic in dogs, cats, swine, horses, and cattle.

(2) When diluted aseptically to the desired concentration and administered intravenously to effect, the average single dose is:

(i) Dogs and cats: 8 milligrams per pound of body weight (when used with a preanesthetic, generally one-half the normal dose).

(ii) Swine: 40 milligrams per 5 pounds of body weight.

(iii) Horses: Light anesthesia, 1 gram per 500 pounds to 1,100 pounds of body weight; deep anesthesia, 1 gram per 300 pounds of body weight (40 milligrams per 12 pounds of body weight).

(iv) Cattle: Short duration, 20 milligrams per 5 pounds of body weight; longer duration, 40 milligrams per 7 pounds of body weight.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(4) NAS/NRC status: The conditions of use specified in this paragraph are NAS/NRC reviewed and found effective. Applications for these uses need not in-

21 CFR Ch. I (4-1-99 Edition)

clude effectiveness data as specified in § 514.111 of this chapter, but may require bioequivalency and safety information.

[40 FR 25812, June 19, 1975, as amended at 49 FR 8434, Mar. 7, 1984; 53 FR 23390, June 22, 1988; 53 FR 40728, Oct. 18, 1988; 62 FR 35076, June 30, 1997]

§ 522.2444 Sodium thiopental implantation or injectable dosage forms.

§ 522.2444a Sodium thiopental for injection.

(a) *Specifications.* The drug contains sodium thiopental sterile powder for dilution with sterile water for injection.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used as an anesthetic for intravenous administration to dogs and cats during short to moderately long surgical and other procedures. It is also used to induce anesthesia in dogs and cats which then have surgical anesthesia maintained by use of a volatile anesthetic.

(2) It is administered as follows:

(i) For brief anesthesia (6 to 10 minutes) a dosage of 6 to 9 milligrams per pound of body weight is suggested.

(ii) To obtain anesthesia of 15 to 25 minutes duration the suggested dosage is 10 to 12 milligrams per pound of body weight.

(iii) Use of a preanesthetic tranquilizer or morphine will decrease the dosage of sodium thiopental required, provide for smoother induction and smoother recovery, and sometimes prolong the recovery period. If morphine is used as a preanesthetic agent the dose of the barbiturate can be reduced as much as 40 to 50 percent. When a tranquilizer is administered the barbiturate dosage can be reduced 10 to 25 percent.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

§ 522.2444b Sodium thiopental, sodium pentobarbital for injection.

(a) *Specifications.* Each gram of the drug contains 750 milligrams of sodium thiopental and 250 milligrams of sodium pentobarbital sterile powder for

dilution with sterile water for injection.

(b) *Sponsor.* See No. 050604 in §510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used as an anesthetic for intravenous administration to dogs and cats during short to moderately long surgical procedures.

(2) It is administered as follows:

(i) For total anesthesia, it is given at approximately 10 to 12 milligrams per pound of body weight over a period of 3.5 to 5 minutes.

(ii) When preanesthetic medication is used, it is important to wait at least an hour before administering thiopental and sodium pentobarbital for injection, and the dosage necessary for anesthesia is reduced. Usually $\frac{1}{2}$ to $\frac{2}{3}$ the normal amount is adequate.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 47 FR 14149, Apr. 2, 1982]

§522.2470 Tiletamine hydrochloride and zolazepam hydrochloride for injection.

(a) *Specifications.* Tiletamine hydrochloride and zolazepam hydrochloride for injection when reconstituted with sterile distilled water provides tiletamine hydrochloride and zolazepam hydrochloride equivalent to 50 milligrams of tiletamine base and 50 milligrams of zolazepam base per milliliter of solution.

(b) *Sponsor.* See No. 000031 in §510.600(c) of this chapter.

(c) *Conditions of use—(1) Indications for use.* It is used for restraint or for anesthesia combined with muscle relaxation in cats and in dogs for restraint and minor procedures of short duration (30 minutes) requiring mild to moderate analgesia.

(2) *Amount.* Expressed as milligrams of the drug combination:

(i) In healthy dogs: An initial intramuscular dosage of 3 to 4.5 milligrams per pound of body weight for diagnostic purposes; 4.5 to 6 milligrams per pound of body weight for minor procedures of short duration such as repair of lacerations and wounds, castrations, and other procedures requiring mild to moderate analgesia. Supplemental doses when required should

be less than the initial dose and the total dose given should not exceed 12 milligrams per pound of body weight. The maximum total safe dose is 13.6 milligrams per pound of body weight.

(ii) In healthy cats: An initial intramuscular dosage of 4.4 to 5.4 milligrams per pound of body weight is recommended for such procedures as dentistry, treatment of abscesses, foreign body removal, and related types of surgery; 4.8 to 5.7 milligrams per pound of body weight for minor procedures requiring mild to moderate analgesia, such as repair of lacerations, castrations, and other procedures of short duration. Initial dosages of 6.5 to 7.2 milligrams per pound of body weight are recommended for ovariohysterectomy and onychectomy. When supplemental doses are required, such individual supplemental doses should be given in increments that are less than the initial dose and the total dose given (initial dose plus supplemental doses) should not exceed the maximum allowable safe dose of 32.7 milligrams per pound of body weight.

(3) *Limitations.* Discard unused reconstituted solution after 48 hours. Not for use in dogs and cats with pancreatic disease, or with severe cardiac or pulmonary dysfunction. Not for use in pregnant animals. Not for use in cats suffering with renal insufficiency. The dosage should be reduced in geriatric dogs and cats. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[47 FR 15328, Apr. 9, 1982, as amended at 51 FR 24142, July 2, 1986]

§522.2471 Tilimicosin phosphate injection.

(a) *Specifications.* Each milliliter contains 300 milligrams of tilimicosin base as tilimicosin phosphate.

(b) *Sponsor.* See 000986 in §510.600(c) of this chapter.

(c) *Related tolerances.* See §556.735 of this chapter.

(d) *Conditions of use—(1) Cattle—(i) Amount.* 10 milligrams per kilogram body weight.

(ii) *Indications for use.* For the treatment of bovine respiratory disease (BRD) associated with *Pasteurella haemolytica*. For the control of respiratory disease in cattle at high risk

§522.2474

of developing BRD associated with *P. haemolytica*.

(iii) *Limitations*. For use only in cattle as a single subcutaneous injection. Not for human use. Use of this antibiotic in humans may prove fatal. Do not use in automatically powered syringes. Do not inject more than 15 milliliters per injection site. If no improvement is noted within 48 hours, the diagnosis should be reevaluated. Do not use intravenously in cattle. Intravenous injection in cattle will be fatal. Do not use in other animal species. Injection of this antibiotic has been found to be fatal in swine and nonhuman primates, and it may be fatal in horses. Safety of use in pregnant and breeding animals has not been established. Do not use in female dairy cattle 20 months of age or older. Use of this antibiotic in this class of cattle may cause milk residues. Do not slaughter within 28 days of last treatment. Federal law restricts this drug to use or on the order of a licensed veterinarian.

(2) [Reserved]

[57 FR 12712, Apr. 13, 1992, as amended at 62 FR 5526, Feb. 6, 1997; 63 FR 7701, Feb. 17, 1998; 63 FR 14818, Mar. 27, 1998]

§522.2474 Tolazoline hydrochloride injection.

(a) *Specifications*. Each milliliter of sterile aqueous solution contains tolazoline hydrochloride equivalent to 100 milligrams of base activity.

(b) *Sponsor*. See No. 061690 in §510.600(c) of this chapter.

(c) *Conditions of use*. It is used as follows:

(1) *Horses*—(i) *Amount*. Administer slowly by intravenous injection 4 milligrams per kilogram of body weight or 1.8 milligrams per pound (4 milliliters per 100 kilograms or 4 milliliters per 220 pounds).

(ii) *Indications for use*. For use in horses when it is desirable to reverse the effects of sedation and analgesia caused by xylazine.

(iii) *Limitations*. The safety of Tolazine™ has not been established in pregnant mares, lactating mares, horses intended for breeding, foals, or horses with metabolically unstable conditions. The safety of Tolazine™ has not been evaluated for reversing

21 CFR Ch. I (4–1–99 Edition)

xylazine used as a preanesthetic to a general anesthetic. This drug is for use in horses only and not for use in food-producing animals. Users with cardiovascular disease (for example, hypertension or ischemic heart disease) should take special precautions to avoid accidental exposure to this product.

Accidental spillage on the skin should be washed off immediately with soap and water. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[61 FR 25785, May 23, 1996]

§522.2476 Trenbolone acetate.

(a) *Specifications*. Each pellet for implanting contains 20 milligrams of trenbolone acetate.

(b) *Sponsor*. See 012579 and 021641 in §510.600(c) of this chapter.

(c) *Related tolerances*. See §556.739 of this chapter.

(d) *Conditions of use*—(1) *Heifers* 200 milligrams trenbolone acetate (10 pellets of 20 milligrams each) for increased rate of weight gain and improved feed efficiency in growing-finishing feedlot heifers, use last 63 days prior to slaughter.

(2) *Steers*. 140 milligrams trenbolone acetate (7 pellets of 20 milligrams each) for improved feed efficiency in growing-finishing feedlot steers, use 126 days prior to slaughter, should be reimplanted once after 63 days.

(3) *Limitations*. Not for use in animals intended for subsequent breeding or in dairy animals. Implant in ear only.

[52 FR 24995, July 2, 1987, as amended at 62 FR 29013, May 29, 1997]

§522.2477 Trenbolone acetate and estradiol.

(a) *Sponsor*. See No. 012579 in §510.600(c) of this chapter for use as in paragraphs (c)(1), (c)(2), and (c)(3) of this section. See No. 021641 in §510.600(c) of this chapter for use as in paragraph (c)(1) of this section.

(b) *Related tolerances*. See §§556.240 and 556.739 of this chapter.

(c) *Conditions of use*—(1) *Feedlot steers*—(i) *Amount*. 120 milligrams of trenbolone acetate and 24 milligrams

of estradiol (6 pellets, each pellet containing 20 milligrams of trenbolone acetate and 4 milligrams of estradiol) per animal.

(ii) *Indications for use.* For increased rate of weight gain and improved feed efficiency in feedlot steers.

(iii) *Limitations.* Implant subcutaneously in ear only. Do not use in animals intended for subsequent breeding or in dairy animals.

(2) *Heifers*—(i) *Amount.* 140 milligrams of trenbolone acetate and 14 milligrams of estradiol (7 pellets, each pellet containing 20 milligrams of trenbolone acetate and 2 milligrams of estradiol) per animal.

(ii) *Indications for use.* For increased rate of weight gain and improved feed efficiency in heifers fed in confinement for slaughter.

(iii) *Limitations.* Implant subcutaneously in ear only. Do not use in animals intended for subsequent breeding or in dairy animals.

(3) *Pasture cattle (slaughter, stocker, feeder steers, and heifers)*—(i) *Amount.* 40 milligrams of trenbolone acetate and 8 milligrams of estradiol (2 pellets, each pellet containing 20 milligrams of trenbolone acetate and 4 milligrams of estradiol) per animal.

(ii) *Indications for use.* For increased rate of weight gain.

(iii) *Limitations.* Implant subcutaneously in ear only. Not for use in animals intended for subsequent breeding or in dairy animals.

[60 FR 4376, Jan. 23, 1995, as amended at 61 FR 29480, June 11, 1996; 61 FR 41499, Aug. 9, 1996; 62 FR 28629, May 27, 1997]

§ 522.2478 Trenbolone acetate and estradiol benzoate.

(a) *Sponsor.* See 000856 in § 510.600(c) of this chapter.

(b) *Related tolerance.* See §§ 556.240 and 556.739 of this chapter.

(c) *Conditions of use*—(1) *Steers*—(i) *Amount.* 200 milligrams of trenbolone acetate and 28 milligrams of estradiol benzoate (one implant consisting of 8 pellets, each pellet containing 25 milligrams of trenbolone acetate and 3.5 milligrams of estradiol benzoate) per animal.

(ii) *Indications for use.* For improved feed efficiency in steers fed in confinement for slaughter.

(iii) *Limitations.* Implant subcutaneously in ear only.

(2) *Heifers*—(i) *Amount.* 200 milligrams of trenbolone acetate and 28 milligrams of estradiol benzoate (one implant consisting of 8 pellets, each pellet containing 25 milligrams of trenbolone acetate and 3.5 milligrams of estradiol benzoate) per animal.

(ii) *Indications for use.* For increased rate of weight gain in heifers fed in confinement for slaughter.

(iii) *Limitations.* Implant subcutaneously in ear only. Not for dairy or beef replacement heifers.

[61 FR 14482, Apr. 2, 1996, as amended at 61 FR 29479, June 11, 1996; 63 FR 63789, Nov. 17, 1998]

§ 522.2483 Sterile triamcinolone acetonide suspension.

(a) *Specifications.* Each milliliter of suspension contains 2 or 6 milligrams triamcinolone acetonide.

(b) *Sponsor.* See 000010 and 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*—(i) *Dogs and cats*—(a) *Intramuscular or subcutaneous.* Single injection of 0.05 to 0.1 milligram (mg.) per pound of body weight in inflammatory, arthritic, or allergic disorders. Single injection of 0.1 mg. per pound of body weight in dermatologic disorders. If symptoms recur, the dose may be repeated, or oral corticosteroid therapy may be instituted.¹

(b) *Intralesional.* 1.2 to 1.8 mg., divided in several injections, spaced around the lesion at 0.5 to 2.5 centimeters apart depending on the size. At any one site the dose injected should not exceed 0.6 mg. and should be well into the cutis to prevent rupture of the epidermis. When treating animals with multiple lesions, do not exceed a total dose of 6 mg.

(c) *Intra-articular and intrasynovial.* Single injection of 1 to 3 mg. dose, dependent on size of joint and severity of symptoms. After 3 or 4 days, repeat dosage if indicated. If initial results are inadequate or too transient, dosage may be increased, not to exceed 3 mg.

(ii) *Horses*—(a) *Intramuscular or subcutaneous.* Single injection of 0.01 to 0.02 mg. per pound of body weight. Usual dose, 12 to 20 mg.

§ 522.2582

21 CFR Ch. I (4–1–99 Edition)

(b) *Intra-articular and intrasynovial.* Single injection of 6 to 18 mg. dose, dependent on size of joint and severity of symptoms. After 3 or 4 days, repeat dosage if indicated. If initial results are inadequate or too transient, dosage may be increased, not to exceed 18 mg.

(2) *Indications for use.* Treatment of inflammation and related disorders in dogs, cats, and horses;¹ and management and treatment of acute arthritis and allergic and dermatologic disorders in dogs and cats.

(3) *Limitations.* (i) Do not use in viral infections. With bacterial infections, appropriate antibacterial therapy should be used.

(ii) Do not use in animals with tuberculosis, chronic nephritis, or cushingoid syndrome, except for emergency therapy.

(iii) Not for use in horses intended for food.

(iv) Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(v) Do not use in the treatment of laminitis.

(vi) Intra-articular injection in equine leg injuries may produce osseous metaplasia.

(vii) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 4976, Feb. 7, 1978, as amended at 50 FR 41490, Oct. 11, 1985; 52 FR 1903, Jan. 16, 1987; 53 FR 40728, Oct. 18, 1988; 62 FR 35077, June 30, 1997]

§ 522.2582 Triflupromazine hydrochloride injection.

(a) *Specifications.* Triflupromazine hydrochloride injection contains 20 milligrams of triflupromazine hydrochloride in each milliliter of sterile aqueous solution.

¹These conditions are NAS/NRC reviewed and are deemed effective. Applications for these uses need not include the effectiveness data specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs, cats, and horses to relieve anxiety and to help control psychomotor overactivity as well as to increase the tolerance of animals to pain and pruritus. The drug is indicated in various office and clinical procedures which require the aid of a tranquilizer, antiemetic, or preanesthetic.¹

(2) The drug is administered to dogs either intravenously at a dosage level of 0.5 to 1 milligram per pound of body weight daily, or intramuscularly at a dosage level of 1 to 2 milligrams per pound of body weight daily. It is administered to cats intramuscularly at a dosage level of 2 to 4 milligrams per pound of body weight daily. It is administered to horses intravenously or intramuscularly at a dosage level of 10 to 15 milligrams per 100 pounds of body weight daily to a maximum dose of 100 milligrams.¹

(3) Not for use in horses intended for food.¹

(4) Do not use in conjunction with organophosphates and/or procaine hydrochloride, because phenothiazines may potentiate the toxicity of organophosphates and the activity of procaine hydrochloride.¹

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13858, Mar. 27, 1975, as amended at 50 FR 41490, Oct. 11, 1985]

§ 522.2610 Trimethoprim and sulfadiazine sterile suspension.

(a)(1) *Specifications.* Each milliliter of sterile aqueous suspension contains 240 milligrams (40 milligrams of trimethoprim and 200 milligrams of sulfadiazine).

(2) *Sponsor.* See 000061 and 000856 in § 510.600(c) of this chapter.

(3) *Conditions of use—(i) Dosage.* One milliliter (40 milligrams of trimethoprim and 200 milligrams of

¹These conditions are NAS/NRC reviewed and are deemed effective. Applications for these uses need not include the effectiveness data specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

Food and Drug Administration, HHS

§ 522.2640a

sulfadiazine) per 20 pounds (9 kilograms) of body weight per day.

(ii) *Indications.* For dogs for treatment of acute urinary tract infections, acute bacterial complications of distemper, acute respiratory tract infections, acute alimentary tract infections, and acute septicemia due to *Streptococcus zooepidemicus*.

(iii) *Limitations.* For subcutaneous use in dogs only; administer once every 24 hours, or for severe infections, after an initial dose, administer half the normal daily dose every 12 hours; continue therapy 2 to 3 days after clinical signs of infection have subsided; if no improvement is seen in 3 to 5 days, reevaluate diagnosis; injection may be used alone or in conjunction with oral dosing; not recommended for use for more than 14 days; a complete blood count should be done for prolonged use; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Specifications.* Each milliliter of sterile aqueous suspension contains 480 milligrams (80 milligrams of trimethoprim and 400 milligrams of sulfadiazine (as the sodium salt)).

(2) *Sponsor.* See 000856 and 011716 in § 510.600(c) of this chapter.

(3) *Conditions of use—(i) Dosage.* Two milliliters (160 milligrams of trimethoprim and 800 milligrams of sulfadiazine) per 100 pounds (45 kilograms) of body weight per day.

(ii) *Indications.* For horses where systemic anti-bacterial action against sensitive organisms is required during treatment of acute strangles, respiratory tract infections, acute urogenital infections, and wound infections and abscesses.

(iii) *Limitations.* For intravenous use; administer as single, daily dose for 5 to 7 days; daily dose may also be halved and given morning and evening; continue acute infection therapy 2 to 3 days after clinical signs have subsided; if no improvement of acute infections is seen in 3 to 5 days, reevaluate diagnosis; a complete blood count should be done periodically for prolonged use; not for use in horses intended for food; Fed-

eral law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 241, Jan. 4, 1983, as amended at 48 FR 23180, May 24, 1983; 48 FR 42809, Sept. 20, 1983; 61 FR 5507, Feb. 13, 1996; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 522.2615 TripeleNNamine hydrochloride injection.

(a) *Specifications.* Each milliliter of aqueous solution contains 20 milligrams of tripeleNNamine hydrochloride.

(b) *Sponsor.* See Nos. 053501 and 059130 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.741 of this chapter.

(d) *Conditions of use—(1) Amount—(i) Dogs, cats, and horses.* For intramuscular use only at a dose of 0.5 milligram per pound of body weight.

(ii) *Cattle.* Administer intravenously or intramuscularly at a dose of 0.5 milligram per pound of body weight.

(2) *Indications for use.* For use in treating conditions in which antihistaminic therapy may be expected to lead to alleviation of some signs of disease.

(3) *Limitations.* Do not use in horses intended for food purposes. Treated cattle must not be slaughtered for food during treatment and for 4 days following the last treatment. Milk that has been taken during treatment and for 24 hours (two milkings) after the last treatment must not be used for food. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[51 FR 44450, Dec. 10, 1986, as amended at 61 FR 29480, June 11, 1996; 62 FR 4164, Jan. 29, 1997]

§ 522.2640 Tylosin injectable dosage forms.

§ 522.2640a Tylosin injection.

(a) *Specifications.* Each milliliter of sterile solution of 50 percent propylene glycol with 4 percent benzyl alcohol contains 50 to 200 milligrams of tylosin activity (as tylosin base). Tylosin conforms to the appropriate antibiotic standard. Tylosin contains at least 95

percent tylosin as a combination of tylosin A, tylosin B, tylosin C, and tylosin D of which at least 80 percent is tylosin A as determined by a method entitled “Determination of Factor Content in Tylosin by High Performance Liquid Chromatography,” which is incorporated by reference. Copies are available from the Dockets Management Branch (HFA–305), Food and Drug Administration, rm. 1–23, 12420 Parklawn Dr., Rockville, MD 20857, or available for inspection at the Office of the Federal Register, 800 North Capitol Street, NW., suite 700, Washington, DC 20001.

(b) *Sponsors.* (1) See No. 000986 in § 510.600(c) of this chapter for use in paragraphs (e)(1), (2), and (3) of this section.

(2) See No. 000010 in § 510.600(c) of this chapter for use as in paragraphs (e)(1) and (2) of this section.

(c) *NAS/NRC status.* These conditions of use are NAS/NRC reviewed and found effective. NADA’s for these uses need not include effectiveness data as specified by § 514.111 of this chapter but may require bioequivalency and safety information.

(d) *Related tolerances.* See § 556.740 of this chapter.

(e) *Conditions of use—(1) Beef cattle and nonlactating dairy cattle—(i) Amount.* 8 milligrams per pound of body weight once daily.

(ii) *Indications for use.* Treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Corynebacterium pyogenes*; foot rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum* and metritis caused by *Corynebacterium pyogenes*.

(iii) *Limitations.* Administer intramuscularly for not more than 5 consecutive days. Continue treatment 24 hours after symptoms disappear. Do not inject more than 10 milliliters per site. Do not use in lactating dairy cattle. Use a 50-milligram-per-milliliter solution for calves weighing less than 200 pounds. Do not administer within 21 days of slaughter.

(2) *Swine—(i) Amount.* 4 milligrams per pound of body weight twice daily.

(ii) *Indications for use.* Treatment of swine arthritis caused by *Mycoplasma*

hyosynoviae; swine pneumonia caused by *Pasteurella* spp.; swine erysipelas caused by *Erysipelothrix rhusiopathiae*; swine dysentery associated with *Treponema hyodysenteriae* when followed by appropriate medication in the drinking water and/or feed.

(iii) *Limitations.* Administer intramuscularly for not more than 3 consecutive days. Continue treatment 24 hours after symptoms disappear. Do not inject more than 5 milliliters per site. Do not administer within 14 days of slaughter. If tylosin medicated drinking water is used as followup treatment for swine dysentery, the animal should thereafter receive feed containing 40 to 100 grams of tylosin per ton for 2 weeks to assure depletion of tissue residues.

(3) *Dogs and cats—(i) Amount.* 3 to 5 milligrams per pound of body weight at 12- to 24-hour intervals.

(ii) *Indications for use—(a) Dogs.* Treatment of upper respiratory infections such as bronchitis, tracheobronchitis, tracheitis, laryngitis, tonsillitis, and pneumonia caused by *Staphylococci* spp., hemolytic *Streptococci* spp., and *Pasteurella multocida*.

(b) *Cats.* Treatment of upper respiratory infections when caused by *Staphylococci* spp. and hemolytic *Streptococci* spp. and for feline pneumonitis when caused by tylosin susceptible organisms.

(iii) *Limitations.* For intramuscular use only. If there is no response to therapy in 5 days, diagnosis and treatment should be reassessed. Use a 50-milligram-per-milliliter solution only. Dogs and cats receiving a dose of less than 50 milligrams (1 milliliter) should be dosed with a tuberculin syringe. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 48643, Oct. 2, 1981, as amended at 47 FR 9398, Mar. 5, 1982; 50 FR 49841, Dec. 5, 1985; 50 FR 50292, Dec. 10, 1985; 53 FR 40728, Oct. 18, 1988; 59 FR 14365, Mar. 28, 1994; 62 FR 35077, June 30, 1997]

§ 522.2640b Tylosin tartrate for injection.

(a) *Specifications.* The drug is a sterile powder containing a mixture of tylosin tartrate and sodium citrate which is

reconstituted to provide 25 milligrams of tylosin activity per milliliter. Tylosin as the tartrate salt, conforms to the appropriate antibiotic standard. Tylosin contains at least 95 percent tylosin as a combination of tylosin A, tylosin B, tylosin C, and tylosin D of which at least 80 percent is tylosin A as determined by a method entitled "Determination of Factor Content in Tylosin by High Performance Liquid Chromatography," which is incorporated by reference. Copies are available from the Dockets Management Branch (HFA-305), Food and Drug Administration, rm. 1-23, 12420 Parklawn Dr., Rockville, MD 20857, or available for inspection at the Office of the Federal Register, 800 North Capitol Street, NW., suite 700, Washington, DC 20001.

(b) *Sponsor.* See No. 000986 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.740 of this chapter.

(d) *Conditions of use*—(1) *Chickens*—(i) *Amount.* 25 milligrams per 2 pounds of body weight.

(ii) *Indications for use.* As an aid in the control and treatment of chronic respiratory disease caused by *Mycoplasma gallisepticum* sensitive to tylosin.

(iii) *Limitations.* Not for use in laying chickens producing eggs for human consumption; inject under the loose skin of the neck behind the head; if no improvement is noted within 5 days, diagnosis should be reevaluated; do not treat within 3 days of slaughter.

(2) *Turkeys*—(i) *Amount.* 6.25 to 12.5 milligrams per sinus.

(ii) *Indications for use.* As an aid in the control and treatment of infectious sinusitis caused by *Mycoplasma gallisepticum* sensitive to tylosin.

(iii) *Limitations.* Do not use in laying turkeys producing eggs for human consumption; inject 6.25 milligrams to 12.5 milligrams per sinus depending on severity of condition; treatment may be repeated in 10 days if the swelling persists; do not treat within 5 days of slaughter; may be used in conjunction with tylosin in drinking water as indicated in § 520.2640(e)(2) of this chapter.

[46 FR 48643, Oct. 2, 1981, as amended at 50 FR 49841, Dec. 5, 1985; 50 FR 49841, Dec. 5, 1985; 59 FR 14365, Mar. 28, 1994]

§ 522.2662 Xylazine hydrochloride injection.

(a) *Specifications.* Xylazine hydrochloride injection is a sterile aqueous solution containing xylazine hydrochloride equivalent to 100 milligrams of xylazine in each milliliter of solution when intended for use in horses, wild deer, and elk, and 20 milligrams of xylazine per milliliter of solution when intended for use in dogs and cats.

(b) *Sponsor.* See 000856 in § 510.600(c) of this chapter for use in horses, wild deer, and elk. See 000859 and 061651 in § 510.600(c) of this chapter for use in horses, wild deer, elk, dogs, and cats. See 061690 in § 510.600(c) of this chapter for use in horses, wild deer, elk, dogs, and cats. See 000010 in § 510.600(c) of this chapter for use in horses only.

(c) *Conditions of use.* (1) The drug is used in horses, wild deer, elk, dogs, and cats to produce sedation, as an analgesic, and a preanesthetic to local anesthesia. It may also be used in horses, dogs, and cats as a preanesthetic to general anesthesia.

(2) It is administered as follows:

(i) To horses from a solution containing 100 milligrams of xylazine per milliliter, intravenously at 0.5 milligram per pound of body weight, or intramuscularly at 1.0 milligram per pound of body weight.

(ii) To dogs and cats from a solution containing 20 milligrams of xylazine per milliliter; intravenously at 0.5 milligram per pound of body weight or intramuscularly or subcutaneously at 1.0 milligram per pound of body weight. In dogs over 50 pounds, a dosage of 0.5 mg. per pound administered intramuscularly may provide sufficient sedation and/or analgesia for most procedures.

(iii) To wild deer and elk from a solution containing 100 milligrams of xylazine (as xylazine hydrochloride) per milliliter, intramuscularly, by hand syringe or syringe dart, in the heavy muscles of the croup or shoulder as follows:

(a) Fallow deer, 2 to 4 milligrams per pound.

(b) Mule deer, sika deer, and whitetailed deer, 1 to 2 milligrams per pound.

(c) Elk, 0.25 to 0.5 milligram per pound.

§ 522.2670

(3) Not to be administered to food-producing animals.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 41 FR 24884, June 21, 1976; 41 FR 28265, July 9, 1976; 53 FR 4848, Feb. 18, 1988; 53 FR 23608, June 23, 1988; 53 FR 40728, Oct. 18, 1988; 55 FR 18724, May 4, 1990; 55 FR 32616, Aug. 10, 1990; 59 FR 14367, Mar. 28, 1994; 60 FR 33110, June 27, 1995; 60 FR 35122 and 35123, July 6, 1995; 61 FR 46548, Sept. 4, 1996; 62 FR 35077, June 30, 1997]

§ 522.2670 Yohimbine injectable.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains either 2 or 5 milligrams of yohimbine (as hydrochloride).

(b) *Sponsor.* See 061690 in § 510.600(c) of this chapter for use of 2 milligrams per milliliter solution in dogs.

(1) *Amount.* 0.05 milligram per pound (0.11 milligram per kilogram) of body weight.

(2) *Indications for use.* To reverse the effects of xylazine in dogs.

(3) *Limitations.* For intravenous use in dogs only. Not for use in food-producing animals. Safety of use in pregnant dogs or in dogs intended for breeding has not been established. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(c) *Sponsor.* See 053923 in § 510.600(c) of this chapter for use of 5 milligrams per milliliter solution in deer and elk.

(1) *Amount.* 0.2 to 0.3 milligram per kilogram of body weight.

(2) *Indications for use.* As an antagonist to xylazine sedation in free ranging or confined members of the family Cervidae (deer and elk).

(3) *Limitations.* For intravenous use only. Do not use in domestic food-producing animals. Do not use for 30 days before or during hunting season. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[58 FR 8543, Feb. 16, 1993, as amended at 60 FR 57832, Nov. 22, 1995]

§ 522.2680 Zeranol.

(a) *Specifications.* Each pellet contains 12 milligrams of zeranol.

(b) *Sponsor.* See 000061 in § 510.600(c) of this chapter.

21 CFR Ch. I (4–1–99 Edition)

(c) *Related tolerances.* See § 556.760 of this chapter.

(d) *Conditions of use.* For use as a subcutaneous ear implant as follows:

(1) *Beef cattle*—(i) *Amount.* 36 milligrams (three 12-milligram pellets) per animal.

(ii) *Indications for use*—(A) For increased rate of weight gain and improved feed conversion in weaned beef calves, growing beef cattle, feedlot steers, and feedlot heifers.

(B) For increased rate of weight gain in suckling calves.

(iii) *Limitations.* Implant subcutaneously in ear only. Do not use in bulls intended for reproduction or in dairy animals. Do not use before 1 month of age or after weaning in heifers intended for reproduction.

(2) *Feedlot lambs*—(i) *Amount.* 12 milligrams (1 pellet) per animal.

(ii) *Indications for use.* For increased rate of weight gain and improved feed conversion.

(iii) *Limitations.* Implant subcutaneously in ear only. Do not use in breeding animals. Do not implant animals within 40 days of slaughter.

(3) *Steers*—(i) *Amount.* 72 milligrams (six 12-milligram pellets) per animal.

(ii) *Indications for use.* For increased rate of weight gain in steers fed in confinement for slaughter.

(iii) *Limitations.* Implant subcutaneously in ear only.

[59 FR 19639, Apr. 25, 1994; 60 FR 26360, May 17, 1995, as amended at 62 FR 61625, Nov. 19, 1997]

PART 524—OPHTHALMIC AND TOPICAL DOSAGE FORM NEW ANIMAL DRUGS

Sec.

524.86 Amitraz liquid.

524.154 Bacitracin or bacitracin zinc-neomycin sulfate-polymyxin B sulfate ophthalmic ointment.

524.155 Bacitracin zinc-polymyxin B sulfate-neomycin sulfate-hydrocortisone or hydrocortisone acetate ophthalmic ointment.

524.321 Cephalonium, polymyxin B sulfate, flumethasone, iodochlorhydroxyquin, piperocaine hydrochloride topical-otic ointment.

524.390 Chloramphenicol ophthalmic and topical dosage forms.

Food and Drug Administration, HHS

Pt. 524

- 524.390a Chloramphenicol ophthalmic ointment.
- 524.390b Chloramphenicol ophthalmic solution.
- 524.390c Chloramphenicol-prednisolone-tetracaine-squalane topical suspension.
- 524.390d Chloramphenicol-prednisolone ophthalmic ointment.
- 524.402 Chlorhexidine diacetate ointment.
- 524.450 Clotrimazole cream.
- 524.463 Copper naphthenate solution.
- 524.520 Cuprimyxin cream.
- 524.575 Cyclosporine ophthalmic ointment.
- 524.660 Dimethyl sulfoxide ophthalmic and topical dosage forms.
- 524.660a Dimethyl sulfoxide solution.
- 524.660b Dimethyl sulfoxide gel.
- 524.770 Doramectin.
- 524.814 Eprinomectin.
- 524.900 Famphur.
- 524.920 Fenthion.
- 524.960 Flumethasone, neomycin sulfate, and polymyxin B sulfate ophthalmic solutions.
- 524.981 Fluocinolone acetonide ophthalmic and topical dosage forms.
- 524.981a Fluocinolone acetonide cream.
- 524.981b Fluocinolone acetonide solution.
- 524.981c Fluocinolone acetonide, neomycin sulfate cream.
- 524.981d Fluocinolone acetonide, dimethyl sulfoxide solution.
- 524.981e Fluocinolone acetonide, dimethyl sulfoxide otic solution.
- 524.1005 Furazolidone aerosol powder.
- 524.1044 Gentamicin sulfate ophthalmic and topical dosage forms.
- 524.1044a Gentamicin ophthalmic solution.
- 524.1044b Gentamicin sulfate, betamethasone valerate otic solution.
- 524.1044c Gentamicin ophthalmic ointment.
- 524.1044d Gentamicin sulfate, betamethasone valerate ointment.
- 524.1044e Gentamicin sulfate spray.
- 524.1044f Gentamicin sulfate, betamethasone valerate topical spray.
- 524.1044g Gentamicin sulfate, betamethasone valerate, clotrimazole ointment.
- 524.1193 Ivermectin pour-on.
- 524.1200 Kanamycin ophthalmic and topical dosage forms.
- 524.1200a Kanamycin ophthalmic ointment.
- 524.1200b Kanamycin ophthalmic aqueous solution.
- 524.1204 Kanamycin sulfate, calcium amphotycin, and hydrocortisone acetate.
- 524.1240 Levamisole.
- 524.1376 2-Mercaptobenzothiazole solution.
- 524.1443 Miconazole nitrate cream; miconazole nitrate lotion; miconazole nitrate spray.
- 524.1451 Moxidectin.
- 524.1465 Mupirocin ointment.
- 524.1484 Neomycin sulfate ophthalmic and topical dosage forms.
- 524.1484a Neomycin sulfate ophthalmic ointment.
- 524.1484b Neomycin sulfate, isoflupredone acetate, tetracaine hydrochloride, and myristyl-gamma-picolinium chloride, topical powder.
- 524.1484c Neomycin sulfate, isoflupredone acetate, tetracaine hydrochloride ointment.
- 524.1484d Neomycin sulfate, hydrocortisone acetate, tetracaine hydrochloride ear ointment.
- 524.1484e Neomycin sulfate and polymyxin B sulfate ophthalmic solution.
- 524.1484f Neomycin sulfate, prednisolone acetate, tetracaine hydrochloride eardrops.
- 524.1484g Neomycin sulfate-thiabendazole-dexamethasone solution.
- 524.1484h Neomycin, penicillin, polymyxin, hydrocortisone suspension.
- 524.1484i Neomycin sulfate, hydrocortisone acetate, sterile ointment.
- 524.1484j [Reserved]
- 524.1484k Neomycin sulfate, prednisolone, tetracaine, and squalane topical-otic suspension.
- 524.1580 Nitrofurazone ophthalmic and topical dosage forms.
- 524.1580a [Reserved]
- 524.1580b Nitrofurazone ointment.
- 524.1580c Nitrofurazone soluble powder.
- 524.1580d [Reserved]
- 524.1580e Nitrofurazone ointment with butacaine sulfate.
- 524.1600 Nystatin ophthalmic and topical dosage forms.
- 524.1600a Nystatin, neomycin, thiostrepton, and triamcinolone acetonide ointment.
- 524.1600b Nystatin, neomycin, thiostrepton, and triamcinolone acetonide ophthalmic ointment.
- 524.1662 Oxytetracycline hydrochloride ophthalmic and topical dosage forms.
- 524.1662a Oxytetracycline hydrochloride and hydrocortisone spray.
- 524.1662b Oxytetracycline hydrochloride, polymyxin B sulfate ophthalmic ointment.
- 524.1742 N-(Mercaptomethyl) phthalimide S-(O,O-dimethyl phosphorodithioate) emulsifiable liquid.
- 524.1880 Prednisolone-neomycin sulfate ophthalmic ointment.
- 524.1881 Prednisolone acetate ophthalmic and topical dosage forms.
- 524.1881a [Reserved]
- 524.1881b Prednisolone acetate-neomycin sulfate sterile suspension.
- 524.1883 Prednisolone sodium phosphate-neomycin sulfate ophthalmic ointment.
- 524.1982 Proparacaine hydrochloride ophthalmic solution.
- 524.2101 Selenium disulfide suspension.
- 524.2140 Squalane, pyrethrins and piperonyl butoxide.
- 524.2350 Tolnaftate cream.
- 524.2481 Triamcinolone acetonide cream.

§ 524.86

524.2620 Liquid crystalline trypsin, Peru balsam, castor oil.

524.2640 Tylosin, neomycin eye powder.

AUTHORITY: 21 U.S.C. 360b.

SOURCE: 40 FR 13873, Mar. 27, 1975, unless otherwise noted.

§ 524.86 Amitraz liquid.

(a) *Specifications.* Amitraz liquid contains 19.9 percent amitraz in an organic solvent.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Indications for use.* For dogs for the treatment of generalized demodicosis (*Demodex canis*).

(2) *Amount.* One 10.6 milliliter bottle per 2 gallons of warm water (250 parts per million) for each treatment, for a total of 3 to 6 treatments, 14 days apart.

(3) *Limitations.* Continue treatment until no viable mites are found in skin scrapings at 2 successive treatments, or until 6 treatments have been applied. Do not use for treatment of localized demodicosis or scabies. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[47 FR 18589, Apr. 30, 1982]

§ 524.154 Bacitracin or bacitracin zinc-neomycin sulfate-polymyxin B sulfate ophthalmic ointment.

(a) *Sponsor.* To firms identified in § 510.600(c) of this chapter as follows:

(1) To 000009; each gram contains 500 units of bacitracin, 3.5 milligrams of neomycin, and 10,000 units of polymyxin B.

(2) To 000061 and 025463; each gram contains 400 units of bacitracin zinc, 3.5 milligrams of neomycin, and 10,000 units of polymyxin B sulfate.

(b) *Conditions of use. Dogs and Cats.* (1) *Amount.* Apply a thin film over the cornea 3 or 4 times daily.

(2) *Indications for use.* Treatment of superficial bacterial infections of the eyelid and conjunctiva of dogs and cats when due to susceptible organisms.

(3) *Limitations.* Laboratory tests should be conducted including in vitro culturing and susceptibility tests on samples collected prior to treatment. Federal law restricts this drug to use

21 CFR Ch. I (4–1–99 Edition)

by or on the order of a licensed veterinarian.

[57 FR 37333, Aug. 18, 1992, as amended at 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 524.155 Bacitracin zinc-polymyxin B sulfate-neomycin sulfate-hydrocortisone or hydrocortisone acetate ophthalmic ointment.

(a) *Sponsor.* To firms identified in § 510.600(c) of this chapter as follows:

(1) To 000061; each gram of ointment contains 400 units of bacitracin zinc, 10,000 units of polymyxin B sulfate, 5 milligrams of neomycin sulfate (equivalent to 3.5 milligrams of neomycin base), and 10 milligrams of hydrocortisone.

(2) To 025463; each gram of ointment contains 400 units of bacitracin zinc, 10,000 units of polymyxin B sulfate, 5 milligrams of neomycin sulfate (equivalent to 3.5 milligrams of neomycin base), and 10 milligrams of hydrocortisone acetate.

(b) *Conditions of use. Dogs and cats.* (1) *Amount.* Apply a thin film over the cornea three or four times daily.

(2) *Indications for use.* For treating acute or chronic conjunctivitis caused by susceptible organisms.

(3) *Limitations.* All topical ophthalmic preparations containing corticosteroids with or without an antimicrobial agent are contraindicated in the initial treatment of corneal ulcers. They should not be used until the infection is under control and corneal regeneration is well underway. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37333, Aug. 18, 1992, as amended at 61 FR 8873, Mar. 6, 1996; 62 FR 61626, Nov. 19, 1997]

§ 524.321 Cephalonium, polymyxin B sulfate, flumethasone, iodochlorhydroxyquin, piperocaine hydrochloride topical-otic ointment.

(a) *Specifications.* Each gram of the drug contains 10 milligrams cephalonium, 5,000 units polymyxin B sulfate, 0.25 milligram flumethasone, 30 milligrams iodochlorhydroxyquin, and 40 milligrams piperocaine hydrochloride in a suitable and harmless ointment base.

Food and Drug Administration, HHS

§ 524.390c

(b) *Sponsor.* See No. 000986 in § 510.600(c) of this chapter.

(c) *Conditions of use.* The drug is recommended for dermal and otic use on dogs and cats for the treatment of the following conditions when complicated by bacteria, yeast, or fungus: Pyodermatitis, allergic dermatitis, dermatophytosis, nonspecific pruritus, and external otitis. For mild inflammations a periodic treatment of applying from once daily to twice weekly may be indicated. In severe conditions apply once or twice daily when continuous treatment may be indicated. Dosage per treatment should not exceed 300 milligrams of the ointment. For otic use treatment should not exceed a total of 12 days. For use only by or on the order of a licensed veterinarian.

§ 524.390 Chloramphenicol ophthalmic and topical dosage forms.

§ 524.390a Chloramphenicol ophthalmic ointment.

(a) *Specifications.* Each gram contains 10 milligrams chloramphenicol in a petrolatum base.

(b) *Sponsor.* See Nos. 000856 and 025463 in § 510.600(c) of this chapter for use as in paragraph (c)(1)(i) of this section. See No. 017030 for use as in paragraph (c)(1)(ii) of this section.

(c) *Conditions of use. Dogs and cats.* (1) *Amount.* Apply as follows:

(i) Every 3 hours around the clock for 48 hours after which night instillations may be omitted.

(ii) Four to six times daily to affected eye for the first 72 hours depending upon the severity of the condition. A small amount of ointment should be placed in the lower conjunctival sac.

(2) *Indications for use.* Treatment of bacterial conjunctivitis caused by pathogens susceptible to chloramphenicol.

(3) *Limitations.* Continue treatment for 48 hours (2 days) after eye appears normal. Therapy for cats should not exceed 7 days. Prolonged use in cats may produce blood dyscrasias. If improvement is not noted in a few days a change of therapy should be considered. When infection may be cause of disease, especially in purulent or catarrhal conjunctivitis, attempts should be made to determine through susceptibility testing, which antibiotics will

be effective prior to applying ophthalmic preparations. This chloramphenicol product must not be used in animals producing meat, eggs, or milk. The length of time that residues persist in milk or tissues has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37333, Aug. 18, 1992]

§ 524.390b Chloramphenicol ophthalmic solution.

(a) *Specifications.* Each milliliter contains 5 milligrams of chloramphenicol.

(b) *Sponsor.* See No. 017030 in § 510.600(c) of this chapter.

(c) *Conditions of use. Dogs and Cats.* (1) *Amount.* Apply one or two drops, 4 to 6 times a day for the first 72 hours, depending upon the severity of the condition. Intervals between applications may be increased after the first 2 days.

(2) *Indications for use.* Treatment of bacterial conjunctivitis caused by organisms susceptible to chloramphenicol. Therapy should be continued for 48 hours after the eye appears normal.

(3) *Limitations.* Therapy for cats should not exceed 7 days. As with other antibiotics, prolonged use may result in overgrowth of nonsusceptible organisms. If superinfection occurs, or if clinical improvement is not noted within a reasonable period, discontinue use, and institute appropriate therapy. Prolonged use in cats may produce blood dyscrasias. Chloramphenicol products must not be used in meat-, egg-, or milk-producing animals. The length of time that residues persist in milk or tissues has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37333, Aug. 18, 1992]

§ 524.390c Chloramphenicol-prednisolone-tetracaine-squalane topical suspension.

(a) *Specification.* Each milliliter contains 4.2 milligrams of chloramphenicol, 1.7 milligrams of prednisolone, 4.2 milligrams of tetracaine, and 0.21 milliliter of squalane.

(b) *Sponsor.* See No. 017030 in § 510.600(c) of this chapter.

§ 524.390d

(c) *Conditions of use. Dogs and cats.* (1) *Amount.* Apply two or three times daily or as needed for not more than 7 days. Severe infections should be supplemented by systemic therapy.

(2) *Indications for use.* Treatment of acute otitis externa and pyoderma (acute moist dermatitis, vulvar fold dermatitis, lip fold dermatitis, interdigital dermatitis, and juvenile dermatitis) in dogs and cats.

(3) *Limitations.* The drug must not be used in the eyes. Prolonged use in cats may produce blood dyscrasias. Laboratory tests should be conducted, including in vitro culturing and susceptibility tests on samples collected prior to treatment. Chloramphenicol products must not be used in meat-, egg-, or milk-producing animals. The length of time that residues persist in milk or tissues has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37334, Aug. 18, 1992]

§ 524.390d Chloramphenicol-prednisolone ophthalmic ointment.

(a) *Specifications.* Each gram contains 10 milligrams of chloramphenicol and 2.5 milligrams of prednisolone acetate.

(b) *Sponsor.* See No. 017030 in § 510.600(c) of this chapter.

(c) *Conditions of use. Dogs and cats.* (1) *Amount.* Apply 4 to 6 times daily to the affected eye for the first 72 hours depending upon the severity of the condition. Continue treatment for 48 hours after the eye appears normal.

(2) *Indications for use.* Treatment of bacterial conjunctivitis and ocular inflammation caused by organisms susceptible to chloramphenicol.

(3) *Limitations.* Therapy for cats should not exceed 7 days, prolonged use in cats may produce blood dyscrasia. As with other antibiotics, prolonged use may result in overgrowth of non-susceptible organisms. If superinfection occurs or if clinical improvement is not noted within a reasonable period, discontinue use and institute appropriate therapy. All topical ophthalmic preparations containing corticosteroids, with or without an antimicrobial agent, are contraindicated in the initial treatment of corneal ulcers. They should not be used until the infection is under control and

21 CFR Ch. I (4–1–99 Edition)

corneal regeneration is well underway. Chloramphenicol products must not be used in meat-, egg-, or milk-producing animals. The length of time that residues persist in milk or tissues has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37334, Aug. 18, 1992]

§ 524.402 Chlorhexidine diacetate ointment.

(a) *Specifications.* The product contains 1 percent of chlorhexidine diacetate in an ointment base.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used as a topical antiseptic ointment for surface wounds on dogs, cats, and horses.¹

(2) The wound area is carefully cleansed and the drug is applied daily.¹

(3) The drug is not to be used in horses intended for use as food.¹

§ 524.450 Clotrimazole cream.

(a) *Specifications.* Each gram of cream contains 10 milligrams of clotrimazole.

(b) *Sponsor.* See 000859 in § 510.600(c).

(c) *Conditions of use—(1) Amount.* Apply ¼-inch ribbon of cream per square inch of lesion once daily for 2 to 4 weeks.

(2) *Indications for use.* For the treatment of fungal infections of dogs and cats caused by *Microsporum canis* and *Trichophyton mentagrophytes*.

(3) *Limitations.* Wash hands thoroughly after use to avoid spread of infection. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 48128, July 18, 1980]

§ 524.463 Copper naphthenate solution.

(a) *Specifications.* The drug contains 37.5 percent copper naphthenate in a suitable solvent.

(b) *Sponsor.* See Nos. 000856 and 017135 in § 510.600(c) of this chapter.

¹These conditions are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

Food and Drug Administration, HHS

§ 524.660a

(c) *Conditions of use—Horses and ponies*—(1) *Amount*. Apply daily to affected hooves until fully healed.

(2) *Indications for use*. As an aid in treating horses and ponies for thrush caused by organisms susceptible to copper naphthenate.

(3) *Limitations*. Use on horses and ponies only. Remove debris and necrotic material before applying. Avoid contact around eyes. Do not use on animals that are raised for food production. Do not contaminate feed. Do not allow runoff of excess drug into hair because contact with the drug may cause some hair loss.

[47 FR 4250, Jan. 29, 1982]

§ 524.520 Cuprimyxin cream.

(a) *Specifications*. The drug contains 0.5 percent cuprimyxin (6-methoxy-1-phenazinol 5, 10-dioxide, cupric complex) in an aqueous cream base.

(b) *Sponsor*. See No. 000004 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) Cuprimyxin is a broad spectrum antibacterial and antifungal cream for the topical treatment of superficial infections in horses, dogs, and cats caused by bacteria, dermatophytes (*Trichophyton* spp., *Microsporum* spp.) and yeast (*Candida albicans*) affecting skin, hair, and external mucosae.

(2) The cream is applied twice daily to affected areas by rubbing into lesions. Treatment should be continued for a few days after clinical recovery to avoid possible relapses.

(3) After application to cutaneous areas, a change in color from dark green to pink is due to the liberation of free myxin from its copper complex.

(4) If no response is seen within seven days, diagnosis and therapy should be reevaluated. If any adverse local reaction is observed after topical application, discontinue treatment.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 45 FR 56799, Aug. 26, 1980]

§ 524.575 Cyclosporine ophthalmic ointment.

(a) *Specifications*. Each gram of ointment contains 2 milligrams of cyclosporine.

(b) *Sponsor*. See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. Apply a 1/4-inch strip of ointment to the affected eye(s) every 12 hours.

(2) *Indications for use*. For management of chronic keratoconjunctivitis sicca (KCS) and chronic superficial keratitis (CSK) in dogs.

(3) *Limitations*. Place ointment directly on cornea or into the conjunctival sac. Safety of use in puppies, pregnant or breeding animals has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[60 FR 48651, Sept. 20, 1995, as amended at 62 FR 48940, Sept. 18, 1997]

§ 524.660 Dimethyl sulfoxide ophthalmic and topical dosage forms.

§ 524.660a Dimethyl sulfoxide solution.

(a) *Specifications*. Dimethyl sulfoxide contains 90 percent of dimethyl sulfoxide and 10 percent of water.

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) It is used or intended for use as a topical application to reduce acute swelling due to trauma:

(i) In horses administered 2 or 3 times daily in an amount not to exceed 100 milliliters per day. Total duration of therapy should not exceed 30 days.

(ii) In dogs administered 3 or 4 times daily in an amount not to exceed 20 milliliters per day. Total duration of therapy should not exceed 14 days.

(2) Not for use in horses and dogs intended for breeding purposes nor in horses slaughtered for food. Other topical medications should only be used when the dimethyl sulfoxide treated area is thoroughly dry. Do not administer by any other route.

(3) For use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 61 FR 5507, Feb. 13, 1996]

§ 524.660b

§ 524.660b Dimethyl sulfoxide gel.

(a) *Specifications.* Dimethyl sulfoxide gel, veterinary contains 90 percent dimethyl sulfoxide in an aqueous gel.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Indications for use.* For use on horses and dogs as a topical application to reduce acute swelling due to trauma.

(2) *Amount*—(i) *Horses.* Administer 2 or 3 times daily in an amount not to exceed 100 grams per day. Total duration of therapy should not exceed 30 days.

(ii) *Dogs.* Administer 3 or 4 times daily in an amount not to exceed 20 grams per day. Total duration of therapy should not exceed 14 days.

(3) *Limitations.* Do not use in horses and dogs intended for breeding purposes or in horses slaughtered for food. Restricted to topical use on horses and dogs only. Due to rapid penetrating ability of dimethyl sulfoxide, rubber gloves should be worn when applying the drug. No other medications should be present on the skin prior to application of the drug. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 48 FR 56205, Dec. 20, 1983; 61 FR 5507, Feb. 13, 1996]

§ 524.770 Doramectin.

(a) *Specifications.* Each milliliter of solution contains 5 milligrams of doramectin.

(b) *Sponsor.* See 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.225 of this chapter.

(d) *Conditions of use*—Cattle—(1) *Amount.* 5 milligrams per 10 kilograms (5 milligrams per 22 pounds).

(2) *Indications for use.* For treatment and control of gastrointestinal roundworms, lungworms, eyeworms, grubs, biting and sucking lice, horn flies, and mange mites, and to control infections and to protect from reinfection with *Cooperia oncophora* and *Dictyocaulus viviparus* for 21 days, and *Ostertagia ostertagia*, *C. punctata*, and *Oesophagostomum radiatum* for 28 days after treatment.

(3) *Limitations.* Administer as a single dose. Do not slaughter cattle within 45

21 CFR Ch. I (4–1–99 Edition)

days of latest treatment. Not for use in female dairy cattle 20 months of age or older. Do not use in calves to be processed for veal. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[62 FR 65753, Dec. 16, 1997, as amended at 63 FR 68183, Dec. 10, 1998]

§ 524.814 Eprinomectin.

(a) *Specifications.* Each milliliter contains 5 milligrams of eprinomectin.

(b) *Sponsor.* See No. 000006 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.227 of this chapter.

(d) *Conditions of use*—(1) *Amount.* One milliliter (5 milligrams) per 10 kilograms of body weight (500 micrograms per kilogram).

(2) *Indications for use.* The drug is used in beef and dairy cattle for treatment and control of gastrointestinal roundworms (*Haemonchus placei* (adult and L4), *Ostertagia ostertagi* (adult and L4, including inhibited L4), *Trichostrongylus axei* (adult and L4), *T. colubriformis* (adult and L4), *T. longispicularis* (adult), *Cooperia oncophora* (adult and L4), *C. punctata* (adult and L4), *C. surnabada* (adult and L4), *Nematodirus helvetianus* (adult and L4), *Bunostomum phlebotomum* (adult and L4), *Oesophagostomum radiatum* (adult and L4), *Strongyloides papillosus* (adults), *Trichuris* spp. (adults)); lungworms (*Dictyocaulus viviparus*, adult and L4); cattle grubs (all parasitic stages *Hypoderma lineatum*, *H. bovis*); lice (*Damalinia bovis*, *Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*); mange mites (*Chorioptes bovis*, *Sarcoptes scabiei*); and horn flies (*Haematobia irritans*). Controls and protects from reinfection of *D. viviparus* for 21 days after treatment and *H. irritans* for 7 days after treatment.

(3) *Limitations.* Apply topically along backbone from withers to tailhead. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[62 FR 33997, June 24, 1997, as amended at 63 FR 59715, Nov. 5, 1998]

§ 524.900 Famphur.

(a) *Chemical name.* O,O- Dimethyl O-[p-(dimethylsulfamoyl)phenyl] phosphorothioate.

(b) *Specifications.* The drug is in liquid form containing 13.2 percent famphur.

(c) *Sponsor.* See Nos. 000061 and 060594 in § 510.600(c) of this chapter.

(d) *Special considerations.* Do not use on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(e) *Related tolerances.* See § 556.273 of this chapter.

(f) *Conditions of use.* (1) The drug is used as a pour-on formulation for the control of cattle grubs and to reduce cattle lice infestations.

(2) It is used at the rate of 1 ounce per 200 pounds body weight, not to exceed a total dosage of 4 ounces, applied from the shoulder to the tail head as a single treatment. It is applied as soon as possible after heel fly activity ceases. Do not use on lactating dairy cows or dry dairy cows within 21 days of freshening, calves less than 3 months old, animals stressed from castration, over-excitement or dehorning, sick or convalescent animals. Animals may become dehydrated and under stress following shipment. Do not treat until they are in good condition. Brahman and Brahman crossbreeds are less tolerant of cholinesterase-inhibiting insecticides than other breeds. Do not treat Brahman bulls.

(3) Do not slaughter within 35 days after treatment. Swine should be eliminated from area where run-off occurs.

[40 FR 13873, Mar. 27, 1975, as amended at 49 FR 34352, Aug. 30, 1984; 57 FR 7652, Mar. 4, 1992; 59 FR 28769, June 3, 1994; 62 FR 55161, Oct. 23, 1997; 62 FR 61626, Nov. 19, 1997]

§ 524.920 Fenthion.

(a) *Chemical name.* O,O-Dimethyl O-[4-(methylthio)-m-tolyl] phosphorothioate.

(b) *Specifications.* (1) The drug is in a liquid form containing 3 percent of fenthion.

(2) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(3) *Special considerations.* Do not use on animals simultaneously or within a few days before or after treatment with

or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(4) *Related tolerances.* See 40 CFR 180.214.

(5) *Conditions of use.* (i) The drug is used as a pour-on formulation for the control of grubs and lice in beef and nonlactating cattle.

(ii) It is used at the rate of one-half fluid ounce per 100 pounds of body weight placed on the backline of the animal. Only one application per season should be made for grub control and this will also provide initial control of lice. A second application for lice control may be made if animals become reinfested, but no sooner than 35 days after the first treatment. Proper timing of treatment is important for grub control; cattle should be treated as soon as possible after heel-fly activity ceases. Cattle should not be slaughtered within 35 days following a single treatment. If a second application is made for lice control, cattle should not be slaughtered within 45 days of the second treatment. The drug must not be used within 28 days of freshening of dairy cattle. If freshening should occur within 28 days after treatment, do not use milk as human food for the balance of the 28-day interval. Do not treat lactating dairy cattle; calves less than 3 months old; or sick, convalescent, or stressed livestock. Do not treat cattle for 10 days before or after shipping, weaning, or dehorning or after exposure to contagious infectious diseases.

(c) *Specifications.* (1) The drug is in a liquid form containing 20 percent fenthion.

(2) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(3) *Special considerations.* Do not use on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(4) *Related tolerances.* See 40 CFR 180.214.

(5) *Conditions of use.* (i) The drug is used for control of cattle grubs and as an aid in controlling lice on beef cattle and on dairy cattle not of breeding age.

(ii) It is applied as a single application placed on the backline of animals as follows:

§ 524.960

21 CFR Ch. I (4–1–99 Edition)

Weight of animal	Dosage (milliliters)
150 to 300 lb	4
301 to 600 lb	8
601 to 900 lb	12
901 to 1,200 lb	16
Over 1,200 lb	20

For most effective results, cattle should be treated as soon as possible after heel-fly activity ceases. Host-parasite reactions such as bloat, salivation, staggering and paralysis may sometimes occur when cattle are treated while the common cattle grub (*Hypoderma lineatum*) is in the gullet, or while the northern cattle grub (*H. bovis*) is in the area of the spinal cord. Cattle should be treated before these stages of grub development. Consult your veterinarian, extension livestock specialist, or extension entomologist regarding the timing of treatment. If it is impossible to determine the area from which the cattle came and/or exact stage of the grubs, it is recommended that the cattle receive only a maintenance ration of low-energy feed during the treatment period. This lessens the likelihood of severe bloat which may occur in cattle on full feed when the common grub is killed while in the gullet. A second application is required for animals heavily infested with lice or for those which become re-infested. A second application should be made no sooner than 35 days after the first treatment.

(iii) Do not treat dairy cattle of breeding age; calves less than 3 months old; sick, convalescent, or severely stressed livestock.

(iv) Do not treat cattle for 10 days before or after shipping, weaning, dehorning, or after exposure to contagious or infectious diseases.

(v) Do not slaughter within 45 days of treatment.

(d) *Specifications.* (1) The drug is a solution containing either 5.6 or 13.8 percent fenthion. Each concentration is available in 2 volumes which are contained in single-dose applicators.

(2) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(3) *Special considerations.* Fenthion is a cholinesterase inhibitor. Do not use this product on dogs simultaneously with or within 14 days before or after treatment with or exposure to cholin-

esterase-inhibiting drugs, pesticides, or chemicals. Do not use with flea or tick collars.

(4) *Conditions of use*—(i) *Amount.* Four to 8 milligrams per kilogram of body weight.

(ii) *Indications for use.* For flea control on dogs only.

(iii) *Limitations.* Apply the contents of the proper size, single-dose tube directly to one spot on the dog's skin. Frequency of repeat treatments depends upon rate of flea reinfestations. Do not use more often than once every 2 weeks. Treatment at 2-week intervals is not to exceed 6 months. Do not use on puppies under 10 weeks of age. Do not use on sick, stressed, or convalescing dogs. Safe use in breeding males has not been established. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 41 FR 16656, Apr. 21, 1976; 42 FR 58741, Nov. 11, 1977; 45 FR 62425, Sept. 19, 1980; 50 FR 19169, May 7, 1985]

§ 524.960 Flumethasone, neomycin sulfate, and polymyxin B sulfate ophthalmic solutions.

(a) *Specifications.* Each milliliter of ophthalmic preparation contains 0.10 milligram flumethasone, 5.0 milligrams neomycin sulfate (3.5 milligrams neomycin base), and 10,000 units of polymyxin B sulfate, with or without hydroxypropyl methylcellulose.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*—(i) *Preparation containing hydroxypropyl methylcellulose.* Dogs: 1 to 2 drops per eye, every 6 hours.

(ii) *Preparation without hydroxypropyl methylcellulose.* Dogs and cats: 2 to 3 drops per eye, every 4 hours.

(2) *Indications for use.* Treatment of the inflammation, edema, and secondary bacterial infections associated with topical ophthalmological conditions of the eye such as corneal injuries, incipient pannus, superficial keratitis, conjunctivitis, acute nongranulomatous anterior uveitis, keratoconjunctivitis, and blepharitis.

(3) *Limitations.* (i) In treating ophthalmological conditions associated with bacterial infections, the drug is contraindicated in those cases in which

Food and Drug Administration, HHS

§ 524.981d

microorganisms are not susceptible to the antibiotics incorporated in the drug.

(ii) The drug is contraindicated in infectious tuberculous lesions of the eye, early acute stages of viral diseases of the cornea and conjunctiva, herpes simplex lesions of the eye, and fungal infections of the conjunctiva and eyelids.

(iii) The usual precautions and contraindications for corticosteroids and adrenocorticoids are applicable with this drug. Corticosteroids may inhibit essential inflammatory responses intrinsic to the fundamental healing mechanism. Adrenocorticoid compounds have been reported to cause an increase in intraocular pressure. Intraocular pressure should be checked frequently. Ocular reexaminations should be made at frequent intervals during long-term therapy.

(iv) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 16012, Mar. 16, 1979, as amended at 61 FR 5507, Feb. 13, 1996]

§ 524.981 Fluocinolone acetonide ophthalmic and topical dosage forms.

§ 524.981a Fluocinolone acetonide cream.

(a) *Specifications.* The drug contains 0.025 percent fluocinolone acetonide.

(b) *Sponsor.* See No. 099207 in §510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is indicated for the relief of pruritus and inflammation associated with certain superficial acute and chronic dermatoses in dogs. It is used in the treatment of allergic and acute moist dermatitis and for the relief of superficial inflammation caused by chemical and physical abrasions and burns.

(2) A small amount is applied to the affected area two or three times daily.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 62 FR 40932, July 31, 1997]

§ 524.981b Fluocinolone acetonide solution.

(a) *Specifications.* The drug contains 0.01 percent fluocinolone acetonide in propylene glycol with citric acid.

(b) *Sponsor.* See No. 099207 in §510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is indicated for the relief of pruritus and inflammation associated with otitis externa and certain superficial acute and chronic dermatoses in the dog. It is also indicated for the relief of pruritus and inflammation associated with acute otitis externa and certain superficial acute and chronic dermatoses in the cat.

(2) A small amount of solution is applied to the affected area 2 or 3 times daily.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 62 FR 40932, July 31, 1997]

§ 524.981c Fluocinolone acetonide, neomycin sulfate cream.

(a) *Specifications.* The drug contains 0.025 percent fluocinolone acetonide and 0.5 percent neomycin sulfate (0.35 percent neomycin base).

(b) *Sponsor.* See No. 099207 in §510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in the relief of pruritus and inflammation associated with superficial acute and chronic dermatoses in dogs. It is used in the treatment of such conditions as allergic and acute moist dermatoses and nonspecific dermatoses in dogs. It is used in the treatment of wound infections in dogs and cats.

(2) A small amount is applied to the infected area two or three times daily.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 62 FR 40932, July 31, 1997]

§ 524.981d Fluocinolone acetonide, dimethyl sulfoxide solution.

(a) *Specifications.* Each milliliter of solution contains 0.01 percent fluocinolone acetonide and 20 percent

§ 524.981e

dimethyl sulfoxide with propylene glycol and citric acid.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs for the relief of impaction commonly present in apparently normal anal sacs, for the reversal of inflammatory changes associated with abnormal anal sacs, and to counteract the offensive odor of anal sac secretions.

(2) It is administered by instillation of 1 to 2 milliliters into each anal sac following expression of anal sac contents. It may be necessary to repeat treatment at 60-day intervals to maintain an odor-free state. The total dosage used should not exceed 2 milliliters per anal sac per treatment.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 61 FR 5507, Feb. 13, 1996]

§ 524.981e Fluocinolone acetonide, dimethyl sulfoxide otic solution.

(a) *Specifications.* Each milliliter of solution contains 0.01 percent of fluocinolone acetonide in 60 percent dimethyl sulfoxide with propylene glycol and citric acid.

(b) *Sponsor.* See No. 000033 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs for the relief of pruritis and inflammation associated with acute and chronic otitis.

(2) It is administered at 4 to 6 drops (0.2 milliliter) twice daily into the ear canal for a maximum period of 14 days. The total dosage used should not exceed 17 milliliters. The ear canal should be cleansed by some appropriate method prior to instillation of the solution and the ear should be massaged gently following instillation.

(3) There should be careful initial evaluation and followup of infected ears. Incomplete response or exacerbation of corticosteroid-responsive lesions may be due to the presence of an infection which requires identification or antibiotic sensitivity testing, and the use of the appropriate antimicrobial agent. As with any corticosteroid, animals with a generalized infection should not be treated with this product without proper sup-

21 CFR Ch. I (4-1-99 Edition)

portive antimicrobial therapy. Preparations with dimethyl sulfoxide should not be used in pregnant animals. For use by or on the order of a licensed veterinarian.

§ 524.1005 Furazolidone aerosol powder.

(a) *Specifications.* The product contains either 4 or 10 percent furazolidone in inert dispersing agent and propellant.

(b) *Sponsors.* (1) See No. 000069 in § 510.600(c) of this chapter for use of the 10 percent product as in paragraphs (c)(2) (i) through (iii) of this section.

(2) See No. 017135 for use of the 4 percent product as in paragraph (c)(2)(iv) of this section.

(c) *Conditions of use*—(1) *Amount.* Hold container about 6 to 12 inches from the eye or affected area and apply only enough powder to impart a light yellow color.

(2) *Indications of use*—(i) *Dogs.* For treatment or prevention of bacterial infection of superficial wounds, abrasions, lacerations, and pyogenic dermatitis.

(ii) *Horses.* For treatment or prevention of bacterial infection of superficial wounds, abrasions, lacerations, and following firing (heat or electrocautery).

(iii) *Cattle.* For treatment of bacterial infections of the bovine eye and for treatment and to reduce the incidence of additional cases of infectious bovine keratoconjunctivitis (pink eye) caused by *Moraxella bovis*.

(iv) *Horses and ponies.* For treatment or prevention of bacterial infection of superficial wounds, abrasions, and lacerations caused by *Staphylococcus aureus*, *Streptococcus spp.* and *Proteus spp.* sensitive to furazolidone.

(3) *Limitations.* For topical application in horses, ponies, and dogs: Clean affected area thoroughly, apply drug once or twice daily, and repeat treatment as required. For treatment of bacterial infections of the bovine eye and infectious bovine keratoconjunctivitis (pink eye) caused by *Moraxella bovis*: Treat affected eyes once daily on each of 3 to 5 consecutive days; to reduce incidence of additional cases of infectious keratoconjunctivitis

also medicate unaffected eyes. Evidence of clinical improvement of bovine eye infections should be noticeable after 5 treatments; if not, reconsult veterinarian. Use only as recommended by a veterinarian in treatment of puncture wounds, wounds requiring surgical debridement or suturing, those of a chronic nature involving proud flesh, generalized and chronic infections of the skin, and those skin conditions associated with intense itching. If redness, irritation, or swelling persists or increases, discontinue use and reconsult veterinarian. Not for use in horses intended for food.

[45 FR 49543, July 25, 1980, as amended at 50 FR 30153, July 24, 1985; 56 FR 50653, Oct. 8, 1991; 57 FR 31314, July 15, 1992; 60 FR 55659, Nov. 2, 1995]

§ 524.1044 Gentamicin sulfate ophthalmic and topical dosage forms.

§ 524.1044a Gentamicin ophthalmic solution.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains gentamicin sulfate equivalent to 3 milligrams of gentamicin.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs and cats for the topical treatment of infections of the conjunctiva caused by susceptible bacteria.

(2) Administer 1 or 2 drops into the conjunctival sac 2 to 4 times a day.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 14189, Apr. 2, 1976, as amended at 52 FR 7832, Mar. 13, 1987]

§ 524.1044b Gentamicin sulfate, betamethasone valerate otic solution.

(a) *Specifications.* Each cubic centimeter of solution contains gentamicin sulfate equivalent to 3 milligrams of gentamicin base and betamethasone valerate equivalent to 1 milligram of betamethasone alcohol.

(b) *Sponsor.* See Nos. 000061 and 051259 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used or indicated for use in dogs in the treatment of acute and chronic otitis

externa caused by bacteria sensitive to gentamicin; the drug is also used or indicated for use in dogs and cats in the treatment of superficial infected lesions caused by bacteria sensitive to gentamicin.

(2)(i) For the treatment of acute and chronic canine otitis externa caused by bacteria sensitive to gentamicin, the drug is administered by instillation of 3 to 8 drops of solution into the ear canal twice daily for 7 to 14 days. Duration of treatment will depend upon the severity of the condition and the response obtained. The duration of treatment and/or frequency of the dosage may be reduced but care should be taken not to discontinue therapy prematurely. The external ear and ear canal should be properly cleaned and dried before treatment. Remove foreign material, debris, crusted exudates, etc., with suitable nonirritating solutions. Excessive hair should be clipped from the treatment area of the external ear.

(ii) For the treatment of canine and feline superficial infected lesions caused by bacteria sensitive to gentamicin, the lesion and adjacent area should be properly cleaned before treatment. Excessive hair should be removed. A sufficient amount of the drug should be applied to cover the treatment area. The drug should be administered twice daily for 7 to 14 days.

(3) If hypersensitivity to any of the components occurs treatment with this product should be discontinued and appropriate therapy instituted. Concomitant use with other drugs known to induce ototoxicity is not recommended. This preparation should not be used in conditions where corticosteroids are contraindicated. Do not administer parenteral corticosteroids during treatment with this drug. The antibiotic sensitivity of the pathogenic organism should be determined prior to use of this preparation.

(4) For use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975. Redesignated at 41 FR 14189, Apr. 2, 1976, and amended at 52 FR 7832, Mar. 13, 1987; 61 FR 48624, Sept. 16, 1996]

§ 524.1044c Gentamicin ophthalmic ointment.

(a) *Specifications.* Each gram of sterile ointment contains gentamicin sulfate

§ 524.1044d

equivalent to 3 milligrams of gentamicin.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used on dogs and cats for topical treatment of conjunctivitis caused by susceptible bacteria.

(2) Apply approximately a ½-inch strip to the affected eye 2 to 4 times a day.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 14188, Apr. 2, 1976, as amended at 52 FR 7832, Mar. 13, 1987]

§ 524.1044d Gentamicin sulfate, betamethasone valerate ointment.

(a) *Specifications.* Each gram of ointment contains gentamicin sulfate equivalent to 3 milligrams of gentamicin base and betamethasone valerate equivalent to 1 milligram of betamethasone.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used or indicated for use in dogs in the treatment of acute and chronic canine otitis externa and canine infected superficial lesions caused by bacteria sensitive to gentamicin.

(2)(i) For the treatment of acute and chronic canine otitis externa the drug is administered by instillation of 3 to 8 drops into the ear canal twice daily for 7 days. The external ear and ear canal should be properly cleaned and dried before treatment. Remove foreign material, debris, crusted exudates, etc., with suitable nonirritating solutions. Excessive hair should be clipped from the treatment area of the external ear.

(ii) For the treatment of canine infected superficial lesions, the lesion and adjacent area should be properly cleaned before treatment. Excessive hair should be removed. A sufficient amount of the drug should be applied to cover the treatment area. The drug should be administered twice daily for 7 to 14 days.

(3) If hypersensitivity to any of the components occurs, treatment should be discontinued and appropriate therapy instituted. Concomitant use of drugs known to induce ototoxicity should be avoided. Observe patients for

21 CFR Ch. I (4–1–99 Edition)

signs of adrenocorticoid overdose. The antibiotic susceptibility of the pathogenic organism should be determined prior to use of this preparation. Administration of recommended doses beyond 7 days may result in delayed wound healing. Animals treated longer than 7 days should be monitored closely.

(4) For use by or on the order of a licensed veterinarian.

[47 FR 26378, June 18, 1982, as amended at 52 FR 7832, Mar. 13, 1987]

§ 524.1044e Gentamicin sulfate spray.

(a) *Specification.* Each milliliter of sterile aqueous solution contains gentamicin sulfate equivalent to 1.07 milligrams of gentamicin.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is indicated for the treatment of pink eye in cattle (infectious bovine keratoconjunctivitis) caused by *Moraxella bovis*.

(2) One actuation of the sprayer delivers 0.7 milliliter containing 0.75 milligram gentamicin. The sprayer should be held upright 3 to 6 inches from the affected eye, with the opening directed towards the eye, and pumped once. It is advisable to treat once a day for up to 3 days.

(3) Conditions other than bacterial infections of the bovine eye and infectious keratoconjunctivitis caused by *Moraxella bovis* may produce similar signs. If conditions persists or increases, discontinue use and consult veterinarian.

[48 FR 41157, Sept. 14, 1983, as amended 52 FR 7833, Mar. 13, 1987]

§ 524.1044f Gentamicin sulfate, betamethasone valerate topical spray.

(a) *Specifications.* Each milliliter of spray contains gentamicin sulfate equivalent to 0.57 milligram of gentamicin base and betamethasone valerate equivalent to 0.284 milligram of betamethasone.

(b) *Sponsor.* See Nos. 000061 and 051259 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs in the treatment of infected superficial lesions caused by bacteria sensitive to gentamicin.

Food and Drug Administration, HHS

§ 524.1193

(2) For the treatment of infected superficial lesions, the lesion and adjacent area should be properly cleaned before treatment. Excessive hair should be removed. Hold bottle upright 3 to 6 inches from the lesion and depress the sprayer head twice. One actuation of the sprayer delivers 0.7 milliliter of the spray. The drug should be administered with two spray actuations 2 to 4 times daily for 7 days.

(3) If hypersensitivity to any of the components occurs, treatment should be discontinued and appropriate therapy instituted. The antibiotic susceptibility of the pathogenic organism should be determined prior to use of this preparation. Administration of recommended doses beyond 7 days may result in delayed wound healing. Animals treated longer than 7 days should be monitored closely.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 740, Jan. 7, 1985, as amended at 52 FR 7833, Mar. 13, 1987; 62 FR 10220, Mar. 6, 1997]

§ 524.1044g Gentamicin sulfate, betamethasone valerate, clotrimazole ointment.

(a) *Specifications.* Each gram (g) of ointment contains gentamicin sulfate equivalent to 3 milligrams (mg) gentamicin base, betamethasone valerate equivalent to 1 mg betamethasone, and 10 mg clotrimazole.

(b) *Sponsor.* See 000061 and 051259 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used for the treatment of canine otitis externa associated with yeast (*Malassezia pachydermatis*, formerly *Pityrosporum canis*) and/or bacteria susceptible to gentamicin.

(2) For 7.5 or 15 g tube, instill 4 drops of ointment twice daily into the ear canal of dogs weighing less than 30 pounds, instill 8 drops twice daily for dogs weighing 30 pounds or more. For 215 g bottle, instill 2 drops of ointment twice daily into the ear canal of dogs weighing less than 30 pounds, instill 4 drops twice daily for dogs weighing 30 pounds or more. Therapy should continue for 7 consecutive days.

(3) The external ear should be cleaned and dried before treatment. Remove

foreign material, debris, crusted exudates, etc., with suitable solutions. Excessive hair should be clipped from the treatment area. If hypersensitivity occurs, treatment should be discontinued and alternate therapy instituted.

(4) Corticosteroids administered to dogs, rabbits, and rodents during pregnancy have resulted in cleft palate in offspring. Other congenital anomalies including deformed forelegs, phocomelia, and anasarca have been reported in offspring of dogs which received corticosteroids during pregnancy. Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition if used during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[58 FR 38973, July 21, 1993, as amended at 63 FR 31932, June 11, 1998]

§ 524.1193 Ivermectin pour-on.

(a) *Specifications.* Each milliliter of solution contains 5 milligrams of ivermectin.

(b) *Sponsors.* (1) See No. 050604 in § 510.600(c) of this chapter for use as in paragraph (d) of this section.

(2) See No. 059130 for use as in paragraphs (d)(1), (d)(2)(i), and (d)(3) of this section.

(c) *Related tolerances.* See § 556.344 of this chapter.

(d) *Conditions of use—*(1) *Amount.* One milliliter per 22 pounds of body weight.

(2) *Indications for use.* (i) For cattle: It is used for the treatment and control of: Gastrointestinal roundworms (adults and fourth-stage larvae) *Ostertagia ostertagi* (including inhibited stage), *Haemonchus placei*, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia* spp., *Oesophagostomum radiatum*; (adults) *O. venulosum*, *Strongyloides papillosus*, *Trichuris* spp.; lungworms (adults and fourth-stage larvae) *Dictyocaulus viviparus*; cattle grubs (parasitic stages) *Hypoderma bovis*, *H. lineatum*; mites *Chorioptes bovis*, *Sarcoptes scabiei* var. *bovis*; lice

§ 524.1200

Linognathus vituli, *Haematopinus eurysternus*, *Damalina bovis*, *Solenoptes capillatus*; horn flies *Haematobia irritans*.

(ii) For cattle: It is also used to control infections of gastrointestinal roundworms *O. ostertagi*, *O. radiatum*, *H. placei*, *T. axei*, *Cooperia punctata*, and *C. oncophora* for 14 days after treatment.

(3) *Limitations*. For use on skin surface only. Do not treat cattle within 48 days of slaughter. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. Drug has been associated with severe adverse reactions in sensitive dogs; therefore drug is not recommended for use in animals other than cattle. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[55 FR 50551, Dec. 7, 1990, as amended at 62 FR 38908, July 21, 1997; 62 FR 63271, Nov. 28, 1997; 63 FR 44385, Aug. 19, 1998]

§ 524.1200 Kanamycin ophthalmic and topical dosage forms.

§ 524.1200a Kanamycin ophthalmic ointment.

(a) *Specifications*. The drug, which is in a suitable and harmless ointment base, contains 3.5 milligrams of kanamycin activity (as the sulfate) per gram of ointment.

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*. It is indicated for use in dogs in various eye infections due to kanamycin sensitive bacteria. It is used treating conditions such as conjunctivitis, blepharitis, dacryocystitis, keratitis, and corneal ulcerations and as a prophylactic in traumatic conditions, removal of foreign bodies, and intraocular surgery. Apply a thin film to the affected eye three or four times daily or more frequently if deemed advisable. Treatment should be continued for at least 48 hours after the eye appears normal. For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 53 FR 27851, July 25, 1988; 64 FR 404, Jan. 5, 1999]

EFFECTIVE DATE NOTE: At 64 FR 404, Jan. 5, 1999, § 524.1200a was amended by removing paragraph (a)(1) and by removing the designation for paragraph (a)(2), effective May

21 CFR Ch. I (4–1–99 Edition)

20, 1999. For the convenience of the user, the superseded text is set forth as follows:

§ 524.1200a Kanamycin ophthalmic ointment.

(a) * * * (1) The kanamycin used conforms to the standards of identity, strength, quality, and purity prescribed by § 444.30 of this chapter.

* * * * *

§ 524.1200b Kanamycin ophthalmic aqueous solution.

(a) *Specifications*. The drug, which is in an aqueous solution including suitable and harmless preservatives and buffer substances, contains 10 milligrams of kanamycin activity (as the sulfate) per milliliter of solution.

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*. It is indicated for use in dogs in various eye infections due to kanamycin sensitive bacteria. It is used in treating conditions such as conjunctivitis, blepharitis, dacryocystitis, keratitis, and corneal ulcerations and as a prophylactic in traumatic conditions, removal of foreign bodies, and intraocular surgery. Instill a few drops into the affected eye every 3 hours or more frequently if deemed advisable. Administer as frequently as possible for the first 48 hours, after which the frequency of applications may be decreased. Treatment should be continued for at least 48 hours after the eye appears normal. For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 53 FR 27851, July 25, 1988; 64 FR 404, Jan. 5, 1999]

EFFECTIVE DATE NOTE: At 64 FR 404, Jan. 5, 1999, § 524.1200b was amended by revising paragraph (a), effective May 20, 1999. For the convenience of the user, the superseded text is set forth as follows:

§ 524.1200b Kanamycin ophthalmic aqueous solution.

(a) *Specifications*. (1) The kanamycin used conforms to the standards of identity, strength, quality, and purity prescribed by § 444.30 of this chapter.

(2) The drug, which is in an aqueous solution including suitable and harmless preservatives and buffer substances, contains 10.0

Food and Drug Administration, HHS

§ 524.1240

milligrams of kanamycin activity (as the sulfate) per cubic centimeter of solution.

* * * * *

§ 524.1204 Kanamycin sulfate, calcium amphomycin, and hydrocortisone acetate.

(a) *Specifications.* (1) Calcium amphomycin is the calcium salt of amphomycin. It conforms to the following specifications:

(i) Its potency is not less than 863 micrograms of amphomycin per milligram;

(ii) Its moisture content is not more than 10 percent; and

(iii) Its pH in a 2-percent aqueous suspension is 6.0 to 7.5.

(2) The drug is in a water-miscible ointment or cream base and each gram of ointment or cream contains: 5.0 milligrams of kanamycin activity as the sulfate, 5.0 milligrams of amphomycin activity as the calcium salt, and 10.0 milligrams of hydrocortisone acetate.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is indicated for use in dogs in the following conditions associated with bacterial infections caused by organisms susceptible to one or both antibiotics: Acute otitis externa, furunculosis, folliculitis, pruritus, anal gland infections, erythema, decubital ulcer, superficial wounds, and superficial abscesses.

(2) The ointment should be applied to the affected areas of the skin at least twice daily. In severe or widespread lesions it may be desirable to apply the ointment more than twice daily. After some improvement is observed, treatment can usually be reduced to once daily. Before application, hair in the affected area should be closely clipped and the area should be thoroughly cleansed of crusts, scales, dirt, or other detritus. When treating infections of the anal gland, the drug should be introduced into the orifice of the gland and not through any fistulous tract. If no response is evident in 7 days, diagnosis and therapy should be reevaluated.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 53 FR 12512, Apr. 15, 1988; 53 FR 27851, July 25, 1988; 64 FR 404, Jan. 5, 1999]

EFFECTIVE DATE NOTE: At 64 FR 404, Jan. 5, 1999, § 524.1204 was amended by removing paragraph (a)(1) and by redesignating paragraphs (a)(2) and (a)(3) as (a)(1) and (a)(2), effective May 20, 1999. For the convenience of the user, the superseded text is set forth as follows:

§ 524.1204 Kanamycin sulfate, calcium amphomycin, and hydrocortisone acetate.

(a) * * * (1) The kanamycin used conforms to the standards of identity, strength, quality, and purity prescribed by § 444.30(a)(1) of this chapter.

* * * * *

§ 524.1240 Levamisole.

(a) *Specifications.* The drug contains 200 milligrams of levamisole per milliliter of diethylene glycol monobutyl ether (DGME) solution.

(b) *Sponsor.* See 000061 and 010042 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.350 of this chapter.

(d) *Conditions of use. Cattle*—(1) *Amount.* 2.5 milliliters per 110 pounds (10 milligrams of levamisole per kilogram) of body weight as a single dose topically to the back of the animal.

(2) *Indications for use.* Anthelmintic effective against stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*, *Chabertia*), and lungworms (*Dictyocaulus*).

(3) *Limitations.* Conditions of constant helminth exposure may require re-treatment within 2 to 4 weeks after the first treatment. Cattle must not be slaughtered within 9 days following last treatment. Do not administer to dairy animals of breeding age. Do not treat animals before dipping or prior to exposure to heavy rain. Consult your

§ 524.1376

veterinarian for assistance in the diagnosis, treatment, and control of parasitism, and before using in severely debilitated animals.

[52 FR 10887, Apr. 6, 1987, as amended at 53 FR 7504, Mar. 9, 1988; 62 FR 61626, Nov. 19, 1997]

§ 524.1376 2-Mercaptobenzothiazole solution.

(a) *Specifications.* The drug contains 1.3 percent 2-mercaptobenzothiazole in a suitable solvent.

(b) *Sponsor.* See 011509 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* Apply twice daily to affected area.

(2) *Indications for use.* For dogs as an aid in treating moist dermatitis and hotspots and as first aid for scrapes and abrasions.

(3) *Limitations.* Clip hair from affected area before applying. If no improvement is seen within 1 week, consult a veterinarian.

[48 FR 15618, Apr. 12, 1983]

§ 524.1443 Miconazole nitrate cream; miconazole nitrate lotion; miconazole nitrate spray.

(a) *Specifications.* (1) The cream contains 23 milligrams of miconazole nitrate (equivalent to 20 milligrams of miconazole base) per gram.

(2) The lotion contains 1.15 percent of miconazole nitrate (equivalent to 1 percent miconazole base).

(3) The spray product consists of a dispensing container, sprayer pump assembly, and lotion which contains 1.15 percent of miconazole nitrate (equivalent to 1-percent miconazole base).

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter for use of cream, lotion, and spray; see No. 051259 in § 510.600(c) of this chapter for use of lotion and spray.

(c) *Conditions of use.* (1) Miconazole nitrate is an antifungal agent for topical treatment of infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

(2) Apply once daily by rubbing into or spraying a light covering on the infected site and the immediate surrounding vicinity. Continue treatment for 2 to 4 weeks until infection is com-

21 CFR Ch. I (4–1–99 Edition)

pletely eradicated as determined by appropriate laboratory examination.

(3) Accurate diagnosis of infecting organism is essential. Identify by microscopic examination of a mounting of infected tissue in potassium hydroxide solution or by culture on an appropriate medium.

(4) If no improvement is observed in 2 weeks, reevaluate diagnosis and therapy.

(5) Avoid contact with eyes since irritation may result.

(6) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 53 FR 26242, July 12, 1988; 62 FR 55161, Oct. 23, 1997; 62 FR 61626, Nov. 19, 1997]

§ 524.1451 Moxidectin.

(a) *Specifications.* Each milliliter contains 5 milligrams of moxidectin (0.5 percent solution).

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.426 of this chapter.

(d) *Conditions of use—(1) Amount.* 0.5 milligrams moxidectin per kilogram (2.2 pounds) of body weight.

(2) *Indications for use.* Beef and non-lactating dairy cattle for treatment and control of internal and external parasites: gastrointestinal roundworms (*Ostertagia ostertagi* (adult and L4, including inhibited larvae), *Haemonchus placei* (adult), *Trichostrongylus axei* (adult and L4), *T. colubriformis* (adult), *Cooperia oncophora* (adult), *C. punctata* (adult), *Bunostomum phlebotomum* (adult), *Oesophagostomum radiatum* (adult), *Nematodirus helvetianus* (adult)); lungworms (*Dictyocaulus viviparus*, adult and L4); cattle grubs (*Hypoderma bovis*, *H. lineatum*); mites (*Chorioptes bovis*, *Psoroptes ovis* (*P. Communis* var. *bovis*)); lice (*Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*, *Damalinia bovis*); and horn flies (*Haematobia irritans*). To control infections and to protect from reinfection with *O. ostertagi* for 28 days after treatment and with *D. viviparus* for 42 days after treatment.

(3) *Limitations.* Apply topically along the top of the back from the withers to the tailhead. Because a withdrawal time for milk has not been established,

Food and Drug Administration, HHS

§ 524.1484b

do not use on female dairy cattle of breeding age. A withdrawal period has not been established for this product on preruminating calves. Do not use on calves to be processed for veal. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[63 FR 14036, Mar. 24, 1998]

§ 524.1465 Mupirocin ointment.

(a) *Specifications.* Each gram contains 20 milligrams of mupirocin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Dogs:*

(i) *Indications for use.* Topical treatment of bacterial infections of the skin, including superficial pyoderma, caused by susceptible strains of *Staphylococcus aureus* and *Staphylococcus intermedius*.

(ii) *Limitations.* Apply twice daily. Treatment should not exceed 30 days. Because of potential hazard of nephrotoxicity due to polyethylene glycol content, care should be exercised in treating deep lesions. Safety of use in pregnant or breeding animals has not been determined. Not for ophthalmic use. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[53 FR 39085, Oct. 5, 1988, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 524.1484 Neomycin sulfate ophthalmic and topical dosage forms.

§ 524.1484a Neomycin sulfate ophthalmic ointment.

(a) *Specifications.* Each gram of the ointment contains 5 milligrams of neomycin sulfate equivalent in activity to 3.5 milligrams of neomycin base.

(b) *Sponsor.* See No. 017030 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is intended for use in dogs and cats for the treatment of superficial ocular bacterial infections limited to the conjunctival or the anterior segment of the eye.

(2) The drug is applied four times each day.

(3) The drug is applied by inserting the tip of the tube beneath the lower lid and by expressing a small quantity

of ointment into the conjunctival sac. The tip of the tube should not come in contact with the eye surface.

(4) Severe infections should be supplemented by systemic therapy.

(5) Prolonged administration of the drug may permit overgrowth of organisms that are not susceptible to neomycin. If new infections due to bacteria or fungi appear during therapy, appropriate measures should be taken.

(6) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 44 FR 49666, Aug. 24, 1979]

§ 524.1484b Neomycin sulfate, isoflupredone acetate, tetracaine hydrochloride, and myristyl-gamma-picolinium chloride, topical powder.

(a) *Specifications.* The product contains 5 milligrams of neomycin sulfate, equivalent to 3.5 milligrams of neomycin base, 1 milligram of isoflupredone acetate, 5 milligrams of tetracaine hydrochloride and .2 milligram of myristyl-gamma-picolinium chloride in each gram of the product in a special adherent powder base.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in horses, dogs, and cats in the treatment or adjunctive therapy of certain ear and skin conditions when such conditions are caused by or associated with neomycin-susceptible organisms and/or allergy. In addition the product is indicated as superficial dressing applied to minor cuts, wounds, lacerations, abrasions, and for postsurgical application where reduction of pain and inflammatory response is deemed desirable. The product may be used as a dusting powder following amputation of tails, claws, and dew-claws and following ear trimming, castrating, and such surgical procedures as ovariohysterectomies. The product may also be used in the treatment of acute otitis externa in dogs, acute moist dermatitis and interdigital dermatitis in dogs.

§ 524.1484c

(2) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 43 FR 18172, Apr. 28, 1978]

§ 524.1484c Neomycin sulfate, isoflupredone acetate, tetracaine hydrochloride ointment.

(a) *Specifications.* The drug contains 5 milligrams of neomycin sulfate (equivalent to 3.5 milligrams of neomycin base), 1 milligram of isoflupredone acetate, and 5 milligrams of tetracaine hydrochloride in each gram of ointment.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in treating such conditions as acute otitis externa in dogs and to a lesser degree, chronic otitis externa in dogs. It also is effective in treating anal gland infections and moist dermatitis in the dog and is a useful dressing for minor cuts, lacerations, abrasions, and post-surgical therapy in the horse, cat, and dog. It may also be used following amputation of dewclaws, tails and claws, following ear trimming and castrating operations.

(2) In treatment of otitis externa and other inflammatory conditions of the external ear canal, a quantity of ointment sufficient to fill the external ear canal may be applied one to three times daily. When used on the skin or mucous membranes, the affected area should be cleansed, and a small amount of the ointment applied and spread or rubbed in gently. The involved area may be treated one to three times a day and these daily applications continued in accordance with the clinical response.

(3) Tetracaine and neomycin have the potential to sensitize. Care should be taken to observe animals being treated for evidence of hypersensitivity or allergy to the drug. If such signs are noted, therapy with the drug should be stopped. Treatment should be limited to the period when local anesthesia is essential to control self-inflicted trauma.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 43 FR 18172, Apr. 28, 1978]

21 CFR Ch. I (4–1–99 Edition)

§ 524.1484d Neomycin sulfate, hydrocortisone acetate, tetracaine hydrochloride ear ointment.

(a) *Specifications.* The product contains 5 milligrams of neomycin sulfate, equivalent to 3.5 milligrams of neomycin base, 5 milligrams of hydrocortisone acetate, and 5 milligrams of tetracaine hydrochloride in each gram of ointment.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is indicated for treating acute otitis externa and, to a lesser degree, chronic otitis externa in dogs and cats. In treatment of ear canker and other inflammatory conditions of the external ear canal, a quantity of ointment sufficient to fill the external ear canal may be applied one to three times daily.¹

(2) Tetracaine and neomycin have the potential to sensitize. Care should be taken to observe animals being treated for evidence of hypersensitivity or allergy to the product. If such signs are noted, therapy with the product should be stopped. Incomplete response or exacerbation of corticosteroid responsive lesions may be due to the presence of nonsusceptible organisms or to prolonged use of antibiotic-containing preparations resulting in overgrowth of nonsusceptible organisms, particularly *Monilia*.¹

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13873, Mar. 27, 1975, as amended at 49 FR 21922, May 24, 1984]

§ 524.1484e Neomycin sulfate and polymyxin B sulfate ophthalmic solution.

(a) *Specifications.* Each milliliter of the ophthalmic preparation contains 5.0 milligrams neomycin sulfate (3.5 milligrams neomycin base), and 10,000 Units of polymyxin B sulfate.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

Food and Drug Administration, HHS

§ 524.1484h

(c) *Conditions of use.* (1) The drug is recommended for the treatment of bacterial infections associated with topical ophthalmological conditions such as corneal injuries, superficial keratitis, conjunctivitis, keratoconjunctivitis, and blepharitis in the dog.

(2) The recommended dosage is 1 to 2 drops per eye every 6 hours.

(3) In treating ophthalmological conditions associated with bacterial infections the drug is contraindicated in those cases in which microorganisms are nonsusceptible to the antibiotics incorporated in the drug.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 61 FR 5507, Feb. 13, 1996]

§ 524.1484f Neomycin sulfate, prednisolone acetate, tetracaine hydrochloride eardrops.

(a) *Specifications.* The product contains 5 milligrams of neomycin sulfate equivalent to 3.5 milligrams of neomycin base, 2.5 milligrams of prednisolone acetate, and 5 milligrams of tetracaine hydrochloride in each milliliter of sterile suspension.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is useful in treating such conditions as acute otitis externa and, to a lesser degree, chronic otitis externa in dogs and cats. It is indicated as treatment or adjunctive therapy of certain ear conditions in dogs and cats caused by or associated with neomycin-susceptible organisms and/or allergy. In otitis externa, 2 to 6 drops may be placed in the external ear canal two or three times daily.¹

(2) Incomplete response or exacerbation of corticosteroid responsive lesions may be due to the presence of nonsusceptible organisms or to prolonged use of antibiotic-containing preparations resulting in overgrowth of nonsusceptible organisms, particularly *Monilia*. Thus, if improvement is not noted within 2 or 3 days, or if redness, irritation, or swelling persists or increases, the diagnosis should be re-determined and appropriate therapeutic measures initiated. Tetracaine and neomycin have the potential to sensitize. Care should be taken to observe ani-

mals being treated for evidence of hypersensitivity or allergy. If such signs are noted, therapy should be stopped.¹

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

§ 524.1484g Neomycin sulfate-thiabendazole-dexamethasone solution.

(a) *Specifications.* Each cubic centimeter of neomycin sulfate-thiabendazole-dexamethasone solution contains: 40 milligrams of thiabendazole, 3.2 milligrams of neomycin (from neomycin sulfate), and 1 milligram of dexamethasone.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is recommended for use as an aid in the treatment of bacterial, mycotic, and inflammatory dermatoses and otitis externa in dogs and cats.

(2) In treating dermatoses affecting areas other than the ear, the surface of the lesions should be well moistened (two to four drops per square inch) twice daily. In treating otitis externa, five to 15 drops of the drug should be instilled in the ear twice daily. The drug is limited to 7 days maximum duration of administration.

(3) For use only by or on order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 62 FR 63271, Nov. 28, 1997]

§ 524.1484h Neomycin, penicillin, polymyxin, hydrocortisone suspension.

(a) *Specifications.* Each milliliter of suspension contains 25 milligrams of neomycin sulfate equivalent to 17.5 milligrams of neomycin, 10,000 international units of penicillin G procaine, 5,000 international units of polymyxin B sulfate, 2 milligrams of hydrocortisone acetate, and 1.25 milligrams of hydrocortisone sodium succinate.

(b) *Sponsor.* See 000009 in § 510.600(c) of this chapter.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(c) *Special considerations.* The labeling shall state: This medication contains penicillin. Allergic reactions in humans are known to occur from topical exposure to penicillin.

(d) *Conditions of use—dogs—(1) Amount.* Rub a small amount into the involved area 1 to 3 times a day. After definite improvement, it may be applied once a day or every other day.

(2) *Indications for use.* Treatment of summer eczema, atopic dermatitis, interdigital eczema, and otitis externa caused by bacteria susceptible to neomycin, penicillin, and polymyxin B.

(3) *Limitations.* For use in dogs only. Shake drug thoroughly and clean lesion before using. If redness, irritation, or swelling persists or increases, discontinue use and reevaluate diagnosis. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[59 FR 5105, Feb. 3, 1994]

§ 524.1484i Neomycin sulfate, hydrocortisone acetate, sterile ointment.

(a) *Specifications.* The drug contains 5 milligrams of neomycin sulfate, equivalent to 3.5 milligrams of neomycin base, and 5 milligrams of hydrocortisone acetate in each gram of ointment.¹

(b) *Sponsor.* No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) *Amount.* Apply three or four times daily into the conjunctival sac. With improvement, frequency may be reduced to two or three times daily. For treatment of ear canker and other inflammatory conditions of the external ear canal, fill external ear canal one to three times daily.¹

(2) *Indications for use.* For treating infections, allergic, and traumatic keratitis, conjunctivitis, acute otitis externa and, to a lesser degree, chronic otitis externa in dogs and cats.¹

(3) *Limitations.* All topical ophthalmic preparations containing corticosteroids, with or without an antimicrobial agent, are contraindicated in the initial treatment of corneal ulcers. They should not be used until infection is under control and corneal regeneration is well underway. Incomplete response or exacerbation of corticosteroid responsive lesions may

be due to the presence of nonsusceptible organisms or to prolonged use on antibiotic-containing preparations resulting in overgrowth of nonsusceptible organisms, particularly *Monilia*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[43 FR 40456, Sept. 12, 1978]

§ 524.1484j [Reserved]

§ 524.1484k Neomycin sulfate, prednisolone, tetracaine, and squalane topical-otic suspension.

(a) *Specifications.* Each milliliter of suspension contains 5 milligrams neomycin sulfate (equivalent to 3.5 milligrams neomycin base), 2 milligrams prednisolone, 5 milligrams tetracaine, and 0.25 milliliter squalane.

(b) *Sponsor.* See 017030 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* 2 to 3 applications daily or as needed.

(2) *Indications for use.* Indicated for use in dogs and cats for treating acute otitis externa and as adjunctive therapy in management of chronic otitis externa. The product may also be used for treating moist dermatitis in dogs.

(3) *Limitations.* Tetracaine and neomycin have the potential to sensitize. If signs of irritation or sensitivity develop, discontinue use. Prolonged use of this product may result in overgrowth of nonsusceptible organisms. If new infections due to bacteria or fungi appear during therapy, appropriate measures should be taken. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 5265, Feb. 4, 1983; 48 FR 8055, Feb. 25, 1983]

§ 524.1580 Nitrofurazone ophthalmic and topical dosage forms.

§ 524.1580a [Reserved]

§ 524.1580b Nitrofurazone ointment.

(a) *Specifications.* The drug contains 0.2 percent nitrofurazone in a water-soluble base.

(b) *Sponsor.* For use on dogs, cats, or horses, see Nos. 000010, 000857, 000864, 000069, 050749, 023851, and 051259 in § 510.600(c) of this chapter. For use on dogs and horses, see No. 017135 in § 510.600(c) of this chapter. For use on

horses, see No. 017153 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Indications for use.* For prevention or treatment of surface bacterial infections of wounds, burns, and cutaneous ulcers of dogs, cats, or horses.¹

(2) *Limitations.* Apply directly on the lesion with a spatula or first place on a piece of gauze. Use of a bandage is optional. The preparation should remain on the lesion for at least 24 hours. The dressing may be changed several times daily or left on the lesion for a longer period. For use only on dogs, cats, and horses (not for food use). In case of deep or puncture wounds or serious burns, use only as recommended by veterinarian. If redness, irritation, or swelling persists or increases, discontinue use; consult veterinarian.¹

[46 FR 43402, June 27, 1980, as amended at 49 FR 6476, Feb. 22, 1984; 50 FR 49373, Dec. 2, 1985; 52 FR 18691, May 19, 1987; 53 FR 32610, Aug. 26, 1988; 53 FR 40728, Oct. 18, 1988; 54 FR 29544, July 13, 1989; 54 FR 37097, Sept. 7, 1989; 55 FR 8462, Mar. 8, 1990; 55 FR 20455, May 17, 1990; 56 FR 37473, Aug. 7, 1991; 56 FR 50653, Oct. 8, 1991; 59 FR 33197, June 28, 1994; 60 FR 55659, Nov. 2, 1995; 62 FR 35077, June 30, 1997]

§ 524.1580c Nitrofurazone soluble powder.

(a) *Specifications.* The drug contains 0.2 percent nitrofurazone in a water-soluble base.

(b) *Sponsor.* See Nos. 000010, 000069, and 050749 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* Apply several times daily to the lesion or affected area from the plastic squeeze bottle.

(2) *Indications for use.* For prevention or treatment of surface bacterial infections of wounds, burns, skin ulcers, and abscesses after incision.¹

(3) *Limitations.* In case of deep or puncture wounds or serious burns, use only as recommended by veterinarian. If redness, irritation, or swelling persists or increases, discontinue use; consult veterinarian. For use only on dogs, cats, and horses (not for food use).¹

[45 FR 43402, June 27, 1980, as amended at 47 FR 43368, Oct. 1, 1982; 48 FR 28984, June 24, 1983; 53 FR 40728, Oct. 18, 1988; 54 FR 30542, July 21, 1989; 56 FR 50653, Oct. 8, 1991; 59 FR 33197, June 28, 1994; 60 FR 55659, Nov. 2, 1995; 62 FR 35077, June 30, 1997]

§ 524.1580d [Reserved]

§ 524.1580e Nitrofurazone ointment with butacaine sulfate.

(a) *Specifications.* The drug contains 0.2 percent nitrofurazone and 0.5 percent butacaine sulfate in a water-soluble base.

(b) *Sponsor.* See No. 051259 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Indications for use.* For prevention or treatment of surface bacterial infections of ears, wounds, burns, and cutaneous ulcers of dogs, cats, and horses.¹

(2) *Limitations.* Apply directly on the lesion with a spatula or first place on a piece of gauze. Use of a bandage is optional. The preparation should remain on the lesion for at least 24 hours. The dressing may be changed several times daily or left on the lesion for a longer period. For use only on dogs, cats, and horses (not for food use). In case of deep or puncture wounds or serious burns, use only as recommended by a veterinarian. If redness, irritation, or swelling persists or increases, discontinue use; consult veterinarian.¹

[49 FR 9417, Mar. 13, 1984]

§ 524.1600 Nystatin ophthalmic and topical dosage forms.

§ 524.1600a Nystatin, neomycin, thiostrepton, and triamcinolone acetonide ointment.

(a) *Specifications.* Each milliliter of petrolatum base or each gram of vanishing cream base ointment contains: 100,000 units of nystatin; neomycin sulfate equivalent to 2.5 milligrams of neomycin base; 2,500 units of thiostrepton; and 1.0 milligram of triamcinolone acetonide.

(b) *Sponsor.* See Nos. 000031, 000069, 000332, 025463, 051259, and 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* (i) For topical dermatological use: Clean affected areas and remove any encrusted discharge or exudate, and apply

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

§ 524.1600b

sparingly either ointment in a thin film.¹

(ii) For otic use: Clean ear canal of impacted cerumen, remove any foreign bodies such as grass awns and ticks, and instill three to five drops of petrolatum base ointment. Preliminary use of a local anesthetic may be advisable.¹

(iii) For infected anal glands and cystic areas: Drain gland or cyst and fill with petrolatum base ointment.¹

(2) *Indications for use.* (i) Topically: Use either ointment in dogs and cats for anti-inflammatory, antipruritic, antifungal, and antibacterial treatment of superficial bacterial infections, and for dermatologic disorders characterized by inflammation and dry or exudative dermatitis, particularly associated with bacterial or candidal (*Candida albicans*) infections.¹

(ii) Otitis, cysts, and anal gland infections: Use petrolatum base ointment in dogs and cats for the treatment of acute and chronic otitis and interdigital cysts, and in dogs for anal gland infections.¹

(3) *Limitations.* For mild inflammations, use once daily to once a week. For severe conditions, apply initially two to three times daily, decreasing frequency as improvement occurs. Not intended for treatment of deep abscesses or deep-seated infections. Not for ophthalmic use. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 43 FR 29770, July 11, 1978; 50 FR 41490, Oct. 11, 1985; 53 FR 39257, Oct. 6, 1988; 54 FR 5431, Feb. 3, 1989; 54 FR 48090, Nov. 21, 1989; 56 FR 50653, Oct. 8, 1991; 60 FR 55660, Nov. 2, 1995; 61 FR 63712, Dec. 2, 1996]

§ 524.1600b Nystatin, neomycin, thiostrepton, and triamcinolone acetate ophthalmic ointment.

(a) *Specifications.* Each cubic centimeter of ointment contains: 100,000 units of nystatin, neomycin sulfate equivalent to 2.5 milligrams of neomycin base, 2,500 units of thiostrepton,

¹These conditions are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

21 CFR Ch. I (4–1–99 Edition)

and 1.0 milligram of triamcinolone acetate.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is recommended for ophthalmic use as an anti-inflammatory, antipruritic, antifungal (*Candida albicans*), and antibacterial ointment for local therapy in keratitis and conjunctivitis in cats and dogs and for infectious keratoconjunctivitis (pink eye) in cattle.

(2) It is to be administered as follows:

(i) For conjunctivitis and keratitis: Apply one drop of ointment to the affected eye(s) two or three times daily. Treatment may be continued for up to 2 weeks if necessary.

(ii) For bovine infectious keratoconjunctivitis: Apply small line of ointment to the affected eye(s) once daily. Treatment may be continued for up to 2 weeks if necessary.

(iii) Frequency of administration is dependent on the severity of the condition. For mild inflammations, applications may range from once daily to once a week; for severe conditions the drug may be applied as often as two to three times daily. Frequency of treatment may be decreased as improvement occurs.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 50 FR 41490, Oct. 11, 1985]

§ 524.1662 Oxytetracycline hydrochloride ophthalmic and topical dosage forms.

§ 524.1662a Oxytetracycline hydrochloride and hydrocortisone spray.

(a) *Specifications.* Each 3-ounce unit of oxytetracycline hydrochloride and hydrocortisone spray contains 300 milligrams of oxytetracycline hydrochloride and 100 milligrams of hydrocortisone with an inert freon propellant such that a 1-second spray treatment will deliver approximately 2.5 milligrams of oxytetracycline hydrochloride and 0.8 milligram of hydrocortisone.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is indicated for relief of discomfort and continued treatment of many allergic,

infectious, and traumatic skin conditions. The indications include prevention of bacterial infections in superficial wounds, cuts, and abrasions, treatment of allergic dermatoses, including urticaria, eczemas, insect bites, and cutaneous drug reactions, infections associated with minor burns and wounds, and nonspecific pruritus in dogs and cats.

(2) A small quantity should be sprayed on the affected surface by holding the container about 6 inches from the area to be treated and pressing the nozzle for 1 or 2 seconds. Only sufficient spray to coat the skin thinly is necessary. The application of small amounts at frequent intervals will give best results. Before treating animals with long or matted hair, it may be necessary to clip the affected area or spread the hairs to allow the medication to contact the skin surface. Relief may be noted following the first or second treatment; however, treatment should not be discontinued too soon after the initial favorable response has been obtained.

(3) Keep away from eyes or other mucous membranes; avoid inhaling; use with adequate ventilation; in case of deep or puncture wounds or serious burns, consult a veterinarian.

§ 524.1662b Oxytetracycline hydrochloride, polymyxin B sulfate ophthalmic ointment.

(a) *Specifications.* Each gram of the ointment contains oxytetracycline hydrochloride equivalent to 5 milligrams of oxytetracycline and 10,000 units of polymyxin B sulfate.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used for the prophylaxis and local treatment of superficial ocular infections due to oxytetracycline- and polymyxin-sensitive organisms. These infections include the following: Ocular infections due to streptococci, rickettsiae, *E. coli*, and *A. aerogenes* (such as conjunctivitis, keratitis, pink-eye, corneal ulcer, and blepharitis in dogs, cats, cattle, sheep, and horses); ocular infections due to secondary bacterial complications associated with distemper in dogs; and ocular infections due to bacterial inflammatory

conditions which may occur secondary to other infectious diseases in dogs, cats, cattle, sheep, and horses.

(2) It is administered topically to the eye two to four times daily.

(3) Allergic reactions may occasionally occur. Treatment should be discontinued if reactions are severe. If new infections due to nonsensitive bacteria or fungi appear during therapy, appropriate measures should be taken.

§ 524.1742 N-(Mercaptomethyl) phthalimide S-(O,O-dimethyl phosphorodithioate) emulsifiable liquid.

(a) *Specifications.* The emulsifiable liquid contains 11.6 percent N-(mercaptomethyl) phthalimide S-(O,O-dimethyl phosphorodithioate).

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Methods of application.* Methods of application to control the following conditions on beef cattle:

To control/method of use	Dilution rate (gal. drug: gal. of water)
Grubs:	
Dip	1:60
Pour-on	1:2
Spray	1:49
Lice:	
Dip	1:60
Pour-on	1:2 or 1:5
Spray	1:49 or 1:100
Hornflies:	
Dip	1:60
Spray	1:49 or 1:100
Cattle Ticks:	
Dip	1:60 or 1:240
Spray	1:49
Southern cattle ticks:	
Dip	1:60 or 1:240
Spray	1:49
Scabies mites:	
Dip	1:60
Lone Star Ticks:	
Dip	1:60
Spray	1:49 or 1:100

(i) *Dip vat procedure.* (a) Prior to charging vat, empty old contents and thoroughly clean the vat. Dip vats should be calibrated to maintain an accurate dilution. Add water, then drug to the vat according to the dilution rate indicated in the table. Add super phosphate at a rate of 100 pounds per 1,000 gallons of vat solution. Super phosphate is added to control the pH of the solution and ensure vat stability.

Super phosphate is usually available at most fertilizer dealers as 0-45-0 or 0-46-0. Stir the dip thoroughly, preferably with a compressed air device; however, any form of thorough mixing is adequate. Re-stir vat contents prior to each use. During the dipping operation, each time the dip's volume is reduced by $\frac{1}{8}$ to $\frac{1}{4}$ of its initial volume, replenish with water and add the drug at a rate of 1 gallon for each 50 or 200 gallons water added—depending on dilution rate 1:60 or 1:240. Also add super phosphate as necessary to maintain pH between 4.5 and 6.5. Stir well and resume dipping. Repeat replenishment process as necessary. For evaporation, add additional water accordingly. For added water due to rainfall, merely replenish dip with the product according to directions. If overflow occurs, either analyze for drug concentration and adjust accordingly or dispose of vat contents and recharge. Check pH after each addition of water or super phosphate to assure proper pH controls.

(b) *Dip maintenance.* (1) With use of dip vat tester, dipping may continue as long as the drug concentration is maintained between 0.15 and 0.25 percent, and the dip is not too foul for satisfactory use as indicated by foul odor or excessive darkening (i.e., color changes from beige to very dark brown).

(2) Without use of dip vat tester, vat should be emptied, cleaned, and recharged each time one of the following occurs: When the dip has been charged for 120 days; when the dip becomes too foul for satisfactory use, within the 120-day limit; if the number of animals dipped equals twice the number of gallons of the initial dip volume, within the 120-day limit.

(ii) *Spray method.* To prepare the spray, mix drug with water according to table and stir thoroughly. Apply the fresh mixture as a high-pressure spray, taking care to wet the skin, not just the hair. Apply to the point of “run-off,” about 1 gallon of diluted spray per adult animal. Lesser amounts will permit runoff for younger animals.

(iii) *Pour-on method.* Dilute the drug with water according to table by slowly adding water to the product while stirring. Apply 1 ounce of the diluted

mixture per 100 pounds of body weight (to a maximum of 8 ounces per head) down the center line of the back.

(2) *Timing of applications for cattle grub control.* For optimum cattle grub control, it is important to treat as soon as possible after the heel fly season, before the grub larvae reach the gullet or spinal canal, as the rapid kill of large numbers of larvae in these tissues may cause toxic side effects, such as bloat, salivation, staggering, and paralysis.

(3) *Treatment regimens.* (i) Control of scabies mites requires two treatments, 10 to 14 days apart.

(ii) Control of Lone Star Ticks and hornflies requires two treatments, 7 days apart.

(4) *Warnings.* The drug is a cholinesterase inhibitor. Do not use this drug on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals. Do not apply within 21 days of slaughter. For use on beef cattle only. Do not treat sick, convalescent, or stressed cattle, or calves less than 3 months old except in Federal or State eradication programs where immediate treatment of all animals in an infested herd is mandatory. Be sure free access to drinking water is available to cattle prior to dipping. Do not dip excessively thirsty animals. Do not dip animals when overheated. Repeat treatment as necessary but not more often than every 7 to 10 days. Treatment for lice, ticks, hornflies, and scabies mites may be made any time of the year except when cattle grub larvae are in the gullet or spinal canal. Treatment for lice, ticks, and scabies mites may be made any time 7 to 10 days following treatment for grubs. Do not treat grubs when the grub larvae are in the gullet or spinal canal. Do not get in eyes, on skin, or on clothing. Do not breathe spray mist. Wear rubber gloves, goggles, and protective clothing. In case of skin contact, wash immediately with soap and water; for eyes, flush with water. Wash all contaminated clothing with soap and hot water before re-use.

Food and Drug Administration, HHS

§ 524.1883

(d) *Related tolerances.* See 40 CFR 180.261.

[40 FR 13873, Mar. 27, 1975, as amended at 46 FR 27914, May 22, 1981; 48 FR 39607, Sept. 1, 1983; 54 FR 51021, Dec. 12, 1989; 61 FR 8873, Mar. 6, 1996; 62 FR 61626, Nov. 19, 1997; 63 FR 5255, Feb. 2, 1998]

§ 524.1880 Prednisolone-neomycin sulfate ophthalmic ointment.

(a) *Specifications.* Prednisolone-neomycin sulfate ophthalmic ointment contains 2 milligrams prednisolone and 5 milligrams neomycin sulfate (equivalent to 3.5 milligrams neomycin base) in each gram of ointment.

(b) *Sponsor.* See No. 017030 in § 510.600(c) of this chapter.

(c) *Conditions of use.* The drug is recommended for use in superficial ocular inflammations or infections limited to the conjunctiva or the anterior segment of the eye of cats and dogs, such as those associated with allergic reactions or gross irritants. A small quantity of the ointment should be expressed into the conjunctival sac four times a day for 7 days. After 7 days, if clinical improvement is not noted, reevaluation of the diagnosis should be considered. All topical ophthalmic preparations containing corticosteroids with or without an antimicrobial agent are contraindicated in the initial treatment of corneal ulcers. They should not be used until the infection is under control and corneal regeneration is well underway. For use only by or on the order of a licensed veterinarian.

§ 524.1881 Prednisolone acetate ophthalmic and topical dosage forms.

§ 524.1881a [Reserved]

§ 524.1881b Prednisolone acetate-neomycin sulfate sterile suspension.

(a) *Specifications.* Prednisolone acetate-neomycin sulfate sterile suspension contains 2.5 milligrams of prednisolone acetate and 5 milligrams of neomycin sulfate (equivalent to 3.5 milligrams of neomycin base) in each milliliter of sterile suspension.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is indicated for treating infectious, allergic and traumatic keratitis and con-

junctivitis, acute otitis externa, and chronic otitis externa in dogs and cats.

(2) For beginning treatment of acute ocular inflammations 1 or 2 drops may be placed in the conjunctival sac 3 to 6 times during a 24 hour period. When improvement occurs, the dosage may be reduced to 1 drop 2 to 4 times daily. In otitis externa, 2 to 6 drops may be placed in the external ear canal 2 or 3 times daily.

(3) All topical ophthalmic preparations containing corticosteroids with or without an anti-microbial agent are contraindicated in the initial treatment of corneal ulcers. They should not be used until infection is under control and corneal regeneration is well underway.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

§ 524.1883 Prednisolone sodium phosphate-neomycin sulfate ophthalmic ointment.

(a) *Specifications.* Prednisolone sodium phosphate-neomycin sulfate ophthalmic ointment contains prednisolone sodium phosphate equivalent to 2.5 milligrams prednisolone 21-phosphate and 5 milligrams neomycin sulfate (equivalent to 3.5 milligrams neomycin base) in each gram of ointment.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is recommended for use in superficial ocular inflammations or infections limited to the conjunctiva or the anterior segment of the eye of cats and dogs, such as those associated with allergic reactions or gross irritants.¹

(2) A small quantity of the ointment should be expressed into the conjunctival sac 4 times a day (at intervals of 1 to 8 hours) for a few days until there is a favorable response, then the frequency of application may be reduced to twice daily as long as the condition remains under control. Treatment may require from a few days to several weeks.¹

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

§ 524.1982

(3) All topical ophthalmic preparations containing corticosteroids with or without an antimicrobial agent are contraindicated in the initial treatment of corneal ulcers. They should not be used until the infection is under control and corneal regeneration is well underway.¹

(4) For use only by or on the order of a licensed veterinarian.¹

[40 FR 13873, Mar. 27, 1975, as amended at 62 FR 63271, Nov. 28, 1997]

§ 524.1982 Proparacaine hydrochloride ophthalmic solution.

(a) *Specifications.* The drug is an aqueous solution containing 0.5 percent proparacaine hydrochloride, 2.45 percent glycerin as a stabilizer, and 0.2 percent chlorobutanol (choral derivative) and 1:10,000 benzalkonium chloride as preservatives.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Special considerations.* The longterm toxicity of proparacaine is unknown. Prolonged use may possibly delay wound healing.

(d) *Conditions of use.* (1) The drug is indicated for use as a topical ophthalmic anesthetic in animals. It is used as an anesthetic in cauterization of corneal ulcers, removal of foreign bodies and sutures from the cornea, and measurement of intraocular pressure (tonometry) when glaucoma is suspected. Local applications may also be used as an aid in the removal of foreign bodies from the nose and ear canal, as an accessory in the examination and treatment of painful otitis, in minor surgery, and prior to catheterization.

(2) It is administered as follows:

(i) For removal of sutures: Instill one to two drops 2 or 3 minutes before removal of stitches.

(ii) For removal of foreign bodies from eye, ear, and nose: For ophthalmic use, instill three to five drops in the eye prior to examination; for otic use, instill five to 10 drops in the ear; for nasal use, instill five to 10 drops in each nostril every 3 minutes for three doses.

(iii) For tonometry: Instill one to two drops immediately before measurement.

21 CFR Ch. I (4–1–99 Edition)

(iv) As an aid in treatment of otitis: Instill two drops into the ear every 5 minutes for three doses.

(v) For minor surgery: Instill one or more drops as required.

(vi) For catheterization: Instill two to three drops with a blunt 20-gauge needle immediately before inserting catheter.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 50 FR 41490, Oct. 11, 1985]

§ 524.2101 Selenium disulfide suspension.

(a) *Specifications.* The product contains 0.9-percent weight in weight (w/w) selenium disulfide (1-percent weight in volume (w/v)).

(b) *NAS/NRC status.* These conditions are NAS/NRC reviewed and found effective. NADA's for similar products for these conditions of use need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(c) *Sponsors.* See 000061, 017135, 023851, and 050604 in § 510.600(c) of this chapter.

(1) *Indications for use.* For use on dogs as a cleansing shampoo and as an agent for removing skin debris associated with dry eczema, seborrhea, and non-specific dermatoses.

(2) *Amount.* One to 2 ounces per application.

(3) *Limitations.* Use carefully around scrotum and eyes, covering scrotum with petrolatum. Allow the shampoo to remain for 5 to 15 minutes before thorough rinsing. Repeat treatment once or twice a week. If conditions persist or if rash or irritation develops, discontinue use and consult a veterinarian.

[47 FR 53351, Nov. 26, 1982, as amended at 48 FR 32762, July 19, 1983; 54 FR 36962, Sept. 6, 1989; 56 FR 9623, Mar. 7, 1991; 58 FR 41025, Aug. 2, 1993; 63 FR 26981, May 15, 1998]

§ 524.2140 Squalane, pyrethrins and piperonyl butoxide.

(a) *Specifications.* The drug contains 25 percent squalane (hexamethyltetracosane), 0.05 percent pyrethrins and 0.50 percent technical piperonyl butoxide.

(b) *Sponsor.* See No. 017030 in § 510.600(c) of this chapter.

Food and Drug Administration, HHS

§ 524.2640

(c) *Conditions of use.* (1) The drug is used for the treatment of ear mites in dogs and cats.

(2) It is administered as follows: Cats and dogs 5–15 pounds body weight, 4 to 5 drops in each ear daily. Dogs 16–30 pounds body weight, 5 to 10 drops in each ear daily. Dogs 30 pounds body weight and over 10 to 15 drops in each ear daily. The recommended treatment is for 7 to 10 days with repeated treatment in 2 weeks if necessary.

§ 524.2350 Tolnaftate cream.

(a) *Specifications.* The drug contains 1 percent tolnaftate (2-naphthyl-*N*-methyl-*N*-(3-tolyl) thionocarbamate) in an anhydrous cream base.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is indicated for treatment of ringworm lesions due to *Microsporum canis* and *Microsporum gypseum* in dogs and cats.

(2) A small amount of the cream is applied to the affected areas once or twice a day for 2 to 4 weeks. The areas to be treated are first cleared of exudate and the hair clipped if the areas are not already denuded. The cream is massaged into each lesion and immediate surrounding area until the cream is no longer visible.

(3) If no response is seen after 2 weeks of treatment with the drug the diagnosis should be reviewed.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 29289, July 7, 1978, as amended at 52 FR 7833, Mar. 13, 1987]

§ 524.2481 Triamcinolone acetonide cream.

(a) *Specifications.* Triamcinolone acetonide cream contains 0.1 percent triamcinolone acetonide in an aqueous vanishing cream base.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is recommended for use on dogs as an anti-inflammatory, antipruritic, and antiallergic agent for topical treatment of allergic dermatitis and summer eczema.

(2) The drug is applied by rubbing into affected areas two to four times daily for 4 to 10 days.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 50 FR 41490, Oct. 11, 1985]

§ 524.2620 Liquid crystalline trypsin, Peru balsam, castor oil.

(a)(1) *Specifications.* The drug is a liquid for direct application or an aerosol preparation formulated so that each gram delivered to the wound site contains 0.12 milligram of crystalline trypsin, 87.0 milligrams of Peru balsam, and 788.0 milligrams of castor oil.

(2) *Sponsor.* See No. 000514 in § 510.600(c) of this chapter.

(b)(1) *Specifications.* The drug is a liquid for direct application or an aerosol preparation formulated so that each gram delivered to the wound site contains 0.1 milligram of crystalline trypsin, 72.5 milligrams of Peru balsam, and 800 milligrams of castor oil.

(2) *Sponsor.* See No. 017135 in § 510.600(c) of this chapter.

(c) *Conditions of use.* The drug is used as an aid in the treatment of external wounds and assists healing by facilitating the removal of necrotic tissue, exudate and organic debris.

[40 FR 13873, Mar. 27, 1975, as amended at 41 FR 56307, Dec. 28, 1976; 50 FR 9800, Mar. 12, 1985; 54 FR 25565, June 16, 1989; 56 FR 37474, Aug. 7, 1991]

§ 524.2640 Tylosin, neomycin eye powder.

(a) *Specifications.* Tylosin is the antibiotic substance produced by growth of *Streptomyces fradiae* or the same antibiotic substance produced by any other means. Tylosin, present as the tartrate salt, conforms to the appropriate antibiotic standard. Tylosin contains at least 95 percent tylosin as a combination of tylosin A, tylosin B, tylosin C, and tylosin D of which at least 80 percent is tylosin A as determined by a method entitled "Determination of Factor Content in Tylosin by High Performance Liquid Chromatography," which is incorporated by reference. Copies are available from the Dockets Management Branch (HFA-305), Food and Drug Administration, rm. 1-23, 12420 Parklawn Dr., Rockville, MD 20857, or available for inspection at the

Office of the Federal Register, 800 North Capitol Street, NW., suite 700, Washington, DC 20001.

(b) *Sponsor.* See No. 000986 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in cattle for the treatment of pinkeye (infectious keratoconjunctivitis).

(2) It is administered by holding the eyelids open and dusting powder into both eyes. The treatment is repeated daily for up to 7 days depending on the severity of the infection. Affected animals should be protected from direct sunlight, dust, and flies. In an affected herd, all animals with or without signs of the disease should receive at least one treatment.

(3) If there is severe eye damage or if the condition persists or increases, discontinue administering the drug and consult a veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 50 FR 49841, Dec. 5, 1985; 59 FR 14365, Mar. 28, 1994]

PART 526—INTRAMAMMARY DOSAGE FORMS

Sec.			
526.88	Amoxicillin trihydrate	for	
	intramammary infusion.		
526.363	Cephapirin benzathine.		
526.365	Cephapirin sodium	for	
	intramammary infusion.		
526.464	Cloxacillin intramammary	dosage	
	forms.		
526.464a	Cloxacillin benzathine	for	
	intramammary infusion.		
526.464b	Cloxacillin benzathine	for	
	intramammary infusion, sterile.		
526.464c	Cloxacillin sodium	for	
	intramammary infusion, sterile.		
526.464d	Cloxacillin sodium	for	
	intramammary infusion.		
526.820	Erythromycin.		
526.1130	Hetacillin potassium	for	
	intramammary infusion.		
526.1590	Novobiocin oil suspension.		
526.1696	Penicillin intramammary	dosage	
	forms.		
526.1696a	Penicillin G procaine in oil.		
526.1696b	Penicillin G procaine-dihydro-		
	streptomycin in soybean oil for		
	intramammary infusion (dry cows).		
526.1696c	Penicillin G procaine-dihydro-		
	streptomycin sulfate for intramammary		
	infusion (dry cows).		
526.1696d	Penicillin G procaine-novobiocin		
	for intramammary infusion.		
526.1810	Pirlimycin hydrochloride aqueous		
	gel.		

AUTHORITY: 21 U.S.C. 360b.

§ 526.88 Amoxicillin trihydrate for intramammary infusion.

(a) *Specifications.* Each single dose syringe contains amoxicillin trihydrate equivalent to 62.5 milligrams of amoxicillin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.38 of this chapter.

(d) *Conditions of use—Lactating cows—*(1) *Amount.* One syringe (equivalent to 62.5 milligrams amoxicillin) per quarter.

(2) *Indications for use.* For the treatment of subclinical infectious bovine mastitis due to *Streptococcus agalactiae* and *Staphylococcus aureus* (penicillin sensitive).

(3) *Limitations.* Administer after milking. Clean and disinfect the teat. Use one syringe per infected quarter every 12 hours for a maximum of 3 doses. Do not use milk taken from treated animals for food purposes within 60 hours (5 milkings) after last treatment. Do not slaughter treated animals for food purposes within 12 days after the last treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37334, Aug. 18, 1992, as amended at 60 FR 55660, Nov. 2, 1995]

§ 526.363 Cephapirin benzathine.

(a) *Specifications.* Each 10 milliliter disposable syringe contains 300 milligrams of cephapirin activity (as cephapirin benzathine) in a peanut-oil gel.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.115 of this chapter.

(d) *Conditions of use—*(1) *Amount.* Infuse contents of one syringe into each infected quarter.

(2) *Indications for use.* Use in dry cows for treatment of mastitis caused by susceptible strains of *Streptococcus agalactiae* and *Staphylococcus aureus*.

(3) *Limitations.* Infuse each infected quarter following last milking or early in the dry period, but no later than 30 days before calving. Milk from treated cows must not be use for food during

Food and Drug Administration, HHS

§ 526.464b

the first 72 hours after calving. Animals infused with this product must not be slaughtered for food until 42 days after the latest infusion. For use in dry cows only.

[43 FR 37174, Aug. 22, 1978, as amended at 53 FR 27851, July 25, 1988]

§ 526.365 Cephapirin sodium for intramammary infusion.

(a) *Specifications.* Each 10-milliliter dose contains 200 milligrams of cephapirin sodium activity in a peanut-oil gel.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.115 of this chapter.

(d) *Conditions of use.* (1) The drug is used for the treatment of lactating cows having bovine mastitis caused by susceptible strains of *Streptococcus agalactiae* and *Staphylococcus aureus*.

(2) Administer one dose into each infected quarter immediately after the quarter has been completely milked out. Do not milk out for 12 hours. Repeat once only in 12 hours. If improvement is not noted within 48 hours after treatment, consult your veterinarian.

(3) Milk that has been taken from animals during treatment and for 96 hours (8 milkings) after the last treatment must not be used for food. Treated animals must not be slaughtered for food until 4 days after the last treatment.

[40 FR 57455, Dec. 10, 1975, as amended at 53 FR 27852, July 25, 1988. Redesignated at 63 FR 8349, Feb. 19, 1998]

§ 526.464 Cloxacillin intramammary dosage forms.

§ 526.464a Cloxacillin benzathine for intramammary infusion.

(a) *Specifications.* Each dose contains cloxacillin benzathine equivalent to 500 milligrams of cloxacillin.

(b) *Related tolerances.* See § 556.165 of this chapter.

(c) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter for use in dairy cows.

(1) *Amount.* Administer aseptically into each infected quarter immediately after last milking or early in dry period.

(2) *Indications for use.* Treatment of mastitis caused by *Staphylococcus aureus* and *Streptococcus agalactiae* including penicillin resistant strains in dairy cows during the dry period.

(3) *Limitations.* For use in dry cows only. Not to be used within 30 days of calving. Animals infused with this product must not be slaughtered for food use for 30 days after the latest infusion. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter for use in dairy cows.

(1) *Amount.* Administer one dose in each quarter immediately after last milking.

(2) *Indications for use.* Treatment and prophylaxis of bovine mastitis in non-lactating cows due to *S. agalactiae* and *S. aureus*.

(3) *Limitations.* For use in dry cows only. Not to be used within 4 weeks (28 days) of calving. Animals infused with this product must not be slaughtered for food use for 4 weeks (28 days) after the latest infusion. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37334, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992, as amended at 60 FR 55660, Nov. 2, 1995]

§ 526.464b Cloxacillin benzathine for intramammary infusion, sterile.

(a) *Specifications.* Each 6 milliliter dose contains cloxacillin benzathine equivalent to 500 milligrams of cloxacillin.

(b) *Related tolerances.* See § 556.165 of this chapter.

(c) *Sponsor.* See No. 055529 in § 510.600(c) of this chapter.

(1) *Amount.* 6 milliliters per infected quarter aseptically immediately after last milking at the time of drying-off of the cow.

(2) *Indications for use.* Treatment of mastitis caused by *Staphylococcus aureus* and *Streptococcus agalactiae* in dairy cows at the time of drying-off of the cow.

(3) *Limitations.* For use in dry cows only. Not to be used within 30 days of calving. Milk taken from treated cows prior to 72 hours (6 milkings) after calving must not be used for human

§ 526.464c

food. Animals infused with this product must not be slaughtered for food from the time of infusion until 72 hours after calving. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(1) *Amount.* One dose per infected quarter immediately after last milking.

(2) *Indications for use.* Treatment and prophylaxis of bovine mastitis in non-lactating cows due to *Streptococcus agalactiae* and *Staphylococcus aureus*.

(3) *Limitations.* For use in dry cows only. Not to be used within 4 weeks (28 days) of calving. Animals infused with this product must not be slaughtered for food use for 4 weeks (28 days) after the latest infusion. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37334, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992, as amended at 58 FR 61016, Nov. 19, 1993; 60 FR 55660, Nov. 2, 1995]

§ 526.464c Cloxacillin sodium for intramammary infusion, sterile.

(a) *Specifications.* Each milliliter contains cloxacillin sodium equivalent to 20.0 milligrams of cloxacillin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.165 of this chapter.

(d) *Conditions of use. Lactating cows—*
(1) *Amount.* 10 milliliters (one dose of 200 milligrams) per infected quarter.

(2) *Indications for use.* Treatment of mastitis in lactating cows due to *Streptococcus agalactiae* and *Staphylococcus aureus*, nonpenicillinase-producing strains.

(3) *Limitations.* Administer after milking, cleaning, and disinfecting, and as early as possible after detection. Treatment should be repeated at 12-hour intervals for a total of three doses. Milk taken from treated animals within 48 hours (four milkings) after the latest treatment should not be used for food. Treated animals should not be slaughtered for food within 10 days after the latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37335, Aug. 18, 1992, as amended at 60 FR 55660, Nov. 2, 1995]

21 CFR Ch. I (4–1–99 Edition)

§ 526.464d Cloxacillin sodium for intramammary infusion.

(a) *Specifications.* Each milliliter contains cloxacillin sodium equivalent 20.0 milligrams of cloxacillin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.165 of this chapter.

(d) *Conditions for use. Lactating cows—*
(1) *Amount.* 10 milliliters (one dose of 200 milligrams) per infected quarter.

(2) *Indications for use.* Treatment of mastitis in lactating cows due to *Streptococcus agalactiae* and *Staphylococcus aureus*, nonpenicillinase-producing strains.

(3) *Limitations.* Administer after milking, cleaning, and disinfecting, and as early as possible after detection. Treatment should be repeated at 12-hour intervals for a total of three doses. Milk taken from treated animals within 48 hours (4 milkings) after the latest treatment should not be used for food. Treated animals should not be slaughtered for food within 10 days after the latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37335, Aug. 18, 1992, as amended at 60 FR 55660, Nov. 2, 1995]

§ 526.820 Erythromycin.

(a) *Specifications.* (1) Each 6-milliliter, single-dose, disposable syringe contains 300 milligrams of erythromycin (as the base), 0.45 milligram of butylated hydroxyanisole, and 0.45 milligram of butylated hydroxytoluene.

(2) Each 12-milliliter, single-dose, disposable syringe contains 600 milligrams of erythromycin (as the base), 0.90 milligram of butylated hydroxyanisole, and 0.90 milligram of butylated hydroxytoluene.

(3) The vehicle is triglyceride of saturated fatty acids from coconut oil.

(4) The drug may or may not be sterile.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use—*(1) *Amount.* (i) Lactating cows: After milking, cleaning, and disinfecting, infuse contents of a single 6-milliliter syringe into each infected quarter; repeat procedure at 12-hour intervals for a maximum of 3 consecutive infusions.

Food and Drug Administration, HHS

§ 526.1590

(ii) Dry cows: After milking, cleaning, and disinfecting, infuse contents of a single 12-milliliter syringe into each infected quarter at the time of drying off.

(2) *Indications for use.* Treatment of mastitis due to *Staphylococcus aureus*, *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, and *Streptococcus uberis* in lactating or dry cows.

(3) *Limitations.* Milk taken from animals during treatment and for 36 hours (3 milkings) after the latest treatment must not be used for food.

[47 FR 15772, Apr. 13, 1982]

§ 526.1130 Hetacillin potassium for intramammary infusion.

(a) *Specifications.* Each 10 milliliter syringe contains hetacillin potassium equivalent of 62.5 milligrams of ampicillin.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use. Lactating cows—*
(1) *Amount.* 10 milliliters of hetacillin potassium equivalent to 62.5 milligrams ampicillin into each infected quarter. Repeat at 24-hour intervals until a maximum of three treatments has been given.

(2) *Indications for use.* Treating acute, chronic, or subclinical bovine mastitis in lactating cows caused by susceptible strains of *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, *Staphylococcus aureus*, and *Escherichia coli*.

(3) *Limitations.* If definite improvement is not noted within 48 hours after treatment, the causal organism should be further investigated. Milk that has been taken from animals during treatment and for 72 hours (6 milkings) after the latest treatment must not be used for food. Treated animals must not be slaughtered for food until 10 days after the latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37335, Aug. 18, 1992]

§ 526.1590 Novobiocin oil suspension.

(a)(1) *Specifications.* Each 10 milliliters of oil suspension contains the equivalent of 400 milligrams of novobiocin (present as sodium novobiocin).

(2) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(3) *Related tolerances.* See § 556.460 of this chapter.

(4) *Conditions of use—*(i) *Amount.* Ten milliliters (equivalent to 400 milligrams of novobiocin) infused in each quarter.

(ii) *Indications for use.* It is used in dry cows for the treatment of mastitis caused by susceptible strains of *Staphylococcus aureus* and *Streptococcus agalactiae*.

(iii) *Limitations.* Infuse each quarter at the time of drying off, but not less than 30 days prior to calving. Do not slaughter treated animals for food use for 30 days following udder infusion. For udder installation for the treatment of mastitis in dry cows only.

(b)(1) *Specifications.* Each 10 milliliters of oil suspension contains the equivalent of 150 milligrams of novobiocin (present as sodium novobiocin).

(2) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(3) *Related tolerances.* See § 556.460 of this chapter.

(4) *Conditions of use—*(i) *Amount.* Infuse 10 milliliters (equivalent to 150 milligrams of novobiocin) in each quarter after milking. Repeat treatment once after 24 hours.

(ii) *Indications for use.* Use in lactating cows for treatment of mastitis caused by susceptible strains of *Staphylococcus aureus*.

(iii) *Limitations.* Do not milk for at least 6 hours after treatment; afterwards, milk at regular intervals. Milk taken from treated animals within 72 hours (6 milkings) after latest treatment must not be used for food. Do not slaughter treated animals for food for 15 days following latest treatment. If redness, swelling, or abnormal milk persists or increases after treatment, discontinue use and consult a veterinarian. For udder instillation in lactating cattle only. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 10554, Mar. 14, 1978]

§ 526.1696

21 CFR Ch. I (4–1–99 Edition)

§ 526.1696 Penicillin intramammary dosage forms.

§ 526.1696a Penicillin G procaine in oil.

(a) *Specifications.* Each milliliter contains penicillin G procaine equivalent to 100,000 units of penicillin G in peanut, sesame, or soybean oils.

(b) *Related tolerances.* See § 556.510 of this chapter.

(c) *Sponsor.* See No. 010515 in § 510.600(c) of this chapter.

(1) *National Academy of Sciences/National Research Council (NAS/NRC) status.* The conditions specified in paragraph (c)(2)(i)(B) of this section were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(2) *Conditions of use.* Treating bovine mastitis caused by *Streptococcus agalactiae*, *S. dysgalactiae*, and *S. uberis* in lactating cows as follows:

(i) *Three-dose regimen.* Administer by intramammary infusion in each infected quarter as follows:

(A) *6-milliliter dose (peanut oil).* Treatment may be repeated at 12-hour intervals. Milk that has been taken from animals during treatment and for 84 hours (7 milkings) after the latest treatment must not be used for food. Animals must not be slaughtered for food during treatment or within 4 days after the latest treatment.

(B) *10-milliliter dose (sesame oil).* Treatment may be repeated at 12-hour intervals. Milk that has been taken from animals during treatment and for 60 hours (5 milkings) after the latest treatment must not be used for food. Animals must not be slaughtered for food during treatment or within 3 days after the latest treatment.

(ii) *Two-dose regimen.* 10-milliliter dose (peanut oil). Administer by intramammary infusion in each infected quarter. Treatment may be repeated at intervals of 12 hours. Milk taken from animals during treatment and for 60 hours (5 milkings) after the latest treatment must not be used for food. Animals must not be slaughtered for food during treatment or within 4 days after latest treatment.

(d) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.

(1) *10-milliliter dose (peanut oil).* Administer by intramammary infusion in each infected quarter. Treatment may be repeated at 12-hour intervals for not more than three doses, as indicated by clinical response.

(2) *Indications for use.* Treating bovine mastitis caused by *Streptococcus agalactiae*, *S. dysgalactiae*, and *S. uberis* in lactating cows as follows:

(3) *Limitations.* Milk that has been taken from animals during treatment and for 60 hours after the latest treatment must not be used for food. Animals must not be slaughtered for food during treatment or within 3 days after the latest treatment.

(e) *Sponsor.* See No. 010515 (sesame oil) and No. 050604 (peanut oil) in § 510.600(c) of this chapter.

(1) *NAS/NRC status.* The conditions of use were NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(2) *Single-dose regimen.* One 10-milliliter dose (sesame oil or peanut oil) in each infected quarter at time of drying-off.

(3) *Indications of use.* Treating bovine mastitis caused by *Streptococcus agalactiae* in dry cows.

(4) *Limitations.* Discard all milk for 72 hours (6 milkings) following calving, or later as indicated by the marketable quality of the milk. Animals must not be slaughtered for food within 14 days postinfusion.

[57 FR 37335, Aug. 18, 1992, as amended at 58 FR 500, Jan. 6, 1993]

§ 526.1696b Penicillin G procaine-dihydrostreptomycin in soybean oil for intramammary infusion (dry cows).

(a) *Specifications.* Each 10 milliliters of suspension contains penicillin G procaine equivalent to 200,000 units of penicillin G and dihydrostreptomycin sulfate equivalent to 300 milligrams of dihydrostreptomycin.

(b) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See §§ 556.200 and 556.510 of this chapter.

(d) *Conditions of use. Dairy cows*—(1) *Amount.* One syringe into each quarter at the last milking prior to drying off.

(2) *Indications for use.* Intramammary treatment of subclinical mastitis in dairy cows at the time of drying off, specifically against infections caused by *Staphylococcus aureus* and *Streptococcus agalactiae*.

(3) *Limitations.* Not to be used within 6 weeks of calving. For use in dry cows only. Milk taken from cows within 24 hours (2 milkings) after calving must not be used for food. Animals infused with this drug must not be slaughtered for food within 60 days of treatment nor within 24 hours after calving.

[57 FR 37336, Aug. 18, 1992]

§ 526.1696c Penicillin G procaine-dihydrostreptomycin sulfate for intramammary infusion (dry cows).

(a) *Specifications.* Each 10 milliliters of suspension contains penicillin G procaine equivalent to 1 million units of penicillin G and dihydrostreptomycin sulfate equivalent to 1 gram of dihydrostreptomycin.

(b) *Sponsor.* See No. 033392 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See §§ 556.200 and 556.510 of this chapter.

(d) *Conditions of use. Dairy cows*—(1) *Amount.* One syringe per quarter at the last milking prior to drying off.

(2) *Indications for use.* Intramammary use to reduce the frequency of existing infection and to prevent new infections with *Staphylococcus aureus* in dry cows.

(3) *Limitations.* Not to be used within 6 weeks of freshening. Not for use in lactating cows. Milk taken from animals within 96 hours (8 milkings) after calving must not be used for feed. Animals infused with this drug must not be slaughtered for food within 60 days from the time of infusion nor within 96 hours after calving. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37336, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992]

§ 526.1696d Penicillin G procaine-novobiocin for intramammary infusion.

(a) *Specifications.* For lactating cattle: each 10-milliliter dose contains 100,000 units of penicillin G procaine

and 150 milligrams of novobiocin as novobiocin sodium. For dry cows: 200,000 units of penicillin G procaine and 400 milligrams of novobiocin as novobiocin sodium.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Lactating cows*—(i) *Amount.* 10 milliliters in each infected quarter after milking. Repeat once after 24 hours.

(ii) *Indications for use.* Treating lactating cows for mastitis caused by susceptible strains of *Staphylococcus aureus*, *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, and *Streptococcus uberis*.

(iii) *Limitations.* For udder instillation in lactating cattle only. Do not milk for at least 6 hours after treatment; thereafter, milk at regular intervals. Milk taken from treated animals within 72 hours (6 milkings) after the latest treatment must not be used for food. Treated animals must not be slaughtered for food for 15 days following the latest treatment. If redness, swelling, or abnormal milk persists, discontinue use and consult a veterinarian.

(2) *Dry cows*—(i) *Amount.* 10 milliliters in each quarter at time of drying off.

(ii) *Indications for use.* Treatment of subclinical mastitis caused by susceptible strains of *Staphylococcus aureus* and *Streptococcus agalactiae*.

(iii) *Limitations.* For udder instillation in dry cows only. Do not use less than 30 days prior to calving. Milk from treated cows must not be used for food during the first 72 hours after calving. Treated animals must not be slaughtered for food for 30 days following udder infusion.

[57 FR 37336, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992]

§ 526.1810 Pirlimycin hydrochloride aqueous gel.

(a) *Specifications.* Each 10-milliliter syringe contains 50 milligrams of pirlimycin (as pirlimycin hydrochloride).

(b) *Sponsor.* See 000009 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.515 of this chapter.

(d) *Conditions of use.* (1) *Dose.* 50 milligrams in each infected quarter, repeated once after 24 hours.

(2) *Indications for use.* For lactating dairy cattle for the treatment of clinical and subclinical mastitis caused by *Staphylococcus* species, such as *Staphylococcus aureus*; and *Streptococcus* species, such as *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, and *Streptococcus uberis*.

(3) *Limitations.* Milk taken from animals during treatment and for 36 hours (three milkings) following the last treatment must not be used for food. Treated animals must not be slaughtered for food use for 28 days following the last treatment. Cows with systemic clinical signs caused by mastitis should receive other appropriate therapy under the direction of a licensed veterinarian. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[58 FR 58486, Nov. 2, 1993]

PART 529—CERTAIN OTHER DOSAGE FORM NEW ANIMAL DRUGS

Sec.

529.50 Amikacin sulfate intrauterine solution.

529.360 Cephalothin discs.

529.400 Chlorhexidine tablets and suspension.

529.469 Competitive exclusion culture.

529.1003 Flurogestone acetate-impregnated vaginal sponge.

529.1030 Formalin solution.

529.1044 Gentamicin sulfate in certain other dosage forms.

529.1044a Gentamicin sulfate intrauterine solution.

529.1044b Gentamicin sulfate solution.

529.1115 Halothane.

529.1186 Isoflurane.

529.1526 Nifurpirinol capsules.

529.2090 Salicylic acid.

529.2464 Ticarcillin powder.

529.2503 Tricaine methanesulfonate.

AUTHORITY: 21 U.S.C. 360b.

SOURCE: 40 FR 13881, Mar. 27, 1975, unless otherwise noted.

§ 529.50 Amikacin sulfate intrauterine solution.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 250 milligrams of amikacin (as the sulfate).

(b) *Sponsor.* See No. 000856 and 059130 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* Two grams (8 milliliters) diluted with 200 milliliters of sterile physiological saline per day for 3 consecutive days.

(2) *Indications for use.* For treating genital tract infections (endometritis, metritis, and pyometra) in mares when caused by susceptible organisms including *E. coli*, *Pseudomonas* spp., and *Klebsiella* spp.

(3) *Limitations.* For intrauterine infusion in the horse only. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 9640, Mar. 8, 1983, as amended at 53 FR 27852, July 25, 1988; 62 FR 15110, Mar. 31, 1997; 62 FR 23358, Apr. 30, 1997]

§ 529.360 Cephalothin discs.

(a) *Specifications.* Cephalothin discs have a uniform potency of 30 micrograms cephalothin per disc.

(b) *Sponsor.* See No. 000986 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The discs are used for determining the in vitro susceptibility of bacteria to cephaloridine and cephalonium.

(2) For veterinary laboratory diagnosis only.

[40 FR 13881, Mar. 27, 1975, as amended at 64 FR 404, Jan. 5, 1999]

EFFECTIVE DATE NOTE: At 64 FR 404, Jan. 5, 1999, § 529.360 was amended by removing the phrase “, comply with the requirements of § 460.1 of this chapter” from paragraph (a) and adding in its place “have a uniform potency of 30 micrograms cephalothin per disc”, effective May 20, 1999.

§ 529.400 Chlorhexidine tablets and suspension.

(a) *Specification.* Each tablet and each 28-milliliter syringe of suspension contain 1 gram of chlorhexidine dihydrochloride.¹

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

Food and Drug Administration, HHS

§ 529.1030

(c) *Conditions of use*—(1) *Amount*. Place 1 or 2 tablets deep in each uterine horn; or infuse a solution of 1 tablet dissolved in an appropriate amount of clean boiled water; or infuse one syringe of suspension into the uterus.¹

(2) *Indications for use*. For prevention or treatment of metritis and vaginitis in cows and mares when caused by pathogens sensitive to chlorhexidine dihydrochloride.¹

(3) *Limitations*. Prior to administration, remove any unattached placental membranes, any excess uterine fluid or debris, and carefully clean external genitalia. Use a clean, sterile inseminating pipette for administering solutions and suspensions. Treatment may be repeated in 48 to 72 hours.¹

[43 FR 10705, Feb. 23, 1979]

§ 529.469 Competitive exclusion culture.

(a) *Specifications*. Each packet of lyophilized culture contains either 2,000 or 5,000 doses in frozen pellets to be reconstituted for use.

(1) For 2,000-dose packet, add contents of one 2,000-dose packet of reconstitution powder to 490 milliliters of deionized water. Mix. Add contents of one 2,000-dose packet of lyophilized culture. Mix thoroughly.

(2) For 5,000-dose packet, add contents of one 5,000-dose packet of reconstitution powder to 1,250 milliliters of deionized water. Mix. Add contents of one 5,000-dose packet of lyophilized culture. Mix thoroughly. Allow to stand for 45 minutes before use. Use within 5 hours of reconstitution.

(b) *Sponsor*. See No. 032761 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*. *Chickens*—(1) *Amount*. Apply 25 milliliters of reconstituted culture as a topical spray on each tray of 100 chicks (0.25 milliliter per chick).

(2) *Indications for use*. For early establishment of intestinal microflora in chickens to reduce *Salmonella* colonization.

(3) *Limitations*. Administer as soon as possible after hatch, preferably at less than 1 day of age. Expose chicks to light for at least 5 minutes after spray treatment to encourage preening for oral uptake of the organisms. Provide

access to feed and water as soon as possible after treatment. Do not administer antibiotics to treated chickens.

[63 FR 25164, May 7, 1998]

§ 529.1003 Flurogestone acetate-impregnated vaginal sponge.

(a) *Specifications*. Each vaginal sponge contains 20 milligrams of flurogestone acetate.

(b) *Sponsor*. See No. 000014 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Indications for use*. For synchronizing estrus/ovulation in cycling adult ewes during their normal breeding season.

(2) *Limitations*. Using applicator provided, insert sponge into ewe's vagina 13 days before desired start of breeding. For intravaginal use in sheep only. Do not use in young ewes that have not had lambs. Use plastic or rubber gloves when handling large numbers of sponges to minimize exposure to drug. Do not leave sponge in the vagina for more than 21 days. Ewes must not be slaughtered for food within 30 days of sponge removal.

[49 FR 45420, Nov. 16, 1984]

§ 529.1030 Formalin solution.

(a) *Specifications*. Formalin solution is an aqueous solution containing approximately 37 percent by weight of formaldehyde gas, U.S.P.

(b) *Sponsor*. Approval to firms identified in § 510.600(c) of this chapter for use as indicated:

(1) No. 050378 for use as in paragraphs (d)(1)(iii), (d)(1)(iv), (d)(1)(v), (d)(2)(iii), (d)(2)(iv), (d)(2)(v), and (d)(3).

(2) Nos. 049968 and 051212 for use as in paragraphs (d)(1)(i), (d)(1)(ii), (d)(2)(i), (d)(2)(ii), and (d)(3).

(c) [Reserved]

(d) *Conditions of use*. It is added to environmental water as follows:

(1) *Indications for use*. (i) Select finfish. For control of external protozoa *Ichthyophthirius* spp., *Chilodonella* spp., *Costia* spp., *Scyphidia* spp., *Epistylis* spp., and *Trichodina* spp., and monogenetic trematodes *Cleidodiscus* spp., *Gyrodactylus* spp., and *Dactylogyrus* spp., on salmon, trout, catfish, largemouth bass, and bluegill.

§ 529.1044

21 CFR Ch. I (4–1–99 Edition)

(ii) Select finfish eggs. For control of fungi of the family Saprolegniaceae on salmon, trout, and esocid eggs.

(iii) Penaeid shrimp. For control of external protozoan parasites *Bodo* spp., *Epistylis* spp., and *Zoothamnium* spp.

(iv) All finfish. For control of external protozoa *Ichthyophthirius* spp., *Chilodonella* spp., *Costia* spp., *Scyphidia* spp., *Epistylis* spp., and *Trichodina* spp., and monogenetic trematodes *Cleidodiscus* spp., *Gyrodactylus* spp., and *Dactylogyrus* spp.

(v) All finfish eggs: For control of fungi of the family Saprolegniaceae.

(2) *Amount*. The drug concentrations required are as follows:

(i) For control of external parasites on select finfish:

Fish	Concentration of formalin (microliters per liter)	
	Tanks and raceways (for up to 1 hour)	Earthen ponds (indefinitely)
Salmon and trout:		
Above 50 ° F	Up to 170	15–25
Below 50 ° F	Up to 250	15–25
Catfish, largemouth bass, and bluegill.	Up to 250	¹ 15–25

¹ Use the lower concentrations when pond is heavily loaded with fish or phytoplankton.

(ii) For control of fungi of the Saprolegniaceae on salmon, trout, and esocid eggs: Apply in constant flow water supply of incubating facilities for 15 minutes. Concentration of formalin used is 1,000 to 2,000 microliters per liter.

(iii) For control of external protozoan parasites on shrimp:

Shrimp	Concentration of formalin (microliters per liter)	
	Tanks and raceways (up to 4 hours daily)	Earthen ponds (single treatment)
Penaeid Shrimp ...	50 to 100 ¹	25 ²

¹ Treat for up to 4 hours daily. Treatment may be repeated daily until parasite control is achieved. Use the lower concentration when the tanks and raceways are heavily loaded.

² Single treatment. Treatment may be repeated in 5 to 10 days if needed.

(iv) For control of external parasites on all finfish:

Aquatic species	Administer in tanks and raceways for up to 1 hour (microliter/liter or part per million (μL/L or ppm))	Administer in earthen ponds indefinitely (μL/L or ppm)
Salmon and trout:		

Aquatic species	Administer in tanks and raceways for up to 1 hour (microliter/liter or part per million (μL/L or ppm))	Administer in earthen ponds indefinitely (μL/L or ppm)
Above 50 °F	Up to 170	15 to 25 ^{1, 2}
Below 50 °F	Up to 250	15 to 25 ^{1, 2}
All other finfish	Up to 250	15 to 25 ^{1, 2}

¹ Use the lower concentration when ponds, tanks, or raceways are heavily loaded with phytoplankton or fish to avoid oxygen depletion due to the biological oxygen demand by decay of dead phytoplankton. Alternatively, a higher concentration may be used if dissolved oxygen is strictly monitored.

² Although the indicated concentrations are considered safe for cold and warm water finfish, a small number of each lot or pond to be treated should always be used to check for any unusual sensitivity to formalin before proceeding.

(v) For control of fungi of the family Saprolegniaceae on all finfish eggs: Eggs of all finfish except Acipenseriformes, 1,000 to 2,000 μL/L (ppm) for 15 minutes; eggs of Acipenseriformes, up to 1,500 μL/L (ppm) for 15 minutes.

(3) *Limitations*. Fish tanks and raceways may be treated daily until parasite control is achieved. Pond treatment may be repeated in 5 to 10 days if needed. However, pond treatments for *Ichthyophthirius* should be made at 2-day intervals until control is achieved. Egg tanks may be treated as often as necessary to prevent growth of fungi. Do not use formalin which has been subjected to temperatures below 40 °F, or allowed to freeze. Do not treat ponds containing striped bass. Treatments in tanks should never exceed 1 hour even if fish show no signs of stress. Do not apply formalin to ponds with water warmer than 27 °C (80 °F), when a heavy bloom of phytoplankton is present, or when the concentration of dissolved oxygen is less than 5 milligrams per liter.

[51 FR 11441, Apr. 3, 1986, as amended at 58 FR 59169, Nov. 8, 1993; 59 FR 60076, Nov. 22, 1994; 63 FR 38304, July 16, 1998]

§ 529.1044 Gentamicin sulfate in certain other dosage forms.

§ 529.1044a Gentamicin sulfate intra-uterine solution.

(a) *Specifications*. Each milliliter of the drug contains 50 or 100 milligrams of gentamicin (as the sulfate) in sterile aqueous solution.

(b) *Sponsor*. See Nos. 000010, 000061, 000856, 000864, 057561, and 059130 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is indicated for use for control of bacterial infections of the uterus in horses (metritis) and as an aid in improving conception in mares with uterine infections caused by bacteria sensitive to gentamicin.

(2) It is administered at a dosage level of 2 to 2.5 grams per day for 3 to 5 days during estrus, each dose being diluted with 200 to 500 milliliters of sterile physiological saline before aseptic infusion into the uterus.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(4) Not for use in horses intended for food.

[40 FR 13881, Mar. 27, 1975, as amended at 40 FR 48676, Oct. 17, 1975; 48 FR 31386, July 8, 1983; 52 FR 7833, Mar. 13, 1987; 58 FR 14314, Mar. 17, 1993; 59 FR 31140, June 17, 1994; 60 FR 45042, Aug. 30, 1995; 60 FR 48894, Sept. 21, 1995; 61 FR 17830, Apr. 23, 1996; 62 FR 611, Jan. 6, 1997; 62 FR 35077, June 30, 1997; 63 FR 6644, Feb. 10, 1998]

§ 529.1044b Gentamicin sulfate solution.

(a) *Specifications.* Each milliliter of solution contains gentamicin sulfate equivalent to 50 milligrams of gentamicin base.

(b) *Sponsor.* See Nos. 000061 and 051259 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is recommended as an aid in the reduction or elimination of the following microorganisms from turkey-hatching eggs: *Arizona hinshawii* (paracolon), *Salmonella st. paul*, and *Mycoplasma meleagridis*.

(2) The drug is added to clean water to provide a dip solution with a gentamicin concentration of 250 to 1,000 parts per million. A concentration of 500 parts per million is recommended. Clean eggs should be held submerged in the gentamicin solution under a vacuum of about 27.5 to 38 centimeters of mercury for 5 minutes followed by additional soaking in gentamicin solution for approximately 10 minutes at atmospheric pressure. Eggs can also be treated by warming them for 3 to 6 hours at approximately 100° F. then immediately submerging them in gentamicin solution maintained at about 40° F., keeping the eggs submerged for 10 to 15 minutes.

(3) For use in the dipping treatment of turkey-hatching eggs only. Eggs which have been dipped in the drug shall not be used for food.

[40 FR 13881, Mar. 27, 1975, as amended at 52 FR 7833, Mar. 13, 1987; 62 FR 22889, Apr. 28, 1997]

§ 529.1115 Halothane.

(a) *Specifications.* The drug is a colorless, odorless, nonflammable, nonexplosive, heavy liquid containing 0.01 percent thymol as a preservative.

(b) *Sponsor.* See 000856 and 012164 in § 510.600(c) of this chapter.

(c) *Conditions of use.*—(1) *Amount.* Two to 5 percent of inhaled atmosphere for induction of anesthesia; 0.5 to 2 percent for maintenance of anesthesia.¹

(2) *Indications for use.* For nonfood animals for the induction and maintenance of anesthesia.¹

(3) *Limitations.* Administered by inhalation. May be administered with either oxygen or a mixture of oxygen and nitrous oxide. Place drug vaporizer between the gas supply and breathing bag to prevent overdosage. Not recommended for obstetrical anesthesia except when uterine relaxation is required. Do not use in pregnant animals; information on possible adverse effects on fetal development is not available. Operating rooms should have adequate ventilation to prevent accumulation of anesthetic gases. Not for use in animals intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[46 FR 27915, May 22, 1981, as amended at 62 FR 29014, May 29, 1997]

§ 529.1186 Isoflurane.

(a) *Specifications.* The drug is a clear, colorless, stable liquid containing no additives or chemical stabilizers. It is nonflammable and nonexplosive.

(b) *Sponsors.* See Nos. 000074, 000209, 010019, 012164, and 059258 in § 510.600(c) of this chapter.

¹These conditions have been reviewed by FDA and found effective. NADA's for similar products for these conditions of use need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

§ 529.1526

(c) *Conditions of use*—(1) *Amount*—(i) *Horses*: For induction of surgical anesthesia: 3 to 5 percent isoflurane (with oxygen) for 5 to 10 minutes. For maintenance of surgical anesthesia: 1.5 to 1.8 percent isoflurane (with oxygen).

(ii) *Dogs*: For induction of surgical anesthesia: 2 to 2.5 percent isoflurane (with oxygen) for 5 to 10 minutes. For maintenance of surgical anesthesia: 1.5 to 1.8 percent isoflurane (with oxygen).

(2) *Indications for use*. For induction and maintenance of general anesthesia in horses and dogs.

(3) *Limitations*. Administer by inhalation; not for use in horses or dogs sensitive to halogenated agents; increasing depth of anesthesia may increase hypotension and respiratory depression; use less than usual amounts of nondepolarizing relaxants; use with vaporizers producing predictable percentage concentrations; not for use in horses intended for food; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[51 FR 594, Jan. 7, 1986, as amended at 54 FR 23472, June 1, 1989; 58 FR 17346, Apr. 2, 1993; 59 FR 44315, Aug. 29, 1994; 60 FR 40456, Aug. 9, 1995; 63 FR 8122, Feb. 18, 1998; 63 FR 24106, May 1, 1998]

§ 529.1526 Nifurpirinol capsules.

(a) *Specifications*. Each capsule contains 3.8 or 7.6 milligrams of nifurpirinol.

(b) *Sponsor*. See No. 000074 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) The drug is used in treating aquarium fish for the control of columnaris disease caused by *Chondrococcus columnaris* susceptible to nifurpirinol.

(2) Use one 3.8 milligram nifurpirinol capsule for each 10 gallons of aquarium water. Empty the contents of the capsule directly into the water and stir briefly. Treat for at least 1 hour. If activated charcoal or carbon filtration is being used, disconnect during treatment, but maintain adequate aeration. Resume water filtration after 1 hour treatment. Usually a single treatment is sufficient. For aquariums with charcoal filters, nifurpirinol can be used once each 24 hours up to 3 consecutive days, discontinuing filtration during treatment. If aquarium does not have

21 CFR Ch. I (4–1–99 Edition)

charcoal filter, do not retreat within 5 days.

(3) Do not use in salt water aquariums.

(4) Do not use while egg bearers or live bearers are reproducing.

[40 FR 60052, Dec. 31, 1975, as amended at 47 FR 20758, May 14, 1982; 56 FR 43699, Sept. 4, 1991]

§ 529.2090 Salicylic acid.

(a) *Specifications*. (1) Each dose contains 0.55 grain of salicylic acid in a gum arabic and dextrin vehicle.

(2) Each dose is incorporated upon a device (teat dilator) suitable for insertion into and subsequent removal from the teat canal.

(b) *Sponsor*. See No. 045087 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) The drug is used for the removal of scar tissue in the teat canal of milk-producing cows.

(2) The labeling bears directions to the user to:

(i) Treat lactating cows initially by inserting dosage and removal of the device;

(ii) Insert second dose and permit device to remain in canal until the next milking; and

(iii) Insert one dose following each milking for not more than 2 days.

(3) Milk that has been drawn from animals within 48 hours of such treatment may not be used for food.

[41 FR 10984, Mar. 15, 1976, as amended at 43 FR 29290, July 7, 1978; 55 FR 29842, July 23, 1990; 55 FR 31481, Aug. 2, 1990; 62 FR 8372, Feb. 25, 1997]

§ 529.2464 Ticarcillin powder.

(a) *Specifications*. Each vial contains ticarcillin disodium equivalent to 6 grams of ticarcillin to be reconstituted with 25 milliliters of sterile water for injection or sterile physiological saline.

(b) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 6 grams per day, intrauterine, for 3 consecutive days during estrus.

(2) *Indications for use*. *Horses*. Intrauterine treatment of endometritis caused by beta-hemolytic streptococci.

(3) *Limitations*. For intrauterine use in horses only. Infuse aseptically. Not

Food and Drug Administration, HHS

§ 530.1

for use in horses raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37336, Aug. 18, 1992, as amended at 60 FR 55660, Nov. 2, 1995]

§ 529.2503 Tricaine methanesulfonate.

(a) *Chemical name.* Ethyl-*m*-amino-benzoate methanesulfonate.

(b) *Sponsor.* See Nos. 050378 and 051212 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used for the temporary immobilization of fish, amphibians, and other aquatic cold-blooded animals (poikilotherms) as an aid in handling during manual spawning (fish stripping), weighing, measuring, marking, surgical operations, transport, photography, and research.

(2) It is used as follows:

(i) For fish the drug is added to ambient water at a concentration of from 15 to 330 milligrams per liter depending upon the degree of anesthetization or sedation desired, the species and size of the fish, and the temperature and softness of the water. Preliminary tests of solutions must be made with small numbers of fish to determine the desired rates of sedation or anesthesia and the appropriate exposure times for the specific lots of fish under prevailing conditions.

(ii) For amphibians and other aquatic coldblooded animals, the drug is added to ambient water in concentrations of from 1:1000 to 1:20,000 depending upon species and stage of development.

(iii) Do not use within 21 days of harvesting fish for food. Use in fish intended for food should be restricted to Ictaluridae, Salmonidae, Esocidae, and Percidae, and water temperature should exceed 10° C. (50° F.). In other fish and in cold-blooded animals, the drug should be limited to hatchery or laboratory use.

[40 FR 13881, Mar. 27, 1975, as amended at 49 FR 5748, Feb. 15, 1984; 51 FR 11439, Apr. 3, 1986; 63 FR 7702, Feb. 17, 1998]

PART 530—EXTRALABEL DRUG USE IN ANIMALS

Subpart A—General Provisions

Sec.

530.1 Scope.

530.2 Purpose.

530.3 Definitions.

530.4 Advertising and promotion.

530.5 Veterinary records.

Subpart B—Rules and Provisions for Extralabel Uses of Drugs in Animals

530.10 Provision permitting extralabel use of animal drugs.

530.11 Limitations.

530.12 Labeling.

530.13 Extralabel use from compounding of approved new animal and approved human drugs.

Subpart C—Specific Provisions Relating to Extralabel Use of Animal and Human Drugs in Food-Producing Animals

530.20 Conditions for permitted extralabel animal and human drug use in food-producing animals.

530.21 Prohibitions for food-producing animals.

530.22 Safe levels and analytical methods for food-producing animals.

530.23 Procedure for setting and announcing safe levels.

530.24 Procedure for announcing analytical methods for drug residue quantification.

530.25 Orders prohibiting extralabel uses for drugs in food-producing animals.

Subpart D—Extralabel Use of Human and Animal Drugs in Animals Not Intended for Human Consumption

530.30 Extralabel drug use in nonfood animals.

Subpart E—Safe Levels for Extralabel Use of Drugs in Animals and Drugs Prohibited From Extralabel Use in Animals

530.40 Safe levels and availability of analytical methods.

530.41 Drugs prohibited for extralabel use in animals.

AUTHORITY: 15 U.S.C. 1453, 1454, 1455; 21 U.S.C. 321, 331, 351, 352, 353, 355, 357, 360b, 371, 379e.

SOURCE: 61 FR 57743, Nov. 7, 1996, unless otherwise noted.

Subpart A—General Provisions

§ 530.1 Scope.

This part applies to the extralabel use in an animal of any approved new animal drug or approved new human drug by or on the lawful order of a licensed veterinarian within the context

§ 530.2

21 CFR Ch. I (4–1–99 Edition)

of a valid veterinary-client-patient relationship.

§ 530.2 Purpose.

The purpose of this part is to establish conditions for extralabel use or intended extralabel use in animals by or on the lawful order of licensed veterinarians of Food and Drug Administration approved new animal drugs and approved new human drugs. Such use is limited to treatment modalities when the health of an animal is threatened or suffering or death may result from failure to treat. This section implements the Animal Medicinal Drug Use Clarification Act of 1994 (the AMDUCA) (Pub. L. 103–396).

§ 530.3 Definitions.

(a) *Extralabel use* means actual use or intended use of a drug in an animal in a manner that is not in accordance with the approved labeling. This includes, but is not limited to, use in species not listed in the labeling, use for indications (disease or other conditions) not listed in the labeling, use at dosage levels, frequencies, or routes of administration other than those stated in the labeling, and deviation from the labeled withdrawal time based on these different uses.

(b) *FDA* means the U.S. Food and Drug Administration.

(c) The phrase *a reasonable probability that a drug's use may present a risk to the public health* means that FDA has reason to believe that use of a drug may be likely to cause a potential adverse event.

(d) The phrase *use of a drug may present a risk to the public health* means that FDA has information that indicates that use of a drug may cause an adverse event.

(e) The phrase *use of a drug presents a risk to the public health* means that FDA has evidence that demonstrates that the use of a drug has caused or likely will cause an adverse event.

(f) A *residue* means any compound present in edible tissues that results from the use of a drug, and includes the drug, its metabolites, and any other substance formed in or on food because of the drug's use.

(g) A *safe level* is a conservative estimate of a drug residue level in edible

animal tissue derived from food safety data or other scientific information. Concentrations of residues in tissue below the safe level will not raise human food safety concerns. A safe level is not a safe concentration or a tolerance and does not indicate that an approval exists for the drug in that species or category of animal from which the food is derived.

(h) *Veterinarian* means a person licensed by a State or Territory to practice veterinary medicine.

(i) A *valid veterinarian-client-patient relationship* is one in which:

(1) A veterinarian has assumed the responsibility for making medical judgments regarding the health of (an) animal(s) and the need for medical treatment, and the client (the owner of the animal or animals or other caretaker) has agreed to follow the instructions of the veterinarian;

(2) There is sufficient knowledge of the animal(s) by the veterinarian to initiate at least a general or preliminary diagnosis of the medical condition of the animal(s); and

(3) The practicing veterinarian is readily available for followup in case of adverse reactions or failure of the regimen of therapy. Such a relationship can exist only when the veterinarian has recently seen and is personally acquainted with the keeping and care of the animal(s) by virtue of examination of the animal(s), and/or by medically appropriate and timely visits to the premises where the animal(s) are kept.

§ 530.4 Advertising and promotion.

Nothing in this part shall be construed as permitting the advertising or promotion of extralabel uses in animals of approved new animal drugs or approved human drugs.

§ 530.5 Veterinary records.

(a) As a condition of extralabel use permitted under this part, to permit FDA to ascertain any extralabel use or intended extralabel use of drugs that the agency has determined may present a risk to the public health, veterinarians shall maintain the following records of extralabel uses. Such records

shall be legible, documented in an accurate and timely manner, and be readily accessible to permit prompt retrieval of information. Such records shall be adequate to substantiate the identification of the animals and shall be maintained either as individual records or, in food animal practices, on a group, herd, flock, or per-client basis. Records shall be adequate to provide the following information:

- (1) The established name of the drug and its active ingredient, or if formulated from more than one ingredient, the established name of each ingredient;
- (2) The condition treated;
- (3) The species of the treated animal(s);
- (4) The dosage administered;
- (5) The duration of treatment;
- (6) The numbers of animals treated; and
- (7) The specified withdrawal, withholding, or discard time(s), if applicable, for meat, milk, eggs, or any food which might be derived from any food animals treated.

(b) A veterinarian shall keep all required records for 2 years or as otherwise required by Federal or State law, whichever is greater.

(c) Any person who is in charge, control, or custody of such records shall, upon request of a person designated by FDA, permit such person designated by FDA to, at all reasonable times, have access to, permit copying, and verify such records.

Subpart B—Rules and Provisions for Extralabel Uses of Drugs in Animals

§ 530.10 Provision permitting extralabel use of animal drugs.

An approved new animal drug or human drug intended to be used for an extralabel purpose in an animal is not unsafe under section 512 of the act and is exempt from the labeling requirements of section 502(f) of the act if such use is:

- (a) By or on the lawful written or oral order of a licensed veterinarian within the context of a valid veterinarian-client-patient relationship; and
- (b) In compliance with this part.

§ 530.11 Limitations.

In addition to uses which do not comply with the provision set forth in § 530.10, the following specific extralabel uses are not permitted and result in the drug being deemed unsafe within the meaning of section 512 of the act:

- (a) Extralabel use in an animal of an approved new animal drug or human drug by a lay person (except when under the supervision of a licensed veterinarian);
- (b) Extralabel use of an approved new animal drug or human drug in or on an animal feed;
- (c) Extralabel use resulting in any residue which may present a risk to the public health; and
- (d) Extralabel use resulting in any residue above an established safe level, safe concentration or tolerance.

§ 530.12 Labeling.

Any human or animal drug prescribed and dispensed for extralabel use by a veterinarian or dispensed by a pharmacist on the order of a veterinarian shall bear or be accompanied by labeling information adequate to assure the safe and proper use of the product. Such information shall include the following:

(a) The name and address of the prescribing veterinarian. If the drug is dispensed by a pharmacy on the order of a veterinarian, the labeling shall include the name of the prescribing veterinarian and the name and address of the dispensing pharmacy, and may include the address of the prescribing veterinarian;

(b) The established name of the drug or, if formulated from more than one active ingredient, the established name of each ingredient;

(c) Any directions for use specified by the veterinarian, including the class/species or identification of the animal or herd, flock, pen, lot, or other group of animals being treated, in which the drug is intended to be used; the dosage, frequency, and route of administration; and the duration of therapy;

(d) Any cautionary statements; and

(e) The veterinarian's specified withdrawal, withholding, or discard time for meat, milk, eggs, or any other food

§ 530.13

which might be derived from the treated animal or animals.

§ 530.13 Extralabel use from compounding of approved new animal and approved human drugs.

(a) This part applies to compounding of a product from approved animal or human drugs by a veterinarian or a pharmacist on the order of a veterinarian within the practice of veterinary medicine. Nothing in this part shall be construed as permitting compounding from bulk drugs.

(b) Extralabel use from compounding of approved new animal or human drugs is permitted if:

(1) All relevant portions of this part have been complied with;

(2) There is no approved new animal or approved new human drug that, when used as labeled or in conformity with criteria established in this part, will, in the available dosage form and concentration, appropriately treat the condition diagnosed. Compounding from a human drug for use in food-producing animals will not be permitted if an approved animal drug can be used for the compounding;

(3) The compounding is performed by a licensed pharmacist or veterinarian within the scope of a professional practice;

(4) Adequate procedures and processes are followed that ensure the safety and effectiveness of the compounded product;

(5) The scale of the compounding operation is commensurate with the established need for compounded products (e.g., similar to that of comparable practices); and

(6) All relevant State laws relating to the compounding of drugs for use in animals are followed.

(c) Guidance on the subject of compounding may be found in guidance documents issued by FDA.

21 CFR Ch. I (4–1–99 Edition)

Subpart C—Specific Provisions Relating to Extralabel Use of Animal and Human Drugs in Food-Producing Animals

§ 530.20 Conditions for permitted extralabel animal and human drug use in food-producing animals.

(a) The following conditions must be met for a permitted extralabel use in food-producing animals of approved new animal and human drugs:

(1) There is no approved new animal drug that is labeled for such use and that contains the same active ingredient which is in the required dosage form and concentration, except where a veterinarian finds, within the context of a valid veterinarian-client-patient relationship, that the approved new animal drug is clinically ineffective for its intended use.

(2) Prior to prescribing or dispensing an approved new animal or human drug for an extralabel use in food animals, the veterinarian must:

(i) Make a careful diagnosis and evaluation of the conditions for which the drug is to be used;—

(ii) Establish a substantially extended withdrawal period prior to marketing of milk, meat, eggs, or other edible products supported by appropriate scientific information, if applicable;

(iii) Institute procedures to assure that the identity of the treated animal or animals is carefully maintained; and

(iv) Take appropriate measures to assure that assigned timeframes for withdrawal are met and no illegal drug residues occur in any food-producing animal subjected to extralabel treatment.

(b) The following additional conditions must be met for a permitted extralabel use of in food-producing animals an approved human drug, or of an animal drug approved only for use in animals not intended for human consumption:

(1) Such use must be accomplished in accordance with an appropriate medical rationale; and

(2) If scientific information on the human food safety aspect of the use of the drug in food-producing animals is not available, the veterinarian must take appropriate measures to assure that the animal and its food products will not enter the human food supply.

(c) Extralabel use of an approved human drug in a food-producing animal is not permitted under this part if an animal drug approved for use in food-producing animals can be used in an extralabel manner for the particular use.

§ 530.21 Prohibitions for food-producing animals.

(a) FDA may prohibit the extralabel use of an approved new animal or human drug or class of drugs in food-producing animals if FDA determines that:

(1) An acceptable analytical method needs to be established and such method has not been established or cannot be established; or

(2) The extralabel use of the drug or class of drugs presents a risk to the public health.

(b) A prohibition may be a general ban on the extralabel use of the drug or class of drugs or may be limited to a specific species, indication, dosage form, route of administration, or combination of factors.

§ 530.22 Safe levels and analytical methods for food-producing animals.

(a) FDA may establish a safe level for extralabel use of an approved human drug or an approved new animal drug when the agency finds that there is a reasonable probability that an extralabel use may present a risk to the public health. FDA may:

(1) Establish a finite safe level based on residue and metabolism information from available sources;

(2) Establish a safe level based on the lowest level that can be measured by a practical analytical method; or

(3) Establish a safe level based on other appropriate scientific, technical, or regulatory criteria.

(b) FDA may require the development of an acceptable analytical method for the quantification of residues above any safe level established under

this part. If FDA requires the development of such an acceptable analytical method, the agency will publish notice of that requirement in the FEDERAL REGISTER.

(c) The extralabel use of an animal drug or human drug that results in residues exceeding a safe level established under this part is an unsafe use of such drug.

(d) If the agency establishes a safe level for a particular species or category of animals and a tolerance or safe concentration is later established through an approval for that particular species or category of animals, for that species or category of animals, the safe level is superseded by the tolerance or safe concentration for that species or category of animals.

§ 530.23 Procedure for setting and announcing safe levels.

(a) FDA may issue an order establishing a safe level for a residue of an extralabel use of an approved human drug or an approved animal drug. The agency will publish in the FEDERAL REGISTER a notice of the order. The notice will include:

(1) A statement setting forth the agency's finding that there is a reasonable probability that extralabel use in animals of the human drug or animal drug may present a risk to the public health;

(2) A statement of the basis for that finding; and

(3) A request for public comments.

(b) A current listing of those drugs for which a safe level for extralabel drug use in food-producing animals has been established, the specific safe levels, and the availability, if any, of a specific analytical method or methods for drug residue detection will be codified in § 530.40.

§ 530.24 Procedure for announcing analytical methods for drug residue quantification.

(a) FDA may issue an order announcing a specific analytical method or methods for the quantification of extralabel use drug residues above the safe levels established under § 530.22 for extralabel use of an approved human drug or an approved animal drug. The agency will publish in the FEDERAL

§ 530.25

21 CFR Ch. I (4–1–99 Edition)

REGISTER a notice of the order, including the name of the specific analytical method or methods and the drug or drugs for which the method is applicable.

(b) Copies of analytical methods for the quantification of extralabel use drug residues above the safe levels established under § 530.22 will be available upon request from the Communications and Education Branch (HFV–12), Division of Program Communication and Administrative Management, Center for Veterinary Medicine, 7500 Standish Pl., Rockville, MD 20855. When an analytical method for the detection of extralabel use drug residues above the safe levels established under § 530.22 is developed, and that method is acceptable to the agency, FDA will incorporate that method by reference.

§ 530.25 Orders prohibiting extralabel uses for drugs in food-producing animals.

(a) FDA may issue an order prohibiting extralabel use of an approved new animal or human drug in food-producing animals if the agency finds, after providing an opportunity for public comment, that:

(1) An acceptable analytical method required under § 530.22 has not been developed, submitted, and found to be acceptable by FDA or that such method cannot be established; or

(2) The extralabel use in animals presents a risk to the public health.—

(b) After making a determination that the analytical method required under § 530.22 has not been developed and submitted, or that such method cannot be established, or that an extralabel use in animals of a particular human drug or animal drug presents a risk to the public health, FDA will publish in the FEDERAL REGISTER, with a 90-day delayed effective date, an order of prohibition for an extralabel use of a drug in food-producing animals. Such order shall state that an acceptable analytical method required under § 530.22 has not been developed, submitted, and found to be acceptable by FDA; that such method cannot be established; or that the extralabel use in animals presents a risk to the public health; and shall:

(1) Specify the nature and extent of the order of prohibition and the reasons for the prohibition;

(2) Request public comments; and

(3) Provide a period of not less than 60 days for comments.

(c) The order of prohibition will become effective 90 days after date of publication of the order unless FDA publishes a notice in the FEDERAL REGISTER prior to that date, that revokes the order of prohibition, modifies it, or extends the period of public comment.

(d) The agency may publish an order of prohibition with a shorter comment period and/or delayed effective date than specified in paragraph (b) of this section in exceptional circumstances (e.g., where there is immediate risk to the public health), provided that the order of prohibition states that the comment period and/or effective date have been abbreviated because there are exceptional circumstances, and the order of prohibition sets forth the agency's rationale for taking such action.

(e) If FDA publishes a notice in the FEDERAL REGISTER modifying an order of prohibition, the agency will specify in the modified order of prohibition the nature and extent of the modified prohibition, the reasons for it, and the agency's response to any comments on the original order of prohibition.

(f) A current listing of drugs prohibited for extralabel use in animals will be codified in § 530.41.

(g) After the submission of appropriate information (i.e., adequate data, an acceptable method, approval of a new animal drug application for the prohibited extralabel use, or information demonstrating that the prohibition was based on incorrect data), FDA may, by publication of an appropriate notice in the FEDERAL REGISTER, remove a drug from the list of human and animal drugs prohibited for extralabel use in animals, or may modify a prohibition.

(h) FDA may prohibit extralabel use of a drug in food-producing animals without establishing a safe level.

Subpart D—Extralabel Use of Human and Animal Drugs in Animals Not Intended for Human Consumption

§ 530.30 Extralabel drug use in nonfood animals.

(a) Because extralabel use of animal and human drugs in nonfood-producing animals does not ordinarily pose a threat to the public health, extralabel use of animal and human drugs is permitted in nonfood-producing animal practice except when the public health is threatened. In addition, the provisions of § 530.20(a)(1) will apply to the use of an approved animal drug.

(b) If FDA determines that an extralabel drug use in animals not intended for human consumption presents a risk to the public health, the agency may publish in the FEDERAL REGISTER a notice prohibiting such use following the procedures in § 530.25. The prohibited extralabel drug use will be codified in § 530.41.

Subpart E—Safe Levels for Extralabel Use of Drugs in Animals and Drugs Prohibited From Extralabel Use in Animals

§ 530.40 Safe levels and availability of analytical methods.

(a) In accordance with § 530.22, the following safe levels for extralabel use of an approved animal drug or human drug have been established:

[Reserved]

(b) In accordance with § 530.22, the following analytical methods have been accepted by FDA:

[Reserved]

§ 530.41 Drugs prohibited for extralabel use in animals.

(a) The following drugs, families of drugs, and substances are prohibited for extralabel animal and human drug uses in food-producing animals.

- (1) Chloramphenicol;
- (2) Clenbuterol;
- (3) Diethylstilbestrol (DES);
- (4) Dimetridazole;
- (5) Iprnidazole;
- (6) Other nitroimidazoles;

(7) Furazolidone (except for approved topical use);

(8) Nitrofurazone (except for approved topical use);

(9) Sulfonamide drugs in lactating dairy cattle (except approved use of sulfadimethoxine, sulfabromomethazine, and sulfaethoxypyridazine);

(10) Fluoroquinolones; and

(11) Glycopeptides.

(b) The following drugs, families of drugs, and substances are prohibited for extralabel animal and human drug uses in nonfood-producing animals: [Reserved]

[62 FR 27947, May 22, 1997]

PART 556—TOLERANCES FOR RESIDUES OF NEW ANIMAL DRUGS IN FOOD

Subpart A—General Provisions

Sec.

556.1 General considerations; tolerances for residues of new animal drugs in food.

Subpart B—Specific Tolerances for Residues of New Animal Drugs

556.20	2-Acetylamino-5-nitrothiazole.
556.30	Aklomide.
556.34	Albendazole.
556.38	Amoxicillin.
556.40	Ampicillin.
556.50	Amprolium.
556.52	Apramycin.
556.60	Arsenic.
556.70	Bacitracin.
556.90	Buquinolate.
556.100	Carbadox.
556.110	Carbomycin.
556.113	Ceftiofur.
556.115	Cephapirin.
556.120	Chlorhexidine.
556.140	Chlorobutanol.
556.150	Chlortetracycline.
556.160	Clopidol.
556.163	Clorsulon.
556.165	Cloxacillin.
556.167	Colistimethate.
556.170	Decoquinat.
556.180	Dichlorvos.
556.200	Dihydrostreptomycin.
556.220	3,5-Dinitrobenzamide.
556.225	Doramectin.
556.227	Eprinomectin.
556.228	Enrofloxacin.
556.230	Erythromycin.
556.240	Estradiol and related esters.
556.260	Ethopabate.
556.270	Ethylenediamine.

§ 556.1

556.273 Famphur.
556.275 Fenbendazole.
556.277 Fenprostalene.
556.283 Florfenicol.
556.286 Flunixin meglumine.
556.290 Furazolidone.
556.300 Gentamicin sulfate.
556.308 Halofuginone hydrobromide.
556.310 Haloxon.
556.320 Hydrocortisone.
556.330 Hygromycin B.
556.344 Ivermectin.
556.347 Lasalocid.
556.350 Levamisole hydrochloride.
556.360 Lincomycin.
556.375 Maduramicin ammonium.
556.380 Melengestrol acetate.
556.390 Methylparaben.
556.400 Methylprednisolone.
556.410 Metoserpate hydrochloride.
556.420 Monensin.
556.425 Morantel tartrate.
556.426 Moxidectin.
556.428 Narasin.
556.430 Neomycin.
556.440 Nequinat.
556.445 Nicarbazin.
556.460 Novobiocin.
556.470 Nystatin.
556.480 Oleandomycin.
556.490 Ormetoprim.
556.495 Oxfendazole.
556.500 Oxytetracycline.
556.510 Penicillin.
556.515 Pirlimycin.
556.520 Prednisolone.
556.530 Prednisone.
556.540 Progesterone.
556.550 Propylparaben.
556.560 Pyrantel tartrate.
556.580 Robenidine hydrochloride.
556.590 Salicylic acid.
556.594 Sarafloxacin.
556.600 Spectinomycin.
556.610 Streptomycin.
556.620 Sulfabromomethazine sodium.
556.625 Sodium sulfachloropyrazine monohydrate.
556.630 Sulfachlorpyridazine.
556.640 Sulfadimethoxine.
556.650 Sulfaethoxyypyridazine.
556.660 Sulfamerazine.
556.670 Sulfamethazine.
556.680 Sulfanitran.
556.685 Sulfaquinoxaline.
556.690 Sulfathiazole.
556.700 Sulfomyxin.
556.710 Testosterone propionate.
556.720 Tetracycline.
556.730 Thiabendazole.
556.735 Tilimicosin.
556.738 Tiamulin.
556.739 Trenbolone.
556.740 Tylosin.
556.741 Tripeleminamine.
556.750 Virginiamycin.
556.760 Zeranol.

21 CFR Ch. I (4–1–99 Edition)

556.770 Zoalene.

AUTHORITY: 21 U.S.C. 342, 360b, 371.

SOURCE: 40 FR 13942, Mar. 27, 1975, unless otherwise noted.

Subpart A—General Provisions

§ 556.1 General considerations; tolerances for residues of new animal drugs in food.

(a) Tolerances established in this part are based upon residues of drugs in edible products of food-producing animals treated with such drugs. Consideration of an appropriate tolerance for a drug shall result in a conclusion either that:

(1) Finite residues will be present in the edible products—in which case a finite tolerance is required; or

(2) It is not possible to determine whether finite residues will be incurred but there is reasonable expectation that they may be present—in which case a tolerance for negligible residue is required; or

(3) The drug induces cancer when ingested by man or animal or, after tests which are appropriate for the evaluation of the safety of such drug, has been shown to induce cancer in man or animal; however, such drug will not adversely affect the animals for which it is intended, and no residue of such drug will be found by prescribed methods of analysis in any edible portion of such animals after slaughter or in any food yielded by or derived from the living animal—in which case the accepted method of analysis shall be published or cited, if previously published and available elsewhere, in this part; or

(4) It may or may not be possible to determine whether finite residues will be incurred but there is no reasonable expectation that they may be present—in which case the establishment of a tolerance is not required; or

(5) The drug is such that it may be metabolized and/or assimilated in such form that any possible residue would be indistinguishable from normal tissue constituents—in which case the establishment of a tolerance is not required.

(b) No tolerance established pursuant to paragraph (a)(1) of this section will

Food and Drug Administration, HHS

§ 556.60

be set at any level higher than that reflected by the permitted use of the drug.

(c) Any tolerance required pursuant to this section will, in addition to the toxicological considerations, be conditioned on the availability of a practicable analytical method to determine the quantity of residue. Such method must be sensitive to and reliable at the established tolerance level or, in certain instances, may be sensitive at a higher level where such level is also deemed satisfactory and safe in light of the toxicity of the drug residue and of the unlikelihood of such residue's exceeding the tolerance.

Subpart B—Specific Tolerances for Residues of New Animal Drugs

§ 556.20 2-Acetylamino-5-nitrothiazole.

A tolerance of 0.1 part per million is established for negligible residues of 2-acetylamino-5-nitrothiazole in the edible tissues of turkeys.

§ 556.30 Aklomide.

Tolerances are established for combined residues of aklomide (2-chloro-4-nitrobenzamide) and its metabolite (4-amino-2-chlorobenzamide) in uncooked edible tissues of chickens as follows:

- (a) 4.5 parts per million in liver and muscle.
- (b) 3 parts per million in skin with fat.

§ 556.34 Albendazole.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of albendazole is 5 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle*. A tolerance is established for albendazole 2-aminosulfone (marker residue) in liver (target tissue) of 0.2 part per million and in muscle of 0.05 part per million.

(2) *Sheep*. A tolerance is established for albendazole 2-aminosulfone (marker residue) in liver (target tissue) of 0.25 part per million and in muscle of 0.05 part per million.

[64 FR 1504, Jan. 11, 1999]

§ 556.38 Amoxicillin.

A tolerance of 0.01 part per million is established for negligible residues of

amoxicillin in milk and in the uncooked edible tissues of cattle.

[49 FR 45422, Nov. 16, 1984]

§ 556.40 Ampicillin.

A tolerance of 0.01 p/m is established for negligible residues of ampicillin in the uncooked edible tissues of swine and cattle and in milk.

§ 556.50 Amprolium.

Tolerances are established as follows for residues of amprolium (1-(4-amino-2-*n*-propyl-5-pyrimidinylmethyl)-2-picolinium chloride hydrochloride):

(a) In the edible tissues and in eggs of chickens and turkeys:

(1) 1 part per million in uncooked liver and kidney.

(2) 0.5 part per million in uncooked muscle tissue.

(3) In eggs:

(i) 8 parts per million in egg yolks.

(ii) 4 parts per million in whole eggs.

(b) In the edible tissues of calves:

(1) 2.0 parts per million in uncooked fat.

(2) 0.5 part per million in uncooked muscle tissue, liver, and kidney.

(c) In the edible tissues of pheasants:

(1) 1 part per million in uncooked liver.

(2) 0.5 part per million in uncooked muscle.

[40 FR 13942, Mar. 27, 1975, as amended at 50 FR 18472, May 1, 1985]

§ 556.52 Apramycin.

A tolerance of 0.1 part per million is established for parent apramycin (marker residue) in kidney (target tissue) of swine. The acceptable daily intake (ADI) for total residues of apramycin is 25 micrograms per kilogram of body weight per day.

[62 FR 40933, July 31, 1997]

§ 556.60 Arsenic.

Tolerances for total residues of combined arsenic (calculated as As) in food are established as follows:

(a) In edible tissues and in eggs of chickens and turkeys:

(1) 0.5 part per million in uncooked muscle tissue.

(2) 2 parts per million in uncooked edible by-products.

(3) 0.5 part per million in eggs.

§ 556.70

(b) In edible tissues of swine:
(1) 2 parts per million in uncooked liver and kidney.

(2) 0.5 part per million in uncooked muscle tissue and by-products other than liver and kidney.

§ 556.70 Bacitracin.

Tolerances for residues of bacitracin from zinc bacitracin or bacitracin methylene disalicylate are established at 0.5 part per million (0.02 unit per gram), negligible residue, in uncooked edible tissues of cattle, swine, chickens, turkeys, pheasants, and quail, and in milk and eggs.

[42 FR 18614, Apr. 8, 1977]

§ 556.90 Buquinolate.

Tolerances are established for residues of buquinolate as follows:

(a) In edible tissues of chickens:

(1) 0.4 part per million in uncooked liver, kidney, and skin with fat.

(2) 0.1 part per million in uncooked muscle.

(b) In eggs:

(1) 0.5 part per million in uncooked yolk.

(2) 0.2 part per million in uncooked whole eggs.

§ 556.100 Carbadox.

A tolerance of 30 parts per billion is established for residues of quinoxaline-2-carboxylic acid (marker residue) in liver (target tissue) of swine.

[63 FR 13337, Mar. 19, 1998]

§ 556.110 Carbomycin.

A tolerance of zero is established for residues of carbomycin in the uncooked edible tissues of chickens.

§ 556.113 Ceftiofur.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of ceftiofur is 30 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Swine, poultry, and sheep*. A tolerance for residues of ceftiofur in edible tissue is not required.

(2) *Cattle*. Tolerances are established for residues of desfuroylceftiofur (marker residue) in edible cattle tissues at 8 parts per million in kidney (target tissue), 2 parts per million in

21 CFR Ch. I (4–1–99 Edition)

the liver, 1 part per million in muscle, and 100 parts per billion in milk.

[63 FR 53579, Oct. 6, 1998]

§ 556.115 Cephapirin.

A tolerance of 0.02 parts per million (ppm) is established for residues of cephapirin in the milk and 0.1 ppm in the uncooked edible tissues of dairy cattle.

[40 FR 57454, Dec. 10, 1975]

§ 556.120 Chlorhexidine.

A tolerance of zero is established for residues of chlorhexidine in the uncooked edible tissues of calves.

§ 556.140 Chlorobutanol.

A tolerance of zero is established for residues of chlorobutanol in milk from dairy animals.

§ 556.150 Chlortetracycline.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of tetracyclines including chlortetracycline, oxytetracycline, and tetracycline is 25 micrograms per kilogram of body weight per day.

(b) *Tolerances*. (1) Tolerances are established for the sum of tetracycline residues in tissues of beef cattle, non-lactating dairy cows, calves, swine, sheep, chickens, turkeys, and ducks, of 2 parts per million (ppm) in muscle, 6 ppm in liver, and 12 ppm in fat and kidney.

(2) A tolerance is established for residues of chlortetracycline in eggs of 0.4 ppm.

[63 FR 52158, Sept. 30, 1998, as amended at 63 FR 57246, Oct. 27, 1998]

§ 556.160 Clopidol.

Tolerances for residues of clopidol (3,5-dichloro-2,6-dimethyl-4-pyridinol) in food are established as follows:

(a) In cereal grains, vegetables, and fruits: 0.2 part per million.

(b) In chickens and turkeys:

(1) 15 parts per million in uncooked liver and kidney.

(2) 5 parts per million in uncooked muscle.

(c) In cattle, sheep, and goats:

(1) 3 parts per million in uncooked kidney.

Food and Drug Administration, HHS

§ 556.220

(2) 1.5 parts per million in uncooked liver.

(3) 0.2 part per million in uncooked muscle.

(d) In swine: 0.2 part per million in uncooked edible tissues.

(e) In milk: 0.02 part per million (negligible residue).

§ 556.163 Clorsulon.

Tolerances are established for residues of clorsulon in cattle as follows:

(a) The tolerance for clorsulon (marker residue) in kidney (target tissue) is 1.0 part per million. A marker residue of 1.0 part per million corresponds to a total residue of 3.0 parts per million in kidney.

(b) The safe concentrations for total clorsulon residues in uncooked edible cattle tissues are: muscle, 1.0 part per million; liver, 2.0 parts per million; kidney, 3.0 parts per million; and fat, 4.0 parts per million.

[50 FR 10221, Mar. 14, 1985]

§ 556.165 Cloxacillin.

A tolerance of 0.01 part per million is established for negligible residues of cloxacillin in the uncooked edible tissues of cattle and in milk.

[40 FR 28792, July 9, 1975]

§ 556.167 Colistimethate.

A tolerance for residues of colistimethate in the edible tissues of chickens is not required.

[63 FR 13123, Mar. 18, 1998]

§ 556.170 Decoquinat.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of decoquinat is 75 micrograms per kilogram of body weight per day.

(b) *Tolerances*. Tolerances are established for residues of decoquinat in the uncooked, edible tissues of chickens, cattle, and goats as follows:

(1) 1 part per million (ppm) in skeletal muscle.

(2) 2 ppm in other tissues.

[64 FR 10103, Mar. 2, 1999]

§ 556.180 Dichlorvos.

A tolerance of 0.1 part per million is established for negligible residues of dichlorvos (2,2-dichlorovinyl dimethyl

phosphate) in the edible tissues of swine.

§ 556.200 Dihydrostreptomycin.

Tolerances are established for residues of dihydrostreptomycin in uncooked, edible tissues of cattle and swine of 2.0 parts per million (ppm) in kidney and 0.5 ppm in other tissues, and 0.125 ppm in milk.

[59 FR 41977, Aug. 16, 1994]

§ 556.220 3,5-Dinitrobenzamide.

No residues of 3,5-dinitrobenzamide may be found in the uncooked edible tissues of chickens as determined by the following method of analysis:

I. *Method of analysis*—3,5-dinitrobenzamide. A method for 3,5-dinitrobenzamide (3,5-DNBA) in chicken tissues is described with a cleanup step that removes most of the interfering materials, thus allowing uncompensated measurements to be read. The 3,5-DNBA is extracted from the sample with acetone and chloroform and prepared for chromatography by removing the aqueous phase in a separatory funnel and the solvents in a flash evaporator. The extract residue is chromatographed on alumina to remove several lipid components and residues of other drugs. The benzamide eluate is passed through a column of Dowex-50 resin, or equivalent, to remove arylamines; for example, 3-amino-5-nitrobenzamide. The 3,5-DNBA fraction is reduced, after removal of alcohol, with TiCl_3 in basic solution to an arylamine, presumably 3,5-diaminobenzamide. The reduced fraction is placed on another Dowex-50 column, most of the interfering substances are removed with washings of alcohol and water, and the arylamine residue is eluted with 4N HCl. Colorimetric measurement is made in a 100-millimeter cell at 530 millimicrons after reacting the residue with Bratton-Marshall reagents.

II. *Reagents*. A. Acetone.

B. Acetyl-(p-nitrophenyl)-sulfanilamide (APNPS) standard—melting point range 264° C.–267° C. Weigh and transfer 10 milligrams of APNPS to a 100-milliliter flask, dissolve and dilute to volume with acetone.

C. Alumina—activated F-20, 80–200 mesh, Aluminum Co. of America, or equivalent substance.

D. Ammonium sulfamate.

E. Ammonium sulfamate solution 1.25 grams of ammonium sulfamate per 100 milliliters of water. Refrigerate when not in use. Prepare fresh weekly.

F. Cation-exchange resin—Dowex 50W-X8, 200–400 mesh, Baker Analyzed Reagent, or equivalent, prepared as follows:

1. Place 500 grams of resin into a 3-liter beaker.

2. Add 2,000 milligrams of 6N HCl.

3. Heat and stir while on a bath at 80° C. for 6 hours. Discontinue heating and continue stirring overnight.

4. Filter the resin on a Buchner funnel (24 cm.) fitted with Whatman No. 1 paper.

5. Wash the resin bed with four 500-milliliter portions of 6N HCl.

6. Wash the resin bed with 500-milliliter portions of deionized water until the effluent has a pH of 5 or higher.

7. Wash the resin bed with three 400-milliliter portions of specially denatured alcohol 3A. Drain thoroughly.

8. Make a slurry of resin in 1,250 milliliters of specially denatured alcohol 3A.

G. Chloroform.

H. Coupling reagent—0.25 gram of *N*-1-naphthyl-ethylenediamine dihydrochloride per 100 milliliters of water. Refrigerate when not in use. Prepare fresh weekly.

I. 3,5-Dinitrobenzamide (3,5-DNBA standard). Add to boiling specially denatured alcohol 3A until a saturated solution is obtained and treat with activated carbon, filtered and crystallize by cooling to room temperature. The 3,5-DNBA therefrom is treated a second time with activated carbon and then recrystallized three more times from specially denatured alcohol 3A. The third crystallization is washed with diethyl ether and dried in a vacuum desiccator, melting point range 185° C.–186° C.

J. Ethyl alcohol—absolute, A.C.S.

K. Eluting reagent A. The formula and volume required in procedure step V-D is dependent on the adsorptive strength of the Al₂O₃. For each lot Al₂O₃, make the following test:

1. Prepare a column (see procedure step V-D for determining formula and volume to eluting reagent A).

2. Transfer 1 milliliter of APNPS standard (100 micrograms per milliliter) in 75 milliliters of chloroform to the column.

3. Wash the column with 100 milliliters of chloroform and discard the eluate.

4. Pass through 100 milliliters of solution consisting of specially denatured alcohol 3A and ethyl alcohol 1:1 (volume to volume). Collect one 50-milliliter and five 10-milliliter portions; these make up the first, second, third, fourth, fifth, and sixth portions of eluate.

5. Place in beakers under a stream of air on a water bath (90° C.) until the solvents are evaporated.

6. Add 10 milliliters of 4N HCl to each, cover with watch glasses and heat (90° C.) for 30 minutes; cool to room temperature.

7. Add the Bratton-Marshall reagents.

8. All fractions show a slight color. Note the portion containing the first significant increase in pink color.

a. If the color increases in the second, third, or fourth portions of eluate, the formula in procedure step V-D is suitable and, depending on the portion, 45, 55, or 65 milliliters, respectively, should be used in procedure step V-D4. Thereby, the APNPS is retained on the column and the benzamides are eluted.

b. If the color increases in the first portion, the eluting strength of the reagent is too strong. Return the test, substituting 1:4 (volume to volume) in procedure step V-D4. If 1:4 (volume to volume) is too strong, rerun with ethyl alcohol in procedure step V-D. If none of these are suitable, another lot of Al₂O₃ should be used.

c. If the color increases in the fifth or sixth portion, the eluting strength of the reagent is too weak. Rerun the test, substituting in procedure step V-D4, respectively, 4:1 (volume to volume), specially denatured alcohol 3A: methyl alcohol, 4:1 (volume to volume), until a suitable formula is found. If none of these are suitable, another lot of Al₂O₃ should be used.

L. Hydrochloric acid, 4N. Add two volumes of water to one volume of HCl.

M. Diatomaceous earth—Hyflo Super Cel, Johns-Manville Co., or equivalent substance.

N. *N*-1-Naphthylethylenediamine dihydrochloride.

O. Sodium hydroxide solution, 10N. Dissolve 100 grams of sodium hydroxide in water and dilute to 25 milliliters.

P. Sodium nitrite solution—0.25 grams of sodium nitrite per 100 milliliters of water. Refrigerate when not in use. Prepare fresh weekly.

Q. Specially denatured alcohol, formula 3A—100 parts of 190-proof ethyl alcohol plus 5 parts of commercial methyl alcohol.

R. Titanium(ous) chloride—20 percent solution.

III. *Special apparatus.* A. Absorption cells—Beckman No. 75195 matched set of two cylindrical silica cells with 100 millimeter optical length, or equivalent cells.

B. Autotransformer—type 500B, or equivalent. To regulate speed of mixer.

C. Centrifuge.

D. Centrifuge tubes—50-milliliter size with glass stopper.

E. Chromatography tubes—Corning No. 38460, 20 millimeters A 400 millimeters and having a tapered 29/42 joint with coarse, fritted disc, or equivalent tubes.

F. Evaporator—vacuum, rotary, thin film.

G. Ion-exchange column—as described by Thiels et al. in “Determination of 3-amino-5-nitro-*o*-toluamide (ANOT) in chicken tissues” published in “Journal of Agricultural and Food Chemistry,” volume 9, pages 201–204 (1961).

H. Glycerol manostat. For regulating pressure on columns: To Al₂O₃ columns, 15-inch head pressure; to ion-exchange columns, 30-inch head pressure.

I. Motor speed control. For regulating speed on 1-quart blender.

J. Volumetric flasks—50 milliliter size, ac-tinic ware.

K. Mixer—Vortex Jr. Model K-500-1, Sci-entific Industries, Inc., or equivalent mixer.

L. One-quart blender.

M. Water bath (45° C.–50° C.).

N. Water bath (90° C.).

IV. *Standard curve.* A. 1. Weigh 100 milli-grams of 3,5-DNBA and transfer to a 1-liter volumetric flask with acetone.

2. Dissolve and dilute with acetone to vol-ume.

3. Dilute 1 milliliter to 100 milliliters.

4. Add 5.0 milliliters of water to each of six centrifuge tubes.

5. Add standard to each of the tubes to con-tain one of the following amounts: 0.0, 1.0, 2.0, 3.0, 5.0, and 10.0 micrograms of 3,5-DNBA.

B. Prepare each tube for colorimetric measurement as follows:

1. Place the tube in a hot water bath (90° C.) until 5.0 milliliters remain. Cool to room temperature.

2. While mixing on Vortex mixer, or equiv-alent, regulated with an autotransformer, add 2 drops of TiCl_3 and 4 drops of 10N NaOH. Continue mixing until chalky-white in ap-pearance.

3. Add 2 milliliters of HCl, mix, and allow to stand for 5 minutes.

4. Transfer to 50-milliliter volumetric flask and dilute with 4N HCl to 40–45 milliliters.

5. Cool to 0° C.–5° C. by placing in a freezer or ice bath.

6. Perform the Bratton-Marshall reaction in subdued light as follows:

a. Add 1 milliliter of sodium nitrite re-agent, mix, and allow to stand for 1 minute.

b. Add 1 milliliter of ammonium sufamate reagent, mix, and allow to stand for 1 minute.

c. Add 1 milliliter of coupling reagent, mix, and allow to stand for 10 minutes.

d. Dilute to volume with 4N HCl.

C. Perform colorimetric measurement at 530 millimicrons as follows:

1. Fill two matched 100-millimeter cells with 4N HCl and place into spectrophotom-eter.

2. Adjust dark current.

3. Adjust to zero absorbance.

4. Replace acid in cell of sample side of compartment with standard to be measured.

5. The standard curve should be run five different times. Plot equivalent concentra-tion in tissue versus mean absorbance at each concentration. If computer is available, a better procedure is to calculate the equa-tion of the standard curve by means of least squares.

V. *Procedure.* A. Extraction. 1. Mince 350 grams of tissue in a 1-quart blending jar for 3 minutes. Use samples obtained from either freshly killed or quickly frozen birds. The latter should be analyzed as soon as thawed.

For fibrous meats (for example, muscle, skin) put through a meat grinder before mincing.

2. Weight 100 ± 0.5 grams of each replicate sample in a 150-milliliter beaker. Analyze each sample in triplicate and average the re-sults. Reproducibility of ± 10 percent between such analyses has been obtained.

3. Transfer the sample to a 1-quart blender jar. For kidney and liver tissues, make a slurry with acetone in the weighing beaker. Transfer with several rinses of acetone.

4. Blend the sample for 5 minutes with 250 milliliters of acetone and a 100-milliliter beakerful of diatomaceous earth.

5. Filter through a Buchner funnel con-taining a wetted Whatman No. 5 filter paper (12.5 cm.) into a 1-liter suction flask.

6. Rinse the blender jar into the funnel with three 25-milliliter portions of acetone.

7. Transfer the pulp and paper from the funnel to the aforementioned blender jar.

8. Add 250 milliliters of chloroform.

9. Blend for 3 minutes.

10. Filter through the aforementioned ap-paratus of procedure step V–A5. For rapid fil-tration of skin and blood samples, prepare funnel by adding diatomaceous earth and tamping evenly over paper to a thickness of 3 to 5 millimeters.

11. Rinse the blender jar into the funnel with three 25-milliliter rinses of chloroform.

B. Phasic separation. 1. Pour the combined filtrates into a 1-liter separatory funnel.

2. Rinse the suction flask twice with 25 milliliters of chloroform.

3. Mix the funnel contents by gently rock-ing and swirling for 30 seconds.

4. Let stand 10 minutes to allow phases to separate.

a. The upper (aqueous) phase (30 to 50 mil-liliters) is not always emulsion-free. Losses from emulsions have not been significant.

b. If an upper (aqueous) phase does not ap-pear, add an additional 100 milliliters of chloroform and 10 milliliters of water and re-peat procedure step V–B3.

5. Withdraw the lower phase into a 1-liter round-bottom flask, and discard upper phase. Withdraw nearly all of the lower phase, let stand for 2 to 3 minutes, then withdraw the remainder.

C. Evaporation. Attach the flask on a thin-film rotary evaporator connected to a vacu-um supply, and place in a water bath main-tained at 45° C.–50° C. until an oily residue remains. Do not overheat the sample or allow to go to dryness.

D. Adsorption chromatography. 1. Prepare a chromatography column using a column with calibrated etchings to indicate appro-priate adsorbent and solvent levels as fol-lows:

a. Fill tube to a depth of 60 millimeters with Al_2O_3 .

b. Tap walls gently with hands.

§ 556.225**21 CFR Ch. I (4–1–99 Edition)**

c. Add anhydrous sodium sulfate to an additional depth of 25 millimeters.

d. Wet and wash column with 50 milliliters of chloroform.

i. During chromatography, make each addition to the tube when the liquid level has reached the top of the sodium sulfate layer.

ii. Increase the percolation rates by applying a slight air pressure to the top of the column.

2. Transfer the residue from procedure step V–C to the column with four 15-milliliter rinses of chloroform. Then rinse the walls of the tube and sodium sulfate layer with three 5-milliliter portions of chloroform. Percolation rate: 15 to 25 milliliters per minute. No color from sample should be seen in sodium sulfate layer after final rinse.

3. Wash column with 100 milliliters of chloroform. Discard eluate.

4. Add 75 milliliters of eluting reagent A and collect eluate A in a 250-milliliter beaker for cation-exchange chromatography.

a. Refer to “Eluting reagent A” under “Reagents” (II–K) for determining formula and volume.

b. Percolation rate: 8 to 12 milliliters per minute.

E. Cation-exchange chromatography—No. 1. 1. Prepare an ion-exchange column as follows:

a. Add a uniform slurry of resin to the column to obtain a 4 to 5 centimeter bed depth after settling.

i. Obtain a uniform slurry using a magnetic stirrer. To add the required amount of resin, calibrate the slurry and transfer it with a 10-milliliter pipette to deliver a reproducible volume.

ii. Increase the flow rate to 2 to 4 milliliters per minute by applying air pressure to the column. A glycerol manostat adjusted to 30 inches and attached between an air supply and column provides adequate pressure.

b. Wash the resin with 10 milliliters of eluting reagent A. Discard eluate.

2. Pass eluate A from procedure step V–D4 through the column. Collect in a 250-milliliter beaker.

3. Pass 50 milliliters of specially denatured alcohol 3A through the column. Combine with the eluate of procedure step V–E2.

F. Reduction. 1. Place the eluate A fraction from procedure step V–E3 on a hot water bath (90° C.) and evaporate with a stream of air until 5 to 10 milliliters remain. Do not overheat the sample or allow the sample to go to dryness.

2. Transfer to centrifuge tube and rinse beaker three times with 3 milliliters of specially denatured alcohol 3A.

3. Evaporate on a hot water bath (90° C.) under a stream of air until alcohol has evaporated. Do not overheat the sample or allow the sample to go to dryness.

4. Remove the tube from the water bath and immediately add 5.0 milliliters of water.

5. While mixing, add 2 drops of titanium chloride and 4 drops of 10N sodium hydroxide. Continue mixing until greyish color disappears.

a. Mix on Vortex Jr. mixer, or equivalent, regulated with autotransformer.

b. Precipitate of insoluble tissue substances and white titanium salts is present after reduction is complete.

6. Dilute to 50 milliliters with specially denatured alcohol 3A and mix.

7. Centrifuge for 5 minutes at 2,000 r.p.m.

G. Cation-exchange chromatography—No. 2. 1. Prepare resin column by procedure step V–E.

2. Pass the centrifugate of procedure step V–F7 through column. Use three rinses of specially denatured alcohol 3A, each 5 milliliters, to aid in transferring of sample.

3. Pass 50 milliliters of specially denatured alcohol 3A through the column.

4. Pass 50 milliliters of deionized water through the column.

5. Elute arylamine residue from the resin with 40 to 43 milliliters of 4N HCl into a 50-milliliter volumetric flask (actinic ware) for 3,5-DNBA analysis. Avoid direct sunlight. The arylamine has been found to be photosensitive.

H. Color development and measurement. 1. Cool to 0° C.–5° C. by placing in a freezer or ice bath.

2. Perform the Bratton-Marshall reaction in subdued light as follows:

a. Add 1 milliliter of sodium nitrite reagent, mix, and allow to stand for 1 minute.

b. Add 1 milliliter of ammonium sulfamate reagent, mix, and allow to stand for 1 minute.

c. Add 1 milliliter of coupling reagent, mix, and allow to stand for 10 minutes.

d. Dilute to volume with 4N HCl.

3. Perform colorimetric measurement at 530 millimicrons as follows:

a. Fill two matched 100-millimeter cells with 4N HCl and place into instrument.

b. Adjust dark current.

c. Adjust to zero absorbance.

d. Replace acid in cell of sample side of compartment with sample to be measured.

e. Record absorbance observed.

I. Calculations. Determine parts per billion (observed) from the standard curve.

§ 556.225 Doramectin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of doramectin is 0.75 microgram per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle*. A tolerance of 100 parts per billion is established for parent doramectin (marker residue) in liver (target tissue) and of 30 parts per billion for parent doramectin in muscle.

Food and Drug Administration, HHS

§ 556.275

(2) *Swine*. A tolerance is established for parent doramectin (marker residue) in liver (target tissue) of 160 parts per billion.

[63 FR 68184, Dec. 10, 1998]

§ 556.227 Eprinomectin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of eprinomectin is 10 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle*. Tolerances are established for residues of eprinomectin B1a (marker residue) in milk of 12 parts per billion, in liver (target tissue) of 4.8 parts per million, and in muscle of 100 parts per billion.

(2) [Reserved]

[63 FR 59715, Nov. 5, 1998]

§ 556.228 Enrofloxacin.

The acceptable daily intake for enrofloxacin is 3 micrograms per kilogram of body weight per day.

(a) *Chickens and turkeys*. A tolerance of 0.3 part per million is established for residues of enrofloxacin (marker residue) in muscle (target tissue) of chickens and turkeys.

(b) *Cattle*. A tolerance of 0.1 part per million for desethyle ciprofloxacin (marker residue) has been established in liver (target tissue) of cattle.

[61 FR 56893, Nov. 5, 1996, as amended at 63 FR 49003, Sept. 14, 1998]

§ 556.230 Erythromycin.

Tolerances for residues of erythromycin in food are established as follows:

(a) 0.1 part per million in uncooked edible tissues of beef cattle and swine.

(b) Zero in milk.

(c) 0.025 part per million in uncooked eggs.

(d) 0.125 part per million (negligible residue) in uncooked edible tissues of chickens and turkeys.

[40 FR 13942, Mar. 27, 1975, as amended at 58 FR 43795, Aug. 18, 1993]

§ 556.240 Estradiol and related esters.

No residues of estradiol, resulting from the use of estradiol or any of the related esters, are permitted in excess of the following increments above the

concentrations of estradiol naturally present in untreated animals:

(a) In uncooked edible tissues of heifers, steers, and calves:

(1) 120 parts per trillion for muscle.

(2) 480 parts per trillion for fat.

(3) 360 parts per trillion for kidney.

(4) 240 parts per trillion for liver.

(b) In uncooked edible tissues of lambs:

(1) 120 parts per trillion for muscle.

(2) 600 parts per trillion for fat, kidney, and liver.

[49 FR 13873, Apr. 9, 1984, as amended at 56 FR 67175, Dec. 30, 1991]

§ 556.260 Ethopabate.

Tolerance for residues of ethopabate converted to metaphenetidine are established in the edible tissues of chickens as follows:

(a) 1.5 parts per million in uncooked liver and kidney.

(b) 0.5 part per million in uncooked muscle.

§ 556.270 Ethylenediamine.

A tolerance of zero is established for residues of ethylenediamine in milk.

§ 556.273 Famphur.

Tolerances are established for residues of famphur including its oxygen analog in or on meat, fat, or meat by-products of cattle at 0.1 part per million.

[62 FR 55161, Oct. 23, 1997]

§ 556.275 Fenbendazole.

(a) *Cattle and goats*. A tolerance¹ of 0.8 part per million is established for parent fenbendazole (the marker residue) in the liver of cattle and goats.

(b) *Swine*. A tolerance¹ for marker residues of fenbendazole in swine is not needed.

(c) *Cattle milk*. A safe concentration of 1.67 parts per million is established

¹As used in this section: “tolerance” refers to a concentration of a marker residue in the target tissue selected to monitor for total residues of the drug in the target animal.

§ 556.277

for total fenbendazole residues. A tolerance of 0.6 part per million is established based on the fenbendazole sulf-oxide metabolite (marker residue).

[59 FR 26943, May 25, 1994, as amended at 61 FR 29478, June 11, 1996]

§ 556.277 Fenprostalene.

A tolerance for marker residue of fenprostalene in cattle is not needed. The safe concentrations for the total residues of fenprostalene in the uncooked edible tissues of cattle are 10 parts per billion in muscle, 20 parts per billion in liver, 30 parts per billion in kidney, 40 parts per billion in fat, and 100 parts per billion in the injection site. As used in this section “tolerance” refers to a concentration of a marker residue in the target tissue selected to monitor for total residues of the drug in the target animal, and “safe concentrations” refer to the concentrations of total residues considered safe in edible tissues.

[49 FR 26716, June 29, 1984]

§ 556.283 Florfenicol.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of florfenicol is 10 micrograms per kilogram of body weight per day.

(b) *Cattle*. A tolerance of 3.7 parts per million (ppm) for florfenicol amine (marker residue) in liver (target tissue) is established. A tolerance of 0.3 ppm for florfenicol amine in cattle muscle is established.

[63 FR 41191, Aug. 3, 1998]

§ 556.286 Flunixin meglumine.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of flunixin is 0.72 micrograms per kilogram of body weight per day.

(b) *Tolerances*. For residues of parent flunixin free acid of 0.125 part per million (ppm) in cattle liver (target tissue) and 0.025 ppm in cattle muscle are established.

[63 FR 38750, July 20, 1998]

§ 556.290 Furazolidone.

A tolerance of zero is established for residues of furazolidone in the uncooked edible tissues of swine.

21 CFR Ch. I (4–1–99 Edition)

§ 556.300 Gentamicin sulfate.

(a) A tolerance of 0.1 part per million is established for negligible residues of gentamicin sulfate in the uncooked edible tissues of chickens and turkeys.

(b) Tolerances are established for total residues of gentamicin in edible tissues of swine as follows: 0.1 part per million in muscle, 0.3 part per million in liver, and 0.4 part per million in fat and kidney. A microbiological determinative procedure and an HPLC confirmatory procedure for gentamicin have been developed to assay gentamicin in kidney at 0.4 ppm. Since residues of gentamicin as the parent compound and total residues are equal, the marker (parent drug) residue concentration of 0.4 ppm in kidney corresponds to 0.4 ppm of total residue.

[48 FR 791, Jan. 7, 1983, as amended at 61 FR 24441, May 15, 1996]

§ 556.308 Halofuginone hydrobromide.

The marker residue selected to monitor for total residues of halofuginone hydrobromide in broilers and turkeys is parent halofuginone hydrobromide and the target tissue selected is liver. A tolerance is established in broilers of 0.16 part per million and in turkeys of 0.13 part per million for parent halofuginone hydrobromide in liver. These marker residue concentrations in liver correspond to total residue concentrations of 0.3 part per million in liver. The safe concentrations for total residues of halofuginone hydrobromide in the uncooked edible tissues of broilers and turkeys are 0.1 part per million in muscle, 0.3 part per million in liver, and 0.2 part per million in skin with adhering fat. As used in this section, “tolerance” refers to a concentration of a marker residue in the target tissue selected to monitor for total residues of the drug in the target animal, and “safe concentrations” refers to the concentrations of total residues considered safe in edible tissues.

[54 FR 28052, July 5, 1989, as amended at 56 FR 8711, Mar. 1, 1991; 57 FR 21209, May 19, 1992]

§ 556.310 Haloxon.

A tolerance of 0.1 part per million is established for negligible residues of

Food and Drug Administration, HHS

§ 556.390

haloxon (3-chloro-7-hydroxy-4-methylcoumarin bis(2-chloroethyl) phosphate) in the edible tissues of cattle.

[40 FR 13942, Mar. 27, 1975, as amended at 45 FR 10333, Feb. 15, 1980]

§ 556.320 Hydrocortisone.

A tolerance is established for negligible residues of hydrocortisone (as hydrocortisone sodium succinate or hydrocortisone acetate) in milk at 10 parts per billion.

§ 556.330 Hygromycin B.

A tolerance of zero is established for residues of hygromycin B in or on eggs and the uncooked edible tissues of swine and poultry.

§ 556.344 Ivermectin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of ivermectin is 1 microgram per kilogram of body weight per day.

(b) *Tolerances*—(1) *Liver*. A tolerance is established for 22,23-dihydroavermectin B_{1a} (marker residue) in liver (target tissue) as follows:

(i) *Cattle*. 100 parts per billion.

(ii) *Swine*. 20 parts per billion.

(iii) *Sheep*. 30 parts per billion.

(iv) *Reindeer*. 15 parts per billion.

(v) *American bison*. 15 parts per billion.

(2) *Muscle*. Muscle residues are not indicative of the safety of other edible tissues. A tolerance is established for 22,23-dihydroavermectin B_{1a} (marker residue) in muscle as follows:

(i) *Swine*. 20 parts per billion.

(ii) [Reserved]

[63 FR 54352, Oct. 9, 1998]

§ 556.347 Lasalocid.

(a) [Reserved]

(b) *Tolerances*—(1) *Chickens*. A tolerance is established for lasalocid residues of 0.3 part per million (ppm) parent lasalocid (marker residue) in skin with adhering fat (target tissue).

(2) *Cattle*. A tolerance is established for lasalocid residues of 0.7 ppm parent lasalocid (marker residue) in liver (target tissue).

(3) *Sheep*. A tolerance for residues of lasalocid is not needed.

(4) *Rabbits*. A tolerance is established for lasalocid residues of 0.7 ppm parent

lasalocid (marker residue) in liver (target tissue).

[64 FR 13069, Mar. 17, 1999]

§ 556.350 Levamisole hydrochloride.

A tolerance of 0.1 part per million is established for negligible residues of levamisole hydrochloride in the edible tissues of cattle, sheep, and swine.

§ 556.360 Lincomycin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of lincomycin is 25 micrograms per kilogram of body weight per day.

(b) *Chickens*. A tolerance for residues of lincomycin in chickens is not required.

(c) *Swine*. Tolerances for lincomycin of 0.6 part per million in liver and 0.1 part per million in muscle are established.

[64 FR 13342, Mar. 18, 1999]

§ 556.375 Maduramicin ammonium.

A tolerance is established for residues of maduramicin ammonium in chickens as follows:

(a) A tolerance for maduramicin ammonium (marker residue) in chickens is 0.38 parts per million in fat (target tissue). A tolerance refers to the concentration of marker residues in the target tissue used to monitor for total drug residues in the target animals.

(b) The safe concentrations for total maduramicin ammonium residues in uncooked edible chicken tissues are: 0.24 parts per million in muscle; 0.72 parts per million in liver; 0.48 parts per million in skin; and 0.48 parts per million in fat. A safe concentration refers to the total residue concentration considered safe in edible tissues.

[54 FR 5229, Feb. 2, 1989]

§ 556.380 Melengestrol acetate.

A tolerance of 25 parts per billion is established for residues of the parent compound, melengestrol acetate, in fat of cattle.

[59 FR 41241, Aug. 11, 1994]

§ 556.390 Methylparaben.

A tolerance of zero is established for residues of methylparaben in milk from dairy animals.

§ 556.400

§ 556.400 Methylprednisolone.

A tolerance is established for negligible residues of methylprednisolone in milk at 10 parts per billion.

§ 556.410 Metoserpate hydrochloride.

A tolerance of 0.02 part per million is established for negligible residues of metoserpate hydrochloride (methyl-*o*-methyl-18-epireserpate hydrochloride) in uncooked edible tissues of chickens.

§ 556.420 Monensin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of monensin is 12.5 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle and goats*. A tolerance of 0.05 part per million is established for negligible residues of monensin in edible tissues of cattle and goats.

(2) *Chickens, turkeys, and quail*. A tolerance for residues of monensin in chickens, turkeys, and quail is not needed.

[64 FR 5159, Feb. 3, 1999]

§ 556.425 Morantel tartrate.

A tolerance of 0.7 part per million is established for *N*-methyl-1,3-propanediamine (MAPA, marker residue) in the liver (target tissue) of cattle and goats. A tolerance for residues of morantel tartrate in milk is not required.

[59 FR 17922, Apr. 15, 1994]

§ 556.426 Moxidectin.

An acceptable daily intake (ADI) of 4 micrograms per kilogram per day in tissue is established. A tolerance is established for parent moxidectin in edible tissues of cattle of 50 parts per billion in muscle and 200 parts per billion in liver.

[63 FR 14036, Mar. 24, 1998]

§ 556.428 Narasin.

A tolerance for narasin residues in chickens is not needed. The safe concentrations for total narasin residues in uncooked edible chicken tissues are: 0.6 part per million in muscle; 1.8 parts per million in liver; 1.2 parts per million in skin with adhering fat and fat. A tolerance refers to the concentration

21 CFR Ch. I (4–1–99 Edition)

of marker residues in the target tissue used to monitor for total drug residues in the target animals. A safe concentration refers to the total residue concentration considered safe in edible tissues.

[51 FR 29097, Aug. 14, 1986]

§ 556.430 Neomycin.

A tolerance of 7.2 parts per million (ppm) is established for residues of parent neomycin (marker residue) in uncooked edible kidney (target tissue), 7.2 ppm in fat, 3.6 ppm in liver, 1.2 ppm in muscle of cattle, swine, sheep, and goats. A tolerance of 0.15 ppm is established for neomycin in milk.

[61 FR 31399, June 20, 1996]

§ 556.440 Nequinat.

A tolerance of 0.1 part per million is established for negligible residues of nequinat in the uncooked edible tissues of chickens.

§ 556.445 Nicarbazin.

A tolerance of 4 parts per million is established for residues of nicarbazin in uncooked chicken muscle, liver, skin, and kidney.

[42 FR 56729, Oct. 28, 1977]

§ 556.460 Novobiocin.

Tolerances for residues of novobiocin are established at 0.1 part per million in milk from dairy animals and 1 part per million in the uncooked edible tissues of cattle, chickens, turkeys, and ducks.

[47 FR 18590, Apr. 30, 1982]

§ 556.470 Nystatin.

A tolerance of zero is established for residues of nystatin in or on eggs and the uncooked edible tissues of swine and poultry.

§ 556.480 Oleandomycin.

Tolerances are established for negligible residues of oleandomycin in uncooked edible tissues of chickens, turkeys, and swine at 0.15 part per million.

§ 556.490 Ormetoprim.

A tolerance of 0.1 part per million is established for negligible residues of

Food and Drug Administration, HHS

§ 556.560

ormetoprim in the edible tissues of chickens, turkeys, ducks, salmonids, and catfish.

[51 FR 18884, May 23, 1986]

§ 556.495 Oxfendazole.

Cattle: A tolerance is established for total oxfendazole residues in edible cattle tissues based on a marker residue concentration of 0.8 part per million (ppm) fenbendazole in the target liver tissue. A fenbendazole concentration of 0.8 ppm in liver corresponds to a total safe concentration of oxfendazole residues of 1.7 ppm in liver. The safe concentrations of total oxfendazole residues in other uncooked edible cattle tissues are: muscle, 0.84 ppm; kidney, 2.5 ppm; and fat, 3.3 ppm. A tolerance refers to the concentration of marker residue in the target tissue selected to monitor for total drug residue in the target animal. A safe concentration is the total residue considered safe in edible tissue.

[55 FR 46943, Nov. 8, 1990]

§ 556.500 Oxytetracycline.

(a) *Acceptable daily intake (ADI).* The ADI for total tetracycline residues (chlortetracycline, oxytetracycline, and tetracycline) is 25 micrograms per kilogram of body weight per day.

(b) *Tolerances.* Tolerances are established for the sum of tetracycline residues in tissues of beef cattle, beef calves, nonlactating dairy cattle, dairy calves, swine, sheep, chickens, turkeys, catfish, lobsters, and salmonids, of 2 parts per million (ppm) in muscle, 6 ppm in liver, and 12 ppm in fat and kidney.

[63 FR 57246, Oct. 27, 1998]

§ 556.510 Penicillin.

Tolerances are established for residues of penicillin and the salts of penicillin in food as follows:

(a) 0.05 part per million (negligible residue) in the uncooked edible tissues of cattle.

(b) Zero in the uncooked edible tissues of chickens, pheasants, quail, swine, and sheep; in eggs; and in milk or in any processed food in which such milk has been used.

(c) 0.01 part per million in the uncooked edible tissues of turkeys.

[40 FR 13942, Mar. 27, 1975, as amended at 43 FR 32749, July 28, 1978]

§ 556.515 Pirlimycin.

A tolerance is established for residues of parent pirlimycin (marker substance) in cattle liver (target tissue) of 0.5 part per million and in milk of 0.4 part per million.

[58 FR 58486, Nov. 2, 1993]

§ 556.520 Prednisolone.

A tolerance of zero is established for residues of prednisolone in milk from dairy animals.

§ 556.530 Prednisone.

A tolerance of zero is established for residues of prednisone in milk from dairy animals.

§ 556.540 Progesterone.

No residues of progesterone are permitted in excess of the following increments above the concentrations of progesterone naturally present in untreated animals:

(a) In uncooked edible tissues of steers and calves:

- (1) 3 parts per billion for muscle.
- (2) 12 parts per billion for fat.
- (3) 9 parts per billion for kidney.
- (4) 6 parts per billion for liver.

(b) In uncooked edible tissues of lambs:

- (1) 3 parts per billion for muscle.
- (2) 15 parts per billion for fat, kidney, and liver.

[49 FR 13873, Apr. 9, 1984]

§ 556.550 Propylparaben.

A tolerance of zero is established for residues of propylparaben in milk from dairy animals.

§ 556.560 Pyrantel tartrate.

Tolerances are established for residues of pyrantel tartrate in edible tissues of swine as follows:

- (a) 10 parts per million in liver and kidney.
- (b) 1 part per million in muscle.

§ 556.580

§ 556.580 Robenidine hydrochloride.

Tolerances are established for residues of robenidine hydrochloride in edible tissues of chickens as follows:

(a) 0.2 part per million in skin and fat.

(b) 0.1 part per million (negligible residue) in edible tissues other than skin and fat.

§ 556.590 Salicylic acid.

A tolerance of zero is established for residues of salicylic acid in milk from dairy animals.

§ 556.594 Sarafloxacin.

A tolerance for residues of sarafloxacin in edible turkey and broiler chickens tissues is not required.

[60 FR 50098, Sept. 28, 1995]

§ 556.600 Spectinomycin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of spectinomycin is 25 micrograms per kilogram of body weight per day.

(b) *Chickens and turkeys*. A tolerance of 0.1 part per million (ppm) for negligible residues of spectinomycin in uncooked edible tissues of chickens and turkeys is established.

(c) *Cattle*. A tolerance of 4 ppm for parent spectinomycin (marker residue) in kidney (target tissue) is established. A tolerance of 0.25 ppm for parent spectinomycin in cattle muscle is established.

[63 FR 24107, May 1, 1998; 63 FR 38304, July 16, 1998]

§ 556.610 Streptomycin.

Tolerances are established for residues of streptomycin in uncooked, edible tissues of chickens, swine, and calves of 2.0 parts per million (ppm) in kidney and 0.5 ppm in other tissues.

[58 FR 47211, Sept. 8, 1993]

§ 556.620 Sulfabromomethazine sodium.

Tolerances for residues of sulfabromomethazine sodium in food are established as follows:

(a) In the uncooked edible tissues of cattle at 0.1 part per million (negligible residue).

21 CFR Ch. I (4–1–99 Edition)

(b) In milk at 0.01 part per million (negligible residue).

[47 FR 30244, July 13, 1982]

§ 556.625 Sodium sulfachloropyrazine monohydrate.

A tolerance of zero is established for residues of sodium sulfachloropyrazine monohydrate in the uncooked edible tissues of chickens.

§ 556.630 Sulfachlorpyridazine.

A tolerance of 0.1 part per million is established for negligible residues of sulfachlorpyridazine in uncooked edible tissues of calves and swine.

§ 556.640 Sulfadimethoxine.

Tolerances are established for residues of sulfadimethoxine in edible products of animals as follows:

(a) In the uncooked edible tissues of chickens, turkeys, cattle, ducks, salmonids, and catfish at 0.1 part per million (negligible residue).

(b) In milk at 0.01 part per million (negligible residue).

[40 FR 13942, Mar. 27, 1975, as amended at 49 FR 46371, Nov. 26, 1984; 51 FR 18884, May 23, 1986]

§ 556.650 Sulfaethoxypyridazine.

Tolerances for residues of sulfaethoxypyridazine in food are established as follows:

(a) Zero in the uncooked edible tissues of swine and in milk.

(b) 0.1 part per million (negligible residue) in uncooked edible tissues of cattle.

§ 556.660 Sulfamerazine.

A tolerance of zero is established for residues of sulfamerazine (N¹-[4-methyl-2-pyrimidinyl]sulfanilamide) in the uncooked edible tissues of trout.

§ 556.670 Sulfamethazine.

A tolerance of 0.1 part per million is established for negligible residues of sulfamethazine in the uncooked edible tissues of chickens, turkeys, cattle, and swine.

[47 FR 25323, June 11, 1982]

§ 556.680 Sulfanitran.

A tolerance of zero is established for residues of sulfanitran (acetyl(*p*-nitro-

Food and Drug Administration, HHS

§ 556.740

phenyl) sulfanilamide) and its metabolites in the uncooked edible tissues of chickens.

§ 556.685 Sulfaquinoxaline.

A tolerance of 0.1 part per million is established for negligible residues of sulfaquinoxaline in the uncooked edible tissues of chickens, turkeys, calves, and cattle.

[61 FR 24443, May 15, 1996]

§ 556.690 Sulfathiazole.

A tolerance of 0.1 part per million is established for negligible residues of sulfathiazole in the uncooked edible tissues of swine.

§ 556.700 Sulfomyxin.

A tolerance of zero is established for residues of sulfomyxin (N-sulfomethyl-polymyxin B sodium salt) in uncooked edible tissues from chickens and turkeys.

§ 556.710 Testosterone propionate.

No residues of testosterone, resulting from the use of testosterone propionate, are permitted in excess of the following increments above the concentrations of testosterone naturally present in untreated animals:

(a) In uncooked edible tissues of heifers:

- (1) 0.64 part per billion in muscle.
- (2) 2.6 parts per billion in fat.
- (3) 1.9 parts per billion in kidney.
- (4) 1.3 parts per billion in liver.

(b) [Reserved]

[52 FR 27683, July 23, 1987]

§ 556.720 Tetracycline.

(a) *Acceptable daily intake (ADI)*. The ADI for total tetracycline residues (chlortetracycline, oxytetracycline, and tetracycline) is 25 micrograms per kilogram of body weight per day.

(b) *Tolerances*. Tolerances are established for the sum of tetracycline residues in tissues of calves, swine, sheep, chickens, and turkeys, of 2 parts per million (ppm) in muscle, 6 ppm in liver, and 12 ppm in fat and kidney.

[63 FR 57246, Oct. 27, 1998]

§ 556.730 Thiabendazole.

Tolerances are established at 0.1 part per million for negligible residues of thiabendazole in uncooked edible tissues of cattle, goats, sheep, pheasants, and swine, and at 0.05 part per million for negligible residues in milk.

[40 FR 13942, Mar. 27, 1975, as amended at 49 FR 29958, July 25, 1984]

§ 556.735 Tilmicosin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of tilmicosin is 25 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle*. A tolerance is established for residues of parent tilmicosin (marker residue) in liver (target tissue) at 1.2 parts per million (ppm).

(2) *Swine*. A tolerance is established for residues of parent tilmicosin (marker residue) in liver (target tissue) at 7.5 ppm and in muscle at 0.1 ppm.

[64 FR 13679, Mar. 22, 1999]

§ 556.738 Tiamulin.

A tolerance of 0.6 part per million is established for 8-*alpha*-hydroxymutilin (marker compound) in liver (target tissue) of swine.

[62 FR 12086, Mar. 14, 1997]

§ 556.739 Trenbolone.

A tolerance for total trenbolone residues in uncooked edible tissues of cattle is not needed. The safe concentration for total trenbolone residues in uncooked edible tissues of cattle is 50 parts per billion (ppb) in muscle, 100 ppb in liver, 150 ppb in kidney, and 200 ppb in fat. A tolerance refers to the concentration of marker residues in the target tissue used to monitor for total drug residues in the target animals. A safe concentration refers to the total residue concentration considered safe in edible tissues.

[52 FR 24995, July 2, 1987, as amended at 54 FR 12595, Mar. 28, 1989]

§ 556.740 Tylosin.

Tolerances are established for residues of tylosin in edible products of animals as follows:

(a) In chickens and turkeys: 0.2 part per million (negligible residue) in

§ 556.741

uncooked fat, muscle, liver, and kidney.

(b) In cattle: 0.2 part per million (negligible residue) in uncooked fat, muscle, liver, and kidney.

(c) In swine: 0.2 part per million (negligible residue) in uncooked fat, muscle, liver, and kidney.

(d) In milk: 0.05 part per million (negligible residue).

(e) In eggs: 0.2 part per million (negligible residue).

§ 556.741 Tripeleennamine.

A tolerance of 200 parts per billion (ppb) is established for residues of tripeleennamine in uncooked edible tissues of cattle and 20 ppb in milk.

[62 FR 4164, Jan. 29, 1997]

§ 556.750 Virginiamycin.

Tolerances are established for negligible residues of virginiamycin in edible tissues of swine as follows:

(a) Swine—

(1) 0.4 ppm in kidney, skin, and fat.

(2) 0.3 ppm in liver.

(3) 0.1 ppm in muscle.

(b) Broiler chickens—

(1) 0.5 ppm in kidney.

(2) 0.3 ppm in liver.

(3) 0.2 ppm in skin and fat.

(4) 0.1 ppm in muscle.

(c) *Cattle*. A tolerance for residues of virginiamycin in cattle is not required.

[46 FR 18966, Mar. 27, 1981, as amended at 59 FR 38902, Aug. 1, 1994]

§ 556.760 Zeranol.

(a) *Cattle*. A tolerance for total zeranol residues in uncooked edible tissues of cattle is not needed. The safe concentration for total zeranol residues in uncooked edible tissues of cattle is 150 parts per billion (ppb) in muscle, 300 ppb in liver, 450 ppb in kidney, and 600 ppb in fat. A tolerance refers to the concentration of marker residues in the target tissue used to monitor for total drug residues in the target animal. A safe concentration refers to the total residue concentration considered safe in edible tissues.

(b) *Sheep*. No residues of zeranol may be found in the uncooked edible tissues of sheep as determined by the following method of analysis:

21 CFR Ch. I (4–1–99 Edition)

I. METHOD OF ANALYSIS—ZERANOL

A gas chromatographic method for the determination of the drug in frozen beef tissues is described. Tissue is frozen and stored in a deep freezer until ready for examination. A weighed portion of wet tissue (with exception of fat) is homogenized and lyophilized to dry solid. The drug is recovered from dry tissue by an extraction with methanol in a Soxhlet extractor. The methanol extract is digested in the presence of hydrochloric acid to hydrolyze conjugates should any be present. Elimination of impurities is brought about by liquid partition transfer successively to chloroform to 1N sodium hydroxide, to carbon tetrachloride, to 1N sodium hydroxide, to ethyl ether, and, finally, to a dry residue. The residue is reacted with a silane mixture to create a volatile derivative which is quantitated by peak area measurements from a flame ionization detector. The drug can be detected at a level of 20 parts per billion with negligible interference from tissues or reagents.

II. REAGENTS

A. Carbon tetrachloride, N.F., Fisher Scientific C-186, or equivalent.

B. Chloroform, N.F., Fisher Scientific C-296, or equivalent.

C. Chromatograph gases, flow rates adjusted to maximize sensitivity for specific chromatograph.

1. Carrier gas, conventional tank helium.

2. Flame makeup gas.

a. Oxygen, conventional tank oxygen.

b. Hydrogen, Linde high purity, or equivalent.

D. Column packing, 3 percent GE SE-52 (Applied Science Laboratories) on P.E. Celite 60–80 mesh (Johns Manville Product No. 154-0048), or equivalent.

E. Ether, anhydrous, Fisher Scientific E-138, or equivalent.

F. Hexamethyldisilazane, Dow-Corning, Peninsular, or equivalent.

G. Hydrochloric acid, analytical reagent grade.

H. Methanol, certified A.C.S., spectranalyzed, Fisher Scientific A-408, or equivalent.

I. Phosphoric acid, analytical reagent grade.

J. Pyridine, anhydrous, A.C.S. reagent grade.

K. Silating reagent mixture: Pipet 8 milliliters each of pyridine and hexamethyldisilazane and 4 milliliters of trimethylchlorosilane into a clean glass vial with a polyethylene cap and mix thoroughly. Let stand overnight and decant supernatant liquid into a vial. Cap and store at room temperature for daily use. If kept dry, the reagent is stable for more than a month. Blanks are scanned by gas chromatography

on each new bottle of J, F, and N material used in the silating reagent mixture for possible peak interference in the region of zeranol derivative.

L. Sodium chloride, analytical reagent grade.

M. Sodium hydroxide, analytical reagent grade.

N. Trimethylchlorosilane, Dow-Corning, Peninsular, or equivalent.

O. Water, distilled in glass.

P. Zeranol, primary standard.

Q. Solutions.

1. 2N Hydrochloric acid in water.

2. 3N Phosphoric acid in water.

3. 2 percent w/v sodium chloride in water.

4. 1N Sodium hydroxide in water.

III. APPARATUS

A. Extraction assemblies, Soxhlet, improved, standard taper grindings, Pyrex brand glass, 1,000 milliliters capacity, Sargent Catalog S-31265D, or equivalent.

B. Flasks, freeze drying, widemouth, 1,000 milliliters capacity, 24/40 standard taper grindings, Pyrex brand glass, Sargent Catalog S-28875-20-F, or equivalent.

C. Flasks, homogenizing, 250 milliliters, Sargent Catalog S-61716, or equivalent.

D. Funnels, separatory, Squibb stopper, with Teflon stopcock plug, Pyrex brand glass, 250- and 500-milliliter capacities, Sargent Catalog S-35815-20-F or G, or equivalent.

E. Gas chromatograph, F and M Model 5750 with flame ionization detector, or equivalent.

F. Gas chromatography column: Stainless steel tubing, 6 feet by $\frac{3}{16}$ inch packed with 3 percent by weight GE SE-52 (Applied Science Laboratories) deposited on P.E. Celite 60-80 mesh (product No. 154-0048), or equivalent. Condition the column by baking for 40-80 hours at 325 °C. with a helium flow, but detached from the detector input. Injections of 1-2 microliters of a 50/50 mixture of hexamethyldisilazane and trimethylchlorosilane will help remove active sites in the column.

1. Prepare a TMS derivative of a 1,000-microgram zeranol standard as described in the procedure section. Inject 1-microliter quantities to determine whether the column is responding to the conditioning. After the column shows a response at the 1,000-microgram level, proceed to smaller quantities to optimize conditions.

2. The column and chromatograph must be conditioned to achieve a minimum sensitivity response so that a peak 5 millimeters in height results from an injection of 5 microliter of standard preparation containing 1 microgram of zeranol in the derivative preparation. This criterion must be met before tissue assay is attempted.

3. The column is brought to 250 °C. after conditioning and held at that temperature for at least 12 hours before making a run.

G. Heating mantle, electric, Glas-Col. Sargent Catalog S-40866H, or equivalent.

H. Hot plate, with gradient rheostat heat control.

I. Meat grinder, manually operated or equivalent.

J. Steam bath.

K. Syringe, Hamilton Micro Syringe Model 701, 10-microliter capacity, or equivalent.

L. Torsion balance, 0.1 gram sensitivity, 500 grams capacity.

M. Vials, 1-dram glass with plastic tops, Owens-Illinois, Opticlear, or equivalent.

N. Virtis freeze drier, Sargent Catalog S-28881-80, or equivalent.

O. Virtis homogenizing mill, macro, Virtis No. 45, Sargent Catalog S-61700, or equivalent.

IV. STANDARD SOLUTIONS

A. Stock solution A: Accurately weigh 0.1000 gram of zeranol, primary standard, into a 250-milliliter beaker. Dissolve the standard in 80 milliliters of methanol and accurately dilute to 100 milliliters in a volumetric flask with methanol. By preparation, the solution contains 1,000 micrograms per milliliter.

B. Stock solution B: Dilute 10.0 milliliters of stock solution A to 100 milliliters with methanol to provide a standard containing 100 micrograms of the drug per milliliter.

C. Stock solution C: Dilute 5.0 milliliters of stock solution B to 100 milliliters with methanol to provide a standard of 5 micrograms per milliliter.

D. Stock solution D: Dilute 2.0 milliliters of stock solution B to 100 milliliters with methanol to provide a standard of 2 micrograms per milliliter. Transfer 1.0 milliliter of stock solution D to a 1-dram glass vial, evaporate to a dry residue in a vacuum desiccator at reduced pressure. The residue contains 2 micrograms of zeranol to be used as a calibration standard in operation of the gas chromatograph.

V. PROCEDURE

A. Preparation of glassware: Glassware should be washed in detergent or chromic acid solution to remove contaminants and rinsed in water to remove traces of cleaning agent. Rinse with methanol before using.

B. Preparation of sample.

1. Collect muscle, liver, kidney, and tripe from a freshly sacrificed animal under the cleanest conditions possible.

2. Grind the fresh tissue in a meat grinder, divide into 100-gram portions, and wrap in aluminum foil. Store wrapped tissue in a deep freeze. Fat should be wrapped in foil and stored in deep freeze.

C. Extraction procedure for muscle, liver, kidney, and tripe.

1. Weight 100 grams of partially thawed tissue into a 250-milliliter homogenizing flask, add 60 milliliters of water, and attach to a Virtis "45" Tissue Mill, or equivalent.

2. Mix the materials at 45,000 r.p.m. for 5 minutes to obtain a thin homogenate.

3. Transfer the homogenate to a 1-liter, widemouth, freeze drying flask using 10-20 milliliters of water for a rinse.

4. Place the flask on its side in a nearly horizontal position in a slurry of dry ice and acetone. Rotate the flask on its side as the homogenate cools to set down a uniform frozen solid layer on the wall of the flask.

5. Mount the flask on a Virtis freeze drier, or equivalent, and lyophilize to dry solids. This operation usually requires 20-24 hours. *Stopping place.*

6. Transfer the solid cake to a clean sheet of paper and crumble by hand to a size convenient for transfer to an extraction thimble.

7. Transfer the solids to a single thickness 60 x 180 milliliter Soxhlet extraction thimble and compact the solids sufficiently to guarantee complete immersion during solid extraction.

8. Transfer 600 milliliters of methanol to a 1-liter pot of a Soxhlet extraction assembly and place the thimble in the extractor. Mount a large glass funnel in the neck of the extractor with the stem extending into the thimble. Rinse the 1-liter freeze drying flask with three 50-milliliter portions of fresh methanol and transfer the rinses through the funnel into the thimble. Mount the condenser in the extractor and extract the solids for 15 hours. The extractor should be heated with the electric heating mantle so that a fill-empty cycle requires 18-24 minutes.

9. Drain the methanol from the thimble. Composite the methanol from the extractor and pot in an 800-milliliter beaker.

10. Rinse the pot with 10 milliliters of methanol and add to the methanol composite. Transfer 50 milliliters of 2N HCl down the pot side wall, and add to methanol composite. Concentrate to 125 milliliters by boiling on a hot plate.

D. Extraction procedure for fat.

1. Cut fat into ¼-inch cubes. The lyophilization of fat is unnecessary since it is essentially water free.

2. Transfer 100 grams of the prepared fat to a 60- x 180-millimeter extraction thimble and extract with 750 milliliters of methanol for 15 hours in the Soxhlet extractor. The extractor should be heated with the electric heating mantle so that a fill-empty cycle requires 18-24 minutes.

3. Drain the methanol from the thimble. Composite the methanol from the extractor and pot into an 800-milliliter beaker.

4. Rinse the pot with 10 milliliters of methanol and add to the methanol composite.

Transfer 50 milliliters of 2N HCl down the pot side wall, and add to methanol composite. Concentrate to 125 milliliters by boiling on a hot plate.

E. Solvent partition.

1. Transfer the methanol concentrate to a 500-milliliter separatory funnel, identified by number as 1, with 70 milliliters of chloroform rinse and mix.

2. Add 300 milliliters of water and without shaking allow liquid phases to separate.

3. Withdraw the chloroform layer into a separatory funnel, identified by number as 2, containing 100 milliliters of 2 percent aqueous sodium chloride.

4. Gently mix the contents of funnel 2 horizontally end to end 30 times and allow phases to separate. Usually about 20 minutes are required to obtain maximum chloroform separation.

5. Withdraw the chloroform layer into a beaker.

6. Extract with shaking the contents of funnels 1 and 2 successively with three more 50-milliliter portions of chloroform.

7. Composite the chloroform extracts and concentrate to 125 milliliters by evaporation on a steam bath and cool to room temperature.

8. Transfer the chloroform composite to a 250-milliliter separatory funnel, fitted with a Teflon stopcock, using 10 milliliters of chloroform as a rinse.

9. Extract the chloroform with three separate 20-milliliter portions of 1N sodium hydroxide solution retaining the emulsion in the sodium hydroxide phase. Agitation of sodium hydroxide with the chloroform extract for the first time is accompanied by the appearance of emulsion.

10. Perform an extraction by gently inverting the closed funnel and returning the funnel to an upright position.

11. Repeat phase mixing 30 times per extraction.

12. Allow phases to separate for 10 minutes. The time delay allows for gradual dissipation of the emulsion to improve phase separation. The zanol transfers from the chloroform to the upper sodium hydroxide phase in this operation.

13. Composite the sodium hydroxide extracts.

14. Wash the sodium hydroxide extract with three 50-milliliter portions of chloroform using the technique as in step 9 and the same 10-minute interval for phase separation. Washing the chloroform removes the emulsion and unwanted impurities from the sodium hydroxide phase.

15. Discard the chloroform washes. Transfer the sodium hydroxide extracts to a 250-milliliter beaker. Rinse each separatory funnel with two 5-milliliter portions of water and add to the sodium hydroxide extract. Wash each funnel twice with tap water and twice with distilled water before next use.

16. Neutralize the washed sodium hydroxide extract to pH 8.0 by dropwise addition of 3N phosphoric acid using a pH meter for pH detection.

17. Transfer the pH 8.0 water extract to a 250-milliliter separatory funnel using 10 to 20 milliliters of water for a rinse.

18. Extract the solution with three separate 50-milliliter portions of carbon tetrachloride. The zeranone transfers to the lower carbon tetrachloride phase. Use the same 30-count phase-mixing technique as in step 9 and allow the mixture to stand 5 minutes for phase separation.

19. Composite the carbon tetrachloride extracts.

20. Extract the carbon tetrachloride composite with two 20-milliliter portions of 1N sodium hydroxide. Zeranone transfers from carbon tetrachloride to the upper sodium hydroxide phase. After phase mixing, allow the mixture to stand 5 minutes for phase separation.

21. Composite the sodium hydroxide extracts.

22. Wash the extract with two 50-milliliter portions of carbon tetrachloride. Allow the mixture to stand 5 minutes for phase separation. Discard the carbon tetrachloride washes.

23. Transfer the sodium hydroxide extract into a 250-milliliter beaker. Rinse the separatory funnel with two 5-milliliter portions of water and add to the sodium hydroxide extract. Wash each funnel twice with tap water and twice with distilled water before next use. Adjust the sodium hydroxide extract to a pH of 9.5 by dropwise addition of 3N phosphoric acid and transfer to a 250-milliliter separatory funnel using 10–20 milliliters of water for a rinse.

24. Extract the pH 9.5 water solution with three separate 30-milliliter portions of anhydrous ethyl ether. Allow the mixture to stand 5 minutes for phase separation. The zeranone transfers to the upper ether phase.

25. Composite the ether extracts in a 125-milliliter Erlenmeyer flask.

26. Reduce the volume of ether to about 1–2 milliliters by evaporation on a hot plate with low heat while removing vapor from top of flask by vacuum aspiration.

27. Transfer ether residue to a 1-dram glass vial. Rinse down flask side wall with 1–2 milliliters of fresh ether and transfer to the glass vial.

28. Continue evaporation of ether to 0.1 milliliter.

29. Place vial in a vacuum desiccator and evaporate residue at line vacuum and room temperature overnight to dryness.

30. Close vial with a plastic cap and submit ether residue for preparation of TMS derivative and gas chromatographic assay. *Stopping place.*

F. Gas liquid chromatography.

1. Start the gas chromatography and maintain the following operational conditions:

Carrier gas pressure: 50 p.s.i. at tank.

Carrier gas flow rate: Sufficient to give zeranone derivative peak a retention time of 4–8 minutes.

Electrometer range: 10^2 or 10^1 .

Detector temperature: 325 °C.

Injection port temperature: 325 °C.

Column temperature: 250°–280 °C., operate isothermally.

Recorder sensitivity: 1 millivolt.

Recorder chart speed: 1 inch per minute.

Sample size: 1 microliter to 5 microliters as necessary to give desired peak area for quantitative measurement.

Septums: Replace each evening and allow to condition overnight at operational temperature.

Flame assembly: Remove silica ash from the flame assembly each week. The flame assembly is removed; the anode, flame jet, and chimney are cleaned with a nylon bristle brush. Water and acetone are drawn through the jet capillary to remove any foreign material.

2. Add 0.2 milliliter of silating reagent to the sample or to the zeranone standard.

3. Stopper the vial and shake vigorously.

4. Warm the vial at 40°–50° C. for a few minutes, then roll the vial on a horizontal plane to insure that all of the interior surfaces of the vial have been in contact with the reagent.

5. Let vial stand for 4 hours or overnight in a warm area (40 °C.) to allow reaction to reach completion.

6. Place vial in a small padded centrifuge tube and centrifuge to settle the precipitate and insure that all the liquid is at the bottom of the vial.

7. Inject 1.0–5.0 microliters of clear solution into the chromatograph. At the beginning of the day's run, make 3–5 injections of a standard to condition the column for that day before taking quantitative data.

8. Run known mixtures at the beginning, middle, and end of the day's run over the concentration range of samples to be analyzed to compensate for day-to-day sensitivity fluctuations and drift. If four or less samples are to be run, calibrating at the beginning and end of the run is sufficient.

VI. CALCULATIONS

Area values are obtained on known mixtures and samples by multiplying the net peak height by the peak width at half height or by counting squares. Area values obtained on knowns are plotted versus zeranone concentration. Calibration plots indicate a near linear function in the 0–10 microgram range. Area values obtained on samples are converted directly to microgram quantities using the curve. Control tests demonstrated a 70 percent recovery of zeranone from spiked

§ 556.770

wet beef liver and muscle necessitating a correction factor.

$$\text{Zeranol, parts per billion} = \frac{\text{Micrograms of zeranol found A1,000}}{W A0.7}$$

Where:

0.7=Correction factor for 70 percent recovery.

W=Grams of tissue examined.

VII. RECOVERY STUDY

A. Fortification of reagent blank.

1. For those using this method for the first time either for recovery study or tissue assay, a solvent blank and solvent fortified with zeranol should be processed through the entire procedure. This preliminary operation will establish whether or not the procedure is free from contamination arising from solvents and glassware and demonstrate the level of recovery of the standard zeranol. Level of recovery should be in the same range as the samples.

2. Transfer 600 milliliters of methanol to a 1-liter beaker. Add 50 milliliters of 2N HCl to the methanol and concentrate to 125 milliliters by boiling on a hot plate.

3. Transfer 600 milliliters of methanol to a 1-liter beaker. Add 50 milliliters of 2N HCl to the methanol and concentrate to 125 milliliters by boiling on a hot plate. Spike the concentrate with 1.0 milliliter of stock solution D.

4. Assay both samples as described in the procedure beginning extraction step V-E1.

B. Fortification of samples.

1. Transfer 100-gram portions of partially thawed tissues into 250-milliliter homogenizing flasks and set half of them aside to serve as tissue blanks.

2. Add to the remaining samples 1 milliliter of stock solution D to serve as fortified samples to which 20 parts per billion zearalanol have been added.

3. Assay both fortified and unfortified tissue as described in the procedure section beginning with V-C1.

[40 FR 13942, Mar. 27, 1975, as amended at 54 FR 31950, Aug. 3, 1989]

§ 556.770 Zoalene.

Tolerances are established for residues of zoalene (3,5-dinitro-*o*-toluamide) and its metabolite 3-amino-5-nitro-*o*-toluamide in food as follows:

(a) In edible tissues of chickens:

(1) 6 parts per million in uncooked liver and kidney.

(2) 3 parts per million in uncooked muscle tissue.

(3) 2 parts per million in uncooked fat.

21 CFR Ch. I (4-1-99 Edition)

(b) In edible tissues of turkeys: 3 parts per million in uncooked muscle tissue and liver.

PART 558—NEW ANIMAL DRUGS FOR USE IN ANIMAL FEEDS

Subpart A—General Provisions

Sec.

558.3 Definitions and general considerations applicable to this part.

558.4 Medicated feed applications.

558.5 New animal drug requirements for liquid Type B feeds.

558.15 Antibiotic, nitrofurans, and sulfonamide drugs in the feed of animals.

Subpart B—Specific New Animal Drugs For Use in Animal Feeds

558.35 Akloamide.

558.55 Amprolium.

558.58 Amprolium and ethopabate.

558.59 Apramycin.

558.60 Arsanilate sodium.

558.62 Arsanilic acid.

558.76 Bacitracin methylene disalicylate.

558.78 Bacitracin zinc.

558.95 Bambermycins.

558.105 [Reserved]

558.115 Carbadox.

558.120 Carbarsones (not U.S.P.).

558.128 Chlortetracycline.

558.140 Chlortetracycline and sulfamethazine.

558.145 Chlortetracycline, procaine penicillin, and sulfamethazine.

558.155 Chlortetracycline, sulfathiazole, penicillin.

558.175 Clopidol.

558.185 Coumaphos.

558.195 Decoquinate.

558.205 Dichlorvos.

558.235 Efratomylin.

558.248 Erythromycin thiocyanate.

558.254 Famphur.

558.258 Fenbendazole.

558.265 Halofuginone hydrobromide.

558.274 Hygromycin B.

558.295 Iodinated casein.

558.300 Ivermectin.

558.305 Laidlomycin propionate potassium.

558.311 Lasalocid.

558.315 Levamisole hydrochloride (equivalent).

558.325 Lincomycin.

558.340 Maduramicin ammonium.

558.342 Melengestrol acetate.

558.348 Mibolerone.

558.355 Monensin.

558.360 Morantel tartrate.

558.363 Narasin.

558.365 Nequinat.

558.366 Nicarbazine.

558.369 Nitarson.

Food and Drug Administration, HHS

§ 558.3

558.376 Nitromide and sulfanitran.
558.415 Novobiocin.
558.430 Nystatin.
558.435 Oleandomycin.
558.450 Oxytetracycline.
558.460 Penicillin.
558.464 Poloxalene.
558.465 Poloxalene free-choice liquid Type C feed.
558.485 Pyrantel tartrate.
558.515 Robenidine hydrochloride.
558.530 Roxarsone.
558.550 Salinomycin.
558.555 Semduramicin.
558.575 Sulfadimethoxine, ormetoprim.
558.579 Sulfaethoxypyridazine.
558.582 Sulfamerazine.
558.586 Sulfaquinoxoline.
558.600 Tiamulin.
558.615 Thiabendazole.
558.618 Tilimicosin.
558.625 Tylosin.
558.630 Tylosin and sulfamethazine.
558.635 Virginiamycin.
558.680 Zovalene.

AUTHORITY: 21 U.S.C. 360b, 371.

SOURCE: 40 FR 13959, Mar. 27, 1975, unless otherwise noted.

Subpart A—General Provisions

§ 558.3 Definitions and general considerations applicable to this part.

(a) Regulations in this part provide for approved uses of drugs and combinations of drugs in animal feeds. Approved combinations of such drugs are specifically identified or incorporated by cross-reference. Unless specifically provided for by the regulations, a combination of two or more drugs is not approved.

(b) The following definitions apply to terms used in this part:

(1) New animal drugs approved for use in animal feed are placed in two categories as follows:

(i) Category I—These drugs require no withdrawal period at the lowest use level in each species for which they are approved.

(ii) Category II—These drugs require a withdrawal period at the lowest use level for at least one species for which they are approved or are regulated on a “no-residue” basis or with a “zero” tolerance because of a carcinogenic concern regardless whether a withdrawal period is required.

(2) A “Type A medicated article” is intended solely for use in the manufac-

ture of another Type A medicated article or a Type B or Type C medicated feed. It consists of a new animal drug(s), with or without carrier (e.g., calcium carbonate, rice hull, corn, gluten) with or without inactive ingredients. The manufacture of a Type A medicated article requires an application approved under § 514.105(a) of this chapter.

(3) A “Type B medicated feed” is intended solely for the manufacture of other medicated feeds (Type B or Type C). It contains a substantial quantity of nutrients including vitamins and/or minerals and/or other nutritional ingredients in an amount not less than 25 percent of the weight. It is manufactured by diluting a Type A medicated article or another Type B medicated feed. The maximum concentration of animal drug(s) in a Type B medicated feed is 200 times the highest continuous use level for Category I drugs and 100 times the highest continuous use level for Category II drugs. The term “highest continuous use level” means the highest dosage at which the drug is approved for continuous use (14 days or more), or, if the drug is not approved for continuous use, it means the highest level used for disease prevention or control. If the drug is approved for multiple species at different use levels, the highest approved level of use would govern under this definition. The manufacture of a Type B medicated feed from a Category II, Type A medicated article requires an application approved under § 514.105(b) of this chapter.

(4) A “Type C medicated feed” is intended as the complete feed for the animal or may be fed “top dressed” (added on top of usual ration) on or offered “free-choice” (e.g., supplement) in conjunction with other animal feed. It contains a substantial quantity of nutrients including vitamins, minerals, and/or other nutritional ingredients. It is manufactured by diluting a Type A medicated article or a Type B medicated feed. A Type C medicated feed may be further diluted to produce another Type C medicated feed. The manufacture of a Type C medicated feed from a Category II, Type A medicated article requires an application approved under § 514.105(b) of this chapter.

§ 558.4

21 CFR Ch. I (4–1–99 Edition)

(5) A Type B or Type C medicated feed manufactured from a drug component (bulk or “drum-run” (dried crude fermentation product)) requires an application approved under § 514.105(a) of this chapter.

[51 FR 7392, Mar. 3, 1986, as amended at 52 FR 2682, Jan. 26, 1987; 54 FR 51386, Dec. 15, 1989; 56 FR 19268, Apr. 26, 1991]

§ 558.4 Medicated feed applications.

(a) The manufacture of a Type B or Type C medicated feed from a Category I, Type A medicated article is exempt from the requirement of an approved medicated feed application.

(b) The manufacture of a Type B or Type C medicated feed from a Category II, Type A medicated article requires an approved medicated feed application.

(c) The use of Type B and Type C medicated feeds shall conform to the conditions of use provided for in subpart B of this part and in §§ 510.515 and 558.15.

(d) This paragraph identifies each drug by category, the maximum level of drug in Type B medicated feeds, and the assay limits for the drug in Type A medicated articles and Type B and Type C medicated feeds, as follows:

CATEGORY I

Drug	Assay limits percent ¹ type A	Type B maximum (200x)	Assay limits percent ¹ type B/C ²
Aklomide	90–110	22.75 g/lb (5.0%)	85–120.
Amprolium with Ethopabate	94–114	22.75 g/lb (5.0%)	80–120.
Bacitracin methylene disalicylate	85–115	25.0 g/lb (5.5%)	70–130.
Bacitracin zinc	84–115	5.0 g/lb (1.1%)	70–130.
Bambermycins	90–110	800 g/ton (0.09%)	80–120/70–130.
Buquinolate	90–110	9.8 g/lb (2.2%)	80–120.
Chlortetracycline	85–115	40.0 g/lb (8.8%)	80–115/70–130.
Coumaphos	95–115	6.0 g/lb (1.3%)	80–120.
Decoquinat	90–105	2.72 g/lb (0.6%)	80–120.
Dichlorvos	100–115	33.0 g/lb (7.3%)	90–120/80–130.
Efrotomycin	94–113	1.45 g/lb (0.32%)	80–120
Erythromycin (thiocyanate salt)	85–115	9.25 g/lb (2.04%)	<20g/ton 70–115/150–50>20g/ton 75–125.
Iodinated casein	85–115	20.0 g/lb (4.4%)	75–125.
Laidlomycin propionate potassium	90–110	1 g/lb (0.22%)	90–115/85–115.
Lasalocid	95–115	40.0 g/lb (8.8%)	Type B (cattle and sheep): 80–120; Type C (all): 75–125.
Lincomycin	90–115	20.0 g/lb (4.4%)	80–130.
Melengestrol acetate	90–110	10.0 g/ton (0.0011%)	70–120
Monensin	90–110	40.0 g/lb (8.8%)	Chickens, turkeys, and quail: 75–125; Cattle: 5–10 g/ton 80–120; Cattle: 10–30 g/ton 85–115; Goats: 20 g/ton 85–115; Liq. feed: 80–120.
Narasin	90–110	7.2 g/lb (1.6%)	85–115/75–125.
Nequinat	95–112	1.83 g/lb (0.4%)	80–120.
Niclosamide	85–120	225g/lb (49.5%)	80–120.
Nystatin	85–125	5.0 g/lb (1.1%)	75–125.
Oleandomycin	85–120	1.125 g/lb (0.25%)	<11.25 g/ton 70–130; >11.25 g/ton 75–125.
Oxytetracycline	90–120	20.0 g/lb (4.4%)	75–125/65–135.
Penicillin	80–120	10.0 g/lb (2.2%)	65–135.
Poloxalene	90–110	54.48 g/lb (12.0%)	Liq. feed: 85–115.
Salinomycin	95–115	6.0 g/lb (1.3%)	80–120.
Semduramicin	90–110	2.25 g/lb (0.50%)	80–110.
Tiamulin	113.4 g/lb, 100–108	3.5 g/lb (0.8%)	90–115.
Tylosin	5 and 10 g/lb, 90–115		70–130.
Virginiamycin	80–120	10.0 g/lb (2.2%)	75–125.
Zoalene	85–115	10.0 g/lb (2.2%)	70–130.
Zoalene	92–104	11.35 g/lb (2.5%)	85–115.

¹Percent of labeled amount.

Food and Drug Administration, HHS

§ 558.4

²Values given represent ranges for either Type B or Type C medicated feeds. For those drugs that have two range limits, the first set is for a Type B medicated feed and the second set is for a Type C medicated feed. These values (ranges) have been assigned in order to provide for the possibility of dilution of a Type B medicated feed with lower assay limits to make Type C medicated feed.

CATEGORY II

Drug	Assay limits percent ¹ Type A	Type B maximum (100x)	Assay limits percent ¹ Type B/C ²
Amprolium	94–114	11.35 g/lb (2.5%)	80–120.
Apramycin	88–112	7.5 g/lb (1.65%)	80–120.
Arsanilate sodium	90–110	4.5 g/lb (1.0%)	85–115/75–125.
Arsanilic acid	90–110	4.5 g/lb (1.0%)	85–115/75–125.
Carbadox	90–110	2.5 g/lb (0.55%)	75–125.
Carbarsone	93–102	17.0 g/lb (3.74%)	85–115.
Clopidol	94–106	11.4 g/lb (2.5%)	90–115/80–120.
Famphur	100–110	5.5 g/lb (1.21%)	90–115/80–120.
Fenbendazole	93–113	8.87 g/lb (1.96%)	75–125
Halofuginone hydrobromide	90–115	272.0 g/ton (.03%)	75–125.
Hygromycin B	90–110	1,200 g/ton (0.13%)	75–125.
Ivermectin	95–105	1,180 g/ton (0.13%)	80–110.
Levamisole	85–120	113.5 g/lb (25%)	85–125.
Maduramicin ammonium	90–110	545 g/ton (.06%)	80–120.
Morantel tartrate	90–110	66.0 g/lb (14.52%)	85–115.
Neomycin	80–120	7.0 g/lb (1.54%)	70–125.
Oxytetracycline	80–120	10.0 g/lb (2.2%)	65–135.
Nicarbazin (granular)	90–110	5.675 g/lb (1.25%)	85–115/75–125
Narasin	90–110	5.675 g/lb (1.25%)	85–115/75–125
Nicarbazin (powder)	98–106	5.675 g/lb (1.25%)	85–115/80–120
Nitarson	90–110	8.5 g/lb (1.87%)	85–120.
Nitromide	90–110	11.35 g/lb (2.5%)	80–120.
Sulfanitrin	85–115	13.6 g/lb (3.0%)	75–125.
Nitromide	90–110	11.35 g/lb (2.5%)	85–115.
Sulfanitrin	85–115	5.65 g/lb (1.24%)	75–125.
Roxarsone	90–110	2.275 g/lb (0.5%)	85–120.
Novobiocin	85–115	17.5 g/lb (3.85%)	80–120.
Pyrantel tartrate	90–110	36 g/lb (7.9%)	75–125.
Robenidine	95–115	1.5 g/lb (0.33%)	80–120.
Ronnel	85–115	27.2 g/lb (6.0%)	80–120.
Roxarsone	90–110	2.275 g/lb (0.5%)	85–120.
Roxarsone	90–110	2.275 g/lb (0.5%)	85–120.
Aklomide	90–110	11.35 g/lb (2.5%)	85–120.
Roxarsone	90–110	2.275 g/lb (0.5%)	85–120.
Clopidol	94–106	11.35 g/lb (2.5%)	80–120.
Bacitracin methylene disalicylate	85–115	5.0 g/lb (1.1%)	70–130.
Roxarsone	90–110	2.275 g/lb (0.5%)	85–120.
Monensin	90–110	5.5 g/lb (1.2%)	75–125.
Sulfadimethoxine	95–115	5.675 g/lb (1.25%)	85–115/75–125.
Ormetoprim (5/3)	95–115	3.405 g/lb (0.75%)	85–115.
Sulfadimethoxine	95–115	85.1 g/lb (18.75%)	85–115/75–125.
Ormetoprim (5/1)	95–115	17.0 g/lb (3.75%)	85–115.
Sulfaethoxypyridazine	95–105	50.0 g/lb (11.0%)	85–115.
Sulfamerazine	85–115	18.6 g/lb (4.0%)	85–115.
Sulfamethazine	85–115	10.0 g/lb (2.2%)	80–120.
Chlortetracycline	85–115	10.0 g/lb (2.2%)	85–125/70–130.
Penicillin	85–115	5.0 g/lb (1.1%)	85–125/70–130.
Sulfamethazine	85–115	10.0 g/lb (2.2%)	80–120.
Chlortetracycline	85–115	10.0 g/lb (2.2%)	85–125/70–130.
Sulfamethazine	85–115	10.0 g/lb (2.2%)	80–120.
Tylosin	80–120	10.0 g/lb (2.2%)	75–125.
Sulfanitrin	85–115	13.6 g/lb (3.0%)	75–125.
Aklomide	90–110	11.2 g/lb (2.5%)	85–120.
Sulfanitrin	85–115	13.6 g/lb (3.0%)	75–125.
Aklomide	90–110	11.2 g/lb (2.5%)	85–120.
Roxarsone	90–110	2.715 g/lb (0.60%)	85–120.
Sulfanitrin	85–115	13.6 g/lb (3.0%)	75–125.
Aklomide	90–110	11.2 g/lb (2.5%)	85–120.
Roxarsone	90–110	2.27 g/lb (0.5%)	85–120.
Sulfaquinoxaline	98–106	11.2 g/lb (2.5%)	85–115.
Sulfathiazole	85–115	10.0 g/lb (2.2%)	80–120.
Chlortetracycline	85–125	10.0g/lb (2.2%)	70–130.
Penicillin	80–120	5.0 g/lb (1.1%)	70–130.
Thiabendazole	94–106	45.4 g/lb (10.0%)	>7% 85–115; <7% 90–110.

CATEGORY II—Continued

Drug	Assay limits percent ¹ Type A	Type B maximum (100x)	Assay limits percent ¹ Type B/C ²
Tilmicosin	90–110	18.2 g/lb (4.0%)	85–115.

¹Percent of labeled amount.

²Values given represent ranges for either Type B or Type C medicated feeds. For those drugs that have two range limits, the first set is for a Type B medicated feed and the second set is for a Type C medicated feed. These values (ranges) have been assigned in order to provide for the possibility of dilution of a Type B medicated feed with lower assay limits to make a Type C medicated feed.

(e) When drugs from both categories are in combination, the Category II requirements will apply to the combination drug product.

[51 FR 7392, Mar. 3, 1986]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.4, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.5 New animal drug requirements for liquid Type B feeds.

(a) Information available to the Commissioner of Food and Drugs shows that certain drugs are unstable when added to some liquid Type B medicated feeds. The demonstrated instability of these drugs gives rise to the question of the stability of other drugs when added to liquid Type B medicated feeds, except where specific approval has been granted for such use. Therefore, the labeling of a drug to provide for its use in a liquid Type B medicated feed causes the drug to be a new animal drug for such use for which an approved new animal drug application is required pursuant to section 512(b) of the Federal Food, Drug, and Cosmetic Act.

(b) The addition of a drug to a liquid Type B medicated feed causes such Type B feed to become an animal feed bearing or containing a new animal drug for which an approved application is required pursuant to section 512(m) of the act.

(c) Each drug product, intended for oral administration to animals, which contains any of the drugs listed in paragraph (d) of this section and which bears labeling for its use in animal feed and/or drinking water shall also include in such labeling the following statement: “FOR USE IN _____ ONLY. NOT FOR USE IN LIQUID TYPE B MEDICATED FEEDS,” the

blank being filled in with the words “DRY FEEDS,” “DRINKING WATER,” “DRY FEEDS AND DRINKING WATER” as applicable, unless:

(1) Such drug product is the subject of an approved new animal drug application providing for its use in liquid Type B medicated feeds, or;

(2) The labeling provisions of this paragraph have been waived on the basis of approval of a petition which includes a copy of the product label; a description of the formulation; and information which establishes that the physical, chemical, or other properties of the particular drug product are such that it cannot reasonably be expected to be diverted for use in liquid Type B medicated feeds. Such petitions shall be submitted to the Food and Drug Administration, Center for Veterinary Medicine, 7500 Standish Pl., Rockville, MD 20855.

(d) The labeling provisions of paragraph (c) of this section apply to all forms of bacitracin, oxytetracycline, and chlortetracycline.

(e) For any drug which is the subject of an approved new animal drug application, the labeling provisions of paragraph (c) of this section may be implemented without prior approval as provided for in § 514.8(d) and (e) of this chapter.

[40 FR 13959, Mar. 27, 1975, as amended at 52 FR 2684, Jan. 26, 1987; 57 FR 6475, Feb. 25, 1992]

§ 558.15 Antibiotic, nitrofurantoin, and sulfonamide drugs in the feed of animals.

(a) The Commissioner of Food and Drugs will propose to revoke currently approved subtherapeutic (increased rate of gain, disease prevention, etc.) uses in animal feed of antibiotic and sulfonamide drugs whether granted by

approval of new animal drug applications, master files and/or antibiotic or food additive regulations, by no later than April 20, 1975, or the nitrofurans by no later than September 5, 1975, unless data are submitted which resolve conclusively the issues concerning their safety to man and animals and their effectiveness under specific criteria established by the Food and Drug Administration based on the guidelines included in the report of the FDA task force on the use of antibiotics in animal feeds. All persons or firms previously marketing identical, related, or similar products except the nitrofurans not the subject of an approved new animal drug application must submit a new animal drug application by July 19, 1973, or by December 4, 1973, in the case of nitrofurans, if marketing is to continue during the interim. New animal drug entities with antibacterial activity not previously marketed, now pending approval or submitted for approval prior to, on, or following the effective date of this publication, shall satisfy such criteria prior to approval.

(b) Any person interested in developing data which will support retaining approval for such uses of such antibiotic, nitrofurans, and sulfonamide drugs pursuant to section 512(l) of the Federal Food, Drug, and Cosmetic Act shall submit to the Commissioner the following:

(1) By July 19, 1973, records and reports of completed, ongoing, or planned studies, including protocols, on the tetracyclines, streptomycin, dihydrostreptomycin, penicillin, and the sulfonamides; for all other antibiotics by October 17, 1973; and for the nitrofurans by March 4, 1974. The Food and Drug Administration encourages sponsors to consult with the Center for Veterinary Medicine on protocol design and plans for future studies.

(2) By April 20, 1974, data from completed studies on the tetracyclines, streptomycin, dihydrostreptomycin, the sulfonamides, and penicillin assessing the effect of the subtherapeutic use of the drug in feed on the salmonella reservoir in the target animal as compared to that in nonmedicated controls. Failure to complete the salmonella studies for any of these drugs

by that time will be grounds for proceeding to immediately withdraw approval.

(3) By April 20, 1975, data satisfying all other specified criteria for safety and effectiveness, including the effect on the salmonella reservoir for any antibiotic or sulfonamide drugs and by September 5, 1975, for the nitrofurans, approved for subtherapeutic use in animal feeds. Drug efficacy data shall be submitted for any feed-use combination product containing such drug and any feed-use single ingredient antibiotic, nitrofurans, or sulfonamide not reviewed by the National Academy of Sciences—National Research Council, Drug Efficacy Study covering drugs marketed between 1938 and 1962.

(4) Progress reports on studies underway every January 1 and July 1 until completion.

(c) Failure on the part of any sponsor to comply with any of the provisions of paragraph (b) of this section for any of the antibacterial drugs included in paragraph (b)(1) of this section, or interim results indicating a health hazard, will be considered as grounds for immediately proceeding to withdraw approval of that drug for use in animal feeds under section 512(l) of the act in the case of failure to submit required records and reports and under section 512(e) where new information shows that such drug is not shown to be safe.

(d) Criteria based upon the guidelines laid down by the task force may be obtained from the Food and Drug Administration, Center for Veterinary Medicine, 7500 Standish Pl., Rockville, MD 20855.

(e) Reports as specified in this section shall be submitted to: Food and Drug Administration, Center for Veterinary Medicine, Office of New Animal Drug Evaluation (HFV-100), 7500 Standish Pl., Rockville, MD 20855.

(f) Following the completion of the requirements of paragraphs (a) and (b) of this section and the studies provided for therein:

(1) Those antibiotic, nitrofurans, and sulfonamide drugs which fail to meet the prescribed criteria for subtherapeutic uses but which are found to be effective for the therapeutic purposes will be permitted in feed only for high-level, short-term therapeutic use and

§ 558.15

21 CFR Ch. I (4–1–99 Edition)

only by or on the order of a licensed veterinarian.

(2) Animal feeds containing antibacterial drugs permitted to remain in use for subtherapeutic purposes shall be labeled to include a statement of the quantity of such drugs.

(g) The submission of applications and data required by paragraphs (a) and (b) of this section is not required for the continued manufacture of any Type A medicated article which is produced solely from a Type A article that is in compliance with the requirements

of this section: *Provided*, That the Type A medicated article contains no drug ingredient whose use in or on animal feed requires an approved application pursuant to section 512(m) of the act and/or where the Type A article is approved by regulation in this part.

(1) The following antibacterial Type A articles manufactured by the designated sponsors are eligible for interim marketing based on their compliance with the requirements of this section:

Drug sponsor	Type A article	Species	Use levels	Indications for use
Pitman-Moore, Inc.	Bacitracin zinc	Chickens, turkeys, swine, pheasants, quail, and cattle.	Sec. 558.78	Sec. 558.78.
A. L. Laboratories, Inc.	do	Chickens, turkeys, pheasants, and quail.	do	Do.
A.L. Laboratories, Inc., Fermenta Animal Health Co..	Bacitracin methylene disalicylate.	Chicken turkeys, swine, and cattle.	Sec. 558.76	Sec. 558.76.
Elanco Products Co	Hygromycin B	Chickens and swine.	Sec. 558.274	Sec. 558.274.
Do	Tylosin	Chickens, swine, and beef cattle.	Sec. 558.625	Sec. 558.625.
Sanofi Animal Health, Inc.	Erythromycin	Chickens, turkeys, and swine.	Sec. 558.248	Sec. 558.248.
The Upjohn Co	Lincomycin	Chickens	Sec. 558.325	Sec. 558.325.
Pfizer, Inc.	Oleandomycin	Chickens, turkeys, and swine.	Sec. 558.435	Sec. 558.435.
Hoechst-Roussel Agri-Vet, Inc ..	Bambermycins	Chickens	Sec. 558.95	Sec. 558.95.
Elanco Products Co	Tylosin and sulfamethazine.	Swine	Sec. 558.630	Sec. 558.630.
American Cyanamid Co., Fermenta Animal Health Co., Feed Specialties Co., Inc., Pfizer, Inc., PennField Oil Co., and VPO, Inc..	Chlortetracycline ..	Chickens, turkeys, swine, and cattle.	Sec. 558.128	Sec. 558.128.
Merck Sharp & Dohme Research Labs., and Solvay Veterinary, Inc.	Procaine Penicillin	Chickens, turkeys, swine, pheasants, and quail.	Sec. 558.460	Sec. 558.460.
Pfizer, Inc., PennField Oil Co. ..	Oxytetracycline	Sec. 558.450	Sec. 558.450	Sec. 558.450.
American Cyanamid Co	Chlortetracycline and sulfamethazine.	Cattle	Sec. 558.128	Sec. 558.128.
Sanofi Animal Health, Inc.	Erythromycin	Cattle	37 mg per head per day. Sec. 558.575	Sec. 558.248.
Hoffman-La Roche, Inc	Sulfadimethoxine and ormetoprim.	Chickens and turkeys..	Sec. 558.575	Sec. 558.575.
Pfizer, Inc.	Oxytetracycline and neomycin.	Chickens, turkeys, swine, and calves.	As provided in paragraph (g)(2) of this section. do	As provided in paragraph (g)(2) of this section. Do.
American Cyanamid Co. and Pfizer, Inc.	Chlortetracycline, sulfamethazine, and penicillin.	Swine	do	Do.
Fermenta Animal Health Co.	Chlortetracycline, sulfathiazole, and penicillin.	do	do	Do.

(2) The following is a list of drug combinations permitted when prepared from antibacterial Type A articles list-

ed in paragraph (g)(1) of this section. Drug combinations listed in subpart B of this part name their sponsors and

Food and Drug Administration, HHS

§ 558.15

are incorporated herein by reference since they are safe and effective by contemporary standards, or such sponsors have been notified of any additional safety or efficacy data required on an individual basis:

Drug sponsor	Type A article	Species	Use levels	Indications for use
Fermenta Animal Health Co.	Chlortetracycline and arsanilic acid.	Swine	10 to 50 g/ton and 0.005 to 0.01 percent.	Enhancement of growth and feed efficiency.
American Cyanamid Co	Chlortetracycline and sulfamethazine.	Cattle	Sec. 558.128	Sec. 558.128.
Pfizer, Inc., PennField Oil Co., and VPO, Inc.	Oxytetracycline and neomycin base.	Chickens	50 g/ton and 35 to 140 g/ton.	Prevention of diseases from oxytetracycline susceptible organisms during periods of stress. As an aid in the prevention of bacterial enteritis and in the control of neomycin-sensitive organisms associated with bluecomb (mud fever or nonspecific enteritis).
Do	do	Chickens (first 2 weeks).	50 to 100 g/ton and 35 to 140 g/ton.	Prevention of early chick mortality due to oxytetracycline-susceptible organisms. As an aid in the prevention of bacterial enteritis and in the control of neomycin-sensitive organisms associated with bluecomb (mud fever or nonspecific enteritis).
Do	do	Chickens	do	To extend period of high egg production, to improve feed efficiency, to improve egg production and feed efficiency in presence of disease and at time of stress. As an aid in maintaining and improving hatchability where birds are suffering stress from moving, vaccinations, culling, extreme temperature changes, and worming; to improve livability of progeny when losses are due to oxytetracycline-susceptible organisms, to improve egg shell quality, prevention of bluecomb (mud fever or nonspecific enteritis). As an aid in the prevention of bacterial enteritis and in the control of neomycin-sensitive organisms associated with bluecomb (mud fever or nonspecific enteritis).
Do	do	do	100 to 200 g/ton and 35 to 140 g/ton.	Prevention of complicated chronic respiratory disease (air-sac infection) and control of complicated chronic respiratory disease by lowering mortality and severity during outbreaks. As an aid in the prevention of bacterial enteritis and in the control of neomycin-sensitive organisms associated with bluecomb (mud fever or nonspecific enteritis).

Drug sponsor	Type A article	Species	Use levels	Indications for use
Do	do	Turkeys	50 g/ton and 35 to 140 g/ton.	As an aid in the prevention of disease from oxytetracycline susceptible organisms during periods of stress. As an aid in the prevention of bacterial enteritis and in the control of neomycin-sensitive organisms associated with bluecomb (mud fever or non-specific enteritis).
Do	do	do	50 to 100 g/ton and 35 to 140 g/ton.	To extend period of high egg production, to improve egg production, to improve feed efficiency, to improve fertility, to improve egg production and feed efficiency in presence of disease and time of stress; as an aid in maintaining and improving hatchability where birds are suffering from stress, exposure, moving, vaccination, culling, extreme losses due to oxytetracycline-susceptible organisms, and to improve egg shell quality prevention of hexamitiasis. As an aid in the prevention of bacterial enteritis and in the control of neomycin-sensitive organisms associated with bluecomb (mud fever or non-specific enteritis).
Do	do	Turkeys (first 4 weeks).	do	As an aid in the prevention of early poult mortality due to oxytetracycline-susceptible organisms. As an aid in the prevention of bacterial enteritis and in the control of neomycin-sensitive organisms associated with bluecomb (mud fever or nonspecific enteritis).
Do	do	do	100 to 150 g/ton and 35 to 105 g/ton.	As an aid in reducing mortality in birds which have suffered an attack of air-sacculitis (it is recommended, wherever possible, to feed from time of attack to marketing).
Do	do	Turkeys	do	As an aid in the prevention of bacterial enteritis and in the control of neomycin-sensitive organisms associated with bluecomb (mud fever or non-specific enteritis).
Do	do	do	100 to 200 g/ton and 35 to 140 g/ton.	Control of bluecomb (mud fever or nonspecific enteritis), infectious sinusitis and hexamitiasis, prevention of infectious synovitis. As an aid in the prevention of bacterial enteritis and in the control of neomycin-sensitive organisms associated with bluecomb (mud fever or non-specific enteritis).
Do	do	do	200 g/ton and 70 to 140 g/ton.	Control of infectious synovitis. For the treatment of bacterial enteritis and bluecomb (mud fever or nonspecific enteritis).

Food and Drug Administration, HHS
§ 558.15

Drug sponsor	Type A article	Species	Use levels	Indications for use
Do	do	Swine	50 g/ton and 35 to 140 g/ton.	As an aid in the prevention of bacterial enteritis (scours), baby pig diarrhea (in baby pigs only), vibronic dysentery, bloody dysentery, and salmonellosis (necro or necrotic enteritis).
Do	do	do	50 to 150 g/ton and 70 to 140 g/ton.	As an aid in the maintenance of weight gains and feed consumption in the presence of atrophic rhinitis. As an aid in the treatment of bacterial enteritis.
Do	do	Calves	50 g/ton and 35 to 140 g/ton.	As an aid in the prevention of bacterial enteritis (scours).
Do	do	do	100 g/ton and 70 to 140 g/ton.	As an aid in the treatment of bacterial enteritis (scours).
Do	do	do	8 to 100 mg/gal and 100 to 200 mg/gal reconstituted milk replacer.	As an aid in the prevention of bacterial diarrhea (scours).
Do	do	do	40 to 200 mg/gal and 200 to 400 mg/gal reconstituted milk replacer.	As an aid in the treatment of bacterial diarrhea (scours).
The Upjohn Co.	Lincomycin, amprolium, and ethopabate.	Chickens	Secs. 558.58 and 558.325.	Secs. 558.58 and 558.325.
Do	Lincomycin and zoalene.	do	Secs. 558.325 and 558.680.	Secs. 558.325 and 558.680.
Do	Lincomycin, amprolium, ethopabate, and roxarsone.	do	Secs. 558.58, 558.325, and 558.530.	Secs. 558.58, 558.325, and 558.530.
Do	Lincomycin, monensin, and roxarsone.	do	Secs. 558.325, 558.355, and 558.530.	Secs. 558.325, 558.355, and 558.530.
Do	Nicarbazine and procaine penicillin.	Chickens	0.01 to 0.02 percent and 2.4 to 50 g/ton.	Do.
Do	Nicarbazine and bacitracin methylene disalicylate.	do	0.01 to 0.02 percent and 4 to 50 g/ton.	Do.
Do	Nicarbazine, bacitracin methylene disalicylate, and roxarsone.	do	0.01 to 0.02 percent, 4 to 50 g/ton, and 0.0025 to 0.005 percent.	Do.
Do	Nicarbazine, procaine penicillin, and roxarsone.	do	0.01 to 0.02 percent, 2.4 to 50 g/ton, and 0.0025 to 0.025 percent.	Do.
Do	Amprolium and bacitracin methylene disalicylate.	Chickens and turkeys.	0.0125 to 0.025 percent and 4 to 50 g/ton.	Secs. 558.55 and 558.76.
Do	Amprolium, ethopabate, and bacitracin methylene disalicylate.	Chickens	0.0125 to 0.025 percent, 0.0004 percent, and 4 to 50 g/ton.	Secs. 558.58 and 558.76.
Do	Amprolium, ethopabate, bacitracin methylene disalicylate, and roxarsone.	do	0.0125 to 0.025 percent, 0.0004 percent, 4 to 50 g/ton, and 0.0025 to 0.005 percent.	Secs. 558.58, 558.76, and 558.530.
Do	Amprolium and procaine penicillin.	Chickens and turkeys.	0.004 to 0.025 percent and 2.4 to 50 g/ton.	Secs. 558.55 and 558.460.

Drug sponsor	Type A article	Species	Use levels	Indications for use
Do	Amprolium, procaine penicillin, and roxarsone.	Chickens	0.004 to 0.025 percent, 2.4 to 50 g/ton, and 0.0025 to 0.005 percent.	Secs. 558.55, 558.460, and 558.530.
Do	Amprolium, ethopabate, procaine penicillin, and erythromycin.	do	0.0125 to 0.025 percent, 0.0004 percent, 2.4 to 50 g/ton, and 4.6 to 18.5 g/ton.	Secs. 558.58 and 558.460.
Do	Amprolium and erythromycin.	do	0.0125 to 0.025 percent and 4.6 to 18.5 g/ton.	Sec. 558.55.
Do	Amprolium and ethopabate.	do	0.0125 to 0.025 percent and 0.0004 percent.	Sec. 558.58.
Do	Amprolium, arsanilic acid, and erythromycin.	do	0.0125 to 0.025 percent, 0.01 percent, and 4.6 to 18.5 g/ton.	Sec. 558.55.
Do	Amprolium, arsanilic acid, and ethopabate.	do	0.0125 to 0.025 percent, 0.01 percent, and 0.0004 percent.	Sec. 558.58.
Do	Amprolium, ethopabate, and bacitracin methylene disalicylate.	do	0.0125 percent, 0.004 percent, and 4 to 50 g/ton.	Do.
Do	Amprolium, ethopabate, bacitracin methylene disalicylate, and roxarsone.	do	0.0125 percent, 0.004 percent, 5 to 35 g/ton, and 0.00375 percent.	Do.
Pitman-Moore, Inc.	Bacitracin zinc, amprolium, and ethopabate.	do	4 to 50 g/ton, 0.0125 to 0.025 percent, and 0.0004 percent.	Prevention of coccidiosis. Growth promotion and feed efficiency. Sec.558.78.
Do	Bacitracin zinc, amprolium, ethopabate, and roxarsone.	do	4 to 50 g/ton, 0.0125 to 0.025 percent, 0.0004 percent, and 0.0025 to 0.005 percent.	Prevention of coccidiosis. Growth promotion and feed efficiency. Improving pigmentation. Sec. 558.78.
Do	Bacitracin zinc and arsanilic acid.	Swine	10 to 50 g/ton and 0.005 to 0.01 percent.	Increased rate of weight gain and improved feed efficiency.
Merck Sharp & Dohme Research Labs.	Amprolium, ethopabate, procaine penicillin, and roxarsone.	Chickens	0.125 to 0.025 percent, 0.0004 percent, 2.4 to 50 g/ton, and 0.0025 to 0.005 percent.	Secs. 558.58, 558.460 and 558.530.
A. L. Laboratories, Inc.	Zoalene and bacitracin methylene disalicylate.	Chickens	0.0125 percent and 4 to 50 g/ton.	Sec. 558.680.
Do	Zoalene, roxarsone, and bacitracin methylene disalicylate.	do	0.0125 percent, 0.005 percent, and 4 to 50 g/ton.	Do.
Do	Zoalene and bacitracin zinc.	do	0.0125 percent and 4 to 50 g/ton.	Do.
Do	Zoalene, roxarsone, and bacitracin zinc.	do	0.0125 percent, 0.0025 to 0.005 percent, and 4 to 50 g/ton.	Do.
Do	Zoalene and penicillin.	do	0.0125 percent and 2.4 to 50 g/ton.	Do.

Food and Drug Administration, HHS

§ 558.35

Drug sponsor	Type A article	Species	Use levels	Indications for use
Do	Zoalene, roxarsone, and penicillin.	do	0.0125 percent, 0.0025 to 0.005 percent, and 2.4 to 50 g/ton.	Do.
Do	Zoalene, arsanilic acid, and bacitracin methylene disalicylate or bacitracin zinc.	do	0.0125 percent, 0.01 percent, and 4 to 50 g/ton.	Do.
Do	Zoalene, arsanilic acid, and penicillin.	do	0.0125 percent, 0.01 percent, and 2.4 to 50 g/ton.	Do.
Do	Zoalene, and bacitracin methylene disalicylate.	do	0.004 to 0.0125 percent and 4 to 50 g/ton.	Do.
Do	Zoalene, roxarsone, and bacitracin methylene disalicylate.	do	0.004 to 0.0125 percent, 0.0025 to 0.005 percent, and 4 to 50 g/ton.	Do.
Whitmoyer Labs, Inc	Carbarsone and bacitracin.	Turkeys	Sec. 558.120	Sec. 558.120.
Elanco Products Co.	Hygromycin B and tylosin.	Chickens	8 to 12 g/ton and 4 to 50 g/ton.	Sec. 558.274.
Do	do	Swine	12 g/ton and 10 to 100 g/ton.	Do.
A. L. Laboratories, Inc.	Nitarson and bacitracin zinc.	Turkeys	0.01875 percent, 4 to 50 g/ton.	As an aid in the prevention of blackhead. To increase rate of weight gain and improve feed efficiency.

[51 FR 8811, Mar. 14, 1986; 51 FR 11014, Apr. 1, 1986, as amended at 51 FR 28547, Aug. 8, 1986; 53 FR 20843, June 7, 1988; 54 FR 37098, Sept. 7, 1989; 54 FR 51386, Dec. 15, 1989; 55 FR 8460, 8462, Mar. 8, 1990; 56 FR 41912, Aug. 23, 1991; 56 FR 64702, Dec. 12, 1991; 57 FR 6476, Feb. 25, 1992; 57 FR 8577, Mar. 11, 1992; 57 FR 14639, Apr. 22, 1992; 58 FR 17515, Apr. 5, 1993; 58 FR 30119, May 26, 1993; 61 FR 51589, Oct. 3, 1996; 64 FR 992, Jan. 7, 1999]

Subpart B—Specific New Animal Drugs for Use in Animal Feeds

§ 558.35 Aklomide.

(a) *Approvals.* Type A medicated articles: to 053501 in §510.600(c) of this chapter, as follows:

- (1) 50 percent aklomide.
- (2) 20 percent sulfanitran and 25 percent aklomide.
- (3) 25 percent aklomide, 20 percent sulfanitran, and 5 percent roxarsone.
- (4) 50 percent aklomide and 10 percent roxarsone.

(b) *Related tolerances.* See §556.30 of this chapter.

(c) *Conditions of use.* It is used in feed for chickens as follows:

- (1) *Amount per ton.* Aklomide, 227 grams (0.025 percent).
- (i) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. tenella* and *E. necatrix*.

(ii) *Limitations.* Not to be fed to birds laying eggs for human consumption.

(2) *Amount per ton.* Aklomide, 227 grams (0.025 percent) combined with sulfanitran, 181.6 grams (0.02 percent).

(i) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, and *E. acervulina*.

(ii) *Limitations.* Not to be fed to laying chickens; withdraw 5 days before slaughter.

(3) *Amount per ton.* Aklomide, 227 grams (0.025 percent) combined with sulfanitran, 181.6 grams (0.02 percent) + roxarsone, 22.7–45.4 grams (0.0025–0.005 percent).

(i) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, and *E. acervulina*; growth promotion and feed efficiency; improving pigmentation.

(ii) *Limitations.* Not to be fed to laying chickens; withdraw 5 days before

§ 558.55

21 CFR Ch. I (4–1–99 Edition)

slaughter; as sole source of organic arsenic; chickens should have access to drinking water at all times.

(4) *Amount per ton.* Aklomide, 227 grams (0.025 percent) combined with roxarsone, 22.7–45.4 grams (0.0025–0.005 percent).

(i) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. tenella*, and *E. necatrix*; growth promotion and feed efficiency; improving pigmentation.

(ii) *Limitations.* Not to be fed to birds laying eggs for human consumption; withdraw 5 days before slaughter; as sole source of organic arsenic; chickens should have access to drinking water at all times.

[40 FR 13959, Mar. 27, 1975, as amended at 41 FR 8312, Feb. 25, 1976; 51 FR 7395, Mar. 3, 1986; 55 FR 8460, Mar. 8, 1990]

§ 558.55 Amprolium.

(a) *Approvals.* Type A medicated articles: 25 percent to 050604 in § 510.600(c) of this chapter for use as in paragraph (d) of this section.

(b) *Special considerations.* Do not use in Type B or Type C medicated feeds containing bentonite.

(c) *Related tolerances.* See § 556.50 of this chapter.

(d) *Conditions of use—(1) Calves.* It is top-dressed on or thoroughly mixed in the daily feed ration as follows:

(i) *Amount.* 227 milligrams per 100 pounds (5 milligrams per kilogram) body weight per day.

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii*.

(b) *Limitations.* Administer from a Type B feed containing from 0.05 to 1.25 percent amprolium with the usual amount of feed consumed in 1 day; feed for 21 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard; withdraw 24 hours before slaughter; as sole source of amprolium.

(ii) *Amount.* 454 milligrams per 100 pounds (10 milligrams per kilogram) body weight per day.

(a) *Indications for use.* As an aid in the treatment of coccidiosis caused by *Eimeria bovis* and *E. zurnii*.

(b) *Limitations.* Administer from a Type B feed containing from 0.05 to 1.25 percent amprolium with the usual amount of feed consumed in 1 day; feed for 5 days; for a satisfactory diagnosis, a microscopic examination of the feces should be done by a veterinarian or diagnostic laboratory before treatment; when treating outbreaks, the drug should be administered promptly after diagnosis is determined; withdraw 24 hours before slaughter; as sole source of amprolium.

(2) *Chickens and turkeys.* It is used as follows:

Amprolium in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 36.3 to 113.5 (0.004% to 0.0125%).		Replacement chickens; development of active immunity to coccidiosis.	Feed as follows—	

Growing conditions	Up to 5 weeks of age	From 5 to 8 weeks of age	Over 8 weeks of age
	<i>Amprolium grams per ton</i>	<i>Amprolium grams per ton</i>	<i>Amprolium grams per ton</i>
Severe exposure to coccidiosis.	113.5 (0.0125%)	72.6–113.5 (0.008%–0.0125%)	36.3–113.5 (0.004%–0.0125%)
Moderate exposure to coccidiosis.	72.6–113.5 (0.008%–0.0125%)	54.5–113.5 (0.006%–0.0125%)	36.3–113.5 (0.004%–0.0125%)
Slight exposure to coccidiosis.	36.3–113.5 (0.004%–0.0125%)	36.3–113.5 (0.004%–0.0125%)	36.3–113.5 (0.004%–0.0125%)

Food and Drug Administration, HHS

§ 558.55

Amprolium in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 36.3 to 113.5 (0.004% to 0.0125%).	Arsanilate sodium 90 (0.01%).	Replacement chickens; development of active immunity to coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic; feed according to subtable in item (i).	
	Arsanilic acid 90 (0.01%).	Replacement chickens; development of active immunity to coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic; feed according to subtable in item (i).	
	Arsanilic acid 90 (0.01%) plus erythromycin 4.6 to 18.5.	Replacement chickens; development of active immunity to coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic. Feed according to subtable in item (i).	
	Arsanilic acid 90 (0.01%) plus erythromycin 92.5.	1. Replacement chickens; development of active immunity to coccidiosis; growth promotion and feed efficiency; improving pigmentation; as an aid in the prevention of chronic respiratory disease during periods of stress. 2. Replacement chickens; development of active immunity to coccidiosis; growth promotion and feed efficiency; improving pigmentation; as an aid in the prevention of infectious coryza.	Feed for 2 d before stress and 3 to 6 d after stress; withdraw 5 d before slaughter; as sole source of organic arsenic. Feed according to subtable in item (i). Feed for 7 to 14 d; withdraw 5 d before slaughter; as sole source of organic arsenic. Feed according to subtable in item (i).	
	Arsanilic acid 90 (0.01%) plus erythromycin 185.	Replacement chickens; development of active immunity to coccidiosis; growth promotion and feed efficiency; improving pigmentation; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease.	Feed for 5 to 8 d; do not use in birds producing eggs for food purposes; withdraw 5 d before slaughter; as sole source of organic arsenic. Feed according to subtable in item (i).	
	Bacitracin 100 to 200	Replacement chickens; development of active immunity to coccidiosis; treatment of chronic respiratory disease (air-sac infection) and blue comb (nonspecific infectious enteritis).	As bacitracin methylene disalicylate or bacitracin zinc. Feed according to subtable in item (i).	
	Chlortetracycline 100 to 200.	Chickens; development of active immunity to coccidiosis; control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption. Feed for 7 to 14 d.	
	Chlortetracycline 200 to 400.	Chickens; development of active immunity to coccidiosis; control of chronic respiratory disease (CRD) and air sac infection caused by <i>M. gallisepticum</i> and <i>E. coli</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption. Feed for 7 to 14 d.	
	Erythromycin 4.6 to 18.5.	Replacement chickens; development of active immunity to coccidiosis; growth promotion and feed efficiency.	As erythromycin thiocyanate. Feed according to subtable in item (i).	
	Erythromycin 92.5	1. Replacement chickens; development of active immunity to coccidiosis; as an aid in the prevention of infectious coryza.	Feed for 7 to 14 d; withdraw 24 h before slaughter. Feed according to subtable in item (i).	

Amprolium in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(ii) 72.6 to 113.5 (0.008% to 0.0125%).	Erythromycin 185	2. Replacement chickens; development of active immunity to coccidiosis; as an aid in the prevention of chronic respiratory disease during periods of stress.	Feed for 2 d before stress and 3 to 6 d after stress; withdraw 24 h before slaughter. Feed according to subtable in item (i).	
	Hygromycin B 8 to 12	Replacement chickens; development of active immunity to coccidiosis; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease.	Feed for 5 to 8 d; do not use in birds producing eggs for food purposes; withdraw 48 h before slaughter. Feed according to subtable in item (i).	
	Penicillin 2.4 to 50	Replacement chickens; development of active immunity to coccidiosis; control of infestation of large round worms (<i>Ascaris galli</i>), cecal worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>).	Feed according to subtable in item (i).	
	Roxarsone 22.7 to 45.4 (0.0025% to 0.005%).	Replacement chickens; development of active immunity to coccidiosis; growth promotion and feed efficiency.	As procaine penicillin. Feed according to subtable in item (i).	
		Replacement chickens; development of active immunity to coccidiosis; growth promotion, and feed efficiency; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic. Feed according to subtable in item (i).	
	Arsanilate sodium 90 (0.01%).	Broiler chickens; prevention of coccidiosis caused by <i>Eimeria tenella</i> only.	Withdraw 5 d before slaughter; as sole source of organic arsenic. Feed according to subtable in item (i).	
	Arsanilic acid 90 (0.01%).	do	Withdraw 5 d before slaughter; as sole source of organic arsenic.	
	Bacitracin 100 to 200	do	do	
	Chlortetracycline 100 to 200.	Broiler chickens; prevention of coccidiosis caused by <i>E. tenella</i> only; treatment of chronic respiratory disease (air-sac infection) and blue comb (nonspecific infectious enteritis).	As bacitracin methylene disalicylate, or zinc bacitracin.	
	Chlortetracycline 200 to 400.	Chickens; prevention of coccidiosis caused by <i>E. tenella</i> only; control of infectious synovitis caused by <i>M. synoviae</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption. Feed for 7 to 14 d.	
	Hygromycin B 8 to 12	Chickens; prevention of coccidiosis caused by <i>E. tenella</i> only; control of chronic respiratory disease (CRD) and air sac infection caused by <i>M. gallisepticum</i> and <i>E. coli</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption. Feed for 7 to 14 d.	
	Penicillin 2.4 to 50	Broiler chickens; prevention of coccidiosis caused by <i>Eimeria tenella</i> only; control of infestation of large round worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>).	Feed according to subtable in item (i).	
		Broiler chickens; prevention of coccidiosis caused by <i>E. tenella</i> only; growth promotion and feed efficiency.	As procaine penicillin	

Food and Drug Administration, HHS

§ 558.55

Amprolium in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(iii) 113.5 (0.0125%) ...	Roxarsone 22.7 to 45.4 (0.0025% to 0.005%).	Broiler chickens; prevention of coccidiosis caused by <i>E. tenella</i> only; growth promotion and feed efficiency; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic.	012799
		1. Laying chickens; prevention of coccidiosis. 2. Laying chickens; treatment of coccidiosis.	 For moderate outbreaks of coccidiosis; administer for 2 weeks.	
(iv) 113.5 to 227 (0.0125% to 0.025%).	Bambermycins 1 to 3 plus roxarsone 22.8 to 34.1 (0.0025% to 0.00375%).	Broiler chickens; as an aid in the prevention of coccidiosis; for increased rate of weight gain, improved feed efficiency, and improved pigmentation.	Feed continuously as the sole ration; as sole source of amprolium and organic arsenic; roxarsone as provided by No. 053501 in §510.600(c) of this chapter, bambermycins by No. 012799; withdraw 5 d before slaughter.	
	Bambermycins 1 to 4	Growing turkeys; prevention of coccidiosis; increased rate of weight gain and improved feed efficiency.	Feed continuously as the sole source of amprolium; bambermycins as provided by No. 012799 in §510.600(c) of this chapter.	
		1. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis. 2. Turkeys; prevention of coccidiosis.	do	
	Arsanilate sodium 90 (0.01%).	1. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation. 2. Turkeys; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic. do	
	Arsanilic acid 90 (0.01%).	1. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation. 2. Turkeys; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation.	do	
	Arsanilic acid 90 (0.01%) plus erythromycin 92.5.	1. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation; as an aid in the prevention of chronic respiratory disease during periods of stress. 2. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation; as an aid in the prevention of infectious coryza.	Feed for 2 d before stress and 3 to 6 d after stress; withdraw 5 d before slaughter; as sole source of organic arsenic. Feed for 7 to 14 d; withdraw 5 d before slaughter; as sole source of organic arsenic.	

Amprolium in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
	Arsanilic acid 90 (0.01%) plus erythromycin 185.	Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease.	Feed for 5 to 8 d; do not use in birds producing eggs for food purposes; withdraw 5 d before slaughter; as sole source of organic arsenic.	1012769
	Arsanilic acid 90 (0.01%) plus erythromycin 4.6 to 18.5.	Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improved pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic.	
	Bacitracin 4 to 50	1. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency. 2. Turkeys; prevention of coccidiosis; growth promotion and feed efficiency.	As bacitracin methylene disacylate or bacitracin zinc. do	
	Bacitracin 100 to 200	1. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; treatment of chronic respiratory disease (air-sac infection), blue comb (non-specific infectious enteritis). 2. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; treatment of chronic ry disease (air-sac infection), blue comb (nonspecific infectious enteritis).	do As bacitracin zinc	
	Bacitracin 100 to 500	Turkeys; prevention of coccidiosis; treatment of infectious sinusitis, blue comb (mud fever). do	As bacitracin zinc Feed contains 50% to 75% of bacitracin but not more than 125 g penicillin; as procaine penicillin; as bacitracin zinc.	000006
	Bacitracin plus penicillin 100 to 500 (of combination).			
	Carbarsone 227 to 340.5.	Turkeys; aid in prevention of coccidiosis (<i>Eimeria adenoides</i> , <i>E. meleagris</i> , and <i>E. gallopavonis</i>) and blackhead.	Feed continuously 2 weeks before coccidiosis and blackhead are expected and continue as long as prevention is needed; withdraw 5 days before slaughter; use as sole source of amprolium and organic arsenic; do not use as a treatment for outbreaks of coccidiosis; carbarsone by 046573 in § 510.600(c) of this chapter.	
	Chlortetracycline 100 to 200.	Chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; control of infectious synovitis caused by <i>M. synoviae</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption. Feed for 7 to 14 d.	

Food and Drug Administration, HHS

§ 558.55

Amprolium in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
	Chlortetracycline 200 to 400.	Chickens where immunity to coccidiosis is not desired; control of chronic respiratory disease (CRD) and air sac infection caused by <i>M. gallisepticum</i> and <i>E. coli</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption. Feed for 7 to 14 d.	
	Erythromycin 4.6 to 18.5.	Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency.	As erythromycin thiocyanate	
	Erythromycin 92.5	1. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; as an aid in the prevention of chronic respiratory disease during periods of stress. 2. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; as an aid in the prevention of infectious coryza.	Feed for 2 d before stress and 3 to 6 d after stress; withdraw 24 h before slaughter. Feed for 7 to 14 d; withdraw 24 h before slaughter.	
	Erythromycin 185	Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease.	Feed for 5 to 8 d, do not use in birds producing eggs for food purposes; withdraw 48 h before slaughter.	
	Hygromycin B 8 to 12	Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; control of infestation of large round worms (<i>Heterakis gallinae</i>) and capillary worms (<i>Capillaria obsignata</i>).	Feed according to subtable in item (i).	
	Penicillin 2.4 to 50	1. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency. 2. Turkeys; prevention of coccidiosis; growth promotion and feed efficiency.	As procaine penicillin do	
	Roxarsone 22.7 to 45.4 (0.0025% to 0.005%).	1. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation. 2. Turkeys; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic. do	
(v) 227 (0.025%)		Laying chickens; treatment of coccidiosis.	For severe outbreaks of coccidiosis; administer for 2 weeks.	

(3) *Pheasants*. It is used as follows:

(i) *Amount*. 0.0175 percent (159 grams per ton).

§ 558.58

21 CFR Ch. I (4–1–99 Edition)

(ii) *Indications for use.* For the prevention of coccidiosis in growing pheasants caused by *Eimeria colchici*, *E. duodenalis*, and *E. phasiani*.

(iii) *Limitations.* Feed continuously as sole ration. Use as sole source of amprolium. Fertility, hatchability, and other reproductive data are not available on amprolium in breeding pheasants. Do not use in feeds containing bentonite.

[41 FR 10985, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.55, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.58 Amprolium and ethopabate.

(a) *Approvals.* Type A medicated articles: (1) 25 percent amprolium and 0.8

percent ethopabate; 25 percent amprolium and 8 percent ethopabate; 5 percent amprolium and 0.16 percent ethopabate; 5 percent amprolium and 1.6 percent ethopabate; to 000006.

(2) 0.15 percent amprolium and 0.004 percent ethopabate and 0.05 gram per pound bacitracin (as bacitracin methylene disalicylate) to 047019 in § 510.600(c) of this chapter.

(b) *Special considerations.* Do not use in Type B or Type C medicated feeds containing bentonite.

(c) *Related tolerances.* See §§ 556.50 and 556.260 of this chapter.

(d) *Conditions of use.* (1) It is used for chickens as follows:

Amprolium and ethopabate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) Amprolium 113.5 (0.0125 pct) and ethopabate 3.6 (0.0004 pct).		Broiler chickens as an aid in the prevention of coccidiosis.	Not for laying hens; as sole source of amprolium.	
(ii) Amprolium 113.5 (0.0125%) and ethopabate 3.6 (0.0004%).	Bambermycins 2 to 3 plus roxarsone 22.8 to 34.1 (0.0025% to 0.00375%).	Broiler chickens; as an aid in the prevention of coccidiosis; for increased rate of weight gain, improved feed efficiency, and pigmentation.	Feed continuously as the sole ration; as sole source of amprolium and organic arsenic; amprolium and ethopabate as provided by No. 050604 in § 510.600(c) of this chapter, roxarsone by No. 046573, bambermycins by No. 012799; withdraw 5 d before slaughter.	
	Lincomycin 2 to 4	Broiler chickens; for increase in rate of weight gain; improved feed efficiency; as an aid in the prevention of coccidiosis.	Not for laying chickens; as lincomycin hydrochloride monohydrate; as sole source of amprolium.	
	Lincomycin 2 to 4 plus roxarsone 45.4 (0.005%).	Broiler chickens; for increase in rate of weight gain; improved feed efficiency and pigmentation; as an aid in the prevention of coccidiosis.	Not for laying chickens; as lincomycin hydrochloride monohydrate; withdraw 5 d before slaughter; as sole source of amprolium and organic arsenic.	
	Roxarsone 45.4 (0.005 pct).	Broiler chickens; to aid in prevention of coccidiosis where severe exposure to coccidiosis from <i>Eimeria acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> is likely to occur; for increased rate of weight gain in broiler chickens raised in floor pens.	Do not feed to laying chickens; withdraw 5 d before slaughter; as sole source of amprolium; do not use as a treatment for outbreaks of coccidiosis; feed as sole ration from time chickens are placed on litter until the past the time when coccidiosis is ordinarily a hazard; roxarsone as provided by No. 046573 in § 510.600(c) of this chapter; combinations as provided by No. 050604.	

Food and Drug Administration, HHS

§ 558.58

Amprolium and ethopabate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(iii) Amprolium 113.5 (0.0125%) and ethopabate 36.3 (0.004%).		Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; as an aid in the prevention of coccidiosis where severe exposure to coccidiosis from <i>Eimeria acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> is likely to occur.	Not for chickens over 16 weeks of age.	046573 and 063238
	Arsanilic acid 90 (0.01 pct) plus erythromycin 4.6 to 18.5.	Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improve pigmentation.	Not for laying hens; withdraw 5 d before slaughter; as sole source of organic arsenic; as erythromycin thiocyanate.	
	Bacitracin 4 to 50	1. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; to aid in prevention of coccidiosis where severe exposure to coccidiosis from <i>Eimeria acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> is likely to occur; for increased rate of weight gain in broiler chickens raised in floor pens.	Not for chickens over 16 weeks of age; do not feed to laying chickens; as sole source of amprolium; not for use as a treatment for outbreaks of coccidiosis; as bacitracin methylene disalicylate as provided by No. 046573 or bacitracin zinc as provided by Nos. 046573 and 063238 in § 510.600(c) of this chapter; feed as the sole ration from the time chickens are placed on litter until past the time when coccidiosis is ordinarily a hazard; combination as provided by No. 050604 in § 510.600(c) of this chapter.	
		2. Broiler chickens; as an aid in prevention of coccidiosis where severe exposure to coccidiosis from <i>Eimeria acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> is likely to occur; improved feed efficiency.	Not for chickens over 16 weeks of age; do not feed to laying chickens; as sole source of amprolium; not for use as a treatment for coccidiosis; bacitracin zinc as provided by Nos. 046573 and 063238 in § 510.600(c) of this chapter; feed as the sole ration from the time chickens are placed on litter until market weight; combination as provided by Nos. 046573 and 063238..	
	Bacitracin 5 to 35 plus roxarsone 34 (0.00375%).	Broiler chickens; for increased rate of weight gain and as an aid in the prevention of coccidiosis where severe exposure to coccidiosis from <i>Eimeria acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> is likely to occur in broiler chickens raised in floor pens.	Do not feed to laying chickens; withdraw 5 d before slaughter; as sole source of amprolium and organic arsenic; do not use as a treatment for outbreaks of coccidiosis; feed as the sole ration from time chickens are placed on litter until past the time when coccidiosis is ordinarily a hazard; amprolium and ethopabate as provided by No. 050604 in § 510.600(c) of this chapter; bacitracin methylene disalicylate as provided by No. 046573 or bacitracin zinc as provided by Nos. 046573 and 063238 in § 510.600(c) of this chapter; roxarsone as provided by No. 046573 in § 510.600(c) of this chapter; combination as provided by No. 050604 in § 510.600(c) of this chapter.	

Amprolium and ethopabate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
	Bacitracin 20 to 35 plus roxarsone 34 (0.00375%).	Broiler chickens; for increased rate of weight gain, improved feed efficiency, and as an aid in the prevention of coccidiosis where severe exposure to coccidiosis from <i>Eimeria acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> is likely to occur in broiler chickens raised in floor pens.	Do not feed to laying chickens; withdraw 5 d before slaughter; as sole source of amprolium and organic arsenic; do not use as a treatment for outbreaks of coccidiosis; feed as the sole ration from time chickens are placed on litter until past the time when coccidiosis is ordinarily a hazard; amprolium and ethopabate as provided by No. 050604 in § 510.600(c) of this chapter; bacitracin methylene disalicylate as provided by No. 046573 in § 510.600(c) of this chapter; roxarsone as provided by No. 046573 in § 510.600(c) of this chapter; combination as provided by No. 050604 in § 510.600(c) of this chapter.	
	Bacitracin 10 to 50 plus roxarsone 15.4 to 45.4 (0.0017% to 0.005%).	Broiler chickens; as an aid in prevention of coccidiosis where severe exposure to coccidiosis from <i>Eimeria acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> is likely to occur; improved feed efficiency.	Do not feed to laying chickens; withdraw 5 d before slaughter; as sole source of amprolium and organic arsenic; do not use as a treatment for outbreaks of coccidiosis; feed as the sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard; amprolium and ethopabate as provided by No. 050604 in § 510.600(c) of this chapter; bacitracin zinc as provided by Nos. 046573 and 063238 roxarsone as provided by No. 046573 combination as provided by Nos. 046573 and 063238.	046573, 063238
	Bacitracin 10 plus roxarsone 30 to 45.4 (0.0033% to 0.005%).	Broiler chickens; as an aid in prevention of coccidiosis where severe exposure to coccidiosis from <i>E. acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> is likely to occur; improved feed efficiency and improved pigmentation.	do	063238
	Bambermycins 1 to 3.	Broiler chickens; as an aid in the prevention of coccidiosis where severe exposure to coccidiosis from <i>Eimeria acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> is likely to occur; for increased rate of weight gain, and improved feed efficiency.	Feed continuously as the sole ration; as sole source of amprolium; amprolium, ethopabate as provided by No. 050604 in § 510.600(c) of this chapter, bambermycins as provided by No. 012799 in § 510.600(c) of this chapter.	
	Bambermycins 1 to 3 plus roxarsone 22.8 to 34.1 (0.0025% to 0.00375%).	Broiler chickens; as an aid in the prevention of coccidiosis where severe exposure to coccidiosis from <i>Eimeria acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> is likely to occur; for increased rate of weight gain; improved feed efficiency, and improved pigmentation.	Feed continuously as the sole ration; as sole source of amprolium and organic arsenic; amprolium and ethopabate as provided by No. 050604 in § 510.600(c) of this chapter, roxarsone by No. 046573 bambermycins by No. 012799. Withdraw 5 days before slaughter.	

Food and Drug Administration, HHS

§ 558.58

Amprolium and ethopabate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(iv) Amprolium 113.5 to 227 (0.0125% to 0.025%) and ethopabate 3.6 (0.0004%).	Erythromycin 4.6 to 18.5.	Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency.	Not for laying hens; withdraw 24 hours before slaughter; erythromycin thiocyanate.	000069
	Virginiamycin, 15	Broiler chickens, as an aid in the prevention of coccidiosis where severe exposure to <i>Eimeria acervulina</i> , <i>E. brunetti</i> , and <i>E. maxima</i> is likely to occur, for increased rate of weight gain and improved feed efficiency.	Feed continuously as sole ration, do not feed to laying hens, not for chickens over 16 weeks of age, as sole source of amprolium, amprolium and ethopabate as provided by 050604 in § 510.600(c), virginiamycin as provided by 000069.	
	Virginiamycin, 5 to 15.	Broiler chickens, as an aid in the prevention of coccidiosis where severe exposure to <i>Eimeria acervulina</i> , <i>E. brunetti</i> , and <i>E. maxima</i> is likely to occur, for increased rate of weight gain.	do	
		For broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis.	Not for laying hens	
	Arsanilic acid 90 (0.01%).	Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation.	As sole source of organic arsenic; withdraw 5 d before slaughter; not for laying hens.	
	Arsanilic acid 90 (0.01%) plus erythromycin 92.5.	1. For broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; as an aid in the prevention of chronic respiratory disease during periods of stress; growth promotion and feed efficiency; improving pigmentation. 2. For broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; as an aid in the prevention of infectious coryza; growth promotion and feed efficiency; improving pigmentation.	Feed for 2 d before stress and 3 to 6 d after stress; withdraw 5 d before slaughter; as sole source of organic arsenic; not for laying hens.	
	Arsanilic acid 90 (0.01%) plus erythromycin 185.	For broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease; growth promotion and feed efficiency; improving pigmentation.	Feed for 7 to 14 d; withdraw 5 d before slaughter; as sole source of organic arsenic; not for laying hens.	
	Bacitracin 4 to 50	For broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency.	As bacitracin methylene disalicylate; not for laying hens.	

Amprolium and ethopabate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
	Bacitracin 100 to 200.	1. Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; treatment of chronic respiratory disease (air-sac infection) and blue comb (nonspecific infectious enteritis). 2. For broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; treatment of chronic respiratory disease (air-sac infection), blue comb (nonspecific infectious enteritis).	do As zinc bacitracin, not for laying hens.	
	Bacitracin 4 to 50 plus roxarsone 22.7 to 45.4 (0.0025% to 0.005%).	Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation.	As bacitracin methylene disalicylate; not for laying hens; as sole source of organic arsenic; withdraw 5 d before slaughter.	
	Chlortetracycline 100 to 200.	For chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption. Feed for 7 to 14 d.	
	Chlortetracycline 200 to 400.	For chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; control of chronic respiratory disease (CRD) and air sac infection caused by <i>M. gallisepticum</i> and <i>E. coli</i> susceptible to chlortetracycline.	In low calcium feed containing 0.8% dietary calcium and 1.5% sodium sulfate; feed continuously as sole ration for 7 to 14 d; do not feed to chickens producing eggs for human consumption.	
	Erythromycin 92.5 ...	1. For broiler chickens and for replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; as an aid in the prevention of chronic respiratory disease during periods of stress. 2. For broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; as an aid in the prevention of infectious coryza.	Feed for 2 d before stress and 3 to 6 d after stress; withdraw 24 h before slaughter; not for laying hens. Feed for 7 to 14 d; withdraw 24 h before slaughter; not for laying hens.	
	Erythromycin 185	For broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease.	Feed for 5 to 8 d; do not use in birds producing eggs for food purposes; withdraw 48 h before slaughter.	
	Penicillin 2.4 to 50 ...	For broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Not for laying hens; as procaine penicillin.	
	Roxarsone 22.7 to 45.4 (0.0025% to 0.005%).	Broiler chickens and replacement chickens where immunity to coccidiosis is not desired; prevention of coccidiosis; growth promotion and feed efficiency; improving pigmentation.	As sole source of organic arsenic; withdraw 5 d before slaughter; not for laying hens.	

Food and Drug Administration, HHS

§ 558.62

Amprolium and ethopabate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(v) Amprolium 136.2 (0.015%) and ethopabate 3.6 (0.0004%).	Bacitracin 10	Broiler chickens; as an aid in the prevention of coccidiosis; growth promotion and feed efficiency.	Feed as sole ration; use as sole source of amprolium; do not feed to laying hens; as bacitracin methylene disalicylate.	047019

(2) [Reserved]

[41 FR 10990, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.58, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.59 Apramycin.

(a) *Approvals.* Type A articles to sponsors identified in § 510.600(c) of this chapter as follows:

(1) 000986 for 75 grams apramycin (as apramycin sulfate) per pound for use as in paragraph (d)(1) of this section.

(2) [Reserved]

(b) [Reserved]

(c) *Related tolerances.* See § 556.52 of this chapter.

(d) *Conditions of use—(1) Swine—(i) Amount.* 150 grams per ton.

(ii) *Indications for use.* For control of porcine colibacillosis (weanling pig scours) caused by susceptible strains of *Escherichia coli*.

(iii) *Limitations.* Use for 14 days. Withdraw 28 days before slaughter.

(2) [Reserved]

[51 FR 9190, Mar. 18, 1986]

§ 558.60 Arsanilate sodium.

(a) *Approvals.* Type A medicated articles: 20, 50, or 100 percent to 015565 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.60 of this chapter.

(c) *Conditions of use.* (1) It is used for chickens and turkeys as follows:

(i) *Grams per ton.* 90 (0.01 percent).

(ii) *Indications for use.* For growth promotion and feed efficiency; improving pigmentation.

(iii) *Limitations.* Withdraw 5 days before slaughter; as sole source of organic arsenic.

(2) Arsanilate sodium may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Amprolium in accordance with § 558.55.

(ii) Zoalene in accordance with § 558.680.

[41 FR 10992, Mar. 15, 1976, as amended at 51 FR 7395, Mar. 3, 1986; 51 FR 33897, Sept. 24, 1986]

§ 558.62 Arsanilic acid.

(a) *Approvals.* Type A medicated articles to sponsors in § 510.600(c) of this chapter as follows:

(1) To 015565: 20, 50, and 100 percent for use as in the table in paragraph (c)(1), entry (ii), item 1; entry (ii), item 2; entry (iii); entry (iv); and entry (v) of this section.

(2) To 015565: 20 percent for use as in paragraph (c)(1), entry (i); entry (ii), item 3 of this section.

(b) *Related tolerances.* See § 556.60 of this chapter.

(c) *Conditions of use.* (1) It is used as follows:

Arsanilic acid in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 45 to 90 (0.005% to 0.01%).		Swine; for increased rate of weight gain and improved feed efficiency in growing swine..	Withdraw 5 days before slaughter; as sole source of organic arsenic.	015565
(ii) 90 (0.01%)		1. Chickens; growth promotion and feed efficiency; improving pigmentation..	do	015565
		2. Turkeys; growth promotion and feed efficiency; improving pigmentation.	do	015565

§ 558.76

21 CFR Ch. I (4–1–99 Edition)

Arsanilic acid in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(iii)	Erythromycin 4.6 to 18.5.	3. Swine; and aid in control of swine dysentery (hemorrhagic enteritis, bloody dysentery). Chickens; growth promotion and feed efficiency; improving pigmentation.	do	015565
(iv)	Erythromycin 92.5 ...	1. Chickens; as an aid in the prevention of chronic respiratory disease during periods of stress; growth promotion and feed efficiency; improving pigmentation. 2. Chickens; as an aid in the prevention of infectious coryza; growth promotion and feed efficiency; improving pigmentation.	As erythromycin thiocyanate; withdraw 5 days before slaughter; as sole source of organic arsenic.	15565
(v)	Erythromycin 185 ...	Chickens; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease; growth promotion and feed efficiency; improving pigmentation.	As erythromycin thiocyanate; feed for 2 days before stress and 3 to 6 days after stress; withdraw 5 days before slaughter; as sole source of organic arsenic.	015565
			As erythromycin thiocyanate; feed for 7 to 14 days; withdraw 5 days before slaughter; as sole source of organic arsenic.	015565
			As erythromycin thiocyanate; feed for 5 to 8 days; do not use in birds producing eggs for food purposes; withdraw 5 days before slaughter; as sole source of organic arsenic.	15565

(2) Arsanilic acid may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Amprolium in accordance with § 558.55.

(ii) Amprolium and ethopabate in accordance with § 558.58.

(iii) Bacitracin zinc in accordance with § 558.78.

(iv) Bacitracin and zoalene in accordance with § 558.680.

(v) Zualene in accordance with § 558.680.

[41 FR 10992, Mar. 15, 1976, as amended at 42 FR 18617, Apr. 8, 1977; 51 FR 7395, Mar. 3, 1986; 51 FR 33897, Sept. 24, 1986; 54 FR 18281, Apr. 28, 1989; 56 FR 19268, Apr. 26, 1991; 60 FR 39847, Aug. 4, 1995]

§ 558.76 Bacitracin methylene disalicylate.

(a) *Approvals.* Type A medicated articles: 10, 25, 30, 40, 50, 60, or 75 grams per pound to 046573 in § 510.600(c) of this chapter.

(b) *Special considerations.* The quantities of antibiotics are expressed in terms of the equivalent amount of antibiotic standard.

(c) *Related tolerances.* See § 556.70 of this chapter.

(d) *Conditions of use.* (1) It is used as follows:

Bacitracin methylene disalicylate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 4 to 50		Chickens, turkeys, and pheasants; increased rate of weight gain and improved feed efficiency ¹ .		046573
(ii) 5 to 20		Quail not over 5 weeks of age; increased rate of weight gain and improved feed efficiency ¹ .		046573
(iii) 10 to 25		Chickens; for increased egg production and improved feed efficiency for egg production.	For first 7 months of production	046573
(iv) 10 to 30		Swine; for increased rate of weight gain and improved feed efficiency.	For growing and finishing swine	000004 and 046573

Food and Drug Administration, HHS

§ 558.76

Bacitracin methylene disalicylate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(v) [Reserved]	Chlortetracycline approximately 400, varying with body weight and food consumption to provide 10 milligrams per pound of body weight per day.	Swine; for increased rate of weight gain and improved feed efficiency; for treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline.	Feed for not more than 14 days to provide 10 milligrams of chlortetracycline per pound of body weight per day; as chlortetracycline provided by Nos. 000004 and 046573 in § 510.600(c) of this chapter. Type C feed may be prepared from Type B feed containing 1 to 3 grams per pound BMD with 400 grams per pound CTC, to 046573 in § 510.600(c).	000004 and 046573
(vi) 50		Broiler chickens; as an aid in the prevention of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin.		046573
(vii)—(viii) [Reserved]		Replacement chickens; as an aid in the prevention of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin	Feed continuously as sole ration	046573
(ix) 100 to 200		Broiler chickens; as an aid in the control of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin.		046573
		Replacement chickens; as an aid in the control of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin	Feed continuously as sole ration. Start at first clinical signs of disease, vary dosage based on severity of infection, administer continuously for 5 to 7 days or as long as clinical signs persist, then reduce medication to prevention level (50 g/t).	046573
(x) 200		Turkeys; as an aid in the control of transmissible enteritis in growing turkeys complicated by organisms susceptible to bacitracin methylene disalicylate.		046573
		Quail; for the prevention of ulcerative enteritis in growing quail due to <i>Clostridium colinum</i> susceptible to bacitracin methylene disalicylate	From Type A medicated articles containing 25, 40, or 50 grams of bacitracin methylene disalicylate. Feed continuously as the sole ration.	046573
(xi) 250		1. Growing/Finishing Swine: For control of swine dysentery associated with <i>Treponema hyodysenteriae</i> on premises with a history of swine dysentery but where signs of the disease have not yet occurred; or following an approved treatment of the disease condition.	As the sole ration. Not for use in swine weighing more than 250 pounds. Diagnosis should be confirmed by a veterinarian when results are not satisfactory.	046573

Bacitracin methylene disalicylate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
		2. Pregnant sows: For control of clostridial enteritis caused by <i>C. perfringens</i> in suckling piglets..	As the sole ration. Feed to sows from 14 days before through 21 days after farrowing on premises with a history of clostridial scours. Diagnosis should be confirmed by a veterinarian when results are not satisfactory.	

¹ These conditions are NAS/NRC reviewed and found effective. Applications for these uses may not require effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(2) It is used as bacitracin methylene disalicylate in feed for animals as follows:

(i) *Amount.* 70 milligrams per head per day.

(a) *Indications for use.* Feedlot beef cattle; reduction in the number of liver condemnations due to abscesses.

(b) *Limitations.* Administer continuously throughout the feeding period.

(ii) *Amount.* 250 milligrams per head per day.

(a) *Indications for use.* Feedlot beef cattle; reduction in the number of liver condemnations due to abscesses.

(b) *Limitations.* Administer continuously for 5 days then discontinue for subsequent 25 days, repeat the pattern during the feeding period.

(3) It is used as bacitracin methylene disalicylate in accordance with the provisions of this section in the combinations provided as follows:

(i) Amprolium in accordance with § 558.55.

(ii) Amprolium with ethopabate in accordance with § 558.58.

(iii) Arsanilic acid with zoalene in accordance with § 558.680

(iv) Carbarsone (not U.S.P.) in accordance with § 558.120.

(v) Nicarbazin as in § 558.366.

(vi) Hygromycin B in accordance with § 588.274.

(vii) Monensin in accordance with § 588.355.

(viii) Lasalocid sodium alone or with roxarsone as in § 558.311.

(ix) Monensin and roxarsone in accordance with § 558.355.

(x) Salinomycin alone or with roxarsone as in § 558.550.

(xi) Halofuginone hydrobromide and roxarsone in accordance with § 558.265.

(xii) Halofuginone in accordance with § 558.265.

(xiii) Narasin alone or in combination with roxarsone as in § 558.363.

(xiv) Semduramicin alone or in combination with roxarsone as in § 558.555.

(xv) Zoalene alone or in combination as in § 558.680.

(xvi) Nitarsone alone or in combination as in § 558.369.

(xvii) Decoquinat and roxarsone as in § 558.195.

(xviii) Decoquinat as in § 558.195.

(xix) Narasin and nicarbazin as in § 558.366.

(xx) Ivermectin as in § 558.300.

[41 FR 10993, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.76, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.78 Bacitracin zinc.

(a) *Approvals.* To sponsors listed in § 510.600(c) of this chapter for use as in paragraph (d) of this section as follows:

(1) To 046573: 50 grams per pound as in paragraphs (d)(1)(i), (d)(1)(ii), (d)(1)(iii), (d)(1)(iv), and (d)(2) of this section.

(2) To 063238: 10, 25, 40, and 50 grams per pound as in paragraphs (d)(1)(i), (d)(1)(ii), (d)(1)(v), (d)(1)(vi), (d)(2), and (d)(3) of this section.

(3) To 000010: 5 and 50 grams per pound for chickens as in paragraph (d)(1)(i) of this section.

(b) *Special considerations.* The quantities of antibiotics are expressed in terms of the equivalent amount of antibiotic standard.

(c) *Related tolerances.* See § 556.70 of this chapter.

(d) *Conditions of use.* (1) It is used as follows:

Food and Drug Administration, HHS

§ 558.95

Bacitracin zinc in grams per ton	Combinations in grams per ton	Indications for use	Limitations	Sponsor
(i) 4 to 50		Chickens, turkeys, and pheasants; for increased rate of weight gain and improved feed efficiency.	Growing chickens, turkeys, and pheasants.	063238, 046573
(ii) 5 to 20		Quail; for increased rate of weight gain and improved feed efficiency.	Growing quail; feed as the Type C feed to starting quail through 5 weeks of age.	063238, 046573
(iii) 10 to 25		Laying chickens; improved feed efficiency and increased egg production.		046573
(iv) 10 to 50		Swine; increased rate of weight gain and improved feed efficiency.	Growing and finishing swine	046573
(v) 20		Growing-finishing swine; increased rate of weight gain.	In Type C feed	063238
(vi) 20 to 40		Growing-finishing swine; improved feed efficiency.	do	063238

(2) It is used in feed for growing cattle at 35 to 70 milligrams per head per day as follows:

(i) To aid in stimulating growth and improving feed efficiency.

(ii) For increased rate of weight gain and improved feed efficiency; see sponsor 063238.

(3) Bacitracin zinc may be used as approved in combination as follows:

(i) Amprolium as in § 558.55.

(ii) Amprolium and roxarsone as in § 558.55.

(iii) Amprolium and ethopabate as in § 558.58.

(iv) Amprolium and ethopabate with roxarsone as in § 558.58.

(v) Carbarsone as in § 558.120.

(vi) Clopidol as in § 558.175.

(vii) Clopidol and roxarsone as in § 558.175.

(viii) Decoquinate as in § 558.195.

(ix) Decoquinate and roxarsone as in § 558.195.

(x) Hygromycin B as in § 558.274.

(xi) Hygromycin B and penicillin as in § 558.274.

(xii) Lasalocid sodium and roxarsone as in § 558.311.

(xiii) Monensin as in § 558.355.

(xiv) Monensin and roxarsone as in § 558.355.

(xv) Robenidine as in § 558.515.

(xvi) Salinomycin as in § 558.550.

(xvii) Salinomycin and roxarsone as in § 558.550.

(xviii) Zoalene as in § 558.680.

(xix) Zoalene and arsanilic acid as in § 558.680.

(xx) Zoalene and roxarsone as in § 558.680.

[41 FR 10994, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.78, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.95 Bambermycins.

(a) *Approvals.* To sponsors identified by drug labeler codes in § 510.600(c) of this chapter for use of bambermycins Type A medicated articles as bambermycins activity per pound in paragraph (d) of this section as follows:

(1) To 012799: 2, 4, and 10 grams for use as in paragraphs (d)(1), (d)(2), (d)(3), and (d)(4) of this section.

(2) To 012799: 0.4 gram for use as in paragraph (d)(2) of this section.

(3) To 011490: 0.4 and 2 grams for use as in paragraph (d)(2) of this section.

(4) To 012286, 016968, and 017790: 0.4 and 2 grams for use as in paragraph (d)(2) and 2 grams for use as in paragraph (d)(3) of this section.

(5) To 012799: 10 grams to make 40 to 800 grams per ton Type B feed for use as in paragraph (d)(4) of this section.

(b) *Special considerations.* (1) Bambermycins liquid Type B feeds may be manufactured from dry bambermycins Type A articles. The liquid Type B feeds must have a pH of 3.8 to 7.5, moisture content of 30 to 45 percent.

(2) The expiration date for the liquid Type B feed is 8 weeks after date of manufacture. The expiration date for

the dry Type C feed made from the liquid Type B feed is 1 week after date of manufacture.

(c) [Reserved]

(d) *Conditions of use*—(1) *Broiler chickens*. It is used as follows:

(i) *Amount per ton*. 1 to 2 grams.

(a) *Indications for use*. For increased rate of weight gain and improved feed efficiency.

(b) *Limitations*. Feed continuously as the sole ration.

(ii) *Amount per ton*. Bambermycins, 1 to 3 grams plus amprolium, 113.5 grams (.0125 percent) plus ethopabate, 36.3 grams (.004 percent).

(a) *Indications for use*. As an aid in the prevention of coccidiosis where severe exposure to coccidiosis from *E. acervulina*, *E. maxima*, and *E. brunetti* is likely to occur. For increased rate of weight gain and improved feed efficiency.

(b) *Limitations*. Feed continuously as the sole ration; as sole source of amprolium; amprolium and ethopabate as provided by No. 050604 in § 510.600(c) of this chapter.

(iii) *Amount per ton*. Bambermycins, 1 to 3 grams plus amprolium, 113.5 grams (.0125 percent) plus ethopabate, 36.3 grams (.004 percent) plus roxarsone, 22.8 to 34.1 grams (.0025–.00375 percent).

(a) *Indications for use*. As an aid in the prevention of coccidiosis where severe exposure to coccidiosis from *E. acervulina*, *E. maxima*, and *E. brunetti* is likely to occur. For increased rate of weight gain, improved feed efficiency, and improved pigmentation.

(b) *Limitations*. Feed continuously as the sole ration; as sole source of amprolium and organic arsenic; amprolium and ethopabate as provided by No. 050604 in § 510.600(c) of this chapter, roxarsone as provided by No. 046573 in § 510.600(c) of this chapter. Withdraw 5 days before slaughter.

(iv) *Amount per ton*. Bambermycins, 1 to 3 grams plus amprolium, 113.5 grams (.0125 percent) plus ethopabate, 3.63 grams (.004 percent) plus roxarsone, 22.8 to 34.1 grams (.0025–.00375 percent).

(a) *Indications for use*. As an aid in the prevention of coccidiosis. For increased rate of weight gain, improved feed efficiency, and improved pigmentation.

(b) *Limitations*. Feed continuously as the sole ration; as sole source of amprolium and organic arsenic; amprolium and ethopabate as provided by No. 050604 in § 510.600(c) of this chapter, roxarsone as provided by No. 046573 in § 510.600(c) of this chapter. Withdraw 5 days before slaughter.

(v) *Amount per ton*. Bambermycins, 1 to 3 grams plus amprolium, 113.5 grams (.0125 percent) plus roxarsone, 22.8 to 34.1 grams (.0025–.00375 percent).

(a) *Indications for use*. As an aid in the prevention of coccidiosis. For increased rate of weight gain, improved feed efficiency, and improved pigmentation.

(b) *Limitations*. Feed continuously as the sole ration; as sole source of amprolium and organic arsenic; amprolium as provided by No. 050604 in § 510.600(c) of this chapter, roxarsone as provided by No. 046573 in § 510.600(c) of this chapter. Withdraw 5 days before slaughter.

(vi) *Amount per ton*. Bambermycins, 1 to 2 grams plus monensin, 90 to 110 grams.

(a) *Indications for use*. For increased rate of weight gain and improved feed efficiency; as an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(b) *Limitations*. Do not feed to laying chickens; feed continuously as sole ration; as monensin sodium provided by No. 000986 in § 510.600(c) of this chapter; as bambermycins provided by No. 012799 in § 510.600(c) of this chapter.

(vii) *Amount per ton*. Bambermycins, 1 gram plus monensin, 90 to 110 grams plus roxarsone, 22.7 to 45.4 grams (.0025 to .005 percent).

(a) *Indications for use*. For increased rate of weight gain and improved feed efficiency; as an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(b) *Limitations*. Do not feed to laying chickens; feed continuously as sole ration; use as sole source of organic arsenic; withdraw 5 days before slaughter; as monensin sodium provided by No. 000986 in § 510.600(c) of this chapter; as bambermycins provided by No. 012799 in § 510.600(c) of this chapter; as

roxarsone provided by No. 046573 in § 510.600(c) of this chapter.

(viii) *Amount per ton.* Bambermycins, 1 gram plus zoalene, 113.4 grams (0.0125 percent).

(a) *Indications for use.* As an aid in the prevention and control of coccidiosis; for increased rate of weight gain and improved feed efficiency.

(b) *Limitations.* Do not feed to chickens over 14 weeks of age; feed continuously as sole ration; zoalene as provided by No. 025700 in § 510.600(c) of this chapter.

(ix) *Amount per ton.* Bambermycins, 1 gram plus zoalene, 113.4 grams (0.0125 percent) plus roxarsone, 22.7 grams (0.0025 percent).

(a) *Indications for use.* As an aid in the prevention and control of coccidiosis; for increased rate of weight gain and improved feed efficiency.

(b) *Limitations.* Do not feed to chickens over 14 weeks of age; feed continuously as sole ration; feed as sole source of organic arsenic; withdraw 5 days before slaughter; zoalene as provided by No. 025700, roxarsone as provided by No. 046573 in § 510.600(c) of this chapter.

(x) *Amount per ton.* Bambermycins, 1 gram plus lasalocid sodium, 68 to 113 grams (0.0075 to 0.0125 percent) plus roxarsone, 45.4 grams (0.005 percent).

(a) *Indications for use.* For prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*; as an aid in the reduction of lesions due to *E. tenella*; and for increased rate of weight gain.

(b) *Limitations.* For broiler chickens only; feed continuously as sole ration; withdraw 5 days before slaughter; roxarsone as provided by Nos. 046573 and 011526 in § 510.600(c) of this chapter, and lasalocid sodium as provided by . 000004 in § 510.600(c) of this chapter.

(xi) *Amount per ton.* Bambermycins 1 to 2 grams, plus roxarsone 45.4 grams, and salinomycin 40 to 60 grams.

(a) *Indications for use.* For prevention of coccidiosis caused by *Eimeria necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, *E. tenella*, and *E. mivati*, including some field strains of *E. tenella* that are more susceptible to roxarsone combined with salinomycin than salinomycin alone; and for improved feed efficiency.

(b) *Limitations.* For broiler chickens only; do not feed to laying chickens; feed continuously as sole ration; as sole source or organic arsenic; withdraw 5 days before slaughter; not approved for use with pellet binders; may be fatal if accidentally fed to adult turkeys or horses; as roxarsone provided by No. 011526 or 046573 in § 510.600(c) of this chapter; as salinomycin sodium biomass provided by Nos. 012799 and 063238 in § 510.600(c) of this chapter.

(xii) *Amount per ton.* Bambermycins 1 to 3 grams, plus salinomycin 40 to 60 grams.

(a) *Indications for use.* For prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*; and improved feed efficiency.

(b) *Limitations.* For broiler chickens only; Do not feed to laying chickens; feed continuously as sole ration; not approved for use with pellet binders; may be fatal if accidentally fed to adult turkeys or horses; as salinomycin sodium biomass provided by Nos. 012799 and 063238 in § 510.600(c) of this chapter.

(xiii) *Amount per ton.* Bambermycins, 1 to 2 grams plus monensin, 90 to 110 grams plus roxarsone, 22.7 to 45.4 grams (0.0025 to 0.005 percent).

(a) *Indications for use.* For increased rate of weight gain; as an aid in prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(b) *Limitations.* See paragraph (b)(1)(vii)(b) of this section.

(2) *Growing-finishing swine.* It is used as follows:

(i) *Amount per ton.* 2 grams.

(a) *Indications for use.* For increased rate of weight gain and improved feed efficiency.

(b) *Limitations.* Feed continuously as sole ration.

(ii) *Amount per ton.* 2 to 4 grams.

(a) *Indications for use.* For increased rate of weight gain.

(b) *Limitations.* Feed continuously as sole ration.

(3) *Growing turkeys.* It is used as follows:

(i) *Amount per ton.* 1 to 2 grams.

(a) *Indications for use.* For improved feed efficiency.

(b) *Limitations.* Feed continuously as the sole ration.

§ 558.95

21 CFR Ch. I (4–1–99 Edition)

(ii) *Amount per ton.* 2 grams.

(a) *Indications for use.* For increased rate of weight gain and improved feed efficiency.

(b) *Limitations.* Feed continuously as sole ration.

(iii) *Amount per ton.* Bambermycins, 1 to 4 grams plus amprolium, 113.5 grams (.0125 percent).

(a) *Indications for use.* For prevention of coccidiosis; for increased rate of weight gain and improved feed efficiency.

(b) *Limitations.* Feed continuously as the sole source of amprolium; amprolium as provided by No. 050604 in § 510.600(c) of this chapter.

(iv) *Amount per ton.* Bambermycins, 1 or 4 grams plus carbarsone, 227 grams (0.025 percent).

(a) *Indications for use.* For improved feed efficiency (1 gram per ton) or increased rate of weight gain (4 grams per ton); as an aid in the prevention of blackhead.

(b) *Limitations.* Feed continuously 2 weeks before blackhead is expected and continue as long as prevention is needed; withdraw 5 days before slaughter; use as sole source of organic arsenic; carbarsone by 046573 in § 510.600(c) of this chapter.

(4) *Cattle*—(i) *Amount per ton.* 1 to 4 grams.

(a) *Indications for use.* For increased rate of weight gain and improved feed efficiency.

(b) *Limitations.* Feed only to cattle being fed in confinement for slaughter. Feed continuously in a Type C medicated feed at a rate of 10 to 20 milligrams of bambermycins per head per day. Liquid Type B feeds containing bambermycins may be used in the preparation of dry complete ration Type C feeds.

(ii) *Amount per ton.* 4 to 20 grams.

(a) *Indications for use.* For increased rate of weight gain.

(b) *Limitations.* Feed continuously to pasture cattle (slaughter, stocker, and feeder cattle, and dairy and beef replacement heifers) at a rate of 10 to 20 milligrams of bambermycins per head per day in at least 1 pound and not more than 10 pounds of Type C medicated feed.

(iii) Used as a free-choice Type C medicated loose mineral feed for pasture cattle (slaughter, stocker, and feeder cattle, and dairy and beef replacement heifers) as follows:

(a) *Specifications.*

Ingredient	International Feed No.	Percent
Deflorinated phosphate (20.5% calcium, 18.5% phosphorus)	6–01–080	42.50
Sodium chloride (salt)	6–04–152	20.10
Calcium carbonate (38% calcium)	6–01–069	15.24
Corn distillers dried grains w/solubles	5–28–236	9.57
Magnesium oxide	6–02–756	5.15
Vitamin and trace mineral premix *		3.72
Mineral oil		1.00
Yeast (primary dehydrated yeast)	7–05–533	0.75
Bambermycins Type A article (10 g/lb)		0.60
Iron oxide	6–02–431	0.50
Magnesium sulfate (67%)	6–02–758	0.32
Selenium premix (270 mg/lb) *		0.21
Copper sulfate	6–01–720	0.18
Potassium sulfate (0.33%)	6–06–098	0.16

*Content of vitamin/trace mineral premix may be varied. However, they should be comparable to those used for other free-choice feeds. Formulation modifications require FDA approval prior to marketing. Selenium must comply with 21 CFR 573.920. Ethylenediamine dihydroiodide (EDDI) should comply with FDA Compliance Policy Guides Sec. 651.100 (CPG 7125.18).

(b) *Amount per ton.* 120 grams.

(c) *Indications for use.* For increased rate of weight gain.

(d) *Limitations.* For free-choice feeding to pasture cattle (slaughter, stocker, and feeder). Feed a nonmedicated commercial mineral product for 6 weeks to stabilize consumption between 2.66 and 5.33 ounces per head per

day. Feed continuously to provide 10- to 20-milligrams bambermycins per head per day. Each use of this free-choice Type C medicated feed must be the subject of an approved Form FDA 1900 as required by 21 CFR 510.455.

(iv) Use free-choice Type C medicated feeds for pasture cattle (slaughter, stocker, and feeder cattle, and dairy

and beef replacement heifers) as follows:

(a) *Amount*. Feed continuously to provide 10 to 20 milligrams of bambermycins per head per day.

(b) *Indications for use*. For increased rate of weight gain.

(c) *Limitations*. Each use in a free-choice Type C medicated feed must be the subject of an approved new animal drug application (NADA) or supplemental NADA as required by 21 CFR 510.455.

(5) Bambermycins may be used as in this section in combination with:

(i) Halofuginone as in § 558.265.

(ii) Narasin as in § 558.363.

(iii) Narasin and roxarsone as in § 558.363.

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.95, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.105 [Reserved]

§ 558.115 Carbadox.

(a) *Approvals*. Type A medicated articles: 2.2 percent (10 grams per pound) to 000069 in § 510.600(c) of this chapter.

(b) *Related tolerances*. See § 556.100 of this chapter.

(c) *Special considerations*. Do not use in Type B or Type C medicated feeds containing bentonite.

(d) *Conditions of use*. It is used for swine as follows:

(1) *Amount per ton*. 10–25 grams (0.0011–0.00275 percent).

(i) *Indications for use*. For increase in rate of weight gain and improvement of feed efficiency.

(ii) *Limitations*. Not for use in pregnant swine or swine intended for breeding purposes. Do not feed to swine within 42 days of slaughter.

(2) *Amount per ton*. 50 grams (0.0055 percent).

(i) *Indications for use*. For control of swine dysentery (vibronic dysentery, bloody scours, or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis*); increased rate of weight gain and improved feed efficiency.

(ii) *Limitations*. Not for use in pregnant swine or swine intended for breed-

ing purposes. Do not feed to swine within 42 days of slaughter.

(3) *Amount per ton*. Carbadox 50 grams (0.0055 percent) plus pyrantel tartrate, 96 grams (0.0106 percent).

(i) *Indications for use*. For control of swine dysentery (vibronic dysentery, bloody scours, or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis*); aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections.

(ii) *Limitations*. Do not feed to swine over 75 pounds; do not feed within 10 weeks of slaughter; consult a veterinarian before feeding to severely debilitated animals; feed continuously as sole ration. Do not use in complete feeds containing less than 15 percent crude protein.

[40 FR 13959, Mar. 27, 1975, as amended at 40 FR 45164, Oct. 1, 1975; 40 FR 57798, Dec. 12, 1975; 42 FR 761, Jan. 4, 1977; 51 FR 7396, Mar. 3, 1986; 63 FR 59216, Nov. 3, 1998]

§ 558.120 Carbarsone (not U.S.P.).

(a) *Approvals*. Type A medicated articles: (1) 37.5 percent to 046573 in § 510.600(c) of this chapter.

(2) 25 percent carbarsone and 5 grams per pound bacitracin (as bacitracin methylene disalicylate) to 046573 in § 510.600(c) of this chapter.

(b) *Related tolerances*. See § 556.60 of this chapter.

(c) [Reserved]

(d) *Conditions of use*. (1) It is used for turkeys as follows:

(i) *Grams per ton*. 227 to 340.5 (0.025 to 0.0375 percent).

(a) *Indications for use*. As an aid in the prevention of blackhead.

(b) *Limitations*. Feed continuously beginning 2 weeks before blackhead is expected and continue as long as prevention is needed; withdraw 5 days before slaughter; as sole source of organic arsenic.

(ii) *Grams per ton*. 227 to 340.5 (0.025 to 0.0375 percent) carbarsone plus 10 grams per ton bacitracin from bacitracin methylene disalicylate.

(a) *Indications for use*. As an aid in the prevention of blackhead; for increased rate of weight gain.

§ 558.128

21 CFR Ch. I (4–1–99 Edition)

(b) *Limitations.* Feed continuously beginning 2 weeks before blackhead is expected and continue as long as prevention is needed; withdraw 5 days before slaughter; as sole source of organic arsenic.

(iii) *Grams per ton.* 227 to 340.5 (0.025 to 0.0375 percent) carbarsone plus 4 to 45 grams per ton bacitracin from bacitracin zinc.

(a) *Indications for use.* As an aid in the prevention of blackhead, increased rate of weight gain, and improved feed efficiency.

(b) *Limitations.* Feed continuously as sole ration. Withdraw 5 days before slaughter. As sole source of organic arsenic; as bacitracin zinc provided by Nos. 046573 and 063238 in §510.600(c) of this chapter.

(2) Carbarsone (not U.S.P.) may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Zoalene in accordance with §558.680.

(ii) Amprolium as in §558.55.

(iii) Bambermycins as in §558.95.

[41 FR 10995, Mar. 15, 1976, as amended at 42 FR 18617, Apr. 8, 1977; 46 FR 46797, Sept. 22, 1981; 48 FR 2758, Jan. 21, 1983; 51 FR 7396, Mar. 3, 1986; 52 FR 2687, Jan. 26, 1987; 53 FR 20843, June 7, 1988; 57 FR 7652, Mar. 4, 1992; 61 FR 515, Jan. 8, 1996; 61 FR 18082, Apr. 24, 1996; 62 FR 61011, Nov. 14, 1997; 63 FR 27845, May 21, 1998]

§ 558.128 Chlortetracycline.

(a) *Approvals.* Type A medicated articles containing the following concentrations of either chlortetracycline calcium complex equivalent to chlortetracycline hydrochloride or, for products intended for use in milk replacer, chlortetracycline hydrochloride:

(1) 50 to 100 grams per pound to 063238 in §510.600(c) of this chapter.

(2) 50 to 100 grams per pound to 000069.

(3) 50 to 100 grams per pound to 046573.

(4) 50 grams per pound to 017519.

(5) 50 to 100 grams per pound to 053389.

(b) *Related tolerances.* See §556.150 of this chapter.

(c) [Reserved]

(d)(1) It is used in feeds as follows:

Chlortetracycline amount	Combination	Indications for use	Limitations	Sponsor
(i) 10 to 50 g/t		1. Chickens; increased rate of weight gain and improved feed efficiency.	Do not feed to chickens producing eggs for human consumption.	063238. 000069, 017519, 046573, 053389.
		2. Growing turkeys; increased rate of weight gain and improved feed efficiency.	Do not feed to turkeys producing eggs for human consumption.	000069, 017519, 046573, 053389, 063238. Do.
		3. Growing swine; increased rate of weight gain and improved feed efficiency.		
(ii) 20 to 50 g/t		Growing sheep; increased rate of weight gain and improved feed efficiency.		000069, 046573, 053389, 063238.
(iii) 50 to 100 g/t		Swine; reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E. <i>Streptococci</i> susceptible to chlortetracycline.		000069, 017519, 046573, 053389, 063238.
(iv) 100 to 200 g/t		Chickens; control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline.	1. Feed continuously for 7 to 14 d. 2. Feed continuously for 7 to 14 d; do not feed to chickens producing eggs for human consumption.	063238. 000069, 017519, 046573, 053389.

Food and Drug Administration, HHS

§ 558.128

Chlortetracycline amount	Combination	Indications for use	Limitations	Sponsor
(v) 200 g/t		Turkeys; control of infectious synovitis caused by <i>M. synoviae</i> susceptible to chlortetracycline.	Feed continuously for 7 to 14 d; do not feed to turkeys producing eggs for human consumption.	000069, 017519, 046573, 053389, 063238.
(vi) 200 to 400 g/t		1. Chickens; control of chronic respiratory disease (CRD) and air sac infection caused by <i>M. gallisepticum</i> and <i>E. coli</i> susceptible to chlortetracycline. 2. Ducks; control and treatment of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline.	1. Feed continuously for 7 to 14 d. 2. Feed continuously for 7 to 14 d; do not feed to chickens producing eggs for human consumption. Feed in complete ration to provide from 8 to 28 milligrams per pound of body weight per day depending upon age and severity of disease, for not more than 21 d. Do not feed to ducks producing eggs for human consumption.	000069, 017519, 046573, 053389, 063238.
(vii) 400 g/t		1. Turkeys; control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to chlortetracycline. 2. Turkey poults not over 4 weeks of age; reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline. 3. Breeding swine; control of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline.	Feed continuously for 7 to 14 d; do not feed to turkeys producing eggs for human consumption. Feed continuously for not more than 14 d.	000069, 017519, 046573, 053389, 063238. Do. Do.
(viii) 500 g/t		Chickens; reduction of mortality due to <i>E. coli</i> infections susceptible to chlortetracycline.	1. Feed for 5 d. 2. Feed for 5 d; do not feed to chickens producing eggs for human consumption.	063238. 000069, 017519, 046573, 053389.
(ix) 10 mg/g of finished feed daily.		Psittacine birds (cockatoos, macaws, and parrots) suspected or known to be infected with psittacosis caused by <i>Chlamydia psittaci</i> sensitive to chlortetracycline.	Feed continuously for 45 d; each bird should consume daily an amount of medicated feed equal to one fifth of its body weight. <i>Warning:</i> "Psittacosis, avian chlamydiosis, or ornithosis is a reportable communicable disease, transmissible between wild and domestic birds, other animals, and man. Contact appropriate public health and regulatory officials."	00004
(x) 0.1 mg/lb of body weight daily.		Calves (up to 250 lb); for increased rate of weight gain and improved feed efficiency.	In milk replacers or starter feed; include on labeling the warning: "A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal."	063238, 000069, 017519, 046573, 053389

Chlortetracycline amount	Combination	Indications for use	Limitations	Sponsor
(xi) 0.5 mg/lb of body weight daily.		Beef cattle (over 700 lb); control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.	Withdraw 48 h prior to slaughter. For sponsor 000004 zero withdrawal time.	Do.
(xii) 10 mg/lb of body weight		1. Calves, beef and nonlactating dairy cattle; treatment of bacterial enteritis caused by <i>E. coli</i> and bacterial pneumonia caused by <i>P. multocida</i> organisms susceptible to chlortetracycline. 2. Calves (up to 250 lb); treatment of bacterial enteritis caused by <i>E. coli</i> susceptible to chlortetracycline. 3. Swine; treatment of bacterial enteritis caused by <i>E. coli</i> and <i>S. choleraesuis</i> and bacterial pneumonia caused by <i>P. multocida</i> susceptible to chlortetracycline.	Feed approximately 400 g/t, varying with body weight and feed consumption to provide 10 mg/lb per day. Treat for not more than 5 d; in feed including milk replacers; withdraw 10 d prior to slaughter except for 24 h for sponsor 046573; zero withdrawal for sponsor 000004; include on labeling the warning: "A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal." In milk replacers or starter feed; include on labeling the warning: "A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal." Feed approximately 400 g/t, varying with body weight and feed consumption to provide 10 mg/lb per day. Feed for not more than 14 d; withdraw 5 d prior to slaughter for sponsor 012286.	Do. 063238, 000069, 017519, 046573, 053389
(xiii) 25 mg/lb of body weight		Turkeys; control of complicating bacterial organisms associated with bluecomb (transmissible enteritis; coronaviral enteritis) susceptible to chlortetracycline.	Feed continuously for 7 to 14 d; do not feed to turkeys producing eggs for human consumption.	Do.
(xiv) 25 to 70 mg/head/day		Calves (250 to 400 lb); increased rate of weight gain and improved feed efficiency.	Include on labeling the warning: "A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal."	063238, 000069, 017519, 046573, 053389
(xv) 70 mg/head/day		Growing cattle (over 400 lb) increased rate of weight gain, improved feed efficiency, and reduction of liver condemnation due to liver abscesses.	do	Do.
(xvi) 80 mg/head/day		Breeding sheep; reducing the incidence of (vibriotic) abortion caused by <i>Campylobacter fetus</i> infection susceptible to chlortetracycline.		063238, 000069, 046573, 053389
(xvii) 350 mg/head/day		1. Beef cattle; control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline.	Withdraw 48 h prior to slaughter. For sponsor 000004 zero withdrawal time.	063238, 000069, 017519, 046573, 053389

Chlortetracycline amount	Combination	Indications for use	Limitations	Sponsor
		2. Beef cattle (under 700 lb); control of active infection of anaplasmosis caused by <i>A. marginale</i> susceptible to chlortetracycline.	do	Do.

(2) For sponsor 000004: it is used in free-choice cattle feeds such as feed blocks or salt-mineral mixes manufactured from approved Type A articles; such feeds are given to beef cattle and nonlactating dairy cattle to provide a daily intake of 0.5 to 2.0 milligrams of chlortetracycline per pound of body weight to aid in the control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline; the use of these Type A articles to make specific free-choice feed formulations must be approved under section 512(b) of the act and be based on a demonstration of drug stability and consumption which is consistent with the effective dose; the specific free-choice feed formulations approved in this paragraph can be manufactured under section 512(m) of the act.

(3) Chlortetracycline may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Amprolium in accordance with § 558.55.

(ii) Amprolium plus ethopabate in accordance with § 558.58.

(iii) Bacitracin methylene disalicylate in accordance with § 558.76.

(iv) Clopidol in accordance with § 558.175.

(v) Decoquinat in accordance with § 558.195.

(vi) Hygromycin B in accordance with § 558.274.

(vii) Monensin in accordance with § 558.355.

(viii) Roxarsone and salinomycin in accordance with § 558.550.

(ix) Robenidine hydrochloride in accordance with § 558.515.

(x) Roxarsone in accordance with § 558.530.

(xi) Salinomycin in accordance with § 558.550.

(xii) Zoalene in accordance with § 558.680.

(xiii) Tiamulin in accordance with § 558.600.

[41 FR 10995, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.128, see the List of Sections Affected in the Finding Aids section of this volume.

§ 558.140 Chlortetracycline and sulfamethazine.

(a) *Approvals.* Type A medicated articles: 35 grams of chlortetracycline per pound with 7.7 percent (35 grams) of sulfamethazine to 063238 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See §§ 556.150 and 556.670 of this chapter.

(c) It is used in feed for beef cattle as follows:

(1) *Amount per head per day.* Chlortetracycline, 350 milligrams plus sulfamethazine, 350 milligrams.

(2) *Indications for use.* Aid in the maintenance of weight gains in the presence of respiratory disease such as shipping fever.

(3) *Limitations.* Feed for 28 days; withdraw 7 days prior to slaughter.

[61 FR 35954, July 9, 1996, as amended at 64 FR 15684, Apr. 1, 1999]

§ 558.145 Chlortetracycline, procaine penicillin, and sulfamethazine.

(a) *Approvals.* Type A medicated articles: (1) 20 grams of chlortetracycline per pound, 4.4 percent (20 grams) of sulfamethazine, and procaine penicillin equivalent in activity to 10 grams of penicillin per pound to 046573 and 063238 in § 510.600(c) of this chapter.

(2) 40 grams of chlortetracycline per pound, 8.8 percent of sulfamethazine, and penicillin procaine equivalent in activity to 20 grams of penicillin per pound to 063238 in § 510.600(c) of this chapter.

(b) *Specifications.* (1) The antibiotic substance refers to the antibiotic or feed-grade antibiotic.

§ 558.155

(2) The antibiotic activities are expressed in terms of the appropriate antibiotic standards.

(3) Type C medicated feed contains in each ton, 100 grams of chlortetracycline, 50 grams of penicillin as procaine penicillin, and 100 grams of sulfamethazine.

(c) *Related tolerances.* See §§ 556.150, 556.510, and 556.670 of this chapter.

(d) *Conditions of use.* (1) It is administered to swine in a Type C feed for reduction of the incidence of cervical abscesses; treatment of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis* and vibronic dysentery); prevention of these diseases during times of stress; maintenance of weight gains in the presence of atrophic rhinitis; growth promotion and increased feed efficiency in swine weighing up to 75 pounds.

(2) Withdraw 15 days prior to slaughter.

[40 FR 13959, Mar. 27, 1975, as amended at 43 FR 19385, May 5, 1978; 47 FR 39814, Sept. 10, 1982; 48 FR 30615, July 5, 1983; 51 FR 7396, Mar. 3, 1986; 52 FR 2684, Jan. 26, 1987; 56 FR 14019, Apr. 5, 1991; 61 FR 18082, Apr. 24, 1996; 62 FR 14300, Mar. 26, 1997; 63 FR 27845, May 21, 1998]

§ 558.155 Chlortetracycline, sulfathiazole, penicillin.

(a) *Approvals.* Type A medicated articles: (1) 20 grams of chlortetracycline hydrochloride, 4.4 percent (20 grams) sulfathiazole, and procaine penicillin equivalent to 10 grams of penicillin per pound to 000004 and 000010 in § 510.600(c) of this chapter.

(2) 40 grams of chlortetracycline hydrochloride, 8.8 percent (40 grams) sulfathiazole and procaine penicillin equivalent in activity to 20 grams of penicillin per pound to Nos. 000004 and 000010 in § 510.600(c) of this chapter.

(b) *Specifications.* (1) The antibiotic substance refers to the antibiotic or feed-grade antibiotic.

(2) The antibiotic activities are expressed in terms of the appropriate antibiotic standards.

(c) *Related tolerances.* See §§ 556.150, 556.510, and 556.690 of this chapter.

(d) *Conditions of use.* It is used for swine as follows:

21 CFR Ch. I (4–1–99 Edition)

(1) *Amount per ton.* Chlortetracycline, 100 grams plus penicillin, 50 grams plus sulfathiazole, 100 grams.

(2) *Indications for use.* For reduction of incidence of cervical abscesses. Treatment of bacterial enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis* and vibronic dysentery). Maintenance of weight gains in the presence of atrophic rhinitis. Swine 10 pounds of body weight to 6 weeks post-weaning: Increased rate of weight gain and improved feed efficiency. Swine 6 to 16 weeks post-weaning: Increased rate of weight gain.

(3) *Limitations.* For swine raised in confinement (dry-lot) or on limited pasture. Feed as sole ration. Withdraw 7 days prior to slaughter.

MINIMUM AMOUNT OF TYPE C FEED WHICH THE ANIMAL SHOULD CONSUME

Type of feed	Approximate body weight in pounds	Minimum desired daily feed intake in pounds
Prestarter (up to 6 weeks postweaning)	20	1
Starter (up to 6 weeks postweaning)	50	1½
Grower (6–16 weeks postweaning)	80	2
Finisher (6–16 weeks postweaning)	150	3

[40 FR 13959, Mar. 27, 1975, as amended at 51 FR 7397, Mar. 3, 1986; 51 FR 28547, Aug. 8, 1986; 52 FR 2684, Jan. 26, 1987; 61 FR 2415, Jan. 26, 1996; 62 FR 35077, June 30, 1997; 62 FR 67725, Dec. 30, 1997; 63 FR 11599, Mar. 10, 1998]

§ 558.175 Clopidol.

(a) *Approvals.* Type A medicated articles: (1) 25 percent to 011526 in § 510.600(c) of this chapter.

(2) 25 percent of clopidol, 10 percent of roxarsone, and 4, 10, 15, or 25 grams of bacitracin methylene disalicylate per pound to 011526 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.160 of this chapter.

(c) [Reserved]

(d) *Conditions of use.* It is used as follows:

(1) *Broiler chickens—*(i) *Amount per ton.* Clopidol 113.5 grams (0.0125 percent).

(a) *Indications for use.* Aid in the prevention of coccidiosis caused by *E.*

tenella, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations*. Do not feed to chickens over 16 weeks of age.

(ii) *Amount per ton*. Clopidol, 113.5 grams (0.0125 percent) plus roxarsone, 45.4 grams (0.005 percent).

(a) *Indications for use*. Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; growth promotion and feed efficiency; improved pigmentation.

(b) *Limitations*. Do not feed to chickens over 16 weeks of age; withdraw 5 days before slaughter; as sole source of organic arsenic.

(iii) *Amount per ton*. Clopidol, 113.5 grams (0.0125 percent) plus roxarsone, 45.4 grams (0.005 percent) plus bacitracin, 4–25 grams.

(a) *Indications for use*. Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; growth promotion and feed efficiency; improved pigmentation; increased rate of weight gain.

(b) *Limitations*. Do not feed to chickens over 16 weeks of age; withdraw 5 days before slaughter; as sole source of organic arsenic; as bacitracin methylene disalicylate, provided by No. 046573 in §510.600(c) of this chapter; or as zinc bacitracin provided by Nos. 046573 and 063238 in §510.600(c) of this chapter.

(iv) *Amount per ton*. Clopidol, 113.5 grams (0.0125%) plus zinc bacitracin, 5 to 25 grams.

(a) *Indications for use*. For increased rate of weight gain and improved feed efficiency; aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations*. Feed continuously as sole ration. Zinc bacitracin as provided by Nos. 046573 and 063238 of §510.600(c) of this chapter.

(v) *Amount per ton*. Clopidol, 113.5 grams (0.0125 percent) plus bacitracin methylene disalicylate, 4 to 50 grams per ton.

(a) *Indications for use*. For increased rate of weight gain; to aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. mivati*, and *E. brunetti*.

(b) *Limitations*. Feed continuously as the sole ration from the time chicks are placed in floor pens until slaughter. Do not feed to chickens over 16 weeks of age. Bacitracin methylene disalicylate as provided by No. 046573 in §510.600(c) of this chapter.

(vi) *Amount per ton*. Clopidol, 113.5 grams (0.0125 percent) plus lincomycin, 2–4 grams.

(a) *Indications for use*. Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; increase in rate of weight gain and improved feed efficiency.

(b) *Limitations*. As lincomycin hydrochloride monohydrate; do not feed to chickens over 16 weeks of age.

(2) *Broiler chickens and replacement chickens*—(i) *Amount per ton*. Clopidol, 113.5 or 227 grams (0.0125 or 0.025 percent).

(a) *Indications for use*. Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations*. Feed up to 16 weeks of age if intended for use as caged layers; feed continuously as the sole ration; withdraw 5 days before slaughter if given at the level of 0.025 percent in feed or reduce level to 0.0125 percent 5 days before slaughter.

(ii) *Amount per ton*. Clopidol, 113.5 grams (0.0125 percent) plus chlortetracycline 100 to 200 grams.

(a) *Indications for use*. Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline.

(b) *Limitations*. Feed continuously as sole ration from the time chicks are placed in floor pens for 7 to 14 days.

(3) [Reserved]

(4) *Replacement chickens*—(i) *Amount per ton*. Clopidol 113.5 grams (0.0125 percent).

(a) *Indications for use*. Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations*. For replacement chickens intended for use as caged layers; do not feed to chickens over 16 weeks of age.

§ 558.185

21 CFR Ch. I (4–1–99 Edition)

(ii) *Amount per ton.* Clopidol 113.5 grams (0.0125 percent) plus roxarsone 45.4 grams (0.005 percent).

(a) *Indications for use.* Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; growth promotion and feed efficiency; improving pigmentation.

(b) *Limitations.* For replacement chickens intended for use as caged layers; do not feed to chickens over 16 weeks of age; withdraw 5 days before slaughter; as sole source of organic arsenic.

(5) *Turkeys—(i) Amount per ton.* Clopidol 113.5 or 227 grams (0.0125 or 0.025 percent).

(ii) *Indications for use.* Aid in the prevention of leucocytozoonosis caused by *Leucocytozoon smithi*.

(iii) *Limitations.* For turkeys grown for meat purposes only; to be administered continuously in feed at 0.0125 or 0.025 percent clopidol as the sole ration depending upon management practices, degree of exposure, and amount of feed eaten; withdraw medication 5 days before slaughter.

[40 FR 13959, Mar. 27, 1975, as amended at 40 FR 31935, July 30, 1975; 40 FR 52723, Nov. 12, 1975; 41 FR 1062, Jan. 6, 1976; 41 FR 53002, Dec. 3, 1976; 44 FR 20676, Apr. 6, 1979; 51 FR 7397, Mar. 3, 1986; 52 FR 2684, Jan. 26, 1987; 53 FR 20843, June 7, 1988; 54 FR 6518, Feb. 13, 1989; 61 FR 35954, July 9, 1996; 62 FR 61626, Nov. 19, 1997; 62 FR 61912, Nov. 20, 1997; 62 FR 62243, Nov. 21, 1997; 63 FR 17948, Apr. 13, 1998; 63 FR 27845, May 21, 1998]

§ 558.185 Coumaphos.

(a) *Approvals.* Type A medicated articles:

(1) [Reserved]

(2) 1.12 and 11.2 percent to 017800 in § 510.600(c) of this chapter for use as in paragraph (d)(1)(ii) of this section.

(b) *Special considerations.* Adequate directions and warnings for use must be given and shall include a statement that coumaphos is a cholinesterase inhibitor and that animals being treated with coumaphos should not be exposed during or within a few days before or after treatment to any other cholinesterase-inhibiting drugs, insecticides, pesticides, or chemicals.

(c) *Related tolerances.* See 40 CFR 180.189.

(d) *Conditions of use.* It is used as follows:

(1) *Beef and dairy cattle—(i) Amount.* Coumaphos 0.00012 lb. (0.054 gram) per 100 lb. body weight per day.

(a) *Indications for use.* As an aid in the reduction of fecal breeding flies through control of fly larvae.

(b) *Limitations.* Feed for the duration of fly season in a Type C feed containing 0.0033 percent or in a feed Type B feed containing not over 0.0066 percent coumaphos; do not feed to animals less than 3 months old; not for use in pelleted feeds.

(ii) *Amount.* Coumaphos, 0.0002 lb. (0.091 gram) per 100 lb. body weight per day.

(a) *Indications for use.* Control of gastrointestinal roundworms (*Haemonchus* spp., *Ostertagia* spp., *Cooperia* spp., *Nematodirus* spp., *Trichostrongylus* spp.).

(b) *Limitations.* Feed 0.0002 lb. (0.091 gram) per 100 lb. body weight per day for 6 consecutive days in the normal grain ration to which the animals are accustomed but not in rations containing more than 0.1 percent coumaphos; do not feed to animals less than 3 months old; do not feed to sick animals or animals under stress, such as those just shipped, dehorned, castrated, or weaned within the last 3 weeks; do not feed in conjunction with oral drenches or with feeds containing phenothiazine. Should conditions warrant, repeat treatment at 30-day intervals.

(2) *Laying chickens—(i) Amount.* Coumaphos 27.2 grams per ton (0.003 percent).

(ii) *Indications for use.* For control of capillary worm (*Capillaria obsignata*) and as an aid in control of common round worm (*Ascaridia galli*) and cecal worm (*Heterakis gallinae*).

(iii) *Limitations.* In Type C feed; administer continuously as the total feed ration for 14 days; when reinfection occurs, treatment may be repeated but not sooner than 3 weeks after the end of the previous treatment; do not feed to chickens within 10 days of vaccination or other conditions of stress; treatment of colored breeds of commercial layers should be avoided while in production since these breeds appear to be more sensitive to coumaphos than

Food and Drug Administration, HHS

§ 558.195

white breeds; as sole medication; medications in general should be avoided while birds are approaching peak production; such interruption of normal feeding practices may upset the flock and lower egg production; diagnosis by competent personnel is essential; flock condition and production records should be carefully evaluated prior to treatment.

(3) *Replacement pullets*—(i) *Amount*. Coumaphos 36.3 grams per ton (0.004 percent).

(ii) *Indications for use*. For control of capillary worm (*Capillaria obsignata*) and as an aid in control of common roundworm (*Ascaridia galli*) and cecal worm (*Heterakis gallinae*).

(iii) *Limitations*. In Type C feed; administer before the onset of production; diagnosis by competent personnel is essential; administer continuously as total feed ration for from 10 to 14 days; do not feed to chickens under 8 weeks of age nor within 10 days of vaccination or other conditions of stress; if birds are maintained on contaminated litter or exposed to infected birds, a second 10 to 14 day treatment is recommended but not sooner than 3 weeks after the end of the previous

treatment; as sole medication; if reinfection occurs after production begins, repeat treatment as recommended for laying flocks.

[40 FR 13959, Mar. 27, 1975, as amended at 42 FR 1463, Jan. 7, 1977; 51 FR 7397, Mar. 3, 1986; 52 FR 2684, Jan. 26, 1987; 61 FR 34729, July 3, 1996]

§ 558.195 Decoquinate.

(a) *Approvals*. Type A medicated articles: 6 percent to 046573 in § 510.600(c) of this chapter.

(b) *Related tolerances in edible products*. See § 556.170 of this chapter.

(c) *Special considerations*. (1) Bentonite should not be used in decoquinate feeds.

(2) Type A medicated articles containing 6 percent decoquinate may be used to make dry or liquid Type B cattle (including veal calf), sheep, and goat feeds as in paragraph (d) of this section.

(3) Type A medicated articles containing 6 percent decoquinate may be used to manufacture dry or liquid Type B cattle feeds as indicated in paragraph (d) of this section.

(d) *Conditions of use*. It is used as follows:

Decoquinate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
13.6 (0.00149 pct)		Young goats; for the prevention of coccidiosis caused by <i>Eimeria christenseni</i> and <i>E. ninakohlyakimovae</i> .	Feed at a rate to provide 22.7 mg per 100 lbs of body weight per day (0.5 mg per kilogram); do not feed to goats producing milk for food; feed for at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard.	046573
13.6 (0.0015 pct)		Young sheep; for the prevention of coccidiosis caused by <i>Eimeria ovinoidalis</i> , <i>E. crandallis</i> , <i>E. parva</i> , <i>E. bakuensis</i> .	Feed Type C feed at a rate to provide 22.7 mg per 100 lb of body weight (0.5 mg per kg) per day. Feed at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard. Do not feed to sheep producing milk for food.	046573

Decoquinate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
13.6 to 27.2 (0.0015 to 0.003 pct).		Cattle; for the prevention of coccidiosis in ruminating and non-ruminating calves and cattle caused by <i>Eimeria bovis</i> and <i>E. zuernii</i> .	Feed Type C feed at a rate to provide 22.7 mg per 100 lb of body weight (0.5 mg per kg) per day. May be prepared from dry or liquid Type B feed containing 0.0125 to 0.5 pct decoquinate. Liquid Type B feed must have a pH range of 5.0 to 6.5 and contain a suspending agent to maintain a viscosity of not less than 500 centipoises. Feed at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard. Do not feed to cows producing milk for food.	
13.6 to 535.7 (0.0015 to 0.059 pct).		Cattle: prevention of coccidiosis in ruminating and nonruminating calves (including veal calves) and cattle caused by <i>Eimeria bovis</i> and <i>E. zuernii</i> .	Feed Type C feed (including dry milk replacer) to provide 22.7 mg per 100 lb body weight (0.5 mg per kg) per day. May be prepared from dry Type B feed containing 0.06 to 0.6 pct decoquinate or liquid Type B feed containing 0.0125 to 0.05 pct decoquinate. The liquid Type B feed must have pH 5.0 to 6.5 and contain a suspending agent to maintain a viscosity of not less than 500 centipoises. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard. Do not feed to animals producing milk for food.	046573
		Young sheep: prevention of coccidiosis caused by <i>Eimeria ovinoidalis</i> , <i>E. parva</i> , <i>E. bakuensis</i> , <i>E. crandallii</i> .	do	do
		Young goats: prevention of coccidiosis caused by <i>Eimeria christensenii</i> , <i>E. ninakohlyakimovae</i> .	do	do
27.2 (0.003 pct)		Broiler chickens; for the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. mivati</i> , <i>E. acervulina</i> , <i>E. maxima</i> and <i>E. brunetti</i> .	Do not feed to laying chickens ..	046573
	Bacitracin 4 to 50	Broiler chickens; for the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. mivati</i> , <i>E. acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> , and for increased rate of weight gain and improved feed efficiency..	Do not feed to laying chickens; feed continuously as sole ration; bacitracin methylene disalicylate as provided by 046573 in §510.600(c) of this chapter..	046573
	Bacitracin 10 to 50 ..	Broiler chickens; for the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. mivati</i> , <i>E. acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> ; for increased rate of weight gain and improved feed efficiency.	Do not feed to laying chickens; feed as sole ration; as zinc bacitracin provided by Nos. 046573 and 011716 in sec. 510.600(c) of this chapter.	046573

Food and Drug Administration, HHS

§ 558.205

Decoquinate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
	Bacitracin methylene disalicylate 50 and roxarsone 22.7–45.4.	Broiler chickens; for prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. mivati</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> ; as an aid in the prevention of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin; for increased rate of weight gain, improved feed efficiency, and improved pigmentation.	Feed continuously as sole ration. Withdraw 5 days before slaughter. Do not feed to laying chickens. Not for use in breeder chickens. Use as sole source of organic arsenic. Poultry should have access to drinking water at all times. Drug overdosage or lack of drinking water may result in leg weakness or paralysis. Decoquinate, bacitracin methylene disalicylate, and roxarsone, as provided by No. 046573 in § 510.600(c) of this chapter.	046573
	Chlortetracycline 100 to 200.	Chickens; for the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. mivati</i> , <i>E. acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> ; control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption; in low calcium feed containing 0.8 pct. of calcium; feed continuously 7 to 14 days.	046573
	Chlortetracycline 200 to 400.	Chickens; for the prevention of coccidiosis caused by <i>E. tenella</i> , <i>E. necatrix</i> , <i>E. mivati</i> , <i>E. acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> ; control of chronic respiratory disease (CRD) and air sac infection caused by <i>M. gallisepticum</i> and <i>E. coli</i> susceptible to chlortetracycline.	do	046573
	Roxarsone 45.4 (0.005 pct.).	Broiler chickens; for the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. mivati</i> , <i>E. acervulina</i> , <i>E. maxima</i> , and <i>E. brunetti</i> ; growth promotion and feed efficiency; improving pigmentation.	Do not feed to laying chickens; withdraw 5 days before slaughter; as sole source of organic arsenic.	046573
	Lincomycin 2	Broiler chickens; for the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. mivati</i> , and <i>E. brunetti</i> ; for increased rate of weight gain and improved feed efficiency.	Do not feed to laying chickens; feed as sole ration; as lincomycin hydrochloride monohydrate provided by No. 000009 in sec. 510.600(c) of this chapter.	000009, 046573
	Roxarsone 11 to 45 (0.0012–0.005 pct.) plus Bacitracin 12 to 50.	Broiler chickens; for the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. mivati</i> , <i>E. maxima</i> , and <i>E. brunetti</i> ; for increased rate of weight gain and improved feed efficiency.	Do not feed to laying chickens; withdraw 5 days before slaughter; as sole source of organic arsenic; as zinc bacitracin provided by Nos. 011716, 046573, and 063238 in sec. 510.600(c) of this chapter; as roxarsone provided by No. 046573 in sec. 510.600(c) of this chapter.	046573

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.195, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.205 Dichlorvos.

(a) *Approvals.* Type A medicated articles: 3.1 and 9.6 percent to 000010 in § 510.600(c) of this chapter.

(b) *Special considerations.* (1) Dichlorvos is to be included in meal or mash or mixed with feed in crumble form only after the crumble feed has been manufactured. Do not mix in feeds to be pelleted nor with pelleted feed. Do not soak the feed or administer as wet mash. Feed must be dry

§ 558.235

when administered. Do not use in animals other than swine. Do not allow fowl access to feed containing this preparation or to feces from treated animals.

(2) Dichlorvos is a cholinesterase inhibitor. Do not use this product in animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals. If human or animal poisoning should occur, immediately consult a physician or a veterinarian. Atropine is antidotal.

(3) Labeling for Type A articles and Type B feeds must include a statement that containers or materials used in packaging such Type A articles and Type B feeds are not to be reused and all such packaging materials must be destroyed after the product has been used.

(c) *Related tolerances.* See § 556.180 of this chapter.

(d) *Conditions of use.* It is used in feed for swine as follows:

(1) *Amount per ton.* Dichlorvos, 348 grams (0.0384 percent).

(i) *Indications for use.* For the removal and control of mature, immature, and/or fourth-stage larvae of the whipworm (*Trichuris suis*), nodular worm (*Oesophagostomum* sp.), large roundworm (*Ascaris suum*) and the thick stomach worm (*Ascarops strongylina*) of the gastrointestinal tract.

(ii) *Limitations.* For swine up to 70 pounds body weight, feed as sole ration for 2 consecutive days. For swine from 70 pounds to market weight, feed as sole ration at the rate of 8.4 pounds of feed per head until the medicated feed has been consumed. For boars, open or bred gilts, and sows, feed as sole ration at the rate of 4.2 pounds per head per day for 2 consecutive days.

(2) *Amount per ton.* Dichlorvos, 479 grams (0.0528 percent).

(i) *Indications for use.* For the removal and control of mature, immature, and/or fourth-stage larvae of the whipworm (*Trichuris suis*), nodular worm (*Oesophagostomum* sp.), large roundworm (*Ascaris suum*), and the thick stomach worm (*Ascarops strongylina*) of the gastrointestinal tract.

21 CFR Ch. I (4–1–99 Edition)

(ii) *Limitations.* For boars, open or bred gilts, and sows, feed as sole ration at the rate of 6 pounds per head for one feeding.

(3) *Amount per ton.* Dichlorvos, 334–500 grams (0.0366–0.0550 percent).

(i) *Indications for use.* An aid in improving litter production efficiency by increasing pigs born alive, birth weights, survival to market, and rate of weight gain. Treatment also removes and controls mature, immature and/or fourth stage larvae of whipworm (*Trichuris suis*), nodular worm (*Oesophagostomum* spp.) large roundworm (*Ascaris suum*), and the thick stomach worm (*Ascarops strongylina*) occurring in the gastrointestinal tract of the sow or gilt.

(ii) *Limitations.* For pregnant swine; mix into a gestation feed to provide 1,000 milligrams per head daily during last 30 days of gestation.

[40 FR 13959, Mar. 27, 1975, as amended at 40 FR 50258, Oct. 29, 1975; 48 FR 46515, Oct. 13, 1983; 51 FR 7397, Mar. 3, 1986; 51 FR 28547, Aug. 8, 1986; 52 FR 2684, Jan. 26, 1987; 62 FR 35077, June 30, 1997]

§ 558.235 Efrotomycin.

(a) *Approvals.* Type A medicated article: 14.5 grams per pound to 050604 in § 510.600(c) of this chapter.

(b) *Conditions of use—(1) Swine.* (i) *Amount.* 3.6 grams per ton.

(A) *Indications for use.* For improved feed efficiency.

(B) *Limitations.* Feed continuously as sole ration. Not to be used in swine weighing more than 250 pounds.

(ii) *Amount.* 3.6 to 14.5 grams per ton.

(A) *Indications for use.* For increased rate of weight gain.

(B) *Limitations.* Feed continuously as sole ration. Not to be used in swine weighing more than 250 pounds.

(2) [Reserved]

[57 FR 38442, Aug. 25, 1992, as amended at 62 FR 63271, Nov. 28, 1997]

§ 558.248 Erythromycin thiocyanate.

(a) *Approvals.* Type A medicated articles: (1) 2.2 percent to 050604 in § 510.600(c) of this chapter for use as in paragraph (d) of this section.

(2) 5 and 10 percent to 050604 for use in paragraphs (d)(1)(i) and (ii) of this section.

Food and Drug Administration, HHS

§ 558.254

(b) *Special considerations.* The levels of antibiotic are expressed in terms of erythromycin master standard. One gram of erythromycin thiocyanate is equivalent to 0.925 gram of erythromycin master standard.

(c) *Related tolerances.* See § 556.230 of this chapter.

(d) *Condition of use.* (1) It is used as follows:

Erythromycin thiocyanate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 4.6 to 18.5		Chickens; growth promotion and feed efficiency.		050604
(ii) 9.25 to 18.5		Turkeys; growth promotion and feed efficiency.	For turkeys not over 12 weeks of age.	050604
(iii) 9.25 to 64.75		Swine; increase in weight gain, improved feed efficiency in starter pigs (9.25 to 64.75) and grower-finishing pigs (9.25).	Starter ration for animals up to 35 lb body weight.	050604
(iv) 18.5		Laying chickens; aids in increasing egg production.		050604
(v) 92.5		1. Chickens; as an aid in the prevention of chronic respiratory disease during periods of stress. 2. Chickens; as an aid in the prevention of infectious coryza. 3. Turkeys; as an aid in the prevention of chronic respiratory disease during periods of stress.	Feed for 2 d before stress and 3 to 6 d after stress; withdraw 24 h before slaughter. Feed for 7 to 14 d; withdraw 24 h before slaughter. Feed for 2 d before stress and 3 to 6 d after stress.	050604
(vi) 185		1. Chickens; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease. 2. Turkeys; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease.	Feed for 5 to 8 d; do not use in birds producing eggs for food purposes; withdraw 48 h before slaughter. Feed for 5 to 8 d; do not use in birds producing eggs for food purposes.	050604 050604

(2) In feed for feedlot beef cattle at 37 milligrams per head per day as an aid in stimulating growth and improving feed efficiency.

(3) Erythromycin thiocyanate may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Amprolium in accordance with § 558.55.

(ii) Amprolium and ethopabate in accordance with § 558.58.

(iii) Arsanilic acid in accordance with § 558.62.

(iv) Zoalene in accordance with § 558.680.

[41 FR 10999, Mar. 15, 1976, as amended at 45 FR 56799, Aug. 26, 1980; 49 FR 31281, Aug. 6, 1984; 51 FR 7397, Mar. 3, 1986; 52 FR 2684, Jan. 26, 1987; 54 FR 12189, Mar. 24, 1989]

§ 558.254 Famphur.

(a) *Approvals.* Type A medicated articles: 13.2 and 33.3 percent to 000061 in § 510.600(c) of this chapter.

(b) *Special considerations.* Famphur is a cholinesterase inhibitor. Do not use this product in animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(c) *Related tolerances.* See § 556.273 of this chapter.

(d) *Conditions of use.* It is used in the feed for cattle as follows:

(1) *Amount.* 1.1 milligrams per pound body weight per day.

(i) *Indications for use.* For control of grubs and as an aid in control of sucking lice.

(ii) *Limitations.* For beef cattle and nonlactating dairy cows; feed for 30 days; withdraw from dry dairy cows

§ 558.258

21 CFR Ch. I (4–1–99 Edition)

and heifers 21 days prior to freshening; withdraw 4 days prior to slaughter.

(2) *Amount.* 2.3 milligrams per pound body weight per day.

(i) *Indications for use.* For control of grubs.

(ii) *Limitations.* For beef cattle and nonlactating dairy cows; feed for 10 days; withdraw from dry dairy cows and heifers 21 days prior to freshening; withdraw 4 days prior to slaughter.

[41 FR 11000, Mar. 15, 1976, as amended at 51 FR 7397, Mar. 3, 1986; 57 FR 7652, Mar. 4, 1992; 62 FR 55161, Oct. 23, 1997; 62 FR 61626, Nov. 19, 1997]

§ 558.258 Fenbendazole.

(a) *Approvals.* Type A medicated articles: 4 percent (18.1 grams per pound), 8 percent (36.2 grams per pound), and 20 percent (90.7 grams per pound) fenbendazole and all combinations provided for in this section to 012799 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.275 of this chapter.

(c) *Conditions of use.* (1) It is used in swine feed as follows:

(i) *Amount.* Fenbendazole, 10 to 80 grams per ton (to provide 9 milligrams per kilogram of body weight) given over a 3- to 12-day period.

(A) *Indications for use.* For the removal of: adult stage lungworms (*Metastrongylus apri* and *M. pudendotectus*); adult and larvae (L3, 4 stages—liver, lung, intestinal forms) large roundworms (*Ascaris suum*); adult stage nodular worms (*Oesophagostomum dentatum*, *O. quadrispinulatum*); small stomach worms (*Hyoststrongylus rubidus*); adult and larvae (L2, 3, 4 stages—intestinal mucosal forms) whipworms (*Trichuris suis*); adult and larvae kidney worms (*Stephanurus dentatus*).

(B) Feed as sole ration.

(ii) *Amount.* Fenbendazole 10 to 80 grams per ton (to provide 9 milligrams per kilogram body weight) and lincomycin 20 grams per ton.

(A) *Indications for use.* As an anthelmintic (as provided in paragraph (c)(1)(i)(A) of this section) and for increased rate of gain in growing-finishing swine.

(B) *Limitations.* Feed as sole ration. Do not feed to swine that weigh more than 250 pounds; as lincomycin pro-

vided by 000009 in § 510.600(c) of this chapter.

(iii) *Amount.* Fenbendazole 10 to 80 grams per ton (to provide 9 milligrams per kilogram body weight) and lincomycin 40 grams per ton.

(A) *Indications for use.* As an anthelmintic (as provided in paragraph (c)(1)(i)(A) of this section) and for control of swine dysentery in animals on premises with a history of swine dysentery, but where symptoms have not yet occurred.

(B) *Limitations.* Feed as sole ration. Do not feed to swine that weigh more than 250 pounds; as lincomycin provided by 000009 in § 510.600(c) of this chapter.

(iv) *Amount.* Fenbendazole 10 to 80 grams per ton (to provide 9 milligrams per kilogram body weight) and lincomycin 100 grams per ton.

(A) *Indications for use.* As an anthelmintic (as provided in paragraph (c)(1)(i)(A) of this section) and for the treatment of swine dysentery.

(B) *Limitations.* Feed as sole ration. Do not use within 6 days of slaughter. Do not feed to swine that weigh more than 250 pounds; as lincomycin provided by 000009 in § 510.600(c) of this chapter.

(v) *Amount.* Fenbendazole 10 to 80 grams per ton (to provide 9 milligrams per kilogram body weight) and lincomycin 200 grams per ton.

(A) *Indications for use.* As an anthelmintic (as provided in paragraph (c)(1)(i)(A) of this section) and for reduction in the severity of swine mycoplasmal pneumonia caused by *Mycoplasma hyopneumoniae*.

(B) *Limitations.* Feed as sole ration. Do not use within 6 days of slaughter. Do not feed to swine that weigh more than 250 pounds; as lincomycin provided by 000009 in § 510.600(c) of this chapter.

(2) It is used in the feed of beef and dairy cattle as follows:

(i) *Amount.* 5 milligrams fenbendazole per kilogram body weight (2.27 milligrams per pound).

(ii) *Indications for use.* For the removal and control of lungworms (*Dictyocaulus viviparus*); barberpole worms (*Haemonchus contortus*); brown stomach worms (*Ostertagia ostertagi*); small stomach worms (*Trichostrongylus*

Food and Drug Administration, HHS

§ 558.265

axei); hookworms (*Bunostomum phlebotomum*); thread-necked intestinal worms (*Nematodirus helvetianus*); small intestinal worms (*Cooperia punctata*

(iii) *Limitations*. Feed as sole ration for one day. Do not use within 13 days of slaughter.

(3) It is used in free-choice beef and dairy cattle feed as follows:

(i) *Amount*. 5 milligrams fenbendazole per kilogram body weight (2.27 milligrams per pound).

(ii) *Indications for use*. For the removal and control of infections of lungworms (*Dictyocaulus viviparus*), barberpole worms (*Haemonchus contortus*), brown stomach worms (*Ostertagia ostertagi*), small stomach worms (*Trichostrongylus axei*), hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small intestinal worms (*Cooperia oncophora* and *C. punctata*), bankrupt worms (*Trichostrongylus colubriformis*), and nodular worms (*Oesophagostomum radiatum*) in cattle.

(iii) *Limitations*. Feed a total of 5 milligrams of fenbendazole per kilogram (2.27 milligrams per pound) of body weight to cattle over a 3 to 6 day period. Retreatment may be needed after 4 to 6 weeks. Do not slaughter within 13 days following last treatment. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(iv) May be fed in a Type C feed as follows:

Ingredient	Percent	Inter-national feed No.
(A) Ingredient:		
Copper sulfate	0.45	6-01-720
Dried Cane Molasses	3.12	4-04-695
Monosodium phosphate	31.16	6-04-288
Salt (sodium chloride)	59.00	6-04-152
Zinc sulfate	0.76	6-05-556
Fenbendazole Type A article (200 grams per kilogram)	5.51	
(B) Ingredient:		
Dicalcium phosphate	32.31	6-00-080
Limestone	17.13	6-02-632
Magnesium oxide	9.79	6-02-756
Zinc sulfate	1.47	6-05-556
Copper sulfate	0.29	6-01-720
Potassium iodide	0.0098	6-03-759
Dried Cane Molasses	0.98	4-04-695
Selenium	0.0002	
Salt	35.93	6-04-152

Ingredient	Percent	Inter-national feed No.
Fenbendazole Type A article (200 grams per kilogram)	2.09	

(C) The content of any added vitamin and trace mineral may be varied; however, they should be comparable to those used by the firm for other free-choice feeds. Formulation modifications require FDA approval prior to marketing. The amount of selenium must comply with published regulations.

(4) *Zoo and wildlife animals*. For removal and control of internal parasites in hoofed zoo and wildlife animals as follows:

(i) *Feral swine* (*Suis scrofa*): 3 milligrams per kilogram per day for 3 days. Treatment for kidney worm (*Stephanurus dentatus*), roundworm (*Ascaris suum*), nodular worm (*Oesophagostomum dentatum*).

(ii) *Ruminants* (subfamily *Antilopinae*, *Hippotraginae*, *Caprinae*): 2.5 milligrams per kilogram per day for 3 days. Treatment for small stomach worm (*Trichostrongylus* spp.), thread necked intestinal worm (*Nematodirus* spp.), barberpole worm (*Haemonchus* spp.), whipworm (*Trichuris* spp.).

(iii) *Rocky mountain bighorn sheep* (*Ovis c. canadensis*): 10 milligrams per kilogram per day for 3 days. Treatment and control of *Protostrongylus* spp.

(iv) *Limitations*. Use as complete feed. Prior withdrawal of feed or water is not necessary. Retreatment may be required in 6 weeks. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Do not use 14 days before or during the hunting season.

[49 FR 3846, Jan. 31, 1984, as amended at 51 FR 7397, Mar. 3, 1986; 53 FR 14788, Apr. 26, 1988; 54 FR 36963, Sept. 6, 1989; 55 FR 48231, Nov. 20, 1990; 57 FR 34516, Aug. 5, 1992; 58 FR 47973, Sept. 13, 1993; 59 FR 35252, July 11, 1994; 61 FR 29478, June 11, 1996]

§ 558.265 Halofuginone hydrobromide.

(a) *Approvals*. Type A medicated articles: 6 grams per kilogram (2.72 grams per pound) to 012579 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.308 of this chapter.

(c) *Conditions of use.* (1) It is used in feed for broiler chickens as follows:

(i) *Amount.* 2.72 grams per ton.

(A) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(B) *Limitations.* Feed continuously as sole ration; withdraw 4 days before slaughter; do not feed to layers; avoid contact with skin, eyes, or clothing; keep out of lakes, ponds, or streams.

(ii) *Amount per ton.* Halofuginone 2.72 grams (0.0003 percent) plus bambermycins 1 to 2 grams.

(A) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*; for increased rate of weight gain and improved feed efficiency.

(B) *Limitations.* Feed continuously as sole ration; withdraw 5 days before slaughter; do not feed to layers.

(iii) *Amount per ton.* Halofuginone 2.72 grams (0.0003 percent) plus virginiamycin 5 grams.

(A) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*; for increased rate of weight gain and improved feed efficiency.

(B) *Limitations.* Feed continuously as sole ration; withdraw 6 days before slaughter; do not feed to layers.

(iv) *Amount per ton.* Halofuginone 2.72 grams (0.0003 percent) plus virginiamycin 5 to 15 grams.

(A) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*; for increased rate of weight gain.

(B) *Limitations.* Feed continuously as sole ration; withdraw 6 days before slaughter; do not feed to layers.

(v) *Amount per ton.* Halofuginone hydrobromide 2.72 grams (0.0003 percent) plus bacitracin methylene disalicylate 10 to 50 grams and roxarsone 22.7 to 45.4 grams.

(A) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*; for

increased rate of weight gain; and for improved feed efficiency.

(B) *Limitations.* Feed continuously as sole ration; withdraw 5 days before slaughter; use as sole source of organic arsenic; do not feed to layers; avoid contact with skin, eyes, or clothing; keep out of lakes, ponds, or streams.

(vi) *Amount per ton.* Halofuginone 2.72 grams (0.0003 percent) plus bacitracin methylene disalicylate 10 to 50 grams.

(A) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, *E. maxima* and for improved feed efficiency.

(B) *Limitations.* Feed continuously as sole ration; withdraw 5 days before slaughter; do not feed to layers; avoid contact with skin, eyes, or clothing; keep out of lakes, ponds, or streams.

(vii) *Amount per ton.* Halofuginone 2.72 grams (0.0003 percent) plus lincomycin 2 to 4 grams.

(A) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima* and for improved feed efficiency.

(B) *Limitations.* Feed continuously as sole ration; withdraw 4 days before slaughter; do not feed to layers; avoid contact with skin, eyes, or clothing; keep out of lakes, ponds, or streams.

(2) It is used in feed for turkeys as follows:

(i) *Amount per ton.* 1.36 to 2.72 grams.

(A) *Indications for use.* For the prevention of coccidiosis in growing turkeys caused by *Eimeria adenoides*, *E. meleagritidis*, and *E. gallopavonis*.

(B) *Limitations.* Feed continuously as sole ration; withdraw 7 days before slaughter; do not feed to layers or water fowl; avoid contact with skin, eyes, or clothing; keep out of lakes, ponds, or streams.

(ii) *Amount per ton.* Halofuginone hydrobromide 1.36 to 2.72 grams plus bacitracin methylene disalicylate 10 to 50 grams.

(A) *Indications for use.* For prevention of coccidiosis caused by *Eimeria adenoides*, *E. meleagritidis*, and *E. gallopavonis*, and for increased rate of weight gain in growing turkeys.

(B) *Limitations.* Feed continuously as sole ration. Withdraw 7 days before

slaughter. Do not feed to laying chickens or water fowl. Keep out of lakes, ponds, and streams. Halofuginone is toxic to fish and aquatic life. Halofuginone is an irritant to eyes and skin. Avoid contact with skin, eyes, or clothing.

(iii) *Amount per ton.* 1.36 to 2.72 grams of halofuginone hydrobromide plus 2 grams of bambermycins.

(A) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria adenoides*, *E. meleagritidis*, and *E. gallopavonis*, and for increased rate of weight gain in growing turkeys.

(B) *Limitations.* Feed continuously as sole ration. Withdraw 7 days before slaughter. Do not feed to laying chickens or waterfowl. Halofuginone hydrobromide is toxic to fish and other aquatic life. Keep out of lakes, ponds, and streams. Halofuginone hydrobromide is an eye and skin irritant. Avoid contact with skin, eyes, and clothing.

(3) It is used in feed for replacement cage laying chickens and replacement broiler breeder chickens as follows:

(i) *Amount per ton.* 2.72 grams.

(A) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. mivati*/E. *mitis*, and *E. brunetti*.

(B) *Limitations.* Feed continuously as sole ration to replacement cage laying chickens until 20 weeks of age. Feed continuously as sole ration to replacement broiler breeder chickens until 16 weeks of age. Withdraw 4 days before slaughter. Do not feed to laying chickens or water fowl. Halofuginone hydrobromide is toxic to fish and aquatic life. Keep out of lakes, ponds, and streams. Halofuginone hydrobromide is an irritant to eyes and skin. Avoid contact with skin, eyes, and clothing.

(ii) [Reserved]

[50 FR 33719, Aug. 21, 1985, as amended at 50 FR 42518, Oct. 21, 1985; 51 FR 7397, Mar. 3, 1986; 51 FR 11439, Apr. 3, 1986; 51 FR 14989, Apr. 22, 1986; 51 FR 23737, July 1, 1986; 53 FR 1018, Jan. 15, 1988; 53 FR 11065, Apr. 5, 1988; 54 FR 11519, Mar. 21, 1989; 54 FR 28052, July 5, 1989; 59 FR 51498, Oct. 12, 1994; 61 FR 21076, May 9, 1996; 61 FR 24694, May 16, 1996]

§ 558.274 Hygromycin B.

(a) *Approvals.* (1) Type A medicated articles: 2.4 and 8 grams per pound to 000986 in § 510.600(c) of this chapter for use as in paragraph (c) of this section.

(2) 2.4 grams per pound to 016968 and 043733 in § 510.600(c) of this chapter for use in swine feed as in paragraph (c)(1)(ii) of this section.

(3) 1.2 grams per pound to 016968 in § 510.600(c) of this chapter for use in swine as in paragraph (c)(1)(ii) of this section.

(4) 0.6 gram per pound to 016968, 017790, 043733, and 050639 in § 510.600(c) of this chapter for use in chickens as in paragraph (c)(1)(i) of this section and in swine as in paragraph (c)(1)(ii) of this section.

(5) 0.48 and 2.4 grams per pound to 026186 in § 510.600(c) of this chapter for use in chickens as in paragraph (c)(1)(i) of this section and in swine as in paragraph (c)(1)(ii) of this section.

(6) [Reserved]

(7) 2.4 grams per pound to 011790 and 017519 in § 510.600(c) of this chapter for use in chickens as in paragraph (c)(1)(i) and in swine as in paragraph (c)(1)(ii) of this section.

(8) 0.6 and 1.6 grams per pound granted to 046573 in § 510.600(c) of this chapter for use in chickens as in paragraph (c)(1)(i) and in swine as in paragraph (c)(1)(ii) of this section.

(b) *Related tolerances.* See § 556.330 of this chapter.

(c) *Conditions of use.* (1) It may be used as follows:

Hygromycin B in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 8 to 12		Chickens: control of infestation of large roundworms (<i>Ascaris galli</i>), cecal worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>).	Withdraw 3 days before slaughter.	000986, 016968, 017519, 017790, 026186, 043733, 046573, 050639

Hygromycin B in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
	Bacitracin 100	Chickens; control of infestation of large roundworms (<i>Ascaris galli</i>), cecal worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>); treatment of chronic respiratory disease (airsac infection), blue comb (nonspecific infectious enteritis).	As bacitracin methylene disalicylate or zinc bacitracin; withdraw 3 days before slaughter.	
	Bacitracin plus penicillin (100 to 200 of combination).	1. Chickens; control of infestation of large roundworms (<i>Ascaris galli</i>), cecal worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>); treatment of chronic respiratory disease (airsac infection), blue comb (nonspecific infectious enteritis). 2. Chickens; control of infestation of large roundworms (<i>Ascaris galli</i>), cecal worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>); treatment of chronic respiratory disease (airsac infection), blue comb (nonspecific infectious enteritis). 3. Chickens; control of infestation of large roundworms (<i>Ascaris galli</i>), cecal worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>); treatment of chronic respiratory disease (airsac infection), blue comb (nonspecific infectious enteritis).	Feed containing not less than 25% of penicillin plus not less than 50% of bacitracin; as procaine penicillin plus bacitracin methylene disalicylate; withdraw 3 days before slaughter. Combination containing not less than 50% nor more than 75% of bacitracin, except that it contains not more than 125 g of penicillin; as procaine penicillin plus zinc bacitracin; withdraw 3 days before slaughter. Combination containing 50% to 75% bacitracin, but not more than 125 g of penicillin, as procaine penicillin; withdraw 3 days before slaughter..	
	Chlortetracycline 100 to 200.	Chickens; control of infestation of large roundworms (<i>Ascaris galli</i>), cecal worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>); control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption; feed for 7 to 14 days; withdraw 3 days before slaughter.	
	Chlortetracycline 200 to 400.	Chickens; control of infestation of large roundworms (<i>Ascaris galli</i>), cecal worms (<i>H. Gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>); control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline.	do	
	Penicillin 100	Chickens; control of infestation of large roundworms (<i>Ascaris galli</i>), cecal worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>); treatment of chronic respiratory disease (airsac infection), blue comb (nonspecific infectious enteritis).	As procaine penicillin; withdraw 3 days before slaughter.	

Food and Drug Administration, HHS

§ 558.300

Hygromycin B in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(ii) 12	Tylosin 4 to 50	Chickens: Control of infestations of large roundworms (<i>Ascaris galli</i>), cecal worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>); growth promotion and feed efficiency.	As tylosin phosphate; withdraw 3 days before slaughter.	000986
		Swine: control of infestation of large roundworms (<i>Ascaris suis</i>), nodular worms (<i>Oesophagostomum dentatum</i>), and whipworms (<i>Trichuris suis</i>).	Withdraw 15 days before slaughter.	000986, 016968, 017519, 017790, 026186, 043733, 046573, 050639
	Chlortetracycline 400..	Swine; control of infestation of large roundworms (<i>Ascaris suis</i>), nodular worms (<i>Oesophagostomum dentatum</i>) and whipworms (<i>Trichuris suis</i>); treatment of bacterial enteritis caused by <i>E. coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>P. multocida</i> susceptible to chlortetracycline..	Withdraw 15 d before slaughter.	
	Tylosin 10 to 100	Swine: Control of infestations of large roundworms (<i>Ascaris suis</i>), nodular worms (<i>Oesophagostomum dentatum</i>), and whipworms (<i>Trichuris suis</i>); growth promotion and feed efficiency.	As tylosin phosphate; withdraw 15 days prior to slaughter; feed continuously as follows: Animal wt.(lbs.): Up to 40.....20 to 100 ¹ 41 to 100.....20 to 40 ¹ 101 to market wt.....10 to 20 ¹	000986

¹ Amount of Tylosin (g/t).

(2) Hygromycin B may also be used in combination with:

- (i) Amprolium in accordance with § 558.55.
- (ii) Zoalene in accordance with § 558.680.

[41 FR 11000, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.274, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.295 Iodinated casein.

(a) *Approvals.* See 017762 in § 510.600(c) of this chapter.

(b) *NAS/NRC status.* The use of this drug is NAS/NRC reviewed and found effective. Applications for these uses need not include efficacy data as required by § 514.111 of this chapter but may require bioequivalency or safety data.

(c) *Conditions of use*—(1) *Ducks*—(i) *Amount per ton.* 100 to 200 grams.

(ii) *Indications for use.* For increased rate of weight gain and improved feathering in growing ducks.

(2) *Dairy cows*—(i) *Amount per pound.* ½ to 1½ grams per 100 lb of body weight.

(ii) *Indications for use.* For increased milk production in dairy cows.

(iii) *Limitations.* This drug is effective for limited periods of time, and the effectiveness is limited to the declining phase of lactation. Administration must be accompanied with increased feed intake; administration may increase heat sensitivity of the animal.

[45 FR 41631, June 20, 1980]

§ 558.300 Ivermectin.

(a) *Approvals.* (1) Type A medicated articles: 0.6 percent (2.72 grams per pound; 6 grams per kilogram) to 050604 in § 510.600(c) of this chapter, and

(2) Type B medicated feeds for ivermectin alone or with lincomycin. See § 558.4 of this chapter for maximum drug levels to 050604 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.344 of this chapter.

(c) [Reserved]

(d) *Conditions of use.* It is used in swine feed as follows:

(1) *Amount per ton.* For weaned, growing-finishing swine, feed 1.8 grams of ivermectin (to provide 0.1 milligram per kilogram of body weight per day). For adult and breeding swine, feed 1.8 to 11.8 grams of ivermectin (to provide 0.1 milligram per kilogram of body weight per day). For adult and breeding swine, may be top-dressed on daily ration for individual treatment at levels of 18.2 to 1180 grams (to provide 0.1 milligram per kilogram of body weight per day).

(i) *Indications for use.* For treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops strongylina*, adults; *Hyostrogylus rubidus*, adults and fourth-stage larvae; *Oesophagostomum* spp., adults and fourth-stage larvae); kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae); lungworms (*Metastrongylus* spp., adults); threadworms (*Strongyloides ransomi*, adults and somatic larvae, and prevention of transmission of infective larvae to piglets, via the colostrum or milk, when fed during gestation); lice (*Haematopinus suis*); and mange mites (*Sarcoptes scabiei* var. *suis*).

(ii) *Limitations.* For use in swine feed only. Feed as only feed for 7 consecutive days. Withdraw 5 days before slaughter. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) *Amount per ton.* 1.8 grams of ivermectin (to provide 0.1 milligram per kilogram of body weight per day) with 20 grams of lincomycin.

(i) *Indications for use.* For treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops strongylina*, adults; *Hyostrogylus rubidus*, adults and fourth-stage larvae; *Oesophagostomum* spp., adults and fourth-stage larvae), kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae), lungworms (*Metastrongylus* spp., adults), lice (*Haematopinus suis*), and mange mites (*Sarcoptes scabiei* var. *suis*). For increased rate of weight gain.

(ii) *Limitations.* For weaned, growing-finishing swine. Feed as only feed for 7 consecutive days. Withdraw 5 days before slaughter. A separate feed containing 20 grams per ton lincomycin may be continued. Not to be fed to swine that weigh more than 250 pounds. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(3) *Amount per ton.* 1.8 grams of ivermectin (to provide 0.1 milligram per kilogram of body weight per day) with 40 grams of lincomycin.

(i) *Indications for use.* For treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops strongylina*, adults; *Hyostrogylus rubidus*, adults and fourth-stage larvae; *Oesophagostomum* spp., adults and fourth-stage larvae), kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae), lungworms (*Metastrongylus* spp., adults), lice (*Haematopinus suis*), and mange mites (*Sarcoptes scabiei* var. *suis*). For control of swine dysentery. For use in swine on premises with a history of swine dysentery, but where symptoms have not yet occurred.

(ii) *Limitations.* For weaned, growing-finishing swine. Feed as only feed for 7 consecutive days. Withdraw 5 days before slaughter. A separate feed containing 40 grams per ton lincomycin may be continued. Not to be fed to swine that weigh more than 250 pounds. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

(4) *Amount per ton.* 1.8 grams of ivermectin (to provide 0.1 milligram per kilogram of body weight per day) with 100 grams of lincomycin.

(i) *Indications for use.* For treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops*

strongylina, adults; *Hyostrogylus rubidus*, adults and fourth-stage larvae; *Oesophagostomum* spp., adults and fourth-stage larvae), kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae), lungworms (*Metastrongylus* spp., adults), lice (*Haematopinus suis*), and mange mites (*Sarcoptes scabiei* var. *suis*). Treatment of swine dysentery.

(ii) *Limitations*. For weaned, growing-finishing swine. Feed as only feed for 7 consecutive days followed by a separate feed containing 100 grams per ton lincomycin for an additional 14 days to complete the lincomycin treatment. Withdraw 6 days before slaughter. Not to be fed to swine that weigh more than 250 pounds. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(5) *Amount per ton*. 1.8 grams of ivermectin (to provide 0.1 milligram per kilogram of body weight per day) with 200 grams of lincomycin.

(i) *Indications for use*. For treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops strongylina*, adults; *Hyostrogylus rubidus*, adults and fourth-stage larvae; *Oesophagostomum* spp., adults and fourth-stage larvae), kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae), lungworms (*Metastrongylus* spp., adults), lice (*Haematopinus suis*), and mange mites (*Sarcoptes scabiei* var. *suis*). For reduction in severity of swine mycoplasmal pneumonia caused by *Mycoplasma hyopneumoniae*.

(ii) *Limitations*. For weaned, growing-finishing swine. Feed as only feed for 7 consecutive days followed by a separate feed containing 200 grams per ton lincomycin for an additional 14 days to complete the lincomycin treatment. Withdraw 6 days before slaughter. Not to be fed to swine that weigh more than 250 pounds. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species

may result in severe gastrointestinal effects. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(6) *Amount per ton*. For weaned, growing-finishing pigs, feed 1.8 grams of ivermectin (to provide 0.1 milligram per kilogram of body weight per day), and 10 to 30 or 250 grams of bacitracin methylene disalicylate. For adult and breeding swine, feed 1.8 grams of ivermectin (to provide 0.1 milligram per kilogram of body weight per day), and 10 to 30 or 250 grams of bacitracin methylene disalicylate.

(i) *Indications for use*. For treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops strongylina*, adults; *Hyostrogylus rubidus*, adults and fourth-stage larvae; *Oesophagostomum* spp., adults and fourth-stage larvae); kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae); lungworms (*Metastrongylus* spp., adults); threadworms (*Strongyloides ransomi*, adults and somatic larvae, and prevention of transmission of infective larvae to piglets, via the colostrum or milk, when fed during gestation); lice (*Haematopinus suis*); and mange mites (*Sarcoptes scabiei* var. *suis*). For increased rate of weight gain and improved feed efficiency in growing and finishing swine. For control of clostridial enteritis caused by *Clostridium perfringens* in suckling piglets. For control of swine dysentery associated with *Treponema hyodysenteriae* on premises with a history of swine dysentery but where signs of disease have not yet occurred, or following an approved treatment of disease condition.

(ii) *Limitations*. For use in swine feed only. Feed as the only feed for 7 consecutive days. For weaned growing and finishing swine, feed bacitracin methylene disalicylate Type C medicated feed from weaning to market weight for increased rate of weight gain and improved feed efficiency. For pregnant sows, feed bacitracin methylene disalicylate to sows from 14 days before through 21 days after farrowing on premises with a history of clostridial

§ 558.305

scours. Withdraw ivermectin-containing feeds 5 days before slaughter.

[58 FR 58653, Nov. 3, 1993, as amended at 60 FR 39848, Aug. 4, 1995; 61 FR 32652, June 25, 1996; 62 FR 63271, Nov. 28, 1997; 63 FR 54353, Oct. 9, 1998; 64 FR 13342, Mar. 18, 1999]

§ 558.305 Laidlomycin propionate potassium.

(a) *Approvals.* Type A medicated articles: 50 grams per pound to 063238 in § 510.600(c) of this chapter.

(b) *Special considerations.* (1) Laidlomycin liquid Type B feeds may be manufactured from dry laidlomycin Type A articles. The liquid Type B feeds must have a pH of 6.0 to 8.0, dry matter of 62 to 75 percent, and bear appropriate mixing directions as follows:

(i) For liquid Type B feeds stored in recirculating tank systems: Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used.

(ii) For liquid Type B feeds stored in mechanical, air, or other agitation type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.

(2) The expiration date for the liquid Type B feed is 21 days after date of manufacture. The expiration date for the dry Type C feed made from the liquid Type B feed is 7 days after date of manufacture.

(c) [Reserved]

(d) *Conditions of use.* Used in cattle feed as follows:

(1) *Amount.* Laidlomycin propionate potassium, 5 grams per ton.

(i) *Indications for use.* For improved feed efficiency and increased rate of weight gain.

(ii) *Limitations.* Feed only to cattle being fed in confinement for slaughter. Feed continuously in a Type C feed at a rate of 30 to 75 milligrams per head per day.

(2) *Amount.* Laidlomycin propionate potassium, 5 to 10 grams per ton.

(i) *Indications for use.* For improved feed efficiency.

21 CFR Ch. I (4–1–99 Edition)

(ii) *Limitations.* Feed only to cattle being fed in confinement for slaughter. Feed continuously in a Type C feed at a rate of 30 to 150 milligrams per head per day.

(3) *Additional limitations.* (i) Do not allow horses or other equines access to feeds containing laidlomycin propionate potassium.

(ii) The safety of laidlomycin propionate potassium in unapproved species has not been established.

(iii) Not for use in animals intended for breeding.

[59 FR 18297, Apr. 18, 1994, as amended at 60 FR 53509, Oct. 16, 1995; 62 FR 9929, Mar. 5, 1997; 63 FR 27845, May 21, 1998]

§ 558.311 Lasalocid.

(a) *Specifications.* A minimum of 90 percent of lasalocid activity is derived from lasalocid A.

(b) *Approvals.* Type A medicated articles approved for sponsors identified in § 510.600(c) of this chapter for use as in paragraph (e) of this section as follows:

(1) 3.0, 3.3, 3.8, 4.0, 4.3, 4.4, 5.0, 5.1, 5.5, 5.7, 6.0, 6.3, 6.7, 7.2, 7.5, 8.0, 8.3, 10.0, 12.5, 15, 20, and 50 percent activity to No. 000004 for use as in paragraphs (e)(1)(i), (ii), (iii), (iv), and (x) of this section.

(2) 15 percent activity to No. 000069 as provided by No. 063238 for use as in paragraph (e)(1)(v) of this section.

(3) 15, 20, 33.1, and 50 percent activity to No. 063238 for use in cattle feeds as in paragraphs (e)(1)(vi), (vii), (ix), (xi), (xii), and (xv) of this section, and for use in sheep as in paragraph (e)(1)(viii) of this section.

(4) 15 percent activity to No. 063238 for use in Type C rabbit feeds as in paragraph (e)(1)(xvi) of this section and for use in ruminant free-choice Type C feeds as in paragraphs (e)(2) and (e)(3) of this section.

(5) 15 percent activity to 021930 (Type A article provided by 063238) for use in free-choice vitamin-mineral Type C cattle feeds as in paragraph (e)(1)(xi).

(6) 20 percent activity as a liquid Type A article to No. 063238 for use in cattle feeds as in paragraphs (e)(1)(vi), (e)(1)(vii), (e)(1)(ix), (e)(1)(xi), (e)(1)(xii), and (e)(3) of this section, and for use in sheep feeds as in paragraph (e)(1)(viii) of this section.

(7) 20 percent activity to No. 063238 for use as follows:

(i) Chukar partridges as in paragraph (e)(1)(xiii).

(ii) Turkeys as in paragraph (e)(1)(xiv).

(iii) Rabbits as in paragraph (e)(1)(xvi).

(c) *Related tolerance.* See § 556.347 of this chapter.

(d) *Special considerations.* (1) Type C cattle and sheep feeds may be manufactured from lasalocid liquid Type B feeds which have a pH of 4.0 to 8.0 and bear appropriate mixing directions as follows:

(i) For liquid Type B feeds stored in recirculating tank systems: Recirculate immediately prior to use for not less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used.

(ii) For liquid Type B feeds stored in mechanical, air, or other agitation-type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.

(2) A positionally stable lasalocid liquid Type B feed will not be subject to the requirements for mixing directions prescribed in paragraph (d)(1) of this section provided it has a pH of 4.0 to 8.0 and contains a suspending agent(s) sufficient to maintain a viscosity of not less than 300 centipoises per second for 3 months. Form FDA 1900 must indicate the pH and centipoises per second for such lasalocid liquid Type B feed.

(3) If a manufacturer is unable to meet the requirements of paragraph (d)(1) or (2) of this section, the manufacturer may secure approval of a positionally stable liquid Type B feed by (i) either filing a new animal drug application for the product or establishing a master file containing data to support the stability of its product; (ii) authorizing the agency to reference and rely upon the data in the master file to support approval of a supplemental new animal drug application to establish positional stability; and (iii) requesting the sponsor of an approved new animal drug application to file a supplement to provide for use of its lasalocid Type A article in the manu-

facture of the liquid Type B feed specified in the appropriate master file. If the data demonstrate the stability of the liquid Type B feed described in the master file, the supplement new animal drug application will be approved. Approval of the supplement will not be published in the FEDERAL REGISTER because such approval will not affect or alter conditions or use of the product in the new animal drug application or the regulation. The approval will, however, provide a basis for the individual liquid feed manufacturer to submit, and for the agency to approve, a medicated feed application under section 512(m) of the act for liquid Type B feed. A manufacturer who seeks to market a positionally unstable lasalocid liquid Type B feed with mixing directions different from the standard directions established in paragraph (d)(1) of this section may also follow this procedure.

(4) If adequate information is submitted to show that a particular liquid Type B feed containing lasalocid is stable outside the pH of 4.0 to 8.0, the pH restriction described in paragraphs (d)(1) and (2) of this section may be waived.

(5) Required label statements:

(i) For liquid Type B feed (cattle and sheep): Mix thoroughly with grain and/or roughage prior to feeding. Feeding undiluted, mixing errors, or inadequate mixing (recirculation or agitation) may result in an excess lasalocid concentration which could be fatal to cattle and sheep. Do not allow horses or other equines access to Type A articles or Type B feeds containing lasalocid as ingestion may be fatal. Safety of lasalocid for use in unapproved species has not been established.

(ii) For Type A articles or Type B feeds (cattle and sheep): Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle and sheep. Do not allow horses or other equines access to Type A articles or Type B feeds containing lasalocid as ingestion may be fatal. Safety of lasalocid for use in unapproved species has not been established.

(6) Lasalocid Type A medicated articles containing lasalocid dried fermentation residue are for use in cattle and sheep feed only.

§ 558.311

21 CFR Ch. I (4–1–99 Edition)

(e)(1) *Conditions of use.* It is used as follows:

Lasalocid sodium activity in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 68 (0.0075 pct) to 113 (0.0125 pct).		For the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> .	For broiler or fryer chickens only; feed continuously as the sole ration.	000004
(ii) 68 (0.0075 pct) to 113 (0.0125 pct).	Roxarsone 45.4 (0.005 pct).	Broiler or fryer chickens; for the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> and as an aid in the reduction of lesions due to <i>E. tenella</i> .	For broiler or fryer chickens only; feed continuously as the sole ration; as sole source of organic arsenic; withdraw 5 d before slaughter; roxarsone provided by Nos. 046573 and 011526 in § 510.600(c) of this chapter.	000004
	Roxarsone 45.4 plus bambermycins 1 (0.00011 pct).	For prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> ; as an aid in the reduction of lesions due to <i>E. tenella</i> ; and for increased rate of weight gain.	For broiler chickens only; feed continuously as sole ration; withdraw 5 days before slaughter; roxarsone provided by Nos. 046573 and 011526 in § 510.600(c) of this chapter, bambermycins provided by No. 012799.	000004
	Roxarsone 45.4 plus lincomycin 2.0.	For prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> ; as an aid in the reduction of lesions due to <i>E. tenella</i> ; and for increased rate of weight gain and improved feed efficiency.	For broiler chickens only; feed continuously as sole ration; withdraw 5 days before slaughter; roxarsone provided by Nos. 046573 and 011526 in § 510.600(c) of this chapter, lincomycin provided by No. 000009.	000004
	Roxarsone 45.4 plus bacitracin 10 to 25.	For prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> ; as an aid in the reduction of lesions due to <i>E. tenella</i> ; and for increased rate of weight gain.	For broiler or fryer chickens only; feed continuously as the sole ration; withdraw 5 days before slaughter; roxarsone provided by Nos. 046573 and 011526 in § 510.600(c) of this chapter, bacitracin methylene disalicylate provided by No. 046573 in § 510.600(c) of this chapter.	000004
	Roxarsone 45.4 plus bacitracin 10 or 30.	For prevention of coccidiosis caused by <i>E. tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> and <i>E. maxima</i> ; as an aid in the reduction of lesions due to <i>E. tenella</i> ; and for increased rate of weight gain (10 grams per ton) or improved feed efficiency (30 grams per ton).	For broiler chickens only; feed continuously as sole ration; withdraw 5 days before slaughter; roxarsone provided by Nos. 046573 and 011526 in § 510.600(c) of this chapter, bacitracin zinc provided by No. 000004.	000004
	Roxarsone 45.5 plus bacitracin methylene disalicylate 50.	Prevention of coccidiosis caused by <i>Eimeria necatrix</i> , <i>E. tenella</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> ; reduction of lesions due to <i>E. tenella</i> ; prevention of necrotic enteritis caused or complicated by <i>Clostridium</i> spp. or other susceptible organisms.	Feed continuously as sole ration; as sole source of organic arsenic; withdraw 5 days before slaughter.	046573
(iii) 68 (0.0075 pct)	Lincomycin 2 (0.00022 pct).	Broiler or fryer chickens; for the prevention of coccidiosis caused by <i>Eimeria mivati</i> , <i>E. brunetti</i> , <i>E. tenella</i> , <i>E. acervulina</i> , <i>E. maxima</i> , and <i>E. necatrix</i> ; for increased rate of weight gain and improved feed efficiency.	For broiler and fryer chickens only; feed continuously as sole ration; withdraw 5 d before slaughter; Type C feed must be used within 4 weeks of manufacture; as lincomycin hydrochloride monohydrate.	000004

Food and Drug Administration, HHS

§ 558.311

Lasalocid sodium activity in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(iv) 68 (0.0075 percent).	Bacitracin 10 to 50 ..	For prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> , and for increased rate of weight gain and improved feed efficiency.	For broiler or fryer chickens only; feed continuously as the sole ration; withdraw 3 days before slaughter bacitracin methylene disalicylate provided by No. 046573 in § 510.600(c) of this chapter.	000004
(v) 68 (0.0075 pct) to 113 (0.0125 pct).	Virginiamycin 20	For prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> , and for increased rate of weight gain and improved feed efficiency.	For broiler and fryer chickens only; feed continuously as sole ration; do not feed to laying chickens; lasalocid sodium provided by No. 063238 in § 510.600(c) of this chapter.	063238
(vi) 10 (0.0011 pct) to 30 (0.0033 pct).		Cattle; for improved feed efficiency.	In Type C feeds; for cattle fed in confinement for slaughter only; feed continuously in complete feed to provide not less than 100 mg nor more than 360 mg of lasalocid sodium activity per head per day.	000004
	Oxytetracycline 7.5 ..	Cattle; for improved feed efficiency and reduction of incidence and severity of liver abscesses.	In Type C feeds, for beef cattle fed in confinement for slaughter; feed continuously at 100 to 360 mg/head/day lasalocid and 75 mg/head/day oxytetracycline. As monoalkyl (C ₈ –C ₁₈) trimethyl ammonium oxytetracycline.	000004
(vii) 25 (0.0027 pct) to 30 (0.0033 pct).		Cattle; for improved feed efficiency and increased rate of weight gain.	In Type C feeds; for cattle fed in confinement for slaughter only; feed continuously in complete feed to provide not less than 250 mg nor more than 360 mg of lasalocid sodium activity per head per day.	000004
	Oxytetracycline 7.5 ..	Cattle; for improved feed efficiency, increased rate of weight gain, and reduction of incidence and severity of liver abscesses.	In Type C feeds, for beef cattle fed in confinement for slaughter; feed continuously at 250 to 360 mg/head/day lasalocid and 75 mg/head/day oxytetracycline. As monoalkyl (C ₈ –C ₁₈) trimethyl ammonium oxytetracycline.	000004
(viii) 20 (0.0022 pct) to 30 (0.0033 pct).		Sheep; for the prevention of coccidiosis caused by <i>Eimeria ovina</i> , <i>E. crandallis</i> , <i>E. ovinoidalis</i> (<i>E. ninakohlyakimovae</i>), <i>E. parva</i> , and <i>E. intricata</i> .	In Type C feeds; for sheep maintained in confinement; feed continuously in complete feed to provide not less than 15 mg nor more than 70 mg of lasalocid sodium activity per head per day depending on body weight.	000004
(ix)		Cattle; for increased rate of weight gain.	For pasture cattle (slaughter, stocker, feeder cattle, and dairy and beef replacement heifers) only; feed continuously at a rate of not less than 60 mg nor more than 200 mg of lasalocid per head per day when on pasture; the drug must be contained in at least 1 pound of feed.	000004
(x) 68 (0.0075 pct) to 113 (0.0125 pct).	Bacitracin 4 to 50	Broiler chickens; for prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> ; and for improved feed efficiency.	For broiler chickens only; feed continuously as the sole ration; withdraw 3 days before slaughter; bacitracin methylene disalicylate provided by No. 046573 in § 510.600(c) of this chapter.	000004

Lasalocid sodium activity in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(xi)		Cattle; for increased rate of weight gain.	For pasture cattle (slaughter, stocker, feeder cattle, and dairy and beef replacement heifers) only; feed continuously on a free-choice basis at a rate of not less than 60 mg nor more than 200 mg of lasalocid per head per day. Each use in a free-choice Type C feed must be the subject of an approved NADA or supplemental NADA as provided in § 510.455 of this chapter.	000004, 021930
(xii)		Cattle; for control of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> .	For cattle; hand feed at a rate of 1 mg of lasalocid per 2.2 pounds body weight per day to cattle weighing up to 800 pounds with a maximum of 360 mg of lasalocid per head per day.	000004
(xiii) 113 (0.0125 pct).		Chukar partridges; for prevention of coccidiosis caused by <i>Eimeria legionensis</i> .	Feed continuously as sole ration up to 8 weeks of age.	000004
(xiv) 68 (0.0075 pct) to 113 (0.0125 pct).		Growing turkeys; for prevention of coccidiosis caused by <i>E. meleagrimitis</i> , <i>E. gallopavonis</i> , and <i>E. adenoeides</i> .	Feed continuously as sole ration	000004
(xv)		Replacement calves; for control of coccidiosis caused by <i>E. bovis</i> and <i>E. zuernii</i> .	In milk replacer powder; hand feed at a rate of 1 mg of lasalocid per 2.2 lb body weight per day; include on labeling warning: "A withdrawal period has not been established for lasalocid in pre-ruminating calves. Do not use in calves to be processed for veal".	000004
(xvi) 113 (0.0125 pct).		Rabbits; for prevention of coccidiosis caused by <i>Eimeria stiedae</i> .	Feed continuously as sole ration up to 6 1/2 weeks of age.	000004

(2) It is used as a free-choice mineral Type C feed as follows:

(i) *Specifications.*

Ingredient	Per-cent	Inter-national feed No.
Defluorinated Phosphate (20.5 percent Calcium, 18.5 percent Phosphorus)	35.9	6–01–080
Sodium Chloride (Salt)	20.0	6–04–152
Calcium Carbonate (38 percent Calcium)	18.0	6–01–069
Cottonseed Meal	10.0	5–01–621
Potassium Chloride	3.0	6–03–755
Selenium Premix (0.02 percent Selenium) ¹	3.0	
Dried Cane Molasses	2.5	4–04–695
Magnesium Sulfate	1.7	6–02–758
Vitamin Premix ¹	1.4	
Magnesium Oxide	1.2	6–02–756
Potassium Sulfate	1.2	6–06–098
Trace Mineral Premix ¹	1.04	

Ingredient	Per-cent	Inter-national feed No.
Bovatec Premix (68 grams per pound) ...	1.06	

¹ Content of the vitamin and trace mineral premixes may be varied; however, they should be comparable to those used by the firm for other free-choice feeds. Formulation modifications require FDA approval prior to marketing. The amount of selenium and ethylenediamine dihydroiodide (EDDI) must comply with published regulations. For selenium (21 CFR 573.920): up to 120 parts per million in a mixture for free-choice feeding at a rate not to exceed an intake of 3 milligrams per head per day.

(ii) *Amount.* 1,440 grams per ton (0.16 percent).

(iii) *Indications for use.* Cattle, for increased rate of weight gain.

(iv) *Limitations.* For pasture cattle (slaughter, stocker, feeder cattle, and dairy and beef replacement heifers); feed continuously on a free-choice basis at a rate of 60 to 200 milligrams lasalocid per head per day; each use of this Type C free-choice feed must be

Food and Drug Administration, HHS

§ 558.315

the subject of an approved FD-1900 as provided in § 510.455 of this chapter.

(v) *Sponsor*. See No. 063238 in § 510.600(c) of this chapter.

(3) It is used as a ruminant free-choice liquid Type C feed as follows:

(i) *Specifications*.

Ingredient	Percent	International feed No.
Cane Molasses	55.167	4-13-241
Condensed Molasses Fermentation Solubles	24.0	N/A
50% Urea Solution (23% N)	12.0	N/A
Ammonium Polyphosphate Solution	1.0	6-08-42
Phosphoric Acid (54%)	3.0	6-03-707
Xanthan Gum	0.05	8-15-818
Water	4.0	N/A
Trace Mineral Mix ¹	0.5	N/A
Vitamin Premix ¹	0.2	N/A
Bovatec Liquid 20 (90.8 gm/lb) ²	0.083	N/A

¹Content of vitamin and trace mineral premixes may be varied. However, they should be comparable to those used for other free-choice liquid feeds. Formulation modifications require FDA approval prior to marketing. The amount of selenium and ethylenediamine dihydroiodide (EDDI) must comply with published regulations (for selenium see 21 CFR 573.920; for EDDI see 51 FR 11483, April 3, 1986).

²To provide 150 gm lasalocid per ton, use 1.652 lb (0.083%) of Bovatec Liquid 20 (90.8 gm/lb). If dry Bovatec-68 (68 gm/lb) is used, use 2.206 lbs per ton (0.111%), replacing molasses.

(ii) *Amount*. 150 grams per ton (.017 percent).

(iii) *Indications for use*. Cattle: for increased rate of weight gain.

(iv) *Limitations*. For pasture cattle (slaughter, stocker, feeder cattle, and dairy and beef replacement heifers). Feed continuously on a free-choice basis at a rate of 60 to 200 milligrams lasalocid per head per day. Each use of this free-choice liquid Type C feed must be the subject of an approved FD-1900 as in § 510.455 of this chapter.

(v) *Sponsor*. See No. 063238 in § 510.600(c) of this chapter.

(4) *Additional combinations*. Lasalocid may be used in accordance with the provisions of this section in combination as follows:

(i) Melengestrol acetate alone or in combination with tylosin in accordance with § 558.342.

(ii) [Reserved]

[41 FR 44382, Oct. 8, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.311, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.315 Levamisole hydrochloride (equivalent).

(a) *Approvals*. Type A medicated articles: 227 grams per pound to 043781 in § 510.600(c) of this chapter.

(b) *Related tolerances*. See § 556.350 of this chapter.

(c) *Conditions of use*. It is used in Type C medicated feed as follows:

(1) *Cattle*—(i) *Amount per pound*. 0.36–3.6 grams (0.08–0.8 percent).

(ii) *Indications for use*. Treatment of the following gastrointestinal worms and lung worm infections; stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*), and lungworms (*Dictyocaulus*).

(iii) *Limitations*. Administer medicated feed mixed thoroughly in one half the usual amount of morning feed; the medicated feed mix should be consumed within 6 hours; when medicated feed is consumed resume normal feeding; medicated feed is to be fed at the rate of 0.36 gram of levamisole hydrochloride (equivalent) per 100 lb. of body weight; conditions of constant helminth exposure may require re-treatment within 2 to 4 weeks after the first treatment; do not slaughter for food within 48 hours of treatment; consult veterinarian before using in severely debilitated animals; do not administer to dairy animals of breeding age; for use in pelleted or meal feeds only; the label shall bear the caution, “Muzzle foam may be observed. However, this reaction will disappear within a few hours. If this condition persists, a veterinarian should be consulted. Follow recommended dosage carefully.”

(2) *Swine*—(i) *Amount per pound*. 0.36 grams (0.08 percent).

§ 558.325

21 CFR Ch. I (4–1–99 Edition)

(ii) *Indications for use.* Treatment of the following nematode infections: large roundworms (*Ascaris suum*), nodular worms (*Oesophagostomum* spp.), lungworms (*Metastrongylus* spp.), intestinal threadworms (*Strongyloides ransomi*), swine kidney worms (*Stephanurus dentatus*).

(iii) *Limitations.* It is recommended that regular feed be withheld overnight and worming feed administered the following morning; dilute supplement with nonmedicated feed as directed; feed the equivalent of 1 lb. of 0.08 percent worming feed per 100 lbs. of body weight of pigs to be treated; may be fed as sole feed or thoroughly mixed with 1 to 2 parts of regular feed prior to feeding; when medicated feed is consumed, resume normal feeding. Pigs maintained under conditions of constant worm exposure may require retreatment within 4 to 5 weeks after the first treatment due to reinfection; do not slaughter for food within 72 hours of treatment; the label shall bear the caution, “Excessive salivation or muzzle foam may be observed. This reaction is occasionally seen and will disappear in a short time after medication. If pigs are infected with mature lungworms, coughing and vomiting may be observed soon after medicated feed is consumed. This reaction is due to the expulsion of worms from the lungs and will be over in several hours.”

[40 FR 13959, Mar. 27, 1975, as amended at 43 FR 11176, Mar. 17, 1978; 43 FR 39351, Sept. 5, 1978; 43 FR 16013, Mar. 16, 1979; 51 FR 7398, Mar. 3, 1986]

§ 558.325 Lincomycin.

(a) *Approvals.* Type A articles and Type B feeds approved for sponsors in § 510.600(c) of this chapter for specific uses as in paragraph (c) of this section as follows:

(1) No. 000009: (i) 4 grams per pound as in paragraphs (c) (1) and (3) of this section.

(ii) 20 grams per pound as in paragraphs (c) (1) through (3) of this section.

(iii) 50 grams per pound as in paragraphs (c) (1) and (2) of this section.

(iv) 10 grams per pound as in paragraphs (c)(1) and (2) (i) through (iv) of this section.

(2)–(4) [Reserved]

(5) No. 043733 for 8 and 20 grams per pound as in paragraphs (c)(2) (i) through (iii) of this section.

(6)–(12) [Reserved]

(13) No. 017800 for 2.5 and 8 grams per pound as in paragraphs (c)(2) (i) through (iv) of this section.

(14)–(15) [Reserved]

(b) *Related tolerances in edible products.* See § 556.360 of this chapter.

(c) [Reserved]

(d) *Conditions of use—(1) Broilers:*

(i) *Amount per ton.* 2 to 4 grams.

(a) *Indications for use.* For increase in rate of weight gain and improved feed efficiency.

(b) *Limitations.* As lincomycin hydrochloride monohydrate.

(ii) *Amount per ton.* 2 grams.

(a) *Indications for use.* For control of necrotic enteritis caused by *Clostridium* spp. or other susceptible organisms.

(b) *Limitations.* As lincomycin hydrochloride monohydrate.

(2) *Swine—(i) Amount per ton.* 40 grams.

(a) *Indications for use.* For control of swine dysentery.

(b) *Limitations.* Feed as sole ration; for use in swine on premises with a history of swine dysentery but where symptoms have not yet occurred; not to be fed to swine that weigh more than 250 pounds.

(ii) *Amount per ton.* 100 grams; 40 grams.

(a) *Indications for use.* For treatment and control of swine dysentery.

(b) *Limitations.* Feed 100 grams per ton for 3 weeks or until signs of disease disappear, followed by 40 grams per ton; feed as sole ration; not to be fed to swine that weigh more than 250 pounds.

(iii) *Amount per ton.* 100 grams.

(a) *Indications for use.* For treatment of swine dysentery.

(b) *Limitations.* Feed as sole ration for 3 weeks or until signs of disease disappear; not to be fed to swine that weigh more than 250 pounds.

(iv) *Amount per ton.* 200 grams.

(a) *Indications for use.* For reduction in the severity of swine mycoplasmal pneumonia caused by *Mycoplasma hyopneumoniae*.

(b) *Limitations.* Feed as sole ration for 21 days; not to be fed to swine that weigh more than 250 pounds.

(v) *Amount per ton.* 20 grams.

(a) *Indications for use.* For increased rate of weight gain in growing-finishing swine.

(b) *Limitations.* Feed as sole ration; not for use in swine weighing over 250 pounds.

(3) Lincomycin may also be used for broilers in combination with:

(i) Amprolium, ethopabate, and roxarsone in accordance with §§ 558.58 and 558.530.

(ii) Amprolium and ethopabate in accordance with § 558.58.

(iii) Clopidol in accordance with § 558.175.

(iv) [Reserved]

(v) Decoquinat in accordance with § 558.195.

(vi) Zoalene in accordance with § 558.680.

(vii) Monensin in accordance with § 558.355.

(viii) Robenidine hydrochloride in accordance with § 558.515.

(ix) Roxarsone, monensin sodium in accordance with §§ 558.355 and 558.530.

(x) Lasalocid sodium in accordance with § 558.311.

(xi) Nicarbazine and roxarsone as in § 558.366.

(xii) Nicarbazine with or without narasin as in § 558.366.

(xiii) Lasalocid sodium and roxarsone as in § 558.311.

(xiv) Halofuginone in accordance with § 558.265.

(xv) Salinomycin with or without roxarsone as in § 558.550.

(4) Lincomycin may also be used for swine in combination with:

(i) Pyrantel tartrate as in § 558.485.

(ii) Fenbendazole as provided in § 558.258.

(iii) Ivermectin as in § 558.300.

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.325, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.340 Maduramicin ammonium.

(a) *Approvals.* Type A medicated articles: 4.54 grams per pound to 063238 in § 510.600(c) of this chapter.

(b) *Tolerances.* See § 556.375 of this chapter.

(c) *Conditions of use.* (1) *Amount.* 4.54 to 5.45 grams per ton (5 to 6 parts per million) (1 to 1.2 pounds per ton).

(i) *Indications for use.* For prevention of coccidiosis caused by *Eimeria acervulina*, *E. tenella*, *E. brunetti*, *E. maxima*, *E. necatrix*, and *E. mivati*.

(ii) *Limitations.* For broiler chickens only. Feed continuously as sole ration. Do not feed to laying hens. Withdraw 5 days before slaughter.

(2) [Reserved]

[54 FR 5229, Feb. 2, 1989, as amended at 54 FR 26732, June 26, 1989; 54 FR 32635, Aug. 9, 1989; 54 FR 33885, Aug. 17, 1989; 55 FR 23, Jan. 2, 1990; 55 FR 8460, Mar. 8, 1990; 55 FR 49616, Nov. 30, 1990; 59 FR 8134, Feb. 18, 1994; 61 FR 18082, Apr. 24, 1996; 63 FR 27845, May 21, 1998]

§ 558.342 Melengestrol acetate.

(a) *Approvals.* Dry Type A medicated articles containing 100 or 200 milligrams per pound or liquid Type A article containing 500 milligrams per pound to 000009 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.380 of this chapter.

(c) *Special considerations.* (1) Type B or C medicated feeds may be manufactured from melengestrol acetate liquid Type A articles or Type B or C medicated feeds which have a pH of 4.0 to 8.0 and bear appropriate mixing directions as follows:

(i) For liquid Type B or C feeds stored in recirculating tank systems: Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used.

(ii) For liquid Type B or C feeds stored in mechanical, air, or other agitation type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.

(2) A positionally stable melengestrol acetate liquid Type B or C feed will not be subject to the requirements for mixing directions prescribed in paragraphs (c)(1) of this section provided it has a pH of 4.0 to 8.0 and contains a suspending agent(s) sufficient to maintain a viscosity of not less than 300 centipoises per second for 3 months.

(d) *Conditions of use.* It is used for heifers as follows:

(1) *Amount.* Melengestrol acetate, 0.25 to 0.50 milligram per head per day.

(i) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and suppression of estrus (heat).

(ii) *Limitations:* Heifers being fed for slaughter; administer 0.5 to 2.0 pounds per head per day of medicated feed containing 0.125 to 1.0 milligram of melengestrol acetate per pound to provide 0.25 to 0.50 milligram of melengestrol acetate per head per day. Melengestrol acetate as provided by No. 000009 in §510.600(c) of this chapter.

(2) *Amount:* Melengestrol acetate, 0.25 to 0.40 milligram per head per day, plus monensin (as monensin sodium), 50 to 360 milligrams per head per day.

(i) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and suppression of estrus (heat).

(ii) *Limitations:* Heifers being fed in confinement for slaughter. Add at the rate of 0.5 to 2.0 pounds per head per day a medicated feed (liquid or dry) containing 0.125 to 0.80 milligram of melengestrol acetate per pound to a feed containing 5 to 30 grams of monensin per ton; or, add at the rate of 0.5 to 2.0 pounds per head per day a medicated feed (liquid or dry) containing 0.125 to 0.80 milligram of melengestrol acetate plus 25 to 720 milligrams of monensin per pound to a nonmedicated feed to provide 0.25 to 0.40 milligram of melengestrol acetate and 50 to 360 milligrams of monensin per head per day. The liquid medicated feeds must be manufactured in accordance with §558.355(f)(3)(i). Melengestrol acetate and monensin as provided by Nos. 000009 and 000986, respectively, in §510.600(c) of this chapter.

(3) *Amount.* Melengestrol acetate, 0.25 to 0.50 milligram per head per day in combination with lasalocid (as lasalocid sodium), 100 to 360 milligrams per head per day.

(i) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and suppression of estrus (heat).

(ii) *Limitations.* Heifers being fed in confinement for slaughter. The liquid medicated feeds are required to be manufactured in accordance with §558.311(d). Add at the rate of 0.5 to 2.0 pounds per head per day a medicated

feed (liquid or dry) containing 0.125 to 1.0 milligram of melengestrol acetate per pound to a feed containing 10 to 30 grams of lasalocid per ton; or, add at the rate of 0.5 to 2.0 pounds per head per day a medicated feed (liquid or dry) containing 0.125 to 1.0 milligram of melengestrol acetate plus 50 to 720 milligrams of lasalocid per pound to a ration of nonmedicated feed to provide 0.25 to 0.5 milligram of melengestrol acetate and 100 to 360 milligrams of lasalocid per head per day. Melengestrol acetate and lasalocid as provided by Nos. 000009 and 063238, respectively, in §510.600(c) of this chapter.

(4) *Amount.* Melengestrol acetate, 0.25 to 0.50 milligram per head per day, plus tylosin (as tylosin phosphate), 90 milligrams per head per day.

(i) *Indications for use.* For increased rate of weight gain, improved feed efficiency, suppression of estrus (heat), and reduced incidence of liver abscesses.

(ii) *Limitations.* Heifers being fed in confinement for slaughter. Melengestrol acetate and tylosin as provided by Nos. 000009 and 000986, respectively, in §510.600(c) of this chapter. To administer 0.25 to 0.50 milligram of melengestrol acetate with 90 milligrams of tylosin per head per day:

(A) Add 0.5 to 2.0 pounds per head per day of a liquid or dry medicated feed containing 0.125 to 1.0 milligram of melengestrol acetate per pound to a medicated feed containing 8 to 10 grams of tylosin per ton; or

(B) Add 0.5 to 2.0 pounds per head per day of a liquid or dry medicated feed containing 0.125 to 1.0 milligram of melengestrol acetate per pound to 4.5 to 18 pounds of a dry medicated feed containing 10 to 40 grams of tylosin per ton; or

(C) Add 0.5 to 2.0 pounds per head per day of a dry medicated feed containing 0.125 to 1.0 milligram of melengestrol acetate (from a dry Type A article) plus 45 to 180 milligrams of tylosin per pound to a ration of nonmedicated feed.

(5) *Amount.* Melengestrol acetate, 0.25 to 0.50 milligram per head per day, plus monensin (as monensin sodium) at 50 to 360 milligrams per head per day, plus

tylosin (as tylosin phosphate), 90 milligrams per head per day.

(i) *Indications for use.* For increased rate of weight gain, improved feed efficiency, suppression of estrus (heat), and reduced incidence of liver abscesses.

(ii) *Limitations.* Heifers being fed in confinement for slaughter. The liquid medicated feeds are required to be manufactured in accordance with § 558.355(f)(3)(i). Melengestrol acetate as provided by No. 000009 and monensin and tylosin as provided by No. 000986 in § 510.600(c) of this chapter. To administer 0.25 to 0.50 milligram of melengestrol acetate to 50 to 360 milligrams of monensin plus 90 milligrams of tylosin per head per day:

(A) Add 0.5 to 2.0 pounds per head per day of a liquid or dry medicated feed containing 0.125 to 1.0 milligram of melengestrol acetate per pound to a medicated feed containing 5 to 30 grams of monensin and 8 to 10 grams of tylosin per ton; or

(B) Add 0.5 to 2.0 pounds per head per day of a liquid or dry medicated feed containing 0.125 to 1.0 milligram of melengestrol acetate plus 25 to 720 milligrams of monensin per pound to 4.5 to 18 pounds of a dry medicated feed containing 10 to 40 grams of tylosin per ton; or

(C) Add 0.5 to 2.0 pounds per head per day of a dry medicated feed containing 0.125 to 1.0 milligram melengestrol acetate (from a dry Type A article), 25 to 600 milligrams of monensin, and 45 to 180 milligrams of tylosin per pound to a ration of nonmedicated feed.

(6) *Amount.* Melengestrol acetate, 0.25 to 0.50 milligram per head per day, plus lasalocid (as lasalocid sodium), 100 to 360 milligrams per head per day, and tylosin (as tylosin phosphate), 90 milligrams per head per day.

(i) *Indications for use.* For increased rate of weight gain, improved feed efficiency, suppression of estrus (heat), and reduced incidence of liver abscesses.

(ii) *Limitations.* Heifers being fed in confinement for slaughter. The liquid medicated feeds are required to be manufactured in accordance with § 558.311(d). Lasalocid, melengestrol acetate, and tylosin as provided by Nos. 000009, 000986, and 063238 respectively,

in § 510.600(c) of this chapter. To administer 0.25 to 0.50 milligram of melengestrol acetate plus 100 to 360 milligrams of lasalocid plus 90 milligrams of tylosin per head per day:

(A) Add 0.5 to 2.0 pounds per head per day of a liquid or dry medicated feed containing 0.125 to 1.0 milligram of melengestrol acetate per pound to a medicated feed containing 10 to 30 grams of lasalocid and 8 to 10 grams of tylosin per ton; or

(B) Add 0.5 to 2.0 pounds per head per day of a liquid or dry medicated feed containing 0.125 to 1.0 milligram of melengestrol acetate plus 50 to 720 milligrams of lasalocid per pound to 4.5 to 18 pounds of a dry medicated feed containing 10 to 40 grams of tylosin per ton; or

(C) Add 0.5 to 2.0 pounds per head per day of a dry pelleted medicated feed containing 0.125 to 1.0 milligram of melengestrol acetate (from a dry Type A article), 50 to 720 milligrams of lasalocid, and 45 to 180 milligrams of tylosin per pound to a ration of non-medicated feed.

(7) *Amount.* 0.5 milligram per head per day.

(i) *Indications for use.* For suppression of estrus (heat).

(ii) *Limitation.* Heifers intended for breeding. Do not exceed 24 days of feeding. Administer 0.5 to 2.0 pounds per head per day of Type C feed containing 0.25 to 1.0 milligram of melengestrol acetate per pound to provide 0.5 milligram of melengestrol acetate per head per day. Melengestrol acetate as provided by No. 000009 in § 510.600(c) of this chapter.

(8) *Amount.* Melengestrol acetate, 0.25 to 0.5 milligram per head per day, plus oxytetracycline, 75 milligrams per head per day.

(i) *Indications for use.* For increased rate of weight gain, improved feed efficiency, suppression of estrus (heat), and reduction of liver condemnation due to liver abscesses.

(ii) *Limitations.* Heifers fed in confinement for slaughter. Add at the rate of 0.5 to 2.0 pounds per head per day a medicated feed (liquid or dry) containing 0.125 to 1.0 milligram of melengestrol acetate per pound to a feed containing 6 to 10 grams of oxytetracycline per ton; or add at the rate of

§ 558.348

0.5 to 2.0 pounds per head per day a dry medicated feed containing 0.125 to 1.0 milligram of melengestrol acetate plus 37.5 to 150 milligrams of oxytetracycline per pound to provide 0.25 to 0.5 milligram of melengestrol acetate and 75 milligrams of oxytetracycline per head per day. Liquid melengestrol acetate may not be mixed with oxytetracycline in a common liquid feed supplement. Melengestrol acetate as provided by 000009, oxytetracycline by 000069, in § 510.600(c) of this chapter.

[42 FR 28535, June 3, 1977, as amended at 46 FR 18967, Mar. 27, 1981; 49 FR 39842, Oct. 11, 1984; 50 FR 1843, Jan. 14, 1985; 50 FR 16228, Apr. 25, 1985; 51 FR 7398, Mar. 3, 1986; 51 FR 36222, Oct. 9, 1986; 52 FR 2685, Jan. 26, 1987; 52 FR 33804, Sept. 8, 1987; 53 FR 27959, July 26, 1988; 55 FR 31828, Aug. 6, 1990; 57 FR 5053, Feb. 12, 1992; 59 FR 41241, Aug. 11, 1994; 60 FR 11028, Mar. 1, 1995; 62 FR 8373, Feb. 25, 1997; 62 FR 14304, Mar. 26, 1997; 63 FR 27845, May 21, 1998; 63 FR 41191, Aug. 3, 1998; 63 FR 66018, Dec. 1, 1998]

§ 558.348 Mibolerone.

(a) *Approvals.* To No. 000009 in § 510.600(c) of this chapter for a canned dog food, each 6½ ounce can containing 30 or 60 micrograms of mibolerone.

(b) *Conditions of use—(1) Amount.* 30 micrograms for animals weighing up to 25 pounds; 60 micrograms for animals weighing 26 to 50 pounds; 120 micrograms for animals weighing 51 to 100 pounds; 180 micrograms for animals weighing over 100 pounds, or German Shepherds or German Shepherd mix weighing 30 to 80 pounds.

(2) *Indications for use.* For the prevention of estrus (heat) in adult female dogs not intended primarily for breeding purposes.

(3) *Limitations.* Administer daily at least 30 days before expected initiation of heat and continue as long as desired, but for not more than 12 months. Mibolerone should not be used in bitches before first estrous period or in purebred Bedlington terriers. It is not intended for animals being used primarily for breeding purposes. Use orally in adult female dogs only. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[47 FR 6617, Feb. 16, 1982]

21 CFR Ch. I (4–1–99 Edition)

§ 558.355 Monensin.

(a) *Specifications.* Monensin, as the base or the sodium salt, contains a minimum of 90 percent monensin activity derived from monensin A and a minimum of 95 percent derived from monensin A plus B. Using thin layer chromatography, the R_f value must be comparable to a reference standard (the R_f value is the distance the spots travel from the starting line divided by the distance the solvent front travels from the starting line), or, using High Performance Liquid Chromatography, the factor distribution of monensin Factor A or B is calculated as the percentage of total biopotency of all peaks. The loss on drying is not more than 10 percent when dried in vacuum at 60 °C for 2 hours.

(b) *Approvals.* Approvals for Type A medicated articles containing the specified levels of monensin activity granted to firms identified by sponsor numbers in § 510.600(c) of this chapter for the conditions of use indicated in paragraph (f) of this section are as follows:

(1) To 000986: 36.3 (for export only), 44, 45, or 60 grams per pound, paragraphs (f) (1)(i) and (4) of this section.

(2) To 000986: 110 grams per lb., paragraphs (f)(1) (i), (iii), (iv), (v), (ix), and (x).

(3) To 000986: 44 grams per lb. with 18 grams per lb. of roxarsone, 110 grams per lb. with 45 grams per lb. of roxarsone, paragraph (f)(1)(ii).

(4) To 000986: 45 and 60 grams per pound, as monensin sodium, paragraph (f)(2) of this section.

(5) To 000069: 45 and 60 grams per pound, as monensin sodium provided by No. 000986, paragraphs (f)(1)(xiii), (xx), and (xxi) of this section.

(6) To 000986: 45 and 60 grams per pound, as monensin sodium, paragraph (f)(5) of this section.

(7) To 000986: 20, 30, 45, 60, 80, and 90.7 grams per pound, as monensin sodium, paragraph (f)(3) of this section.

(8) To 063238: 45 and 60 grams per pound, as monensin sodium provided by No. 000986, paragraph (f)(1)(xiv) of this section.

(9) To 063238: 45 and 60 grams per pound, as monensin sodium provided by No. 000986, paragraphs (f)(1)(xv) and (xvi) of this section.

(10) To 012799: 45 and 60 grams per pound, as monensin sodium, paragraph (f)(1)(xvii) of this section.

(11) To 046573: 45 and 60 grams per pound, as monensin sodium provided by No. 000986, paragraphs (f)(1)(xiv), (xviii), (xix), (xxiii), (xxiv), (xxv), (xxvi), and (xxvii) of this section.

(12) To 000069: 45 and 60 grams per pound, as monensin sodium provided by No. 000986, paragraph (f)(1)(xxii) of this section.

(13) To 021930: 60 and 80 grams per pound, paragraph (f)(3)(v) of this section.

(14) To 000986: 60, 80, and 90.7 grams per pound, as monensin sodium, paragraph (f)(6) of this section.

(c) [Reserved]

(d) *Special considerations.* (1) Type C chicken feed containing monensin as the mycelial cake shall bear an expiration date of 90 days after its date of manufacture.

(2) Type C cattle feeds containing 30 grams or less monensin sodium per ton shall bear an expiration date of 30 days after its date of manufacture.

(3) Type C goat feeds shall bear an expiration date of 30 days after date of manufacture.

(4) [Reserved]

(5) Liquid Type B feeds shall bear an expiration date of 8 weeks after its date of manufacture.

(6) The labeling of all formulations containing monensin shall bear the following caution statement: Do not allow horses or other equines access to formulations containing monensin. Ingestion of monensin by equines has been fatal.

(7) The labeling of all Type A articles and Type B feeds (liquid and dry) containing monensin intended for use in cattle shall bear, in addition to the caution statement in paragraph (d)(6) of this section, the following caution statements:

(i) Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions;

(ii) Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle.

(iii) Must be thoroughly mixed in feeds before use;

(iv) Do not feed undiluted;

(v) Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

(vi) Do not feed to lactating dairy cows.

(8) The labeling of complete feeds containing monensin intended for use in cattle shall bear the caution statements specified in paragraphs (d) (6) and (7) (i) and (v) of this section.

(9) The labeling of Type A articles containing monensin intended for use in chickens shall bear the caution statements specified in paragraphs (d) (6) and (7) (iii) and (iv) of this section.

(10) The labeling of all Type A articles and Type B feeds (liquid and dry) containing monensin intended for use in goats shall bear, in addition to the caution statement in paragraph (d)(6) of this section, the following caution statements:

(i) Monensin medicated goat feed is safe for use in goats only. Consumption by unapproved species may result in toxic reactions.

(ii) Feeding undiluted or mixing errors resulting in high concentrations of monensin could be fatal to goats.

(iii) Must be thoroughly mixed in feeds before use.

(iv) Do not feed undiluted.

(v) Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

(11) The labeling of complete feeds containing monensin intended for use in goats shall bear the caution statements specified in paragraphs (d)(6) and (d)(10) (i) and (v) of this section.

(e) *Related tolerances.* See § 556.420 of this chapter.

(f) *Conditions of use.* It is used as follows:

(1) *Broiler chickens*—(i) *Amount per ton.* Monensin, 90–110 grams.

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as the sole ration; in the absence of coccidiosis, the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain; as monensin or monensin sodium.

(ii) *Amount per ton.* Monensin, 90–110 grams, plus roxarsone 45.4 grams (0.005 percent).

(a) *Indications for use.* Growth promotion and feed efficiency, improving pigmentation; as an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati* and *E. maxima*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as the sole ration; withdraw 5 days before slaughter; as sole source of organic arsenic; as monensin or monensin sodium.

(iii) *Amount per ton.* Monensin, 90–110 grams plus bacitracin, 5–25 grams.

(a) *Indications for use.* For increased rate of weight gain and improved feed efficiency; as an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as sole ration; in the absence of coccidiosis, the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain; as bacitracin methylene disalicylate provided by No. 046573 in § 510.600(c) of this chapter; as monensin sodium.

(iv) *Amount per ton.* Monensin, 90–110 grams plus bacitracin, 10 grams.

(a) *Indications for use.* For increased rate of weight gain and improved feed efficiency; as an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as sole ration; in the absence of coccidiosis, the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain; as zinc bacitracin provided by No. 063238 in § 510.600(c) of this chapter; as monensin sodium.

(v) *Amount per ton.* Monensin, 90–110 grams plus bacitracin, 10–30 grams.

(a) *Indications for use.* For improved feed efficiency; as an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as sole ration; in the absence of coccidiosis, the use of monensin with no withdrawal

period may limit feed intake resulting in reduced weight gain; as zinc bacitracin provided by No. 063238 in § 510.600(c) of this chapter; as monensin sodium.

(vi) *Amount per ton.* Monensin, 90 to 110 grams plus bambermycins, 1 to 2 grams. See § 558.95(b)(1)(vi).

(vii) *Amount per ton.* Monensin, 90 to 110 grams plus bambermycins, 1 gram plus roxarsone, 22.7 to 45.4 grams (.0025 to .005 percent). See § 558.95(b)(1)(vii).

(viii) *Amount per ton.* Monensin, 90 to 110 grams plus oxytetracycline, 200 grams.

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*; and for the control of complicated chronic respiratory disease (CRD or air-sac infection) caused by *Mycoplasma gallisepticum* and *Escherichia coli*.

(b) *Limitations.* In the absence of coccidiosis, the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain; do not feed to laying chickens; feed continuously as sole ration; as monensin sodium.

(ix) *Amount per ton.* Monensin, 90–110 grams plus lincomycin, 2 grams.

(a) *Indications for use.* For increase in rate of weight gain and improved feed efficiency; as an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(b) *Limitations.* Do not feed to laying chickens; to be fed as a sole ration; in the absence of coccidiosis, the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain; as monensin sodium.

(x) *Amount per ton.* Monensin, 90–110 grams plus lincomycin, 2 grams and roxarsone, 15–45 grams.

(a) *Indications for use.* For increase in rate of weight gain; as an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as the sole ration; withdraw 5 days before slaughter; as sole source of organic arsenic; as roxarsone provided by No. 046573,

§ 510.600(c) of this chapter; as monensin sodium provided by No. 000986, § 510.600(c) of this chapter; as lincomycin provided by No. 000009, § 510.600(c) of this chapter; as a combination provided by No. 000009, § 510.600(c) of this chapter.

(xi) *Amount per ton.* Monensin, 90 to 110 grams, plus lincomycin, 2 grams and roxarsone, 15 to 30 grams.

(a) *Indications for use.* For increase in rate of weight gain, improved feed efficiency, improved pigmentation, and as an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati* and *E. maxima*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as the sole ration; withdraw 5 days before slaughter; as sole source of organic arsenic; as roxarsone provided by No. 046573 in § 510.600(c) of this chapter; as monensin sodium provided by No. 000986 in § 510.600(c) of this chapter; as lincomycin provided by No. 000009 in § 510.600(c) of this chapter; as a combination provided by No. 000009 in § 510.600(c) of this chapter.

(xii) *Amount per ton.* Monensin, 90 to 110 grams, plus bacitracin methylene disalicylate, 10 to 25 grams, and roxarsone, 11.3 to 45.4 grams.

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. maxima*, and *E. mivati*; for increased rate of weight gain and for improved feed efficiency.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as sole ration; withdraw 5 days before slaughter; as sole source of organic arsenic; as monensin sodium provided by No. 000986 in § 510.600 of this chapter; as bacitracin methylene disalicylate provided by No. 046573 in § 510.600 of this chapter; as roxarsone provided by No. 011526 or 046573 in § 510.600 of this chapter.

(xiii) *Amount per ton.* Monensin, 90 to 110 grams, plus 5 grams virginiamycin.

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. maxima*, and *E. mivati*; for increased rate of weight gain and improved feed efficiency.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as sole ration; as monensin sodium provided by No. 000986 in § 510.600 of this chapter; virginiamycin provided by No. 000007 in § 510.600 of this chapter.

(xiv) *Amount per ton.* Monensin, 90 to 110 grams, plus 500 grams chlortetracycline.

(a) *Indications for use.* As an aid in the reduction of mortality due to *Escherichia coli* infections susceptible to such treatment. As an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(b) *Limitations.* Do not feed to laying chickens; feed for 5 days as the sole ration; withdraw 24 hours before slaughter; in the absence of coccidiosis, the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain; not to be fed continuously for more than 5 days; as monensin sodium; as chlortetracycline hydrochloride provided by Nos. 046573 and 063238 in § 510.600(c) of this chapter.

(xv) *Amount per ton.* Monensin, 90 to 110 grams, plus bacitracin zinc, 10 grams, and roxarsone, 15 grams (0.0017 percent).

(a) *Indications for use.* For increase in rate of weight gain; for the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as the sole ration; withdraw 5 days before slaughter; as sole source of organic arsenic; feed must be used within 4 weeks of manufacture; as monensin sodium; as bacitracin zinc provided by Nos. 063238 and 046573 in § 510.600(c) of this chapter; as roxarsone provided by No. 046573 in § 510.600(c) of this chapter.

(xvi) *Amount per ton.* Monensin, 90 to 110 grams, plus bacitracin zinc, 4 to 50 grams, and roxarsone, 15 to 45.4 grams (0.0017 percent to 0.005 percent).

(a) *Indications for use.* For improved feed efficiency; for improved pigmentation by enhancing carotenoid and xanthophyll utilization; for the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as the sole

ration; withdraw 5 days before slaughter; as sole source of organic arsenic; feed must be used within 4 weeks of manufacture; as monensin sodium; as bacitracin zinc provided by Nos. 046573 and 063238 in § 510.600(c) of this chapter; as roxarsone provided by No. 046573 in § 510.600(c) of this chapter.

(xvii) *Amount per ton.* Monensin, 90 to 110 grams plus bambermycins, 1 to 2 grams plus roxarsone, 22.7 to 45.4 grams (0.0025 to 0.005 percent). See § 558.95(b)(1)(xiii) of this chapter.

(xviii) *Amount per ton.* Monensin, 90 to 110 grams, plus bacitracin methylene disalicylate, 50 grams, and roxarsone, 22.7 to 34.0 grams (0.0025 percent to .00375 percent).

(a) *Indications for use.* For increase in rate of weight gain and improved feed efficiency; as an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; as an aid in the prevention of necrotic enteritis caused or complicated by *Clostridium spp* or other organisms susceptible to bacitracin methylene disalicylate.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as the sole ration; withdraw 5 days before slaughter; as sole source of organic arsenic; as monensin sodium provided by No. 000986 in § 510.600(c) of this chapter; as bacitracin methylene disalicylate provided by No. 046573 in § 510.600(c) of this chapter; as roxarsone provided by No. 046573 in § 510.600(c) of this chapter.

(xix) *Amount per ton.* Monensin, 90 to 110 grams, plus bacitracin methylene disalicylate, 50 grams, and roxarsone, 22.7 to 45.4 grams (0.0025 percent to .005 percent).

(a) *Indications for use.* For increased rate of weight gain; as an aid in the prevention of necrotic enteritis caused or complicated by *Clostridium spp* or other organisms susceptible to bacitracin methylene disalicylate; as an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as the sole ration; withdraw 5 days before slaughter; as sole source of organic arsenic; as monensin sodium provided by No.

000986 in § 510.600(c) of this chapter; as bacitracin methylene disalicylate provided by No. 046573 in § 510.600(c) of this chapter; as roxarsone provided by No. 046573 in § 510.600(c) of this chapter.

(xx) *Amount per ton.* Monensin, 90 to 110 grams, plus virginiamycin, 5 to 15 grams, and roxarsone, 22.7 grams (0.0025 percent).

(a) *Indications for use.* For increase in rate of weight gain; as an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as the sole ration; withdraw 5 days before slaughter; as sole source of organic arsenic; as monensin sodium; as roxarsone provided by Nos. 046753 and 011526 in § 510.600(c) of this chapter.

(xxi) *Amount per ton.* Monensin, 90 to 110 grams, plus virginiamycin, 5 to 15 grams.

(a) *Indications for use.* For increase in rate of weight gain; as an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as sole ration; as monensin sodium provided by No. 000986 in § 510.600 of this chapter; virginiamycin provided by No. 000007 in § 510.600 of this chapter.

(xxii) *Amount per ton.* Monensin, 90 to 110 grams plus oxytetracycline, 500 grams.

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. Mivati*, and *E. maxima*; as an aid in the reduction of mortality due to air-sacculities (air-sac infection) caused by *Escherichia coli* sensitive to oxytetracycline.

(b) *Limitations.* Feed for 5 days as sole ration. Do not feed to laying chickens. Withdraw 24 hours before slaughter. As monensin sodium provided by No. 000986 in § 510.600(c) of this chapter. As mono-alkyl (C₈-C₁₈) trimethylammonium oxytetracycline provided by No. 000069 in § 510.600(c) of this chapter.

(xxiii) *Amount per ton.* Monensin, 90 to 110 grams, plus bacitracin zinc, 4 to

50 grams, and roxarsone, 22.7 to 45.4 grams (0.0025 percent to 0.005 percent).

(a) *Indications for use.* For improved feed efficiency; as an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as the sole ration; withdraw 5 days before slaughter; as sole source of organic arsenic; as monensin sodium provided by No. 000986 in § 510.600(c) of this chapter; as bacitracin zinc provided by No. 046573 in § 510.600(c) of this chapter; as roxarsone provided by No. 046573 in § 510.600(c) of this chapter.

(xxiv) *Amount per ton.* Monensin, 90 to 110 grams, plus bacitracin methylene disalicylate, 4 to 50 grams.

(xxv) *Amount per ton.* Monensin, 90 to 110 grams plus bacitracin, 4 to 50 grams.

(a) *Indications for use.* For increased rate of weight gain and improved feed efficiency; as an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(b) *Limitations.* Do not feed to laying chickens; feed continuously as sole ration; in the absence of coccidiosis, the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain; as bacitracin zinc provided by No. 046573 in § 510.600(c) of this chapter, as monensin sodium.

(xxvi) *Amount per ton.* Monensin 90 to 110 grams plus bacitracin 100 to 200 grams and roxarsone 22.7 to 34.0 grams.

(a) *Indications for use.* As an aid in the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin methylene disalicylate; as an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; for increased rate of weight gain and improved feed efficiency.

(b) *Limitations.* For broiler chickens only. Feed continuously as sole ration. Use as sole source of organic arsenic. Withdraw 5 days before slaughter. Do not feed to laying hens. To control necrotic enteritis, start medication at first clinical signs of disease. The dosage range permitted provides for dif-

ferent levels based on the severity of infection. Use continuously for 5 to 7 days or as long as clinical signs persist, then reduce dosage to prevention level. Animals should have access to drinking water at all times. Drug overdosage or lack of water may result in leg weakness. As roxarsone and bacitracin methylene disalicylate provided by No. 046573 in § 510.600(c) of this chapter.

(xxvii) *Amount per ton.* Monensin 90 to 110 grams plus bacitracin 100 to 200 grams and roxarsone 22.7 to 45.4 grams.

(a) *Indications for use.* As an aid in the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin methylene disalicylate; as an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; for increased rate of weight gain.

(b) *Limitations.* For broiler chickens only. Feed continuously as sole ration. Use as sole source of organic arsenic. Withdraw 5 days before slaughter. Do not feed to laying hens. To control necrotic enteritis, start medication at first clinical signs of disease. The dosage range permitted provides for different levels based on the severity of infection. Use continuously for 5 to 7 days or as long as clinical signs persist, then reduce dosage to prevention level. Animals should have access to drinking water at all times. Drug overdosage or lack of water may result in leg weakness. As roxarsone and bacitracin methylene disalicylate provided by No. 046573 in § 510.600(c) of this chapter.

(2) *Turkeys—(i) Amount per ton.* Monensin, 54 to 90 grams.

(a) *Indications for use.* For the prevention of coccidiosis in turkeys caused by *E. adenoides*, *E. meleagritidis*, and *E. gallopavonis*.

(b) *Limitations.* For growing turkeys only; as monensin sodium; feed continuously as sole ration. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis.

(ii) *Amount per ton.* Monensin, 54 to 90 grams, and bacitracin methylene disalicylate, 4 to 50 grams.

(a) *Indications for use.* For prevention of coccidiosis caused by *Eimeria adenoides*, *E. meleagritidis*, and *E. gallopavonis*, for increased rate of weight gain, and for improved feed efficiency.

(b) *Limitations.* For growing turkeys only; as monensin sodium; feed continuously as sole ration. Do not allow horses, other equines, mature turkeys or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis. Bacitracin methylene disalicylate as provided by No. 046573 in § 510.600(c) of this chapter.

(iii) *Amount per ton.* Monensin, 54 to 90 grams, and bacitracin methylene disalicylate, 200 grams.

(a) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria adenoides*, *E. meleagritidis*, and *E. gallopavonis*, and as an aid in the control of transmissible enteritis complicated by organisms susceptible to bacitracin methylene disalicylate.

(b) *Limitations.* For growing turkeys only; as monensin sodium; feed continuously as sole ration. Do not allow horses, other equines, mature turkeys or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis. Bacitracin methylene disalicylate as provided by No. 046573 in § 510.600(c) of this chapter.

(iv) *Amount per ton.* Monensin, 54 to 90 grams, with virginiamycin, 10 to 20 grams.

(a) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria adenoides*, *E. meleagritidis*, and *E. gallopavonis*, and for increased rate of weight gain and improved feed efficiency in growing turkeys.

(b) *Limitations.* For growing turkeys only. Feed continuously as sole ration. Do not allow horses, other equines, ma-

ture turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses, mature turkeys, and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis. Virginiamycin as provided by No. 000069 in § 510.600(c) of this chapter.

(3) *Cattle—(i) Amount per ton.* Monensin, 5–30 grams.

(a) *Indications for use.* Improved feed efficiency.

(b) *Limitations.* (1) Feed only to cattle being fed in confinement for slaughter. Feed continuously in complete feed at a rate of 50 to 360 milligrams of monensin per head per day; as monensin sodium. Complete feeds may be manufactured from monensin liquid Type B feeds. The liquid Type B feeds have a pH of 4.3 to 7.1 and their labels must bear appropriate mixing directions. Mixing directions for liquid Type B feeds stored in recirculating tank systems are: Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily, as directed in this paragraph, even when Type B feed is not used. Mixing directions for liquid Type B feeds stored in mechanical, air, or other agitation-type tank systems are: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily, as directed in this paragraph, even when Type B feed is not used. The liquid Type B feed must bear directions to mix thoroughly with grain and/or roughage prior to feeding and must bear caution statements as follows: Inadequate mixing, (recirculation or agitation), of liquid Type B feeds has resulted in increased monensin concentration which has been fatal to cattle.

(2) An approved positionally stable monensin liquid Type B feed will not be subject to the requirements for mixing directions and cautionary labeling prescribed in paragraph (f)(3)(i)(b)(1) of this section. A manufacturer may secure approval of a positionally stable liquid Type B feed by (i) either filing

an NADA for the product or by establishing a master file containing data to support the stability of its product; (ii) authorizing the agency to reference and rely upon the data in the master file to support approval of a supplemental NADA to establish positional stability; and (iii) requesting No. 000986 in § 510.600(c) of this chapter to file a supplemental NADA to provide for the use of its monensin Type A article in the manufacture of the liquid Type B feed specified in the appropriate master file. If the data demonstrate the stability of the liquid Type B feed described in the master file, the agency will approve the supplemental NADA. Approval of the Type B feed need not be published in the FEDERAL REGISTER because approval will not affect or alter the content of the regulation. The approval will, however, provide a basis for the individual liquid feed manufacturer to submit, and for the agency to approve, a medicated feed application under section 512(m) of the act for the liquid Type B feed. A manufacturer who seeks to market a positionally unstable monensin liquid Type B feed with mixing directions different from the standard established in paragraph (f)(3)(i)(b)(1) of this section may also follow this procedure.

(ii) *Amount per ton.* Monensin, 5 to 30 grams, plus tylosin, 8 to 10 grams.

(a) *Indications for use.* Improved feed efficiency; for reduction of incidence of liver abscesses caused by *Fusobacterium necrophorum* and *Actinomyces (Corynebacterium) pyogenes*.

(b) *Limitations.* Feed only to cattle being fed in confinement for slaughter. Feed continuously as sole ration at the rate of 50 to 360 milligrams of monensin and 60 to 90 milligrams of tylosin per head per day; as monensin sodium; as tylosin phosphate.

(iii) *Amount per ton.* Monensin, 25 to 400 grams.

(a) *Indications for use.* For increased rate of weight gain; for prevention and control of coccidiosis due to *Eimeria bovis* and *E. zuernii*.

(b) *Limitations.* Feed to pasture cattle (slaughter, stocker, feeder, and dairy and beef replacement heifers). For increased rate of weight gain, feed at a rate of 50 to 200 milligrams monensin per head per day in not less than 1

pound of feed or, after the 5th day, feed at a rate of 400 milligrams per head per day every other day in not less than 2 pounds of feed. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending on severity of challenge, up to 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per day in not less than 1 pound of feed.

(iv) *Amount per ton.* Monensin, 5 to 30 grams per ton (to provide 50 to 360 milligrams per head per day), plus melengestrol acetate, 0.25 to 1.6 grams per ton (to provide 0.25 to 0.40 milligram per head per day). See § 558.342(c)(2).

(a) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and suppression of estrus (heat).

(b) *Limitations.* Heifers being fed in confinement for slaughter: Administer melengestrol acetate and monensin by: (1) Adding melengestrol acetate from a separate Type B feed containing 0.125 to 0.8 milligram per pound to Type C medicated feeds containing monensin at 5 to 30 grams per ton, (2) adding melengestrol acetate from a separate Type B feed containing 0.125 to 0.8 milligram per pound and monensin from a separate Type B feed containing 50 to 1,200 grams per ton to Type C medicated feeds, (3) adding melengestrol acetate and monensin which are contained in the same dry Type B feed at the ranges in paragraph (f)(3)(iv)(b) (1) and (2) of this section to Type C medicated feeds, or (4) using a liquid Type B feed containing 0.125 to 0.8 milligram melengestrol acetate per pound and 25 to 600 milligrams monensin per pound (50 to 1,200 grams per ton) to make Type C medicated feeds. Type C medicated feeds in paragraph (f)(3)(iv)(b) (1) and (2) of this section may be manufactured from monensin liquid Type B feeds in accordance with paragraph (f)(3)(i)(b) of this section.

(v) *Amount.* 150 milligrams per pound (0.033 percent).

(a) *Indications for use.* Increased rate of weight gain.

(b) *Limitations.* As protein-mineral blocks to be fed free choice to cattle (slaughter, stocker, feeder, and dairy

and beef replacement heifers) on pasture which may require supplemental feed. Provide 50 to 200 milligrams of monensin (0.34 to 1.33 pounds) per head per day, at least 1 block per 10 to 12 head of cattle. Roughage must be available at all times. Do not allow animals access to other protein blocks, salt or mineral, while being fed this product. Do not allow horses or other equines access to formulations containing monensin (ingestion of monensin by equines has been fatal). Block's effectiveness in cull cows and bulls has not been established. Approval must comply with § 510.455 of this chapter.

(vi) *Amount per ton.* Monensin, 25 to 400 grams.

(a) *Indications for use.* For improved feed efficiency; for prevention and control of coccidiosis due to *E. bovis* and *E. zuernii*.

(b) *Limitations.* Feed to mature reproducing beef cows. Feed as supplemental feed, either hand-fed in a minimum of 1 pound of feed or mixed in a total ration. For improved feed efficiency, feed continuously at a rate of 50 to 200 milligrams monensin per head per day. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending upon severity of challenge, up to a maximum of 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per head per day.

(vii) *Amount per ton.* Monensin, 10 to 30 grams.

(a) *Indications for use.* For improved feed efficiency; for prevention and control of coccidiosis due to *E. bovis* and *E. zuernii*.

(b) *Limitations.* For feedlot cattle, feed continuously to provide 50 to 360 milligrams monensin per head per day. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending upon the severity of challenge, up to maximum of 360 milligrams per head per day.

(viii) *Additional combinations.* Monensin may be used for heifers being fed in confinement for slaughter with melengestrol acetate with or without tylosin as in § 558.342.

(ix) *Amount.* To 000986: To make liquid Type B medicated feed containing 400 grams per ton monensin sodium with 150 grams per ton tylosin phosphate used to make a dry Type C medicated feed containing 21.4 to 26.8 grams per ton monensin plus 8 to 10 grams per ton tylosin.

(a) *Indications for use.* Improved feed efficiency; for reduction of incidence of liver abscesses caused by *Fusobacterium necrophorum* and *Actinomyces (Corynebacterium) pyogenes*.

(b) *Limitations.* Feed only to cattle being fed in confinement for slaughter. Feed continuously at the rate of 8.2 to 10.2 kilograms (18 to 22.5 pounds) of Type C medicated feed per head per day to supply 240 milligrams of monensin and 90 milligrams of tylosin per head per day; as monensin sodium; as tylosin phosphate. Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by equines has been fatal. Safe use in unapproved species and breeding cattle has not been established. The liquid Type B medicated feed must bear an expiration date of 14 days after date of manufacture. The mixing directions for this liquid Type B medicated feed stored in recirculation or agitation tank systems are: Recirculate or agitate immediately prior to use for not less than 10 minutes, moving at least 1 percent of the tanks contents per minute from the bottom of the tank to the top. Recirculate or agitate as directed daily, even when the Type B medicated feed is not used. Inadequate mixing (recirculation or agitation) of liquid Type B medicated feeds may result in increased monensin concentrations which have been fatal to cattle. Both an approved NADA and an approved medicated feed application are required to make this liquid Type B medicated feed.

(x) *Amount per ton.* Monensin, 1,620 grams as monensin sodium (810 milligrams per pound).

(a) *Indications for use.* For increased rate of weight gain.

(b) *Specifications.* Use as free-choice Type C medicated feed formulated as mineral granules as follows:

Ingredient	Percent	International feed no.
Monocalcium phosphate (21% phosphorus, 15% calcium)	29.49	6-01-082
Sodium chloride (salt)	24.25	6-04-152
Dried cane molasses	20.0	4-04-695
Ground limestone (33% calcium) or calcium carbonate (38% calcium)	13.75	6-02-632
Cane molasses	3.0	4-04-696
Processed grain by-products (as approved by AAFCO)	5.0	
Vitamin/trace mineral premix ¹	2.5	
Monensin Type A article, 80 grams per pound	1.01	
Antidusting oil	1.0	

¹Content of the vitamin/trace mineral premix may be varied. However, they should be comparable to those used for other free-choice feeds. Formulation modifications require FDA approval prior to marketing. The amount of selenium and ethylenediamine dihydroiodide (EDDI) must comply with the published requirements. (For selenium see 21 CFR 573.920; for EDDI see 51 FR 11483 (April 3, 1986).)

(c) *Limitations.* Medicated mineral granules to be fed free-choice to pasture cattle (slaughter, stocker, feeder, and dairy and beef replacement heifers). Feed continuously on a free-choice basis at the rate of 50 to 200 milligrams per head per day. During the first 5 days of feeding, cattle should receive no more than 100 milligrams per day. Do not feed additional salt or minerals. Do not mix with grain or other feeds. Monensin is toxic to cattle when consumed at higher than approved levels. Stressed and/or water deprived cattle should be adapted to the pasture and to unmedicated mineral supplement before using this product. Do not feed to lactating dairy cattle. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. The product's effectiveness in cull cows and bulls has not been established. Each use of this free-choice Type C feed must be the subject of an approved medicated feed application (MFA or Form FDA 1900) or supplemental MFA as required by §510.455 of this chapter.

(xi) *Amount per ton.* Monensin, 10 to 200 grams.

(a) *Indications for use.* For prevention and control of coccidiosis due to *E. bovis* and *E. zuernii*.

(b) *Limitations.* For calves excluding veal calves. Feed at a rate of 0.14 to 1.0 milligram monensin per pound of body weight per day, depending upon the severity of challenge, up to maximum of 200 milligrams per head per day.

(4) *Replacement chickens intended for use as cage layers—(i) Amount per ton.* Monensin, 90 to 110 grams.

(ii) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(iii) *Limitations.* Do not feed to laying chickens; feed continuously as sole ration; as monensin sodium; do not feed to chickens over 16 weeks of age.

(5) *Bobwhite quail—(i) Amount per ton.* Monensin, 73 grams.

(ii) *Indications for use.* For the prevention of coccidiosis in growing bobwhite quail caused by *Eimeria dispersa* and *E. Lettyae*.

(iii) *Limitations.* Feed continuously as the sole ration; do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin.

(6) *Goats—(i) Amount per ton.* Monensin, 20 grams.

(a) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria crandallis*, *E. christenseni*, and *E. ninakohlyakimovae*.

(b) *Limitations.* (1) Feed only to goats being fed in confinement. Do not feed to lactating goats. Feed continuously in Type C feed as monensin sodium. Type C feeds may be manufactured from monensin liquid Type B feeds. The liquid Type B feeds have a pH of 4.3 to 7.1 and their labels must bear appropriate mixing directions. Mixing directions for liquid Type B feeds stored in recirculating tank systems are: Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily, as directed in this paragraph, even when Type B feed is not used. Mixing directions for liquid Type B feeds stored in mechanical, air, or other agitation-type tank

§ 558.360

systems are: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily, as directed in this paragraph, even when Type B feed is not used. The liquid Type B feed must bear directions to mix thoroughly with grain and/or roughage prior to feeding and must bear caution statement as follows: Inadequate mixing (recirculation or agitation), of liquid Type B feeds has resulted in increased monensin concentration which could be fatal to goats.

(2) An approved positionally stable monensin liquid Type B feed will not be subject to the requirements for mixing directions and cautionary labeling prescribed in paragraph (f)(6)(i)(b)(1) of this section. A manufacturer may secure approval of a positionally stable liquid Type B feed by (i) either filing an NADA for the product or by establishing a master file containing data to support the stability of its product; (ii) authorizing the agency to reference and rely upon the data in the master file to support approval of a supplemental NADA to establish positional stability; and (iii) requesting No. 000986 in § 510.600(c) of this chapter to file a supplemental NADA to provide for the use of its monensin Type A article in the manufacture of the liquid Type B feed specified in the appropriate master file. If the data demonstrate the stability of the liquid Type B feed described in the master file, the agency will approve the supplemental NADA. Approval of the Type B feed need not be published in the FEDERAL REGISTER because approval will not affect or alter the content of the regulation. The approval will, however, provide a basis for the individual liquid feed manufacturer to submit, and for the agency to approve, a medicated feed application under section 512(m) of the act for the liquid Type B feed. A manufacturer who seeks to market a positionally unstable monensin liquid Type B feed with mixing directions different from the standard established in paragraph (f)(6)(i)(b)(1) of this section may also follow this procedure.

(ii) [Reserved]

[40 FR 13959, Mar. 27, 1975]

21 CFR Ch. I (4–1–99 Edition)

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.355, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.360 Morantel tartrate.

(a) *Approvals.* Type A medicated articles: 88 grams per pound to 000069 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.425 of this chapter.

(c) *Special considerations.* (1) Do not use in Type B or Type C medicated feeds containing bentonite.

(2) Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.

(d) *Conditions of use—(1) Amount.* 0.44 to 4.4 grams of morantel tartrate per pound of feed.

(2) *Indications for use—(i) Cattle.* For removal and control of mature gastrointestinal nematode infections of cattle including stomach worms (*Haemonchus* spp., *Ostertagia* spp., *Trichostrongylus* spp.), worms of the small intestine (*Cooperia* spp., *Trichostrongylus* spp., *Nematodirus* spp.), and worms of the large intestine (*Oesophagostomum radiatum*).

(ii) *Goats.* For removal and control of mature gastrointestinal nematode infections of goats including *Haemonchus contortus*, *Ostertagia* (*Teladorsagia*) *circumcincta*, and *Trichostrongylus axei*.

(3) *Limitations.* Feed as a single therapeutic treatment at 0.44 gram of morantel tartrate per 100 pounds of body weight. Fresh water should be available at all times. When medicated feed is consumed, resume normal feeding. Conditions of constant worm exposure may require retreatment in 2 to 4 weeks. Do not treat cattle within 14 days of slaughter; do not treat goats within 30 days of slaughter.

[46 FR 50950, Oct. 16, 1981, as amended at 47 FR 53352, Nov. 26, 1982; 51 FR 7399, Mar. 3, 1986; 51 FR 9005, Mar. 17, 1986; 52 FR 11642, Apr. 10, 1987; 59 FR 17922, Apr. 15, 1994]

§ 558.363 Narasin.

(a) *Approvals.* Type A medicated articles containing specified levels of narasin approved for sponsors identified in § 510.600(c) of this chapter for use as in paragraph (d) of this section are as follows:

(1) To 000986: 36, 45, 54, 72, and 90 grams per pound, paragraph (d)(1)(i) of this section.

(2) To 000986: 36, 45, 54, 72, and 90 grams per pound, with 10, 20, 50, and 80 percent roxarsone, paragraph (d)(1)(ii) of this section.

(3) To 000986: 36 grams per pound, with 36 grams per pound nicarbazin, paragraph (d)(1)(iii) of this section.

(4) To 012799: 36, 45, 54, 72, and 90 grams per pound, with 2 and 10 grams per pound bambermycins, paragraph (d)(1)(iv) of this section.

(5) To 012799: 45 grams per pound, with 4 and 10 grams per pound bambermycins, and 45.4, 90, and 227 grams per pound roxarsone, paragraph (d)(1)(vii) of this section.

(6) To 046573: 45 grams per pound with 10, 25, 30, 40, 50, 60, or 75 grams per pound bacitracin methylene disalicylate and 45.4, 90, or 227 grams per pound roxarsone, paragraphs (d)(1)(viii) and (d)(1)(ix) of this section.

(b) *Tolerances.* See § 556.428 of this chapter.

(c) [Reserved]

(d) *Conditions of use.* It is used as follows:

(1) *Broiler chickens*—(i) *Amount per ton.* Narasin, 54 to 72 grams.

(A) *Indications for use.* For prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(B) *Limitations.* For broiler chickens only. Feed continuously as sole ration. Do not allow adult turkeys, horses, or other equines access to narasin formulations. Ingestion of narasin by these species has been fatal.

(ii) *Amount per ton.* Narasin, 54 to 72 grams, plus roxarsone 45.4 grams (0.005 percent).

(A) *Indications for use.* For the prevention of coccidiosis in broiler chickens caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati* including some field strains of *E. tenella* which are more susceptible to roxarsone combined with narasin than to narasin alone.

(B) *Limitations.* For broiler chickens only; feed continuously as the sole ration; do not feed to laying chickens; may be fatal if accidentally fed to adult turkeys or to horses; withdraw 5

days before slaughter; as sole source of organic arsenic; not approved for use with pellet binders.

(iii) *Amount per ton.* Narasin, 27 to 45 grams, plus nicarbazin, 27 to 45 grams.

(A) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

(B) *Limitations.* For broiler chickens only. Feed continuously as the sole ration. Do not feed to laying hens. Do not allow adult turkeys, horses, or other equines access to formulations containing narasin. Ingestion of narasin by these animals has been fatal. Withdraw 5 days before slaughter. The 2 drugs can be combined only at a 1:1 ratio for the 27 to 45 grams per ton range. Only granular nicarbazin as provided by No. 000986 in § 510.600(c) of this chapter may be used in the combination.

(iv) *Amount per ton.* Narasin, 54 to 72 grams, plus bambermycins, 1 to 2 grams.

(A) *Indications for use.* For prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, and for increased rate of weight gain and improved feed efficiency.

(B) *Limitations.* For broiler chickens only. Feed continuously as the sole ration. May be fatal if fed to adult turkeys, horses, or other equines.

(v) *Amount per ton.* Narasin 54 to 72 grams, roxarsone 22.7 to 45.4 grams, and bacitracin methylene disalicylate 10 to 50 grams.

(A) *Indications for use.* For prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, for increased rate of weight gain, and for improved feed efficiency.

(B) *Limitations.* For broiler chickens only. Feed continuously as sole ration. Withdraw 5 days before slaughter. Do not feed to laying hens. Use as sole source of organic arsenic. Drug overdose or lack of water may result in leg weakness. Do not allow adult turkeys, horses, or other equines access to narasin formulations. Ingestion of narasin by these species has been fatal. Narasin as provided by 000986,

roxarsone by 046573, bacitracin methylene disalicylate by 046573 in § 510.600(c) of this chapter.

(vi) *Amount per ton.* Narasin 54 to 72 grams, and bacitracin methylene disalicylate 10 to 50 grams.

(A) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix*, and *E. tenella*, for increased rate of weight gain, and for improved feed efficiency.

(B) *Limitations.* For broiler chickens only. Feed continuously as sole ration. Do not feed to laying hens. Do not allow adult turkeys, horses, or other equines access to narasin formulations. Ingestion of narasin by these species has been fatal. Narasin as provided by 000986, bacitracin methylene disalicylate by 046573 in § 510.600(c) of this chapter.

(vii) *Amount per ton.* Narasin 54 to 72 grams, bambermycins 1 to 2 grams, and roxarsone 22.7 to 45.4 grams.

(A) *Indications for use.* For prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. mivati*, *E. acervulina*, *E. maxima*, and *E. brunetti*, and for increased rate of weight gain, improved feed efficiency, and improved pigmentation in broiler chickens.

(B) *Limitations.* For broiler chickens only. Feed continuously as sole ration. Do not feed to laying hens. Do not allow adult turkeys or horses or other equines access to formulations containing narasin. Ingestion of narasin by these animals has been fatal. Use as sole source of organic arsenic. Poultry should have access to drinking water at all times. Drug overdosage or lack of water intake may result in leg weakness or paralysis. Withdraw 5 days before slaughter. Narasin as provided by 000986 in § 510.600(c) of this chapter, bambermycins by 012799, and roxarsone by 046573.

(viii) *Amount per ton.* Narasin, 54 to 72 grams, and bacitracin methylene disalicylate, 50 grams, with roxarsone, 22.7 to 45.4 grams.

(A) *Indications for use.* For prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, as an aid in the prevention of necrotic enteritis caused or complicated by *Clos-*

tridium spp. or other organisms susceptible to bacitracin, and for increased rate of weight gain, improved feed efficiency, and improved pigmentation.

(B) *Limitations.* For broiler chickens only. Feed continuously as sole ration. Withdraw 5 days before slaughter. Do not feed to laying hens. Do not allow adult turkeys, horses, or other equines access to narasin formulations. Ingestion of narasin by these species has been fatal. Use as sole source of organic arsenic. Poultry should have access to drinking water at all times. Drug overdose or lack of water intake may result in leg weakness or paralysis. Narasin as provided by 000986, bacitracin methylene disalicylate and roxarsone by 046573 in § 510.600(c) of this chapter.

(ix) *Amount per ton.* Narasin, 54 to 72 grams, and bacitracin methylene disalicylate, 100 to 200 grams, with roxarsone, 22.7 to 45.4 grams.

(A) *Indications for use.* For prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, as an aid in the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin, and for increased rate of weight gain, improved feed efficiency, and improved pigmentation.

(B) *Limitations.* For broiler chickens only. Feed continuously as sole ration. Withdraw 5 days before slaughter. Do not feed to laying hens. Do not allow adult turkeys, horses, or other equines access to narasin formulations. Ingestion of narasin by these species has been fatal. Use as sole source of organic arsenic. Poultry should have access to drinking water at all times. Drug overdose or lack of water intake may result in leg weakness or paralysis. Narasin as provided by 000986, bacitracin methylene disalicylate and roxarsone by 046573 in § 510.600(c) of this chapter.

(2) Narasin may also be used for broilers in combination with:

(i) Nicarbazin with lincomycin as in § 558.366.

Food and Drug Administration, HHS

§ 558.366

(ii) Nicarbazin and bacitracin methylene disalicylate as in § 558.366.

[51 FR 29098, Aug. 14, 1986, as amended at 53 FR 24260, June 28, 1988; 54 FR 1928, Jan. 18, 1989; 54 FR 5930, Feb. 7, 1989; 54 FR 9430, Mar. 7, 1989; 54 FR 29335, July 12, 1989; 55 FR 8460, Mar. 8, 1990; 57 FR 8578, Mar. 11, 1992; 57 FR 32176, July 21, 1992; 63 FR 13124, Mar. 18, 1998; 63 FR 51823, Sept. 29, 1998; 63 FR 57248, Oct. 27, 1998; 64 FR 4966, Feb. 2, 1999]

§ 558.365 Nequinat.

(a) *Approvals.* Type A medicated articles: 4 percent to 017800 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.440 of this chapter.

(c) *Special considerations.* Do not use in Type B or Type C medicated feeds containing bentonite.

(d) *Conditions of use.* It is used as follows:

(1) *Broiler or fryer chickens—(i) Amount per ton.* Nequinat, 18.16 grams.

(a) *Indications for use.* An aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations.* Feed continuously as the sole ration.

(ii) *Amount per ton.* Nequinat, 18.16 grams (0.002 percent) plus roxarsone, 45.4 grams (0.005 percent).

(a) *Indications for use.* An aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, and *E. mivati*; growth promotion and feed efficiency; for improving pigmentation.

(b) *Limitations.* Feed continuously as sole ration throughout the starting pe-

riod; withdraw 5 days before slaughter; as sole source of organic arsenic.

(iii) *Amount per ton.* Nequinat, 18.16 grams (0.002 percent) plus oxytetracycline, 200 grams.

(a) *Indications for use.* For control of complicated chronic respiratory disease (air-sac infection), infectious synovitis, and treatment of blue comb (nonspecific infectious enteritis).

(b) *Limitations.* As monoalkyl (C₈–C₁₈) trimethylammonium oxytetracycline as provided by No. 000069 in § 510.600(c) of this chapter.

(2) *Roaster chickens or replacement chickens for caged layers—(i) Amount per ton.* Nequinat, 18.16 grams (0.002 percent).

(ii) *Indications for use.* An aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(iii) *Limitations.* Feed continuously as the sole ration; do not feed to chickens over 16 weeks of age.

[40 FR 13959, Mar. 27, 1975, as amended at 51 FR 7399, Mar. 3, 1986; 52 FR 2685, Jan. 26, 1987]

§ 558.366 Nicarbazin.

(a) Type A medicated articles: 25 percent to 000986, 060728, and 063271 in § 510.600(c) of this chapter for use as indicated in the table in paragraph (c) of this section.

(b) *Related tolerances.* See § 556.445 of this chapter.

(c) *Conditions of use.* It is used in chicken feed as follows:

Nicarbazin in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
27 to 45	Narasin 27 to 45	Broiler chickens; prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , <i>E. mivati</i> .	Sec. 558.363(d)(1)(iii)	000986
	Narasin 27 to 45 and bacitracin methylene disalicylate 4 to 50.	Broiler chickens; prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , <i>E. mivati</i> ; for increased rate of weight gain and improved feed efficiency.	Feed continuously as sole ration. Withdraw 5 days before slaughter. Do not allow turkeys, horses, or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Do not feed to laying hens. Narasin and nicarbazin as provided by 000986, bacitracin methylene disalicylate by 046573.	000986

Nicarbazin in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
113.5 (0.0125 pct)	Narasin 27 to 45 and Lincomycin 2 to 4.	Broiler chickens; prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , <i>E. mivati</i> , for increased rate of weight gain and improved feed efficiency.	Feed continuously as sole ration. Withdraw 5 days before slaughter. Do not allow turkeys, horses, or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Do not feed to laying hens. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Narasin and nicarbazin as provided by 000986, lincomycin by 000009.	000986
		Chickens; aid in preventing outbreaks of cecal (<i>Eimeria tenella</i>) and intestinal (<i>E. acervulina</i> , <i>E. maxima</i> , <i>E. necatrix</i> , and <i>E. brunetti</i>) coccidiosis.	Feed continuously as sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard; do not use as a treatment for coccidiosis; do not use in flushing mashers; do not feed to laying hens; withdraw 4 days before slaughter.	000986 060728 063271
	Bacitracin methylene disalicylate 30.	Broiler chickens; aid in preventing outbreaks of cecal (<i>Eimeria tenella</i>) and intestinal (<i>E. acervulina</i> , <i>E. maxima</i> , <i>E. necatrix</i> , and <i>E. brunetti</i>) coccidiosis; for increased rate of weight gain and improved feed efficiency.	do	060728 063271
	Lincomycin 2 (0.00044 pct).	Broiler chickens; aid in preventing outbreaks of cecal (<i>Eimeria tenella</i>) and intestinal (<i>E. acervulina</i> , <i>E. maxima</i> , <i>E. necatrix</i> and <i>E. brunetti</i>) coccidiosis; for increased rate of weight gain.	do	060728 063271
	Roxarsone 22.7 (0.0025)..	do	Feed continuously as sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard; as sole source of organic arsenic; do not use a treatment for coccidiosis; do not use in flushing mashers; do not feed to laying hens; withdraw 5 days before slaughter.	060728 063271
	Roxarsone 22.7 (0.0025) plus lincomycin 2 (0.0004)..	do	do	060728 063271

[42 FR 56729, Oct. 28, 1977; 43 FR 1942, Jan. 13, 1978, as amended at 44 FR 40887, July 13, 1979; 50 FR 13562, Apr. 5, 1985; 51 FR 7399, Mar. 3, 1986; 54 FR 1928, Jan. 18, 1989; 60 FR 29483, June 5, 1995; 61 FR 1832, Jan. 24, 1996; 61 FR 14021, Mar. 29, 1996; 61 FR 14483, Apr. 2, 1996; 62 FR 29011, May 29, 1997; 63 FR 13124, Mar. 18, 1998; 63 FR 57248, Oct. 27, 1998; 64 FR 4966, Feb. 2, 1999]

§ 558.369 Nitarsone.

(a) *Approvals.* Type A medicated articles: 50 percent to 046573 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.60 of this chapter.

(c) [Reserved]

(d) *Conditions of use.* It is used as follows:

(1) *Chickens and turkeys.*—(i) *Amount.* Nitarsone, 0.01875 percent.

(ii) *Indications for use.* As an aid in the prevention of blackhead.

(iii) *Limitations.* Early medication is essential to prevent spread of disease. Adequate drinking water must be provided near feeder at all times. The drug is not effective in preventing blackhead in birds infected more than 4 or 5 days. Discontinue use 5 days before slaughtering animals for human consumption to allow elimination of the drug from edible tissues. The drug is dangerous for ducks, geese, and dogs. Overdosage or lack of water may result in leg weakness or paralysis. Use as sole source of arsenic.

(2) *Turkeys*—(i) *Amount.* Nitarsone 0.01875 percent, plus bacitracin methylene disalicylate 4 to 50 grams per ton.

(ii) *Indications for use.* As an aid in the prevention of blackhead, and for increased rate of weight gain and improved feed efficiency.

(iii) *Limitations.* For growing turkeys. Feed continuously as sole ration. Early medication is essential to prevent spread of disease. Adequate drinking water must be provided near feeders at all times. Overdosage or lack of water may result in leg weakness or paralysis. The drug is not effective in preventing blackhead in birds infected more than 4 or 5 days. Discontinue use 5 days before slaughtering animals for human consumption to allow elimination of the drug from edible tissues. The drug is dangerous for ducks, geese, and dogs. Use as sole source of arsenic.

[46 FR 47535, Sept. 29, 1981, as amended at 47 FR 14152, Apr. 2, 1982; 51 FR 7399, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 55 FR 8460, Mar. 8, 1990; 57 FR 8578, Mar. 11, 1992; 63 FR 39028, July 21, 1998]

§ 558.376 Nitromide and sulfanitran.

(a) *Approvals.* Type A medicated articles: 25 percent nitromide, 30 percent sulfanitran, with or without 5 percent roxarsone to 053501 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See §§ 556.220 and 556.680 of this chapter.

(c) *Conditions of use.* It is used for chickens as follows:

(1) *Amount.* 227 grams per ton nitromide (0.025 percent) and 272 grams per ton sulfanitran (0.03 percent).

(i) *Indications for use.* As an aid in the prevention of coccidiosis caused by

Eimeria tenella, *E. necatrix*, and *E. acervulina*.

(ii) *Limitations.* Not to be fed to laying chickens; withdraw 5 days before slaughter; from Type A articles containing not more than 25 percent nitromide and 30 percent sulfanitran.

(2) *Amount.* 227 grams per ton nitromide (0.025 percent) and 272 grams per ton sulfanitran (0.03 percent), plus 45.4 grams per ton roxarsone (0.005 percent).

(i) *Indications for use.* Prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, and *E. acervulina*; growth promotion and feed efficiency; improving pigmentation.

(ii) *Limitations.* Not to be fed to laying chickens; withdraw 5 days before slaughter; from Type A articles containing not more than 25 percent nitromide, 30 percent sulfanitran, and 5 percent roxarsone; as sole source of organic arsenic.

[41 FR 11002, Mar. 15, 1976, as amended at 51 FR 7399, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 55 FR 8460, Mar. 8, 1990]

§ 558.415 Novobiocin.

(a) *Approvals.* Type A medicated articles: 25 grams of activity per pound to 000009 in § 510.600(c) of this chapter. Type B medicated feeds: 17.5 grams per pound to 000009 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.460 of this chapter.

(c) *Conditions of use.* It is used in animal feeds as follows:

(1) *Chickens*—(i) *Amount.* Novobiocin, 6–7 mgs. per lb. body weight per day.

(a) *Indications for use.* Aid in the treatment of breast blisters associated with staphylococcal infections susceptible to novobiocin.

(b) *Limitations.* Administer, as sole ration, feed which contains not less than 200 grams of novobiocin activity per ton of feed; not for laying chickens; feed 5 to 7 days; withdraw 4 days before slaughter.

(ii) *Amount.* Novobiocin, 10–14 mgs. per lb. body weight per day.

(a) *Indications for use.* Treatment of staphylococcal synovitis and generalized staphylococcal infections susceptible to novobiocin.

(b) *Limitations.* Administer, as sole ration, feed which contains not less than

§ 558.430

350 grams of novobiocin activity per ton of feed; not for laying chickens; feed 5 to 7 days; withdraw 4 days before slaughter.

(2) *Turkeys*—(i) *Amount*. Novobiocin, 4–5 mgs. per lb. body weight per day.

(a) *Indications for use*. Aid in the treatment of breast blisters associated with staphylococcal infections susceptible to novobiocin.

(b) *Limitations*. Administer, as sole ration, feed which contains not less than 200 grams of novobiocin activity per ton of feed; not for laying turkeys; feed 5 to 7 days; withdraw 4 days before slaughter.

(ii) *Amount*. Novobiocin, 5–8 mgs. per lb. body weight per day.

(a) *Indications for use*. Aid in the control of recurring outbreaks of fowl cholera caused by strains of *Pasteurella multocida* susceptible to novobiocin following initial treatment with 7–8 mgs. per pound body weight per day.

(b) *Limitations*. Administer, as sole ration, feed which contains not less than 200 grams of novobiocin activity per ton of feed; feed 5 to 7 days; not for laying turkeys; withdraw 4 days before slaughter.

(iii) *Amount*. Novobiocin, 7–8 mgs. per lb. body weight per day.

(a) *Indications for use*. Treatment of staphylococcal synovitis and generalized staphylococcal infection susceptible to novobiocin; treatment of acute outbreaks of fowl cholera caused by strains of *Pasteurella multocida* susceptible to novobiocin.

(b) *Limitations*. Administer, as sole ration, feed which contains not less than 350 grams of novobiocin activity per ton of feed; feed 5 to 7 days; not for laying turkeys; withdraw 4 days before slaughter.

(3) *Mink*—(i) *Amount*. 20 mgs. per lb. body weight per day.

(ii) *Indications for use*. For treatment of generalized infections, abscesses, or urinary infections caused by staphylococcal or other novobiocin sensitive organisms.

(iii) *Limitations*. Administer, as sole ration, feed which contains not less than 200 grams of novobiocin activity per ton of feed; feed for 7 days.

(4) *Ducks*—(i) *Amount*. Novobiocin, 350 grams per ton.

(ii) *Indications for use*. Control of infectious serositis and fowl cholera in ducks caused by *Pasteurella anatipestifer* and *P. multocida*, susceptible to novobiocin.

(iii) *Limitations*. Administer, as sole ration, for 5 to 7 days, continue medication for 14 days if necessary, repeat if reinfection occurs; discontinue use at least 3 days before slaughter; not for use in laying ducks.

[40 FR 13959, Mar. 27, 1975, as amended at 45 FR 42263, June 24, 1980; 51 FR 7399, Mar. 3, 1986; 52 FR 36402, Sept. 29, 1987]

§ 558.430 Nystatin.

(a) *Approvals*. Type A medicated articles: 20 grams of activity per pound to 046573 in § 510.600(c) of this chapter.

(b) *Related tolerances*. See § 556.470 of this chapter.

(c) *Conditions of use*. It is used for chickens and turkeys as follows:

(1) *Amount*. 50 grams per ton.

(i) *Indications for use*. Chickens and turkeys; aid in control of crop mycosis and mycotic diarrhea (*Candida albicans*).

(ii) *Limitations*. Growing and laying chickens; growing turkeys.

(2) *Amount*. 100 grams per ton.

(i) *Indications for use*. Chickens and turkeys; treatment of crop mycosis and mycotic diarrhea (*Candida albicans*).

(ii) *Limitations*. Growing and laying chickens; growing turkeys; to be fed for 7 to 10 days.

[41 FR 11002, Mar. 15, 1976, as amended at 51 FR 7399, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 53 FR 40729, Oct. 18, 1988; 55 FR 8461, Mar. 8, 1990; 57 FR 8578, Mar. 11, 1992]

§ 558.435 Oleandomycin.

(a) *Approvals*. Type A medicated articles: 5 grams of activity per pound to 000069 in § 510.600(c) of this chapter.

(b) *Related tolerances*. See § 556.480 of this chapter.

(c) *Special considerations*. Do not use bentonite in Type B or Type C medicated feeds containing oleandomycin. Oleandomycin refers to oleandomycin or feed-grade oleandomycin.

(d) *Conditions of use*. It is used in animal feed as follows:

(1) *Chickens and turkeys*—(i) *Amount per ton*. Oleandomycin, 1 to 2 grams.

(ii) *Indications for use*. For increased rate of weight gain and improved feed

Food and Drug Administration, HHS

§ 558.450

efficiency for broiler chickens and growing turkeys.

(2) *Swine*—(i) *Amount per ton*. Oleandomycin, 5 to 11.25 grams.

(ii) *Indications for use*. For increased rate of weight gain and improved feed efficiency in growing-finishing swine.

[40 FR 13959, Mar. 27, 1975, as amended at 44 FR 40283, July 10, 1979; 51 FR 7399, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987]

§ 558.450 Oxytetracycline.

(a) *Approvals*. Type A medicated articles:

(1) 10, 20, 30, 50, 100, and 200 grams per pound to 000069 in § 510.600(c) of this chapter.

(2) 50 and 100 grams per pound to 053389 in § 510.600(c) of this chapter.

(b) *Special considerations*. (1) In accordance with § 558.5 labeling shall bear the statement: “FOR USE IN DRY

ANIMAL FEED ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS.”

(2) The articles in paragraph (a)(1) of this section contain an amount of mono-alkyl (C₈–C₁₈) trimethylammonium oxytetracycline expressed in terms of an equivalent amount of oxytetracycline hydrochloride or an amount of oxytetracycline dihydrate base expressed in terms of an equivalent amount of oxytetracycline hydrochloride.

(3) The articles in paragraph (a)(2) of this section contain an amount of mono-alkyl (C₈–C₁₈) trimethylammonium oxytetracycline expressed in terms of an equivalent amount of oxytetracycline hydrochloride.

(c) *Related tolerances*. See § 556.500 of this chapter.

(d)(1) *Conditions of use*. It is used in feed as follows:

TABLE 1

Oxytetracycline amount	Combination	Indications for use	Limitations	Sponsor
(i) 10 to 20 grams per ton (g/ton)	Nequinat 18.16 g/ton (0.002%)	Sheep; increased rate of weight gain and improved feed efficiency.	do	000069, 053389
(ii) 10 to 50 g/ton		1. Chickens; increased rate of weight gain and improved feed efficiency.	Do not feed to chickens producing eggs for human consumption.	Do.
		2. Growing turkeys; increased rate of weight and improved feed efficiency.	Do not feed to turkeys producing eggs for human consumption.	Do.
		3. Swine; increased rate of weight and improved feed efficiency.	do	Do.
(iii) 100 g/ton	Nequinat 18.16 g/ton (0.002%)	Turkeys; control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to oxytetracycline.	Feed continuously for 7 to 14 days (d); do not feed to turkeys producing eggs for human consumption.	Do.
(iv) 100 to 200 g/ton		Chickens; control of infectious synovitis caused by <i>Mycoplasma synoviae</i> ; control of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline.	Feed continuously for 7 to 14 d; do not feed to chickens producing eggs for human consumption; in low calcium feed, withdraw 3 d before slaughter.	Do.
		Chickens; control of infectious synovitis caused by <i>M. synoviae</i> ; control of fowl cholera caused by <i>P. multocida</i> susceptible to oxytetracycline; as an aid in the control of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , and <i>E. mivati</i> .	do	000069

TABLE 1—Continued

Oxytetracycline amount	Combination	Indications for use	Limitations	Sponsor
(v) 200 g/ton		Turkeys; control of infectious synovitis caused by <i>M. synoviae</i> susceptible to oxytetracycline.	Feed continuously for 7 to 14 d; withdraw 5 d before slaughter; do not feed to turkeys producing eggs for human consumption.	000069, 053389
(vi) 400 g/ton	Monensin 90 to 110 g/ton	Chickens; control of chronic respiratory disease (CRD) and air sac infection caused by <i>M. gallisepticum</i> and <i>Escherichia coli</i> susceptible to oxytetracycline.	Feed continuously for 7 to 14 d; do not feed to chickens producing eggs for human consumption; in low calcium feeds, withdraw 3 d before slaughter.	Do.
		Chickens; control of CRD and air sac infection caused by <i>M. gallisepticum</i> and <i>E. coli</i> susceptible to oxytetracycline; and as an aid in the prevention of coccidiosis caused by <i>E. necatrix</i> , <i>E. tenella</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> .	do	000069
	Nequinatate 18.16 g/ton (0.002%)	Chickens; control of CRD and air sac infection caused by <i>M. gallisepticum</i> and <i>E. coli</i> susceptible to oxytetracycline; as an aid in prevention of coccidiosis caused by <i>E. tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , and <i>E. mivati</i> .	do	Do.
(vii) 500 g/ton		Chickens; reduction of mortality due to air sacculitis (air-sac-infection) caused by <i>E. coli</i> susceptible to oxytetracycline.	Feed continuously for 5 d; do not feed to chickens producing eggs for human consumption; withdraw 24 hours before slaughter; in low calcium feeds withdraw 3 d before slaughter.	000069, 053389
	Monensin 90 to 110 g/ton	Chickens; reduction of mortality due to air sacculitis (air-sac-infection) caused by <i>E. coli</i> susceptible to oxytetracycline; as an aid in the prevention of coccidiosis caused by <i>E. necatrix</i> , <i>E. tenella</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> .	do	000069
	Salinomycin 40 to 60 g/ton	Chickens; reduction of mortality due to air sacculitis (air-sac-infection) caused by <i>E. coli</i> susceptible to oxytetracycline; prevention of coccidiosis caused by <i>E. necatrix</i> , <i>E. tenella</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. mivati</i> , and <i>E. maxima</i> .	do	000069, 012799

TABLE 1—Continued

Oxytetracycline amount	Combination	Indications for use	Limitations	Sponsor
(viii) 0.05 to 0.1 milligram/pound (mg/lb) of body weight daily.		Calves (up to 250 lb); for increased rate of weight gain and improved feed efficiency.	Feed continuously; in milk replacers or starter feed.	000069, 053389
(ix) 10 mg/lb of body weight daily.		1. Calves and beef and nonlactating dairy cattle; treatment of bacterial enteritis caused by <i>E. coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>P. multocida</i> susceptible to oxytetracycline.	Feed continuously for 7 to 14 d; in feed or milk replacers; withdraw 5 d before slaughter.	Do.
		2. Calves (up to 250 lb); treatment of bacterial enteritis caused by <i>E. coli</i> susceptible to oxytetracycline.	Feed continuously for 7 to 14 d; in milk replacers or starter feed; withdraw 5 d before slaughter.	Do.
		3. Sheep; treatment of bacterial enteritis caused by <i>E. coli</i> and bacterial pneumonia caused by <i>P. multocida</i> susceptible to oxytetracycline.	Feed continuously for 7 to 14 d; withdraw 5 d before slaughter.	Do.
		4. Swine; treatment of bacterial enteritis caused by <i>E. coli</i> and <i>Salmonella choleraesuis</i> susceptible to oxytetracycline and treatment of bacterial pneumonia caused by <i>P. multocida</i> susceptible to oxytetracycline.	Feed continuously for 7 to 14 d; withdraw 5 d before slaughter.	Do.
		5. Breeding swine; control and treatment of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to oxytetracycline.	Feed continuously for not more than 14 d; withdraw 5 d before slaughter.	Do.
(x) 25 mg/lb of body weight		Turkeys; control of complicating bacterial organisms associated with bluecomb (transmissible enteritis; coronaviral enteritis) susceptible to oxytetracycline.	Feed continuously for 7 to 14 d; withdraw 5 d before slaughter; do not feed to turkeys producing eggs for human consumption.	Do.
(xi) 25 mg/head/day		Calves (250 to 400 lb); increased rate of weight gain and improved feed efficiency.		Do.
(xii) 75 mg/head/day		Growing cattle (over 400 lb); increased rate of weight gain; improved feed efficiency, and reduction of liver condemnation due to liver abscesses.		Do.
(xiii) 0.5 to 2.0 g/head/day		Cattle; prevention and treatment of the early stages of shipping fever complex.	Feed 3 to 5 d before and after arrival in feedlots.	Do.

TABLE 1—Continued

Oxytetracycline amount	Combination	Indications for use	Limitations	Sponsor
(xiv) 200 mg/colony		Honey bees; control of American foulbrood caused by <i>Bacillus larvae</i> and European foulbrood caused by <i>Streptococcus pluton</i> susceptible to oxytetracycline.	Remove at least 6 weeks prior to main honey flow.	Do.

(2) It is used in fish feed as follows:

TABLE 2

Oxytetracycline amount	Combination	Indications for use	Limitations	Sponsor
(i) 250 mg/kilogram of fish/d (11.35 g/100 lb of fish/d).		Pacific salmon for marking of skeletal tissue.	For salmon not over 30 g body weight; administer as sole ration for 4 consecutive days in feed containing oxytetracycline hydrochloride or mono-alkyl (C ₈ –C ₁₈) trimethyl ammonium oxytetracycline; fish not to be liberated for at least 7 d following the last administration of medicated feed.	000069
(ii) 2.5 to 3.75 g/100 lb of fish/d.		1. Salmonids; control of ulcer disease caused by <i>Hemophilus piscium</i> , furunculosis caused by <i>Aeromonas salmonicida</i> , bacterial hemorrhagic septicemia caused by <i>A. liquefaciens</i> , and pseudomonas disease.	Administer as mono-alkyl (C ₈ –C ₁₈) trimethyl ammonium oxytetracycline in mixed ration for 10 d; do not liberate fish or slaughter fish for food for 21 d following the last administration of medicated feed; do not administer when water temperature is below 9 °C (48.2 °F).	000069
		2. Catfish; control of bacterial hemorrhagic septicemia caused by <i>A. liquefaciens</i> and pseudomonas disease.	Administer as mono-alkyl (C ₈ –C ₁₈) trimethyl ammonium oxytetracycline in mixed ration for 10 d; do not liberate fish or slaughter fish for food for 21 d following the last administration of medicated feed; do not administer when water temperature is below 16.7 °C (62 °F).	000069
(iii) 1 g/lb of medicated feed.		Lobsters; control of gaffkemia caused by <i>Aerococcus viridans</i> .	Administer as sole ration for 5 consecutive days in feed containing monoalkyl (C ₈ –C ₁₈) trimethyl ammonium oxytetracycline; withdraw medicated feed 30 d before harvesting lobsters.	000069

(3) Oxytetracycline may be used in accordance with the provisions of this section in the combinations provided as follows:

- (i) Robenidine hydrochloride in accordance with § 558.515.
(ii) Lasalocid as in § 558.311.

Food and Drug Administration, HHS

§ 558.465

(iii) Melengestrol acetate as in § 558.342.

[61 FR 51590, Oct. 3, 1996, as amended at 63 FR 41192, Aug. 3, 1998]

§ 558.460 Penicillin.

(a) *Specifications.* As penicillin procaine G or feed grade penicillin procaine.

(b) *Sponsors.* Type A medicated articles: To 000069, 100 grams per pound. To 046573, 100 and 227 grams per pound.

(c) Related tolerances. See § 556.510 of this chapter.

(d) *Conditions of use.* (1) It is used as follows:

Penicillin in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 2.4 to 50		Chickens, turkeys, and pheasants; for increased rate of weight gain and improved feed efficiency..	Do not feed to poultry producing eggs for human consumption..	000069, 046573.
(ii) 5 to 20		Quail; for increased rate of weight gain and improved feed efficiency..	Quail; not over 5 weeks of age.	Do.
(iii) 10 to 50		Swine; for increased rate of weight gain and improved feed efficiency..		Do.

(2) Penicillin may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Amprolium in accordance with § 558.55.

(ii) Amprolium plus ethopabate in accordance with § 558.58.

(iii)-(v) [Reserved]

(vi) Hygromycin B in accordance with § 558.274.

(vii) Roxarsone and zoalene in accordance with § 558.680.

(viii) Zonalene in accordance with § 558.680.

[41 FR 11004, Mar. 15, 1976, as amended at 42 FR 18618, Apr. 8, 1977; 42 FR 36995, July 19, 1977; 47 FR 42103, Sept. 24, 1982; 51 FR 7399, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 58 FR 30120, May 26, 1993; 60 FR 39847, Aug. 4, 1995; 63 FR 36179, July 2, 1998]

§ 558.464 Poloxalene.

(a) *Approvals.* (1) Dry Type A medicated articles: 53 percent to 000069 in § 510.600(c) of this chapter.

(2) Liquid Type A medicated articles: 99.5 percent to 000069 in § 510.600(c) of this chapter.

(b) *Conditions of use.* (1) For prevention of legume (alfalfa, clover) and wheat pasture bloat in cattle.

(2) Poloxalene dry Type A article and liquid Type A article must be thoroughly blended and evenly distributed in feed prior to use. This may be accomplished by adding the Type A arti-

cle to a small quantity of feed, mixing thoroughly, then adding this mixture to the remaining feed and again mixing thoroughly. Dosage is 1 gram of poloxalene per 100 pounds of body weight daily and continued during exposure to bloat producing conditions. If bloating conditions are severe, the dose is doubled. Treatment should be started 2 to 3 days before exposure to bloat-producing conditions. Repeat dosage if animals are exposed to bloat-producing conditions more than 12 hours after the last treatment. Do not exceed the higher dosage levels in any 24-hour period.

[40 FR 39857, Aug. 29, 1975, as amended at 51 FR 7399, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 56 FR 50654, Oct. 8, 1991; 60 FR 55660, Nov. 2, 1995]

§ 558.465 Poloxalene free-choice liquid Type C feed.

(a) *Approvals.* Type A medicated articles: 99.5 percent to 000069 in § 510.600(c) of this chapter.

(b) *Conditions of use.* (1) For control of legume (alfalfa, clover) and wheat pasture bloat in cattle, use 7.5 grams of poloxalene per pound of liquid Type C feed (1.65 percent weight/weight). Each animal must consume 0.2 pound of Type C feed per 100 pounds of body weight daily for adequate protection.

(2) For control of legume (alfalfa, clover) bloat in cattle grazing of prebloom legumes, use 10.00 grams of poloxalene

§ 558.485**21 CFR Ch. I (4–1–99 Edition)**

per pound of liquid Type C feed (2.2 percent weight/weight). Each animal must consume 0.15 pound of Type C feed per 100 pounds of body weight daily for adequate protection. If consumption exceeds 0.2 pound of Type C feed per 100 pounds of body weight daily, cattle should be changed to a Type C feed containing 7.5 grams of poloxalene per pound.

(3) Poloxalene liquid Type A article must be thoroughly blended and evenly distributed into a liquid Type C feed and offered to cattle in a covered liquid Type C feed feeder with lick wheels. The formula for the liquid Type C feed, on a weight/weight basis, is as follows: Ammonium polyphosphate 2.66 percent, phosphoric acid (75 percent) 3.37 percent, sulfuric acid 1.00 percent, water 10.00 percent, and molasses sufficient to make 100.00 percent, vitamins A and D and/or trace minerals may be added. One free-turning lick wheel per 25 head of cattle must be provided.

(4) The medicated liquid Type C feed must be introduced at least 2 to 5 days before legume consumption to accustom the cattle to the medicated liquid Type C feed and to lick wheel feedings. If the medicated liquid wheel Type C feed feeding is interrupted, this 2- to 5-day introductory feeding should be repeated.

[40 FR 13959, Mar. 27, 1975, as amended at 42 FR 21281, Apr. 26, 1977; 51 FR 7399, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 56 FR 50654, Oct. 8, 1991; 60 FR 55660, Nov. 2, 1995]

§ 558.485 Pyrantel tartrate.

(a) *Approvals.* Type A medicated articles containing pyrantel tartrate to firms identified by drug labeler codes in § 510.600(c) of this chapter for the specific usage indicated in paragraph (e) of this section:

(1) To 000069: 9.6, 19.2, 48 and 80 grams per pound, paragraph (e)(1) of this section; 48 grams per pound, paragraph (e)(2) of this section.

(2) To 017800: 19.2 and 48 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(3) To 016968: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(4) [Reserved]

(5) To 017790: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(6) [Reserved]

(7) To 051359: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(8) To 011490: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(9) To 011749: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(10) To 043733: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(11) To 017519: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(12) To 046987: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) and (e)(1)(ii) of this section.

(13) To 034936: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) and (e)(1)(ii) of this section.

(14) [Reserved]

(15) To 049685: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(16)–(17) [Reserved]

(18) To 001800: 9.6 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(19) To 050568: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(20) To 050639: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(21) [Reserved]

(22) To 017473: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(23) To 021676: 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(24)–(26) [Reserved]

(27) To 010439: 9.6 and 19.2 grams per pound, paragraphs (e)(1)(i) through (e)(1)(iii) of this section.

(28) To 062240: 48 grams per pound, paragraph (e)(2) of this section.

(b) [Reserved]

(c) *Related tolerances.* See § 556.560 of this chapter.

(d) *Special considerations.* (1) Consult veterinarian before using in severely debilitated animals.

(2) Do not mix in Type B or Type C medicated feeds containing bentonite.

(e) *Conditions of use.* It is used as follows:

(1) *Swine*—(i) *Amount per ton.* 96 grams (0.0106 percent).

(A) *Indications for use.* Aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections.

(B) *Limitations.* Feed continuously as the sole ration in a Type C feed; withdraw 24 hours prior to slaughter.

(ii) *Amount per ton.* 96 grams (0.0106 percent).

(A) *Indications for use.* For the removal and control of large roundworm (*Ascaris suum*) infections.

(B) *Limitations.* Feed for 3 days as the sole ration in a Type C feed; withdraw 24 hours prior to slaughter.

(iii) *Amount per ton.* 800 grams (0.0881 percent).

(A) *Indications for use.* For the removal and control of large roundworm (*Ascaris suum*) and nodular worm (*Oesophagostomum*) infections.

(B) *Limitations.* As sole ration for a single therapeutic treatment in Type C feed; feed at the rate of 1 lb of feed per 40 lb of body weight for animals up to 200 lb, and 5 lb of feed per head for animals 200 lb or over; withdraw 24 hours prior to slaughter.

(iv) *Amount per ton.* Pyrantel tartrate, 96 grams (0.0106 percent) and carbadox, 50 grams (0.0055 percent).

(A) *Indications for use.* For control of swine dysentery (vibrionic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis*); aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections.

(B) *Limitations.* Do not feed to swine weighing over 75 pounds; do not feed within 10 weeks of slaughter; consult a veterinarian before feeding to severely debilitated animals; feed continuously as sole ration. Do not use in Type C feeds containing less than 15 percent crude protein.

(v) *Amount per ton.* Pyrantel tartrate, 96 grams (0.0106 percent) and tylosin, 40 to 100 grams, as tylosin phosphate.

(A) *Indications for use.* For prevention of swine dysentery (vibrionic); aid in the prevention of migration and establishment of large roundworms (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum* spp.) infections.

(B) *Limitations.* Use 100 grams tylosin per ton for at least 3 weeks followed by 40 grams tylosin per ton until market weight; withdraw 24 hours before slaughter. Consult your veterinarian before feeding to severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.

(vi) *Amount per ton.* Pyrantel tartrate, 96 grams (0.0106 percent) and tylosin 40 to 100 grams, as tylosin phosphate.

(A) *Indications for use.* Treatment and control of swine dysentery (vibrionic); aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum* spp.) infections.

(B) *Limitations.* Administer tylosin in feed as tylosin phosphate after treatment with tylosin in drinking water as tylosin base; 0.25 grams per gallon in drinking water for 3 to 10 days, 40 to 100 grams tylosin per ton in feed for 2 to 6 weeks; withdraw 24 hours before slaughter. Consult your veterinarian before feeding to severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.

(vii) *Amount per ton.* Pyrantel tartrate, 96 grams (0.0106 percent) and lincomycin, 40 grams, as lincomycin hydrochloride monohydrate.

(A) *Indications for use.* For control of swine dysentery; aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum* spp.) infections.

(B) *Limitations.* Feed as sole ration; for use in swine on premises with a history of swine dysentery but where symptoms have not yet occurred; not to be fed to swine that weigh more

than 250 pounds; withdraw 6 days before slaughter. Consult your veterinarian before feeding to severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.

(viii) *Amount per ton.* Pyrantel tartrate, 96 grams (0.0106 percent) and lincomycin, 100 grams, then 40 grams, as lincomycin hydrochloride monohydrate.

(A) *Indications for use.* For treatment and control of swine dysentery; aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum* spp.) infections.

(B) *Limitations.* Feed 100 grams per ton for 3 weeks or until signs of disease disappear, followed by 40 grams per ton; feed as sole ration; not to be fed to swine that weigh more than 250 pounds; withdraw 6 days before slaughter. Consult your veterinarian before feeding to severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.

(ix) *Amount per ton.* Pyrantel tartrate, 96 grams (0.0106 percent) and lincomycin, 100 grams, as lincomycin hydrochloride monohydrate.

(A) *Indications for use.* For treatment of swine dysentery; aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum* spp.) infections.

(B) *Limitations.* Feed 100 grams per ton 3 weeks or until signs of disease disappear, followed by 40 grams per ton; feed as sole ration; not to be fed to swine that weigh more than 250 pounds; withdraw 6 days before slaughter. Consult your veterinarian before feeding to severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.

(x) *Amount per ton.* Pyrantel tartrate, 96 grams (0.0106 percent) and lincomycin, 100 or 40 grams.

(A) *Indications for use.* For treatment and/or control of swine dysentery; for removal and control of large roundworm (*Ascaris suum*) infections.

(B) *Limitations.* Administer in accordance with paragraph (c)(2)(i), (c)(2)(ii),

or (c)(2)(iii) of § 558.325 and paragraph (e)(1)(ii)(B) of this section.

(xi) *Amount per ton.* Pyrantel tartrate, 800 grams (0.0881 percent) and lincomycin, 100 or 40 grams.

(A) *Indications for use.* For treatment and/or control of swine dysentery; for removal and control of large roundworm (*Ascaris suum*) and nodular worm (*Oesophagostomum* spp.) infections.

(B) *Limitations.* Administer in accordance with paragraph (c)(2)(i), (c)(2)(ii), or (c)(2)(iii) of § 558.325 and paragraph (e)(1)(iii)(B) of this section.

(xii) *Amount per ton.* Pyrantel tartrate, 96 grams (0.0106 percent) and lincomycin, 200 grams as lincomycin hydrochloride monohydrate.

(A) *Indications for use.* For the reduction in severity of swine mycoplasma pneumonia caused by *Mycoplasma hyopneumoniae*; aid in the prevention of migration and establishment of large roundworms (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum* spp.) infections.

(B) *Limitations.* Feed as sole ration for 21 days; not to be fed to swine that weigh more than 250 pounds; withdraw 6 days before slaughter; consult your veterinarian before feeding to severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.

(C) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(2) *Horses*—(i) *Amount.* 1.2 milligrams per pound (2.64 milligrams per kilogram) of body weight.

(A) *Indications for use.* Prevention of *Strongylus vulgaris* larval infections; control of adult large strongyles (*S. vulgaris*, *S. edentatus*, and *Triodontophorus* spp.), adult and 4th stage larvae small strongyles (*Cyathostomum* spp., *Cylicocyclus* spp., *Cylicostephanus* spp., *Cylicodontophorus* spp., *Poteriostomum* spp.), adult and 4th stage larvae pinworms (*Oxyuris equi*), and adult and 4th stage larvae ascarids (*Parascaris equorum*).

(B) *Limitations.* Administer either as a top-dress (not to exceed 12,000 grams per ton) or mixed in the horse's daily grain ration (not to exceed 1,200 grams per ton) during the time that the animal is at risk of exposure to internal

parasites. Not for use in horses intended for food. Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.485, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.515 Robenidine hydrochloride.

(a) *Approvals.* Type A medicated articles: 30 grams per pound to 063238 in § 510.600(c) of this chapter.

(b) *Special considerations.* Type C feed containing robenidine hydrochloride must be fed within 50 days from the date of manufacture. Do not use in Type B or Type C medicated feeds containing bentonite.

(c) *Related tolerances in edible products.* See § 556.580 of this chapter.

(d) *Conditions of use.* It is used in feed for chickens as follows:

(1) *For broiler and fryer chickens—(i) Amount per ton.* Robenidine hydrochloride, 30 grams (0.0033 percent).

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima*, and *E. necatrix*.

(b) *Limitations.* Do not feed to layers; feed continuously as the sole ration; withdraw 5 days prior to slaughter.

(ii) *Amount per ton.* Robenidine hydrochloride, 30 grams (0.0033 percent) plus roxarsone, 22.5–45.4 grams (.005 percent).

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima*, and *E. necatrix* and increased rate of weight gain.

(b) *Limitations.* Do not feed to layers; feed continuously as the sole ration; withdraw 5 days prior to slaughter; as sole source of organic arsenic. Roxarsone provided by No. 046573, § 510.600(c) of this chapter.

(iii) *Amount per ton.* Robenidine hydrochloride, 30 grams (0.0033 percent) plus chlortetracycline, 100 to 200 grams.

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima*, and *E. necatrix*;

control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline.

(b) *Limitations.* Withdraw 5 days prior to slaughter; do not feed to chickens producing eggs for human consumption; feed continuously as sole ration up to 14 days.

(iv) *Amount per ton.* Robenidine hydrochloride, 30 grams (0.0033 percent) plus chlortetracycline, 200 to 400 grams.

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima*, and *E. necatrix*; control of chronic respiratory disease (CRD) and air sac infection caused by *M. gallisepticum* and *E. coli* susceptible to chlortetracycline.

(b) *Limitations.* Withdraw 5 days prior to slaughter; do not feed to chickens producing eggs for human consumption; feed continuously as sole ration up to 14 days.

(v) *Amount per ton.* Robenidine hydrochloride, 30 grams (0.0033 percent) plus chlortetracycline, 500 grams.

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima*, and *E. necatrix*; as an aid in the reduction of mortality due to *E. coli* susceptible to chlortetracycline.

(b) *Limitations.* Withdraw 5 days prior to slaughter; do not feed to chickens producing eggs for human consumption; feed continuously up to 5 days.

(vi) *Amount per ton.* Robenidine hydrochloride, 30 grams (.0033 percent) plus bacitracin, 4 to 50 grams (as zinc bacitracin).

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima*, and *E. necatrix*, and: in the presence of 4 to 30 grams per ton of bacitracin, for increased rate of weight gain; in the presence of 27 to 50 grams per ton of bacitracin, for improved feed efficiency.

(b) *Limitations.* Feed continuously as sole ration; do not feed to laying chickens; withdraw 5 days prior to slaughter; as zinc bacitracin provided by Nos. 063238 and 046573 in § 510.600(c) of this chapter.

§ 558.530

21 CFR Ch. I (4–1–99 Edition)

(vii) *Amount per ton.* Robenidine hydrochloride, 30 grams (.0033 percent) plus bacitracin, 4 to 50 grams (as bacitracin methylene disalicylate).

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by *E. mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima*, and *E. necatrix*, and; in the presence of 4 to 30 grams per ton of bacitracin, for increased rate of weight gain; in the presence of 27 to 50 grams per ton of bacitracin, for improved feed efficiency.

(b) *Limitations.* Feed continuously as sole ration; do not feed to laying chickens; withdraw 5 days prior to slaughter; as bacitracin methylene disalicylate provided by No. 046573 in § 510.600(c) of this chapter.

(viii) *Amount per ton.* Robenidine hydrochloride, 30 grams (0.0033 percent) plus lincomycin, 2 grams.

(a) *Indications for use.* For increase in rate of weight gain and improved feed efficiency and as an aid in prevention of coccidiosis caused by *E. mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima*, and *E. necatrix*.

(b) *Limitations.* Do not feed to laying hens; feed continuously as the sole ration; withdraw 5 days before slaughter; lincomycin as provided by No. 000009, § 510.600(c) of this chapter; approval for this combination granted to No. 000009 as identified in § 510.600(c) of this chapter.

(2) *For broiler chickens—(i) Amount per ton.* Robenidine hydrochloride, 30 grams (0.0033 percent) plus oxytetracycline, 400 grams.

(ii) *Indications for use.* As an aid in the prevention of coccidiosis caused by *Eimeria mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima*, and *E. necatrix*; control of CRD and air sac infection caused by *Mycoplasma gallisepticum* and *Escherichia coli* susceptible to oxytetracycline.

(iii) *Limitations.* Feed continuously for 7 to 14 days; do not feed to chickens producing eggs for human consumption; withdraw 5 days before slaughter; do not use in feeds containing bentonite; feed must be used within 50 days of manufacture; oxytetracycline

as provided by No. 000069 of this chapter.

[40 FR 13959, Mar. 27, 1975, as amended at 40 FR 33444, Aug. 8, 1975; 40 FR 39859, Aug. 29, 1975; 41 FR 1062, Jan. 6, 1976; 41 FR 28514, July 12, 1976; 41 FR 53002, Dec. 3, 1976; 51 FR 7400, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 53 FR 20843, June 7, 1988; 55 FR 8461, Mar. 8, 1990; 57 FR 8578, Mar. 11, 1992; 61 FR 18082, Apr. 24, 1996; 61 FR 35956, July 9, 1996; 61 FR 51593, Oct. 3, 1996; 62 FR 61626, Nov. 19, 1997; 62 FR 66986, Dec. 23, 1997; 63 FR 17948, Apr. 13, 1998; 63 FR 27845, May 21, 1998]

§ 558.530 Roxarsone.

(a) *Approvals.* Type A medicated articles: (1) 10, 20, and 50 percent to 046573 in § 510.600(c) of this chapter for use as in paragraph (d)(1) of this section.

(2) 10, 20, 50, and 80 percent to 046573 in § 510.600(c) of this chapter for use as in paragraphs (d)(1), (d)(2), (d)(3), and (d)(4) of this section.

(b) *Related tolerances.* See § 556.60 of this chapter.

(c) [Reserved]

(d) *Conditions of use—(1) Growing chickens and growing turkeys—(i) Grams per ton.* Roxarsone 22.7 and 45.4 (0.0025 to 0.005 percent).

(ii) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and improved pigmentation.

(iii) *Limitations.* Withdraw 5 days before slaughter; as sole source of organic arsenic; drug overdose or lack of water may result in leg weakness; feed continuously throughout growing period.

(2) *Growing chickens—(i) Grams per ton.* Roxarsone, 22.7 to 45.4 (0.0025 to 0.005 percent) plus chlortetracycline, 10 to 50.

(A) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and improved pigmentation.

(B) *Limitations.* Do not feed to chickens producing eggs for human consumption; withdraw 5 days before slaughter; as sole source of organic arsenic; drug overdose or lack of water may result in leg weakness; feed continuously throughout growing period.

(ii) *Grams per ton.* Roxarsone 22.7 to 45.4 (0.0025 to 0.005 percent) plus chlortetracycline, 100 to 200.

(A) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and improved pigmentation; control of infectious synovitis caused

by *Mycoplasma synoviae* susceptible to chlortetracycline.

(B) *Limitations.* See paragraph (d)(2)(i)(B) of this section except feed continuously for 7 to 14 days.

(iii) *Grams per ton.* Roxarsone 22.7 to 45.4 (0.0025 to 0.005 percent) plus chlortetracycline, 200 to 400.

(A) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and improved pigmentation; control of chronic respiratory disease (CRD) and air sac infection caused by *M. gallisepticum* and *Escherichia coli* susceptible to chlortetracycline.

(B) *Limitations.* See paragraph (d)(2)(i)(B) of this section except feed continuously for 7 to 14 days.

(iv) *Grams per ton.* Roxarsone 22.7 to 45.4 (0.0025 to 0.005 percent) plus chlortetracycline, 500.

(A) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and improved pigmentation; reduction of mortality due to *E. coli* infections susceptible to chlortetracycline.

(B) *Limitations.* See paragraph (d)(2)(i)(B) of this section except feed for 5 days.

(3) *Growing turkeys*—(i) *Grams per ton.* Roxarsone 22.7 to 45.4 (0.0025 to 0.005 percent) plus chlortetracycline, 10 to 50.

(A) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and improved pigmentation.

(B) *Limitations.* Do not feed to turkeys producing eggs for human consumption; withdraw 5 days before slaughter; as sole source of organic arsenic; drug overdose or lack of water may result in leg weakness; feed continuously throughout growing season.

(ii) *Grams per ton.* Roxarsone 22.7 to 45.4 (0.0025 to 0.005 percent) plus chlortetracycline 200.

(A) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and improved pigmentation; control of infectious synovitis caused by *M. synoviae* susceptible to chlortetracycline.

(B) *Limitations.* See paragraph (d)(3)(i)(B) of this section except that the drug should only be fed continuously for 7 to 14 days.

(iii) *Grams per ton.* Roxarsone 22.7 to 45.4 (0.0025 to 0.005 percent) plus chlortetracycline, 400.

(A) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and improved pigmentation; control of hexamitiasis caused by *Hexamita meleagridis* susceptible to chlortetracycline. Turkey poulters not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by *Salmonella typhimurium* susceptible to chlortetracycline.

(B) *Limitations.* See paragraph (d)(3)(i)(B) of this section except that the drug should only be fed continuously for 7 to 14 days.

(iv) *Amount.* Roxarsone 22.7 to 45.4 grams per ton (0.0025 to 0.005 percent) plus chlortetracycline, 25 milligrams per pound of body weight daily.

(A) *Indications for use.* For increased rate of weight gain, improved feed efficiency, and improved pigmentation; control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline.

(B) *Limitations.* See paragraph (d)(3)(i)(B) of this section except that the drug should only be fed continuously for 7 to 14 days.

(4) *Growing-finishing swine*—(i) *Grams per ton.* Roxarsone 22.7 to 34.1 (0.0025 to 0.00375 percent).

(A) *Indications for use.* For increased rate of weight gain and improved feed efficiency.

(B) *Limitations.* Withdraw 5 days before slaughter; as sole source of organic arsenic; feed continuously throughout growing season.

(ii) *Grams per ton.* Roxarsone 22.7 to 34.1 (0.0025 to 0.00375 percent) plus chlortetracycline, 400 (to administer 10 milligrams per pound of body weight).

(A) *Indications for use.* For increased rate of weight gain and improved feed efficiency; treatment of bacterial enteritis caused by *E. coli* and *S. choleraesuis* and bacterial pneumonia caused by *P. multocida* susceptible to chlortetracycline.

(B) *Limitations.* Withdraw 5 days before slaughter; as sole source of organic arsenic; feed for not more than 14 days.

(iii) *Grams per ton.* Roxarsone 181.5 (0.02 percent).

§ 558.550

21 CFR Ch. I (4–1–99 Edition)

(A) *Indications for use.* For the treatment of swine dysentery.

(B) *Limitations.* Feed for not more than 6 consecutive days; if improvement is not observed, consult a veterinarian; withdraw 5 days before slaughter; as a sole source or organic arsenic; animals must consume enough medicated feed to provide a therapeutic dose.

(iv) *Grams per ton.* Roxarsone, 181.5 (0.02 percent) plus chlortetracycline, 10 to 50.

(A) *Indications for use.* For the treatment of swine dysentery; increased rate of weight gain and improved feed efficiency.

(B) *Limitations.* See paragraph (d)(4)(iii)(B) of this section.

(v) *Grams per ton.* Roxarsone, 181.5 (0.02 percent) plus chlortetracycline, 400.

(A) *Indications for use.* For the treatment of swine dysentery; treatment of bacterial enteritis caused by *E. coli* and *S. choleraesuis* and bacterial pneumonia caused by *P. multocida* susceptible to chlortetracycline.

(B) *Limitations.* See paragraph (d)(4)(iii)(B) of this section.

(5) *Permitted combinations.* It may be used in accordance with this section in combination as follows:

(i) Aklomide as in § 558.35.

(ii) Amprolium as in § 558.55.

(iii) Amprolium and ethopabate as in § 558.58.

(iv) Bacitracin methylene disalicylate as in § 558.76.

(v) Bacitracin zinc as in § 558.78.

(vi) Bambermycins and bambermycins plus certain anticoccidials as in § 558.95.

(vii) [Reserved]

(viii) Chlortetracycline as in § 558.128.

(ix) Clopidol as in § 558.175.

(x) Decoquinatone alone or in combination as in § 558.195.

(xi) Monensin as in § 558.355.

(xii) Nequinatone as in § 558.365.

(xiii) Nicarbazine as in § 558.366.

(xiv) Nitroimidazole and sulfanitran as in § 558.376.

(xv) Robenidine hydrochloride as in § 558.515.

(xvi) Sulfadimethoxine, ormetoprim as in § 558.575.

(xvii) Zoalene as in § 558.680.

(xviii) Penicillin and zoalene as in § 558.680.

(xix) Lasalocid as in § 558.311.

(xx) Lasalocid sodium and lincomycin as in § 558.311.

(xxi) Monensin and virginiamycin as in § 558.355.

(xxii) Monensin and bacitracin zinc as in § 558.355.

(xxiii) Narasin with bacitracin methylene disalicylate as in § 558.363.

(xxiv) Semduramicin with bacitracin methylene disalicylate as in § 558.555.

(6) *Additional combinations.* (i) Roxarsone may be used in combination “as an aid in the reduction of lesions due to *E. tenella*” as follows:

(A) Lasalocid as in § 558.311.

(B) Lasalocid plus bacitracin methylene disalicylate as in § 558.311.

(C) Lasalocid plus lincomycin as in § 558.311.

(D) Lasalocid and bacitracin zinc as in § 558.311.

(ii) Roxarsone may be used in combination with salinomycin as in § 558.550.

(iii) Roxarsone may be used in combination with salinomycin and bacitracin methylene disalicylate as in § 558.550.

(iv) Roxarsone may be used in combination with salinomycin and bacitracin zinc as in § 558.550.

(v) Roxarsone may be used in combination with halofuginone hydrobromide and bacitracin methylene disalicylate as in § 558.265.

(vi) Roxarsone may be used in combination with narasin as in § 558.363 of this part.

(vii) Roxarsone may be used in combination with salinomycin and chlortetracycline as in § 558.550.

[46 FR 52331, Oct. 27, 1981]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.530, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.550 Salinomycin.

(a) *Approvals.* Type A medicated articles containing 30 or 60 grams of salinomycin activity per pound (as salinomycin sodium biomass) as follows:

(1) To 063238 in § 510.600(c) of this chapter for use as in paragraph (d) of this section.

(2) To 012799 for use as in paragraphs (d)(1)(i), (d)(1)(iii) through (d)(1)(xvi), and (d)(3)(i) through (d)(3)(iii) of this section.

(3) To 046573 for use as in paragraphs (d)(1)(xv) and (d)(1)(xvi) of this section.

(b)–(c) [Reserved]

(d) *Conditions of use.* (1) Broilers: It is used as follows:

(i)(a) *Amount per ton.* Salinomycin 40 to 60 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(c) *Limitations.* Feed continuously as sole ration. Do not feed to layers. Not approved for use with pellet binders. May be fatal if accidentally fed to adult turkeys or horses.

(ii)(a) *Amount per ton.* Salinomycin 40 to 60 grams and roxarsone 45.4 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, including some field strains of *E. tenella* which are more susceptible to roxarsone combined with salinomycin than to salinomycin alone.

(c) *Limitations.* Feed continuously as sole ration. Use as sole source of organic arsenic. Not approved for use with pellet binders. Do not feed to layers. May be fatal if accidentally fed to adult turkeys or horses. Withdraw 5 days before slaughter. Roxarsone as provided by No. 011526 or 046573 in § 510.600(c) of this chapter.

(iii)(a) *Amount per ton.* Salinomycin 40 to 60 grams and bacitracin methylene disalicylate 4 to 30 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati* for increased rate of weight gain and improved feed efficiency.

(c) *Limitation.* Feed continuously as sole ration. Not approved for use with pellet binders. Do not feed to layers. May be fatal if accidentally fed to adult turkeys or horses. Bacitracin methylene disalicylate as provided by No. 046573 in § 510.600(c) of this chapter.

(iv)(a) *Amount per ton.* Salinomycin 40 to 60 grams with roxarsone 45.4 grams and bacitracin methylene disalicylate 4 to 50 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, including some field strains of *E. tenella* which are more susceptible to roxarsone combined with salinomycin than to salinomycin alone; for increased rate of weight gain.

(c) *Limitations.* Feed continuously as sole ration. Use as sole source of organic arsenic. Not approved for use with pellet binders. Do not feed to layers. May be fatal if accidentally fed to adult turkeys or horses. Withdraw 5 days before slaughter. Roxarsone and bacitracin as provided by No. 046573 in § 510.600(c) of this chapter.

(v)(a) *Amount per ton.* Salinomycin 40 to 60 grams per ton with roxarsone 22.7 to 45.4 grams per ton.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima* and for improved feed efficiency.

(c) *Limitations.* Feed continuously as sole ration. Use as sole source of organic arsenic. Not approved for use with pellet binders. Do not feed to layers. May be fatal if accidentally fed to adult turkeys or to horses. Withdraw 5 days before slaughter. Roxarsone as provided by No. 046573 in § 510.600(c) of this chapter.

(vi)(a) *Amount per ton.* Salinomycin 40 to 60 grams and bacitracin methylene disalicylate 4 to 50 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and for improved feed efficiency.

(c) *Limitations.* Feed continuously as sole ration. Not approved for use with pellet binders. Do not feed to layers. May be fatal if accidentally fed to adult turkeys or horses. Bacitracin MD as provided by No. 046573 in § 510.600(c) of this chapter.

(vii)(a) *Amount per ton.* Salinomycin 40 to 60 grams and bacitracin zinc 10 to 50 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and for increased rate of weight gain.

(c) *Limitations.* Feed continuously as sole ration. Not approved for use with pellet binders. Do not feed to layers. May be fatal if accidentally fed to adult turkeys or horses. Bacitracin zinc as provided by No. 063238 in § 510.600(c) of this chapter.

(viii)(a) *Amount per ton.* Salinomycin 40 to 60 grams with roxarsone 45.4 grams and bacitracin zinc 4 to 50 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, including some field strains of *E. tenella* which are more susceptible to roxarsone combined with salinomycin than to salinomycin alone; for increased rate of weight gain and improved feed efficiency.

(c) *Limitations.* See paragraph (b)(1)(iv)(c) of this section.

(ix)(a) *Amount per ton.* Salinomycin 40 to 60 grams with roxarsone 34.1 grams and bacitracin zinc 10 to 50 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and for increased rate of weight gain and improved feed efficiency.

(c) *Limitations.* Feed continuously as sole ration. Use as sole source of organic arsenic. Do not feed to layers. May be fatal if accidentally fed to adult turkeys or horses. Withdraw 5 days before slaughter. Roxarsone as provided by No. 046573 and bacitracin as provided by No. 063238 in § 510.600(c) of this chapter.

(x)(a) *Amount per ton.* Salinomycin 40 to 60 grams and virginiamycin 5 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and for increased rate of weight gain and improved feed efficiency.

(c) *Limitations.* Feed continuously as sole ration. Not approved for use with pellet binders. Do not feed to layers or to chickens over 16 weeks of age. May be fatal if accidentally fed to adult turkeys or horses. Virginiamycin as provided by No. 053571 in § 510.600(c) of this chapter.

(xi)(a) *Amount per ton.* Salinomycin 40 to 60 grams and virginiamycin 5 to 15 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and for increased rate of weight gain.

(c) *Limitations.* See paragraph (b)(1)(x)(c) of this section.

(xii)(a) *Amount per ton.* Salinomycin 40 to 60 grams, virginiamycin 5 grams, and roxarsone 45.4 grams.

(b) *Indications for use.* For prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, including some field strains of *E. tenella* which are more susceptible to roxarsone combined with salinomycin than to salinomycin alone, and for improved feed efficiency.

(c) *Limitations.* Feed continuously as sole ration. Withdraw 5 days prior to slaughter. Use as sole source of organic arsenic. Not approved for use with pellet binders. Do not feed to layers. May be fatal if accidentally fed to adult turkeys or horses. Virginiamycin as provided by No. 053571 in § 510.600(c) of this chapter. Roxarsone as provided by No. 046573 in § 510.600(c) of this chapter.

(xiii)(a) *Amount per ton.* Salinomycin 40 to 60 grams and lincomycin 2 to 4 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati* and for improved feed efficiency.

(c) *Limitations.* Feed continuously as sole ration. Not approved for use with pellet binders. Do not feed to layers. Do not allow horses, adult turkeys, guinea pigs, rabbits, hamsters, or ruminants access to this feed. Ingestion by these species may result in severe gastrointestinal effects or may be fatal. Lincomycin hydrochloride monohydrate as provided by No. 000009 in § 510.600(c) of this chapter.

(xiv)(a) *Amount per ton.* Salinomycin 40 to 60 grams, roxarsone 45.4 grams, and lincomycin 2 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, including some field strains of *E. tenella*

that are more susceptible to roxarsone combined with salinomycin than to salinomycin alone, and for improved feed efficiency.

(c) *Limitations.* Feed continuously as sole ration. Not approved for use with pellet binders. Drug overdose or lack of water may result in leg weakness. Do not feed to layers. Do not allow horses, adult turkeys, guinea pigs, rabbits, hamsters, or ruminants access to this feed. Ingestion by these species may result in severe gastrointestinal effects or may be fatal. Withdraw 5 days before slaughter. Lincomycin hydrochloride monohydrate as provided by No. 000009 in §510.600(c) of this chapter. Roxarsone as provided by No. 046573 in §510.600(c) of this chapter.

(xv)(a) *Amount per ton.* Salinomycin 40 to 60 grams, chlortetracycline 500 grams, and roxarsone 45.4 grams.

(b) *Indications for use.* For prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, including some field strains of *E. tenella* which are more susceptible to roxarsone combined with salinomycin than to salinomycin alone, and as an aid in the reduction of mortality due to *E. coli* infections susceptible to such treatment.

(c) *Limitations.* Do not feed to layers. In feeds containing 0.8 percent dietary calcium, not to be fed for more than 5 days. Not approved for use with pellet binders. Withdraw 5 days before slaughter. May be fatal if accidentally fed to adult turkeys or to horses. Chlortetracycline as provided by Nos. 046573 and 063238 and roxarsone as provided by No. 046573 in §510.600(c) of this chapter.

(xvi)(a) *Amount per ton.* Salinomycin 40 to 60 grams and chlortetracycline 500 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and as an aid in the reduction of mortality due to *E. coli* infections susceptible to such treatment.

(c) *Limitations.* Do not feed to layers. In feeds containing 0.8 percent dietary calcium. Not to be fed for more than 5 days. Not approved for use with pellet binders. Withdraw 24 hours before slaughter. May be fatal if accidentally fed to adult turkeys or horses. Chlor-

tetracycline as provided by No. 063238 in §510.600(c) of this chapter.

(xvii)(A) *Amount per ton.* Salinomycin 40 to 60 grams with roxarsone 34.1 or 45.4 grams and bacitracin methylene disalicylate 4 to 50 grams.

(B) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, including some field strains of *E. tenella* that are more susceptible to roxarsone combined with salinomycin than to salinomycin alone; for increased rate of weight gain. Use of 34.1 or 45.4 grams per ton roxarsone is indicated to meet the *E. tenella* challenge which varies with environmental and management conditions.

(C) *Limitations.* Feed continuously as sole ration. Use as sole source of organic arsenic. Not approved for use with pellet binders. Do not feed to laying chickens. May be fatal if accidentally fed to adult turkeys or horses. Poultry should have access to drinking water at all times. Overdosage or lack of water may result in leg weakness or paralysis. Withdraw 5 days before slaughter. Salinomycin as provided by No. 000004 in §510.600(c) of this chapter. Roxarsone and bacitracin as provided by No. 046573 in §510.600(c) of this chapter.

(2) *Quail*—(i)(a) *Amount per ton.* Salinomycin 50 grams.

(b) *Indications for use.* For the prevention of coccidiosis caused by *E. dispersa* and *E. lettyae*.

(c) *Limitations.* Feed continuously as sole ration. Not approved for use with pellet binders. May be fatal if accidentally fed to adult turkeys or horses.

(ii) [Reserved]

(3) *Roaster and replacement (breeder and layer) chickens:* It is used as follows:

(i) *Amount per ton.* Salinomycin 40 to 60 grams.

(ii) *Indications for use.* For prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(iii) *Limitations.* Feed continuously as sole ration. Do not feed to laying hens producing eggs for human consumption. Not approved for use with pellet binders. May be fatal if accidentally fed to horses or adult turkeys.

§ 558.555

21 CFR Ch. I (4–1–99 Edition)

(4) *Permitted combinations.* Salinomycin may be used as in this section in combinations as follows:

(i) Bambermycins and roxarsone as in § 558.95.

(ii) Bambermycins as in § 558.95.

(iii) Oxytetracycline as in § 558.450.

[48 FR 30616, July 5, 1983]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.550, see the List of CFR Sections Affected in the Finding Aids section of this volume.

EDITORIAL NOTE: At 62 FR 29014, May 29, 1997, in § 558.550, paragraph (b) was redesignated as paragraph (d). At 62 FR 60781, Nov. 13, 1997, paragraph (b)(1)(vii)(c) was amended by removing “No. 000004” and adding in its place “Nos. 000004 and 046573” and at 62 FR 66985, Dec. 23, 1997, paragraph (b)(1)(ix)(c) was amended by removing “No. 000004” and adding in its place “Nos. 000004 and 046573”. Since § 558.550(b) does not exist, these amendments could not be incorporated.

§ 558.555 Semduramicin.

(a) *Approvals.* Type A medicated article containing 5.13 percent semduramicin sodium (equivalent to 50 grams semduramicin per kilogram or 22.7 grams per pound) to 000069 in § 510.600(c) of this chapter.

(b) *Conditions of use.* (1) Broilers:

(i) *Amount.* Semduramicin: 25 parts per million.

(ii) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria tenella*, *E. acervulina*, *E. maxima*, *E. brunetti*, *E. necatrix*, and *E. mivati/E. mitis*.

(iii) *Limitations.* Do not feed to laying hens.

(2) *Amount.* Semduramicin 22.7 grams with bacitracin methylene disalicylate 10 to 50 grams and roxarsone 45.4 grams per ton.

(i) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati/E. mitis*, *E. necatrix*, and *E. tenella*, including some field strains of *E. tenella* that are more susceptible to semduramicin combined with roxarsone than semduramicin alone, and for improved feed efficiency.

(ii) *Limitations.* Feed continuously as sole ration. Use feed within 2 weeks of production. Withdraw 5 days before slaughter. Do not feed to laying hens. Use as sole source of organic arsenic. Poultry should have access to drinking

water at all times. Drug overdosage or lack of water intake may result in leg weakness or paralysis. Semduramicin as provided by 000069 in § 510.600(c) of this chapter, bacitracin methylene disalicylate and roxarsone as provided by 046573.

(3) *Amount.* Semduramicin 22.7 grams with bacitracin methylene disalicylate 10 to 50 grams per ton.

(i) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati/E. mitis*, *E. necatrix*, and *E. tenella*, and for improved feed efficiency in broiler chickens.

(ii) *Limitations.* Feed continuously as sole ration. Use feed within 2 weeks of production. Do not feed to laying hens. Semduramicin as provided by 000069, bacitracin methylene disalicylate as provided by 046573 in § 510.600(c) of this chapter.

(4) *Amount.* Semduramicin 22.7 grams with roxarsone 45.4 grams per ton.

(i) *Indications for use.* For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati/E. mitis*, *E. necatrix*, and *E. tenella*, including some field strains of *E. tenella* that are more susceptible to semduramicin combined with roxarsone than semduramicin alone.

(ii) *Limitations.* Feed continuously as sole ration. Withdraw 5 days before slaughter. For broiler chickens only. Do not feed to laying hens. Use as sole source of organic arsenic. Roxarsone as provided by 046573, semduramicin as provided by 000069 in § 510.600(c) of this chapter.

[59 FR 17477, Apr. 13, 1994, as amended at 60 FR 57928, Nov. 24, 1995; 61 FR 29481, June 11, 1996; 61 FR 43451, Aug. 23, 1996; 61 FR 66584, Dec. 18, 1996; 62 FR 66985, Dec. 23, 1997]

§ 558.575 Sulfadimethoxine, ormetoprim.

(a) *Approvals.* Type A medicated articles to sponsors as identified in § 510.600(c) of this chapter for uses as in paragraph (c) of this section as follows:

(1) 25 percent sulfadimethoxine and 15 percent ormetoprim to 063238 for use for poultry as in paragraphs (c)(1), (2), (3), and (4) of this section.

(2) 25 percent sulfadimethoxine and 5 percent ormetoprim to 063238 for use

for fish as in paragraphs (c)(5) and (6) of this section.

(b) *Related tolerances.* See §§ 556.490 and 556.640 of this chapter.

(c) *Conditions of use.* It is used in feeds for animals as follows:

(1) *Broiler chickens*—(i) *Amount per ton.* Sulfadimethoxine, 113.5 grams (0.0125 percent) plus ormetoprim, 68.1 grams (0.0075 percent).

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by all *Eimeria* species known to be pathogenic to chickens, namely, *E. tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, and bacterial infections due to *H. gallinarum* (infectious coryza), *E. coli* (colibacillosis) and *P. multocida* (fowl cholera).

(b) *Limitations.* Feed as sole ration; withdraw 5 days before slaughter.

(ii) *Amount per ton.* Sulfadimethoxine, 113.5 grams (0.0125 percent) plus ormetoprim, 68.1 grams (0.0075 percent) plus roxarsone, 22.7 grams (0.0025 percent).

(a) *Indications for use.* As an aid in the prevention of coccidiosis caused by all *Eimeria* species known to be pathogenic to chickens, namely *E. tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, and bacterial infections due to *H. gallinarum* (infectious coryza), *E. coli*, (colibacillosis); and *P. multocida* (fowl cholera); growth promotion and feed efficiency; improving pigmentation.

(b) *Limitations.* Withdraw 5 days before slaughter; as sole source of organic arsenic.

(2) *Replacement chickens*—(i) *Amount per ton.* Sulfadimethoxine, 113.5 grams (0.0125 percent) plus ormetoprim, 68.1 grams (0.0075 percent).

(ii) *Indications for use.* As an aid in the prevention of coccidiosis caused by all *Eimeria* species known to be pathogenic to chickens, namely *E. tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, and bacterial infections due to *H. galmaxima*, and bacterial infections due to *H. gallinarum* (infectious coryza), *E. coli* (colibacillosis) and *P. multocida* (fowl cholera).

(iii) *Limitations.* Feed as a sole ration; do not feed to chickens over 16 weeks (112 days) of age; withdraw 5 days before slaughter.

(3) *Turkeys*—(i) *Amount per ton.* Sulfadimethoxine, 56.75 grams (0.00625 percent) plus ormetoprim, 34.05 grams (0.00375 percent).

(ii) *Indications for use.* As an aid in the prevention of coccidiosis caused by all *Eimeria* species known to be pathogenic to turkeys, namely, *E. adenoides*, *E. gallopavonis*, and *E. meleagritidis* and bacterial infection due to *P. multocida* (fowl cholera).

(iii) *Limitations.* Do not feed to turkeys producing eggs for food; withdraw 5 days before slaughter.

(4) *Ducks*—(i) *Amount per ton.* Sulfadimethoxine, 227 grams (0.025 percent) plus ormetoprim, 136.2 grams (0.015 percent).

(a) *Indications for use.* As an aid in the control of bacterial infections due to *P. multocida* (fowl cholera) in ducks, including breeding ducks.

(b) *Limitations.* Feed as sole ration for 7 days; withdraw 5 days before slaughter; medication should be started at the first signs of infection; do not feed to ducks producing eggs for food.

(ii) *Amount per ton.* Sulfadimethoxine, 454 grams (0.05 percent) plus ormetoprim, 272.4 grams (0.03 percent).

(a) *Indications for use.* As an aid in the control of bacterial infections due to *E. coli*, *P. anatipestifer*, and severe challenge of *P. multocida* (fowl cholera) in ducks.

(b) *Limitations.* Feed as a sole ration for 7 days; withdraw 5 days before slaughter; medication should be started at the first signs of infection; not for breeding ducks; do not feed to ducks producing eggs for food.

(5) *Salmonids*—(i) *Amount.* 50 milligrams of active ingredients per kilogram of body weight per day.

(ii) *Indications of use.* For the control of furunculosis in salmonids (trout and salmon) caused by *Aeromonas salmonicida* strains susceptible to sulfadimethoxine and ormetoprim combination.

(iii) *Limitations.* Administer for 5 consecutive days; withdraw 42 days before release as stocker fish or slaughter.

(6) *Catfish*—(i) *Amount.* 50 milligrams of active ingredients per kilogram of body weight per day.

(ii) *Indications for use.* For control of enteric septicemia of catfish caused by

§ 558.579

Edwardsiella ictaluri strains susceptible to sulfadimethoxine and ormetoprim combination.

(iii) *Limitations*. Administer for 5 consecutive days; withdraw 3 days before slaughter or release as stocker fish.

[40 FR 13959, Mar. 27, 1975, as amended at 42 FR 13550, Mar. 11, 1977; 49 FR 33442, Aug. 23, 1984; 49 FR 46371, Nov. 26, 1984; 51 FR 7400, Mar. 3, 1986; 51 FR 18884, May 23, 1986; 52 FR 2686, Jan. 26, 1987; 54 FR 1686, Jan. 17, 1989; 63 FR 27846, May 21, 1998]

§ 558.579 Sulfaethoxyypyridazine.

(a) *Approvals*. Type A medicated articles: 5.5 percent for swine, and 5.5 and 11 percent for cattle to 010042 in § 510.600(c) of this chapter.

(b) *Related tolerances*. See § 556.650 of this chapter.

(c) *Conditions of use*. It is used in animal feed as follows:

(1) *Swine*—(i) *Amount*. 25 milligrams per pound body weight per day.

(ii) *Indications for use*. For treatment of bacterial scours, pneumonia, enteritis, bronchitis, septicemia accompanying *Salmonella choleraesuis* infection.

(iii) *Limitations*. Administer 1,000 grams per ton (0.11 percent) for not less than 4 days nor more than 10 days; do not treat within 10 days of slaughter; as sole source of sulfonamide; for use by or on the order of a licensed veterinarian.

(2) *Cattle*—(i) *Amount*. 25 milligrams per pound body weight per day.

(ii) *Indications for use*. For treatment of respiratory infections (pneumonia, shipping fever), foot rot, calf scours; as adjunctive therapy in septicemia accompanying mastitis and metritis.

(iii) *Limitations*. Administer as a top dressing or in mixed feed for 4 days; do not treat within 16 days of slaughter; as sole source of sulfonamide; milk that has been taken from animals during treatment and for 72 hours (6 milkings) after the latest treatment must not be used for food; for use by or on the order of a licensed veterinarian.

[41 FR 11005, Mar. 15, 1976, as amended at 51 FR 7400, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987]

§ 558.582 Sulfamerazine.

(a) *Approvals*. Type A medicated articles: 99 percent to 063238 in § 510.600(c) of this chapter.

21 CFR Ch. I (4–1–99 Edition)

(b) *Related tolerances*. See § 556.660 of this chapter.

(c) *Conditions of use*. It is used in fish feed for rainbow trout, brook trout, and brown trout as follows:

(1) *Amount*. 10 grams of sulfamerazine per 100 pounds of fish per day.

(2) *Indications for use*. Control of furunculosis.

(3) *Limitations*. Treat for not more than 14 days; do not treat within 3 weeks of marketing or stocking in stream open to fishing.

[41 FR 11005, Mar. 15, 1976, as amended at 51 FR 7400, Mar. 3, 1986; 61 FR 18082, Apr. 24, 1996; 63 FR 27846, May 21, 1998]

§ 558.586 Sulfaquinoxoline.

(a) *Approvals*. Type A medicated articles: 40 percent to 050749 in § 510.600(c) of this chapter.

(b)–(c) [Reserved]

(d) *NAS/NRC status*. The conditions of use specified in this section have been reviewed by NAS/NRC and are found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency information. Applications must be accompanied by a written commitment to undertake the human safety studies required by FDA.

(e) *Special considerations*. (1) For control of outbreaks of disease, medication should be initiated as soon as the diagnosis is determined. Medicated chickens, turkeys, and rabbits must actually consume enough medicated feed which provides a recommended dose of approximately 3.5 to 60 milligrams per pound per day in chickens, 2.5 to 100 milligrams per pound per day in turkeys, and 2.8 to 68 milligrams per pound per day in rabbits depending upon age and class of animal, ambient temperature, and other factors. Consult a veterinarian or poultry pathologist for diagnosis.

(2) [Reserved]

(f) *Conditions of use*. It is used as follows:

(1) *Chickens*—(i) *Amount*. 0.015 percent.

(a) *Indications for use*. As an aid in preventing outbreaks of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, and *E. brunetti* under average conditions of exposure.

(b) *Limitations.* Feed continuously from the time birds are placed on litter and continue past the age when coccidiosis is ordinarily a hazard. If death losses exceed 0.5 percent in a 2-day period, obtain a laboratory diagnosis. If coccidiosis is the cause, use the sulfaquinoxaline levels recommended for control of outbreaks, returning to the original dosage schedule after the outbreak has subsided. Losses may result from intercurrent disease, other conditions affecting drug intake, or variant strains of coccidia species which can contribute to the virulence of coccidiosis under field conditions. Do not treat chickens within 10 days of slaughter. Do not medicate chickens producing eggs for human consumption.

(ii) *Amount.* 0.0175 percent.

(a) *Indications for use.* As an aid in preventing outbreaks of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, and *E. brunetti* where excessive exposure to coccidia is increased due to overcrowding or other management factors.

(b) *Limitations.* Feed continuously from the time birds are placed on litter and continue past the age when coccidiosis is ordinarily a hazard. If death losses exceed 0.5 percent in a 2-day period, obtain a laboratory diagnosis. If coccidiosis is the cause, use the sulfaquinoxaline levels recommended for control of outbreaks, returning to the original dosage schedule after the outbreak has subsided. Losses may result from intercurrent disease, other conditions affecting drug intake, or variant strains of coccidia species which can contribute to the virulence of coccidiosis under field conditions. Do not treat chickens within 10 days of slaughter. Do not medicate chickens producing eggs for human consumption.

(iii) *Amount.* 0.1 to 0.05 percent.

(a) *Indications for use.* As an aid in controlling outbreaks of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, and *E. brunetti*.

(b) *Limitations.* Feed at 0.1 percent level for first 48 to 72 hours. Skip 3 days; 0.05 percent for 2 days, skip 3 days; 0.05 percent for 2 days. If bloody droppings recur, give 0.05 percent for another 2 days. Do not treat chickens

within 10 days of slaughter. Do not medicate chickens producing eggs for human consumption.

(2) *Turkeys*—(i) *Amount.* 0.0175 percent.

(a) *Indications for use.* As an aid in preventing outbreaks of coccidiosis caused by *Eimeria meleagritidis* and *E. adenoides*.

(b) *Limitations.* Feed 0.0175 percent continuously during time birds are closely confined. May be continued for week to 10 days after flock is transferred to range to reduce danger of an outbreak following moving of the flock. Do not treat turkeys within 10 days of slaughter. Do not medicate turkeys producing eggs for human consumption.

(ii) *Amount.* 0.05 percent.

(a) *Indications for use.* As an aid in controlling outbreaks of coccidiosis caused by *Eimeria meleagritidis*, and *E. adenoides*.

(b) *Limitations.* Feed 0.05 percent for 2 days. Follow with 3 days on regular feed and 2 more days on 0.05 percent sulfaquinoxaline feed. Again follow with 3 days on regular feed and 2 more days on 0.05 percent sulfaquinoxaline feed. Continue this schedule if necessary till all signs of the outbreaks have subsided. Do not treat turkeys within 10 days of slaughter. Do not medicate turkeys producing eggs for human consumption.

(3) *Chickens and turkeys*—(i) *Amount.* 0.05 or 0.1 percent.

(a) *Indications for use.* As an aid in the control of acute fowl cholera caused by *Pasteurella multocida* susceptible to sulfaquinoxaline and fowl typhoid caused by *Salmonella gallinarum* susceptible to sulfaquinoxaline.

(b) *Limitations.* Feed 0.1 percent for 48 to 72 hours. Mortality should be brought under control. After medication, move birds to clean ground or to a clean house. If disease recurs, use 0.05 percent in feed again for 2 days. Do not treat chickens or turkeys within 10 days of slaughter for food. Do not medicate chickens or turkeys producing eggs for human consumption.

(ii) [Reserved]

(4) *Rabbits*—(i) *Amount.* 0.025 percent.

(a) *Indications for use.* As an aid in preventing coccidiosis caused by *Eimeria stiedae*.

§ 558.600

21 CFR Ch. I (4–1–99 Edition)

(b) *Limitations.* Treatment to be started after weaning. Feed continuously for 30 days or feed medicated feed for 2 days out of every week until marketing. Do not treat within 10 days of slaughter.

(ii) *Amount.* 0.1 percent.

(a) *Indications for use.* As an aid in controlling outbreaks of coccidiosis caused by *Eimeria stiedae*.

(b) *Limitations.* Feed for 2 weeks. Do not treat within 10 days of slaughter.

[48 FR 3965, Jan. 28, 1983, as amended at 51 FR 7400, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 55 FR 29843, July 23, 1990; 59 FR 33197, June 28, 1994]

§ 558.600 Tiamulin.

(a) *Approvals.* Type A article containing 5, 10, or 113.4 grams of tiamulin (as tiamulin hydrogen fumarate) per pound to 000010 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.738 of this chapter.

(c) *Conditions of use in swine*—(1) *Amount.* 35 grams of tiamulin per ton.

(i) *Indications for use.* For control of swine dysentery associated with *Serpulina (Treponema) hyodysenteriae* susceptible to tiamulin.

(ii) *Limitations.* Feed continuously as sole ration on premises with a history of swine dysentery but where signs of disease have not yet occurred or following approved treatment of disease. Withdraw 2 days before slaughter. Not for use in swine over 250 pounds body weight. Use as only source of tiamulin. Swine being treated with tiamulin should not have access to feeds containing polyether ionophores (e.g., lasalocid, monensin, narasin, or salinomycin) as adverse reactions may occur.

(2) *Amount.* 10 grams of tiamulin per ton.

(i) *Indications for use.* For increased rate of weight gain and improved feed efficiency.

(ii) *Limitations.* Feed continuously as the sole ration. Not for use in swine weighing over 250 pounds. Use as sole source of tiamulin. Swine being treated with tiamulin should not have access to feeds containing polyether ionophores (e.g., lasalocid, monensin, narasin, or salinomycin) as adverse reactions may occur.

(3) *Amount.* 200 grams of tiamulin per ton.

(i) *Indications for use.* Treatment of swine dysentery associated with *Serpulina (Treponema) hyodysenteriae* susceptible to tiamulin.

(ii) *Limitations.* Feed continuously as the sole feed for 14 consecutive days. Withdraw feed 7 days before slaughter. Not for use in swine over 113.40 kilograms (250 pounds) body weight. Use as the only source of tiamulin. Swine being treated with tiamulin should not have access to feeds containing polyether ionophores (e.g., monensin, lasalocid, narasin, semduramicin, or salinomycin) as adverse reactions may occur.

(4) *Amount per ton.* 35 grams of tiamulin (as tiamulin hydrogen fumarate), plus the equivalent of approximately 400 grams of chlortetracycline hydrochloride varying with body weight and feed consumption to provide 10 milligrams of chlortetracycline per pound of body weight daily.

(i) *Indications for use.* Treatment of swine bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to chlortetracycline, and control of swine dysentery associated with *Serpulina (Treponema) hyodysenteriae* susceptible to tiamulin.

(ii) *Limitations.* Feed continuously as sole ration for 14 days. Not for use in swine weighing over 250 pounds. Use as only source of chlortetracycline and tiamulin. Swine being treated with tiamulin should not have access to feeds containing polyether ionophores (e.g., monensin, salinomycin, narasin, semduramicin, and lasalocid) as adverse reactions may occur. If signs of toxicity occur, discontinue use. Withdraw 2 days before slaughter. As chlortetracycline calcium complex, Type A medicated articles containing the equivalent of 50 to 100 grams per pound of chlortetracycline hydrochloride provided by 046573, 053389, and 063238 in § 510.600(c) of this chapter.

[52 FR 26956, July 17, 1987, as amended at 53 FR 39257, Oct. 6, 1988; 59 FR 41975, Aug. 16, 1994; 62 FR 12086, Mar. 14, 1997; 62 FR 35077, June 30, 1997; 63 FR 27846, May 21, 1998; 63 FR 52969, Oct. 2, 1998]

§ 558.615 Thiabendazole.

(a) *Approvals.* Dry Type A medicated articles: 22, 44.1, 66.1, and 88.2 percent to 050604 in § 510.600(c) of this chapter. The 66.1 percent Type A is solely for the manufacture of cane molasses liquid Type B feed which is mixed in dry feeds. The 88.2 percent Type A is used solely for the manufacture of an aqueous slurry for adding to a Type C dry cattle feed.

(b) *Special considerations.* Do not use in Type B or Type C medicated feed containing bentonite.

(c) *Related tolerances.* See § 556.730 of this chapter.

(d) *Conditions of use.* It is used in feed for animals as follows:

(1) *Cattle—(i) Amount.* 3 grams per 100 lb. body weight.

(a) *Indications for use.* Control of infections of gastrointestinal roundworms (*Trichostrongylus* spp., *Haemonchus* spp., *Ostertagia* spp., *Nematodirus* spp., *Oesophagostomum radiatum*).

(b) *Limitations.* Use 3 grams per 100 lb. body weight at a single dose; may repeat once in 2 to 3 weeks; do not treat animals within 3 days of slaughter; milk taken from treated animals within 96 hours (8 milkings) after the latest treatment must not be used for food.

(ii) *Amount.* 5 grams per 100 lb. body weight.

(a) *Indications for use.* Control of severe infections of gastrointestinal roundworms (*Trichostrongylus* spp., *Haemonchus* spp., *Ostertagia* spp., *Nematodirus* spp., *Oesophagostomum radiatum*); control of infections of *Cooperia* spp.

(b) *Limitations.* 5 grams per 100 lb. body weight at a single dose or divided into 3 equal doses, administered 1 dose each day, on succeeding days; may repeat once in 2 to 3 weeks; do not treat animals within 3 days of slaughter; milk taken from treated animals within 96 hours (8 milkings) after the latest treatment must not be used for food.

(2) *Goats—(i) Amount.* 3 grams per 100 lb. body weight.

(ii) *Indications for use.* Control of severe infections of gastrointestinal roundworms (*Trichostrongylus* spp., *Haemonchus* spp., *Ostertagia* spp., *Cooperia* spp., *Nematodirus* spp.,

Bunostomum spp., *Strongyloides* spp., *Chabertia* spp., and *Oesophagostomum* spp.).

(iii) *Limitations.* 3 grams per 100 lb. body weight at a single dose; do not treat animals within 30 days of slaughter; milk taken from treated animals within 96 hours (8 milkings) after the latest treatment must not be used for food.

(3) *Sheep and goats—(i) Amount.* 2 grams per 100 lb. body weight.

(ii) *Indications for use.* Control of infections of gastrointestinal roundworms (*Trichostrongylus* spp., *Haemonchus* spp., *Ostertagia* spp., *Cooperia* spp.; *Nematodirus* spp., *Bunostomum* spp., *Strongyloides* spp., *Chabertia* spp., and *Oesophagostomum* spp.); also active against ova and larvae passed by sheep from 3 hours to 3 days after the feed is consumed (good activity against ova and larvae of *T. colubriformis* and *axeii*, *Ostertagia* spp., *Nematodirus* spp., *Strongyloides* spp.; less effective against those of *Haemonchus contortus* and *Oesophagostomum* spp.).

(iii) *Limitations.* Use 2 grams per 100 lb. body weight at a single dose; do not treat animals within 30 days of slaughter; milk taken from treated animals within 96 hours (8 milkings) after the latest treatment must not be used for food.

(4) *For swine—(i) Amount.* 45.4–908 grams per ton (0.005–0.1 percent).

(ii) *Indications for use.* Aid in the prevention of infections of large roundworms (genus *Ascaris*).

(iii) *Limitations.* Administer continuously feed containing 0.05–0.1 percent thiabendazole per ton for 2 weeks followed by feed containing 0.005–0.02 percent thiabendazole per ton for 8–14 weeks; do not treat animals within 30 days of slaughter.

(5) *Pheasants—(i) Amount.* 454 grams per ton (0.05 percent) continuously for 2 weeks (14 days).

(ii) *Indications for use.* For the treatment of gapeworms (*Syngamus trachea*) in pheasants.

(iii) *Limitations.* Do not use treated pheasants for food for 21 days after last day of treatment. Fertility, hatchability, and other reproductive data

§ 558.618

are not available on use in breeding animals.

[40 FR 13959, Mar. 27, 1975, as amended at 47 FR 49641, Nov. 2, 1982; 49 FR 29958, July 25, 1984; 51 FR 7400, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 62 FR 63271, Nov. 28, 1997]

§ 558.618 Tilmicosin.

(a) *Approvals.* Type A medicated articles: 90.7 grams of tilmicosin (as tilmicosin phosphate) per pound (200 grams per kilogram) to 000986 in § 510.600(c) of this chapter.

(b) *Special considerations.* Do not use in any feed containing bentonite.

(c) *Related tolerances.* See § 556.735 of this chapter.

(d) *Conditions of use.* It is used in swine feed as follows:

(1) *Amount per ton.* 181 grams to 363 grams tilmicosin.

(2) *Indications for use.* For the control of swine respiratory disease associated with *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

(3) *Limitations.* For use in swine feed only. The safety of tilmicosin has not been established in pregnant swine or swine intended for breeding purposes. Do not allow horses or other equine access to feeds containing tilmicosin. Feed continuously as the sole ration for 21-day period, beginning approximately 7 days before an expected disease outbreak. Withdraw 7 days before slaughter. Federal law restricts this drug to use under the professional supervision of a licensed veterinarian. Any animal feed bearing or containing this drug shall be fed to animals only by or upon a lawful veterinary feed directive (VFD) issued by a licensed veterinarian in the course of the veterinarian's professional practice. VFD's for tilmicosin phosphate shall not be refilled.

(4) *VFD Requirements.* This drug and any article or feed manufactured from it shall bear the following cautionary statements: "Caution: Federal law limits this drug to use under the professional supervision of a licensed veterinarian. Animal feed bearing or containing this veterinary feed directive drug shall be fed to animals only by or upon a lawful veterinary feed directive issued by a licensed veterinarian in the course of the veterinarian's professional practice." A VFD shall contain

21 CFR Ch. I (4–1–99 Edition)

the following information: The name, address, and phone number of the veterinarian and the client; identification of the animals to be treated, including, identification of the species, number of animals, and the location of the animals; date of treatment and, if different, date of prescribing the VFD drug; the condition or disease being diagnosed or treated; name of the animal drug; level of animal drug in feed and amount of feed; feeding instructions with withdrawal time; any special instructions and cautionary statements necessary for use of the drug in conformance with the approval; expiration date of VFD; number of refills, if permitted by approval; signature of the veterinarian; veterinarian's license number and name of the State issuing the license.

[61 FR 68148, Dec. 27, 1996; 62 FR 15391, Apr. 1, 1997, as amended at 64 FR 13679, Mar. 22, 1999]

§ 558.625 Tylosin.

(a) *Specifications.* Tylosin is the antibiotic substance produced by growth of *Streptomyces fradiae* or the same antibiotic substance produced by any other means. Tylosin, present as the phosphate salt, conforms to the appropriate antibiotic standard. Tylosin contains at least 95 percent tylosin as a combination of tylosin A, tylosin B, tylosin C, and tylosin D of which at least 80 percent is tylosin A as determined by a method entitled "Determination of Factor Content in Tylosin by High Performance Liquid Chromatography," which is incorporated by reference. Copies are available from the Dockets Management Branch (HFA-305), Food and Drug Administration, rm. 1-23, 12420 Parklawn Dr., Rockville, MD 20857, or available for inspection at the Office of the Federal Register, 800 North Capitol Street, NW., suite 700, Washington, DC 20001.

(b) *Approvals.* Type A medicated article levels of tylosin granted to firms as sponsor(s) and identified by drug listing numbers in § 510.600(c) of this chapter for the specific usage indicated in paragraph (f) of this section.

(1) To 000986: 10, 40, 100 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

Food and Drug Administration, HHS

§ 558.625

(2) To 051359: 1, 2, 4, 5, 8, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(3) To 043733: 20 and 40 grams per pound, paragraphs (f)(1)(i) through (vi) of this section.

(4) [Reserved]

(5) To 017800: 0.4, 0.8, 1, and 8 grams per pound, paragraph (f)(1)(vi)(a) of this section; 10 and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(6)–(7) [Reserved]

(8) To 035369: 4 and 10 grams per pound, paragraph (f)(1)(vi)(a) of this section; 10 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(9) To 043727: 4 grams per pound, paragraph (f)(1)(vi)(a) of this section; 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(10) To 017519: 0.4, 0.8, and 1.6 grams per pound, paragraph (f)(1)(vi)(a) of this section; 20, 40, and 100 grams per pound, paragraphs (f)(1)(i) through (f)(1)(vi) of this section.

(11) [Reserved]

(12) To 021930: 2 grams per pound, paragraph (f)(1)(vi)(a) of this section; 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(13) To 035393: 0.4 and 2 grams per pound; paragraph (f)(1)(vi)(a) of this section.

(14) To 016968: 1, 2, 4, 8, and 10 grams per pound, paragraphs (f)(1) (i), (iii), (iv), and (vi) of this section; 20, 25, 40, and 100 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(15) [Reserved]

(16) To 049768: 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(17) To 021780: 0.8 gram per pound, paragraph (f)(1)(vi)(a) of this section; 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(18) [Reserved]

(19) To 033999: 10 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(20)–(21) [Reserved]

(22) To 026282: 10 grams per pound; paragraph (f)(1)(vi)(a) of this section.

(23)–(24) [Reserved]

(25) To 000069: 4, 8, and 10 grams per pound, paragraph (f)(1)(vi)(a) of this section; 20 and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(26)–(30) [Reserved]

(31) To 029341: 5 grams per pound; paragraph (f)(1)(vi)(a) of this section.

(32) [Reserved]

(33) To 034936: 0.8 and 2 grams per pound, paragraph (f)(1)(vi)(a) of this section; 4, 8, and 10 grams per pound, paragraphs (f)(1)(i), (iii), (iv), and (vi) of this section; 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section; 100 grams per pound, paragraphs (f)(1) (i), (ii), (iii), (iv), and (vi) of this section.

(34) To 028260: 0.8 gram per pound, paragraph (f)(1)(vi)(a) of this section.

(35) To 039741: 2 and 10 grams per pound, paragraph (f)(1)(vi)(a) of this section.

(36)–(37) [Reserved]

(38) To 053740: 1 gram per pound, paragraph (f)(1)(vi)(a) of this section; 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(39) To 050604: 10 grams per pound, paragraph (f)(1)(vi)(a) of this section.

(40) To 035955: 10 grams per pound, paragraph (f)(1)(vi)(a) of this section.

(41) [Reserved]

(42) To 010439: 0.4, 0.5, and 2 grams per pound, paragraph (f)(1)(vi) (a) of this section; 5, 10, 20, and 40 grams per pound, paragraph (f)(1)(i) through (vi) of this section.

(43)–(44) [Reserved]

(45) To 017139: 4 and 10 grams per pound, paragraph (f)(1)(vi)(a) of this section.

(46)–(47) [Reserved]

(48) To 017790: 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(49)–(51) [Reserved]

(52) To 017519: 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(53) [Reserved]

(54) To 046573: 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(55)–(56) [Reserved]

(57) To 028459: 0.4 and 10 grams per pound; paragraph (f)(1)(vi)(a) of this section.

(58)–(62) [Reserved]

§ 558.625

21 CFR Ch. I (4–1–99 Edition)

(63) To 046987: 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(64)–(65) [Reserved]

(66) To 024174: 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(67) To 027190: 2 grams per pound; paragraph (f)(1)(vi)(a) of this section.

(68) To 017473: 10 and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(69)–(72) [Reserved]

(73) To 035098: 0.33 and 0.67 gram per pound, paragraph (f)(1)(vi)(a) of this section; 0.8, 1, 2, and 10 grams per pound, paragraphs (f)(1) (i) and (vi) (a), (b) and (d) of this section; 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(74)–(76) [Reserved]

(77) To 050639: 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(78) To 050972: 0.36, 0.4, 0.72, and 0.8 gram per pound, paragraph (f)(1)(vi)(a) of this section; 1 gram per pound, paragraphs (f)(1)(vi) (a), (b), and (d) of this section.

(79) To 012286: 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(80) To 049685: 5, 10, 20, and 40 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(81)–(82) [Reserved]

(83) To 046573: 5-, 10-, 20-, and 40-grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(84) [Reserved]

(85) To 047126: 10, 40, and 100 grams per pound, paragraphs (f)(1) (i) through (vi) of this section.

(86)–(88) [Reserved]

(89) To 053389: 5, 10, 20, and 40 grams per pound, paragraph (f)(1) (i) through (vi) of this section.

(c)–(d) [Reserved]

(e) *Related tolerances.* See § 556.740 of this chapter.

(f) *Conditions of use.* (1) It is used in animal feeds as follows:

(i) *For beef cattle—(a) Amount per ton.* 8–10 grams.

(b) *Indications for use.* For reduction of incidence of liver abscesses caused by *Fusobacterium necrophorum* and *Actinomyces (Corynebacterium) pyogenes*.

(c) *Limitations.* As tylosin phosphate; each animal must receive not more than 90 milligrams per day and not less than 60 milligrams per day; feed continuously as sole ration.

(ii) *Broiler chickens—(a) Amount per ton.* Tylosin, 800–1000 grams.

(b) *Indications for use.* To aid in the control of chronic respiratory disease caused by *Mycoplasma gallisepticum*.

(c) *Limitations.* As tylosin phosphate; withdraw 5 days before slaughter; administer in feed to chickens 0 to 5 days of age, follow with second administration in feed for 24–48 hours at 3 to 5 weeks of age.

(iii) *Chickens—(a) Amount per ton.* Tylosin, 4–50 grams.

(1) *Indications for use.* For increased rate of weight gain and improved feed efficiency.

(2) *Limitations.* As tylosin phosphate.

(iv) *Laying chickens—(a) Amount per ton.* Tylosin, 20–50 grams.

(b) *Indications for use.* For improved feed efficiency.

(c) *Limitations.* As tylosin phosphate.

(v) *Replacement chickens—(a) Amount per ton.* Tylosin, 1,000 grams.

(b) *Indications for use.* To aid in the control of chronic respiratory disease caused by *Mycoplasma gallisepticum*.

(c) *Limitations.* As tylosin phosphate; withdraw 5 days before slaughter; administer in feed to chickens 0 to 5 days of age, follow with second administration in feed for 24 to 48 hours at 3 to 5 weeks of age.

(vi) *Swine—(a) Amount per ton.* Tylosin, 10–100 grams.

(1) *Indications for use.* For increased rate of weight gain and improved feed efficiency.

(2) *Limitations.* As tylosin phosphate; continuous use as follows: *Grams per ton:* 20–100, prestarter or starter; 20–40, grower; 10–20, finisher.

(b) *Amount per ton.* Tylosin, 40–100 grams.

(1) *Indications for use.* Prevention of swine dysentery (vibrionic).

(2) *Limitations.* Use 100 grams per ton for at least 3 weeks followed by 40 grams per ton until market weight; as tylosin phosphate.

(c) *Amount per ton.* Tylosin, 40–100 grams.

(1) *Indications for use.* Treatment and control of swine dysentery (vibrionic).

(2) *Limitations.* Administer in feed as tylosin phosphate after treatment with tylosin in drinking water as tylosin base; 0.25 gram per gallon in drinking water for 3–10 days, 40–100 grams per ton in feed for 2–6 weeks.

(d) *Amount per ton.* Tylosin, 100 grams.

(1) *Indications for use.* Maintaining weight gains and feed efficiency in presence of atrophic rhinitis.

(2) *Limitations.* As tylosin phosphate.

(2) Tylosin may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Hygromycin B as in § 558.274.

(ii) Melengestrol acetate alone or in combination with certain ionophores in accordance with § 558.342.

(iii) [Reserved]

(iv) Monensin in accordance with § 558.355.

(v) Pyrantel tartrate in accordance with § 558.485.

(e) *Amount per ton.* Tylosin 100 grams. (1) *Indications for use.* Prevention and/or control of porcine proliferative enteropathies (ileitis) associated with *Lawsonia intracellularis*.

(2) *Limitations.* As tylosin phosphate, administer for 21 days.

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.625, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.630 Tylosin and sulfamethazine.

(a) [Reserved]

(b) *Approvals.* Type A medicated article levels, a combination of equal amounts of tylosin and sulfamethazine, granted to firms as sponsor(s) and identified by drug listing numbers in § 510.600(c) of this chapter for the conditions of use indicated in paragraph (f) of this section.

(1) To 000986: 40 grams per pound each, paragraph (f)(2)(i).

(2) To 000986: 10 grams per pound each, paragraph (f)(2)(i).

(3) To 017519: 10 grams per pound each, paragraph (f)(2)(ii) of this section.

(4) To 021780: 2 grams per pound each, paragraph (f)(2)(ii) of this section.

(5) To 017800: 40 grams per pound each, paragraph (f)(2)(ii) of this section.

(6) To 017139: 4, 10, or 20 grams per pound each, paragraph (f)(2)(ii) of this section.

(7) To 021930: 2 grams per pound each, paragraph (f)(2)(i) of this section; 5, 10, 20, or 40 grams per pound each, paragraph (f)(2)(ii) of this section.

(8) To 017519, 026186: 5 or 10 grams per pound each, paragraph (f)(2)(ii) of this section.

(9) [Reserved]

(10) To 010439, 011749, 016968, 017473, 017519, 017790, 021780, 024174, 030841, 034936, 035098, 043727, 043733, 046573, 046987, 050568, 050639, and 051359, 053389: 5, 10, 20, or 40 grams per pound each, paragraph (f)(2)(ii) of this section.

(c)–(d) [Reserved]

(e) *Related tolerances.* See §§ 556.670 and 556.740 of this chapter.

(f) *Conditions of use.* It is used in feed for swine as follows:

(1) *Amount per ton.* Tylosin, 100 grams plus sulfamethazine, 100 grams.

(2) *Indications for use.* (i) Maintaining weight gains and feed efficiency in the presence of atrophic rhinitis; lowering the incidence and severity of *Bordetella bronchiseptica* rhinitis; prevention of swine dysentery (vibriotic); control of swine pneumonias caused by bacterial pathogens (*P. multocida* and/or *C. pyogenes*); for reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E Streptococci. Only the sulfamethazine portion of this combination is active in controlling jowl abscesses.

(ii) Maintaining weight gains and feed efficiency in the presence of atrophic rhinitis; lowering the incidence and severity of *Bordetella bronchiseptica* rhinitis; prevention of swine dysentery (vibriotic); control of swine pneumonias caused by bacterial pathogens (*Pasteurella multocida* and/or *Corynebacterium pyogenes*).

(3) *Limitations.* As tylosin phosphate; withdraw 15 days before slaughter.

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.630, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.635 Virginiamycin.

(a) [Reserved]

(b) *Approvals.* Type A medicated articles. (1) 1.1 percent activity (5 grams per pound), 2.2 percent activity (10 grams per pound), 4.4 percent activity (20 grams per pound), 11 percent activity (50 grams per pound), and 50 percent activity (227 grams per pound) used as in paragraph (f) of this section; and 30 percent activity (136.2 grams per pound) for the manufacture of Type C medicated feed for cattle used as in paragraph (f)(3); to 000069 in § 510.600(c) of this chapter.

(2) 2.2 percent activity (10 grams per pound) to 046573, 016968, and 017790 in § 510.600(c) of this chapter for use as in paragraphs (f)(1)(iv) and (f)(1)(v) of this section.

(c) [Reserved]

(d) *Related tolerances.* See § 556.750 of this chapter.

(e) *Special considerations.* (1) Not for use in breeding swine over 120 pounds.

(2) Dilute Type A article with at least 10 pounds of a feed ingredient prior to final mixing in 1 ton of Type C feed.

(3)–(4) [Reserved]

(f) *Conditions of use*—(1) *Swine.* It is used as follows:

(i) 100 grams per ton for 2 weeks, for treatment of swine dysentery in non-breeding swine over 120 pounds.

(ii) 100 grams per ton for 2 weeks, 50 grams per ton thereafter, for treatment and control of swine dysentery in swine up to 120 pounds.

(iii) 25 grams per ton, as an aid in control of dysentery in swine up to 120 pounds. For use in animals or on premises with a history of swine dysentery but where symptoms have not yet occurred.

(iv) 10 grams per ton from weaning up to 120 pounds for increased rate of weight gain and improved feed efficiency, followed by 5 grams per ton to market weight for increased rate of weight gain and improved feed efficiency. For continuous use from weaning to market weight.

(v) 10 grams per ton from weaning up to 120 pounds for increased rate of weight gain and improved feed efficiency, followed by 5 to 10 grams per ton to market weight for increased

rate of weight gain. For continuous use from weaning to market weight.

(2) *Poultry.* It is used as follows:

(i) 5 to 15 grams per ton for increased rate of weight gain, for use in broiler chickens, not for use in layers.

(ii) 5 grams per ton for increased rate of weight gain and improved feed efficiency in broiler chickens, not for use in layers.

(iii) 20 grams per ton for prevention of necrotic enteritis caused by *Clostridium perfringens* susceptible to virginiamycin in broiler chickens; not for use in layers.

(iv) 10 to 20 grams per ton for increased rate of weight gain and improved feed efficiency in growing turkeys.

(3) *Cattle.* It is used as follows:

(i) 16.0 to 22.5 grams per ton to provide 100 to 340 milligrams per head per day for increased rate of weight gain.

(ii) 13.5 to 16.0 grams per ton to provide 85 to 240 milligrams per head per day for reduction of incidence of liver abscesses.

(iii) 11.0 to 16.0 grams per ton to provide 70 to 240 milligrams per head per day for improved feed efficiency.

(iv) Feed continuously as sole ration to cattle fed in confinement for slaughter. Not for use in animals intended for breeding.

(4) Virginiamycin may be used in accordance with the provisions of this section in the combinations provided, as follows:

(i) Monensin sodium in accordance with § 558.355.

(ii) Lasalocid sodium in accordance with § 558.311.

(iii) Monensin and roxarsone as in § 558.355.

(iv) Amprolium and ethopabate as in § 558.58.

(v) Halofuginone as in § 558.265.

(vi) Salinomycin alone or with roxarsone as in § 558.550.

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.635, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.680 Zoalene.

(a) *Approvals.* Type A medicated articles: 25 percent to 046573 in § 510.600(c) of this chapter.

Food and Drug Administration, HHS

§ 558.680

(b) *Related tolerances.* See § 556.770 of this chapter. (c) *Conditions of use—(1) Chickens and turkeys:*

Zoalene in grams/ton	Combination in grams/ton	Indications for use	Limitations
(i) 36.3–113.5 (0.004–0.0125%).		Replacement chickens; development of active immunity to coccidiosis.	Grower ration not to be fed to birds over 14 weeks of age; as follows:

Growing conditions	Starter ration Grams per ton	Grower ration Grams per ton
Severe exposure	113.5 (0.0125%)	75.4–113.5 (0.0083%–0.0125%)
Light to moderate exposure	75.4–113.5 (0.0083%–0.0125%)	36.3–75.4 (0.004%–0.0083%)

Zoalene in grams/ton	Combination in grams/ton	Indications for use	Limitations
	Arsanilate sodium 90 (0.01%).	Replacement chickens; development of active immunity to coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Grower ration not to be fed to birds over 14 weeks of age; withdraw 5 days (d) before slaughter; as sole source of organic arsenic; feed as in subtable in item (i).
	Arsanilic acid 90 (0.01%).	Replacement chickens; development of active immunity to coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Grower ration not to be fed to birds over 14 weeks of age; withdraw 5 d before slaughter; as sole source of organic arsenic; feed as in subtable in item (i).
	Arsanilic acid 90 (0.01%) plus erythromycin 4.6 to 18.5.	Replacement chickens; growth promotion and feed efficiency; development of active immunity to coccidiosis; improving pigmentation.	As erythromycin thiocyanate; grower ration not to be fed to birds over 14 weeks of age; withdraw 5 d before slaughter; as sole source of organic arsenic; feed as in subtable item (i).
	Arsanilic acid 90 (0.01%) plus erythromycin 92.5.	1. Replacement chickens; as an aid in the prevention of chronic respiratory disease during periods of stress; development of active immunity to coccidiosis; growth promotion and feed efficiency; improving pigmentation. 2. Replacement chickens; as an aid in the prevention of infectious coryza; development of active immunity to coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Feed for 2 d before stress and 3 to 6 d after stress; as erythromycin thiocyanate; grower ration not to be fed to birds over 14 weeks of age; withdraw 5 d before slaughter; as sole source of organic arsenic; feed as in subtable in item (i). Feed for 7 to 14 d; as erythromycin thiocyanate; grower ration not to be fed to birds over 14 weeks of age; withdraw 5 d before slaughter; as sole source of organic arsenic; feed as in subtable in item (i).
	Arsanilic acid 90 (0.01%) plus erythromycin 185.	Replacement chickens; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease; growth promotion and feed efficiency; improving pigmentation and development of active immunity to coccidiosis.	Feed for 5 to 8 d; do not use in birds producing eggs for food purposes; withdraw 5 d before slaughter; as erythromycin thiocyanate; as sole source of organic arsenic; feed as in subtable in item (i).
	Arsanilic acid 90 (0.01%) plus penicillin 2.4 to 50.	Replacement chickens; growth promotion and feed efficiency; development of active immunity to coccidiosis; improving pigmentation.	As procaine penicillin; grower ration not to be fed to birds over 14 weeks of age; withdraw 5 d before slaughter; as sole source of organic arsenic; feed as in subtable in item (i).

Zoalene in grams/ton	Combination in grams/ton	Indications for use	Limitations
	Bacitracin 100 to 500.	Replacement chickens; treatment of chronic respiratory disease (air-sac infection); blue comb (nonspecific infectious enteritis); development of active immunity to coccidiosis.	As bacitracin zinc; grower ration not to be fed to birds over 14 weeks of age; feed as in subtable in item (i).
	Chlortetracycline 100 to 200.	Replacement chickens; development of active immunity to coccidiosis; control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption; grower ration not to be fed to birds over 14 weeks of age; feed as in subtable in item (i).
	Chlortetracycline 200 to 400.	Replacement chickens; development of active immunity to coccidiosis; control of chronic respiratory disease (CRD) and air sac infection caused by <i>M. gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption; grower ration not to be fed to birds over 14 weeks of age; feed as in subtable in item (i).
	Erythromycin 4.6 to 18.5.	Replacement chickens; growth promotion and feed efficiency; development of active immunity to coccidiosis.	As erythromycin thiocyanate; grower ration not to be fed to birds over 14 weeks of age; feed as in subtable in item (i).
	Erythromycin 92.5.	1. Replacement chickens, as an aid in the prevention of chronic respiratory disease during periods of stress; development of active immunity to coccidiosis. 2. Replacement chickens; as an aid in the prevention of infectious coryza; development of active immunity to coccidiosis.	Feed for 2 d before stress and 3 to 6 after stress; withdraw 24 hours (h) before slaughter; as erythromycin thiocyanate; grower ration not to be fed to birds over 14 weeks of age; feed as in subtable in item (i). Feed for 7 to 14 d; withdraw 24 h before slaughter; as erythromycin thiocyanate; grower ration not to be fed to birds over 14 weeks of age; feed as in subtable in item (i).
	Erythromycin 185.	Replacement chickens; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease; development of active immunity to coccidiosis.	Feed for 5 to 8 d; do not use in birds producing eggs for food purposes; withdraw 48 h before slaughter; grower ration not to be fed to birds over 14 weeks of age; feed as in subtable in item (i).
	Hygromycin B 8 to 12.	Replacement chickens; development of active immunity to coccidiosis; control of infestation of large round worms (<i>Ascaris galli</i>) cecal worms (<i>Heterakis gallinae</i>) and capillary worms (<i>Capillaria obsignate</i>).	Grower ration not to be fed to birds over 14 weeks of age; feed as in subtable in item (i).
	Penicillin 2.4 to 50.	Replacement chickens; growth promotion and feed efficiency; development of active immunity to coccidiosis.	As procaine penicillin; grower ration not to be fed to birds over 14 weeks of age; feed as in subtable in item (i).
	Penicillin 2.4 to 50 plus roxarsone 22.7 to 45.4 (0.0025% to 0.005%).	Replacement chickens; growth promotion and feed efficiency; development of active immunity to coccidiosis; improving pigmentation.	As procaine penicillin; grower ration not to be fed to birds over 14 weeks of age; withdraw 5 d before slaughter; as sole source of organic arsenic; feed as in subtable in item (i).
	Roxarsone 22.7 to 45.5 (0.0025% to 0.005%).	Replacement chickens; development of active immunity to coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Grower ration not to be fed to birds over 14 weeks of age; withdraw 5 d before slaughter; as sole source of organic arsenic; feed as in subtable in item (i).
(ii) 113.5 (0.0125%).		Broiler chickens; prevention and control of coccidiosis.	

Zoalene in grams/ton	Combination in grams/ton	Indications for use	Limitations
	Arsanilate sodium 90 (0.01%).	Broiler chickens; prevention and control of coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic.
	Arsanilic acid 90 (0.01%).	Broiler chickens; growth promotion and feed efficiency; prevention and control of coccidiosis; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic.
	Arsanilic acid 90 (0.01%) plus erythromycin 4.6 to 18.5.	Broiler chickens; growth prevention and control of coccidiosis; improving pigmentation.	As erythromycin thiocyanate; withdraw 5 d before slaughter; as sole source of organic arsenic.
	Arsanilic acid 90 (0.01%) plus erythromycin 92.5.	1. Broiler chickens; as an aid in the prevention of chronic respiratory disease during periods of stress; growth promotion and feed efficiency; improving pigmentation; control of coccidiosis. 2. Broiler chickens; prevention and control of coccidiosis; growth promotion and feed efficiency; improving pigmentation; as an aid in the prevention of infectious coryza.	Do. Do.
	Arsanilic acid 90 (0.01%) plus erythromycin 185.	Broiler chickens; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease; prevention and control of coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Feed for 5 to 8 d; do not use in birds producing eggs for food purposes; as erythromycin thiocyanate; withdraw 5 d before slaughter; as sole source of organic arsenic.
	Arsanilic acid 90 (0.01%) plus penicillin 2.4 to 50.	Broiler chickens; growth promotion and feed efficiency; prevention and control of coccidiosis; improving pigmentation.	As procaine penicillin; withdraw 5 d before slaughter; as sole source of organic arsenic.
	Arsanilic acid 90 (0.01%) plus bacitracin 4 to 50.	Broiler chickens; prevention and control of coccidiosis; improving pigmentation; growth promotion and feed efficiency.	Withdraw 5 d before slaughter; as sole source of organic arsenic; as bacitracin methylene disalicylate.
	Bacitracin 4 to 50.	Broiler chickens; growth promotion and feed efficiency; prevention and control of coccidiosis.	As bacitracin methylene disalicylate or zinc bacitracin.
	Bacitracin 4 to 50 plus roxarsone 22.7 to 45.4 (0.0025 to 0.005%).	Broiler chickens; growth promotion and feed efficiency; prevention and control of coccidiosis; improving pigmentation.	As bacitracin methylene disalicylate or zinc bacitracin; withdraw 5 d before slaughter; as sole source of organic arsenic.
	Bacitracin 100 to 500.	Broiler chickens; treatment of chronic respiratory disease (air-sac infection); blue comb (nonspecific infectious enteritis); prevention and control of coccidiosis.	As zinc bacitracin.
	Chlortetracycline 100 to 200	Broiler chickens; prevention and control of coccidiosis; control of infectious synovitis caused by <i>M. synoviae</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption; feed continuously for 7 to 14 d.
	Chlortetracycline 200 to 400	Broiler chickens; prevention and control of coccidiosis; control of chronic respiratory disease (CRD) and air sac infection caused by <i>M. gallisepticum</i> and <i>E. coli</i> susceptible to chlortetracycline.	Do not feed to chickens producing eggs for human consumption; feed continuously for 7 to 14 d.

Zoalene in grams/ton	Combination in grams/ton	Indications for use	Limitations
(iii) 113.5 to 170.3 (0.0125 to 0.01875%).	Erythromycin 4.6 to 18.5.	Broiler chickens; growth promotion and feed efficiency; prevention and control of coccidiosis.	As erythromycin thiocyanate.
	Erythromycin 92.5.	1. Broiler chickens; as an aid in the prevention of chronic respiratory disease during period of stress; prevention and control of coccidiosis. 2. Broiler chicken; as an aid in the prevention of infectious coryza; prevention and control of coccidiosis.	Feed for 2 d before stress and 3 to 6 d after stress; withdraw 24 h before slaughter; as erythromycin thiocyanate. Feed for 7 to 14 d; withdraw 24 h before slaughter; as erythromycin thiocyanate.
	Erythromycin 185.	Broiler chickens; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease; prevention and control of coccidiosis.	Feed for 5 to 8 d; do not use in birds producing eggs for food purposes; withdraw 48 h before slaughter; as erythromycin thiocyanate.
	Hygromycin B 8 to 12.	Broiler chickens; prevention and control of coccidiosis; control of infestation of large round worms (<i>Ascaris galli</i>) cecal worms (<i>Heterakis gallinae</i>) and capillary worms (<i>Capillaria obsignate</i>).	
	Lincomycin 2.	Broiler chickens; increase in rate of weight gain; improved feed efficiency; as an aid in the prevention and control of coccidiosis.	Do not feed to laying chickens; to be fed as the sole ration; as lincomycin hydrochloride monohydrate provided by No. 000009 in § 510.600(c) of this chapter.
	Penicillin 2.4 to 50.	Broiler chickens; growth promotion and feed efficiency; prevention and control of coccidiosis.	As procaine penicillin.
	Penicillin 2.4 to 50 plus roxarsone 22.7 to 45.4 (0.0025 to 0.005%).	Broiler chickens; prevention and control of coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic; as procaine penicillin.
	Roxarsone 22.7 to 45.4 (0.0025 to 0.005%).	Broiler chickens; prevention and control of coccidiosis; growth promotion and feed efficiency; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic.
	Arsanilate sodium 90 (0.01%).	Turkeys; prevention and control of coccidiosis. Turkeys; growth promotion and feed efficiency; improving pigmentation.	For turkeys grown for meat purposes only. For turkeys grown for meat purposes only; withdraw 5 d before slaughter; as sole source of organic arsenic.
	Arsanilic acid 90(0.01%). Bacitracin methylene disalicylate 4–50.	do. Turkeys; prevention and control of coccidiosis, and increased rate of weight gain and improved feed efficiency.	Do. For turkeys grown for meat purposes only, not to be fed to laying birds, feed continuously as sole ration until 14 to 16 weeks of age.
	Carbarsone (not U.S.P.) 277 to 340.5 (0.025% to 0.0375%).	Turkeys; prevention and control of coccidiosis; aid in the prevention of blackhead.	For turkeys grown for meat purposes only; feed continuously beginning 2 weeks before blackhead and coccidiosis are expected and continue as long as prevention of blackhead and prevention and control of coccidiosis is needed; withdraw 5 d before slaughter; as sole source of organic arsenic.
	Roxarsone 22.7 to 45.4 (0.0025% to 0.005%).	Turkeys; growth promotion and feed efficiency; improving pigmentation.	Withdraw 5 d before slaughter; as sole source of organic arsenic.

Food and Drug Administration, HHS

§ 570.3

(2) *Permitted combinations.* It may be used in accordance with the provisions of this section in the combinations provided, as follows:

(i) Bambermycins in accordance with § 558.95.

(ii) Roxarsone in accordance with § 558.530.

[41 FR 11005, Mar. 15, 1976, as amended at 42 FR 18618, Apr. 8, 1977; 42 FR 20817, Apr. 22, 1977; 42 FR 36995, July 19, 1977; 51 FR 7401, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 55 FR 8461, Mar. 8, 1990; 57 FR 8403, Mar. 10, 1992; 57 FR 8578, Mar. 11, 1992; 61 FR 35957, July 9, 1996; 63 FR 38750, July 20, 1998]

PART 564 [RESERVED]

PART 570—FOOD ADDITIVES

Subpart A—General Provisions

Sec.

570.3 Definitions.

570.6 Opinion letters on food additive status.

570.13 Indirect food additives resulting from packaging materials prior sanctioned for animal feed and pet food.

570.14 Indirect food additives resulting from packaging materials for animal feed and pet food.

570.15 Adoption of regulation on initiative of Commissioner.

570.17 Exemption for investigational use and procedure for obtaining authorization to market edible products from experimental animals.

570.18 Tolerances for related food additives.

570.19 Pesticide chemicals in processed foods.

Subpart B—Food Additive Safety

570.20 General principles for evaluating the safety of food additives.

570.30 Eligibility for classification as generally recognized as safe (GRAS).

570.35 Affirmation of generally recognized as safe (GRAS) status.

570.38 Determination of food additive status.

AUTHORITY: 21 U.S.C. 321, 341, 342, 346a, 348, 371.

SOURCE: 41 FR 38644, Sept. 10, 1976, unless otherwise noted.

Subpart A—General Provisions

§ 570.3 Definitions.

(a) *Secretary* means the Secretary of Health and Human Services.

(b) *Department* means the Department of Health and Human Services.

(c) *Commissioner* means the Commissioner of Food and Drugs.

(d) As used in this part, the term *act* means the Federal Food, Drug, and Cosmetic Act approved June 25, 1936 (52 Stat. 1040 *et seq.*, as amended; 21 U.S.C. 301–392).

(e) *Food additives* includes all substances not exempted by section 201(s) of the act, the intended use of which results or may reasonably be expected to result, directly or indirectly, either in their becoming a component of food or otherwise affecting the characteristics of food. A material used in the production of containers and packages is subject to the definition if it may reasonably be expected to become a component, or to affect the characteristics, directly or indirectly, of food packed in the container. *Affecting the characteristics of food* does not include such physical effects, as protecting contents of packages, preserving shape, and preventing moisture loss. If there is no migration of a packaging component from the package to the food, it does not become a component of the food and thus is not a food additive. A substance that does not become a component of food, but that is used, for example, in preparing an ingredient of the food to give a different flavor, texture, or other characteristic in the food, may be a food additive.

(f) *Common use in food* means a substantial history of consumption of a substance by a significant number of animals in the United States.

(g) The word *substance* in the definition of the term *food additive* includes a food or feed or a component of a food or feed consisting of one or more ingredients.

(h) *Scientific procedures* include those human, animal, analytical, and other scientific studies, whether published or unpublished, appropriate to establish the safety of a substance.

(i) *Safe* or *safety* means that there is a reasonable certainty in the minds of competent scientists that the substance is not harmful under the intended conditions of use. It is impossible in the present state of scientific knowledge to establish with complete certainty the absolute harmlessness of

§570.6

21 CFR Ch. I (4–1–99 Edition)

the use of any substance. Safety may be determined by scientific procedures or by general recognition of safety. In determining safety, the following factors shall be considered:

(1) The probable consumption of the substance and of any substance formed in or on food because of its use;

(2) The cumulative effect of the substance in the diet, taking into account any chemically or pharmacologically related substance or substances in such diet;

(3) Safety factors which, in the opinion of experts qualified by scientific training and experience to evaluate the safety of food and food ingredients, are generally recognized as appropriate.

(j) The term *nonperishable processed food* means any processed food not subject to rapid decay or deterioration that would render it unfit for consumption. Not included are hermetically sealed foods and other processed foods requiring refrigeration.

(k) *General recognition of safety* shall be determined in accordance with §570.30.

(l) *Prior sanction* means an explicit approval granted with respect to use of a substance in food prior to September 6, 1958, by the Food Drug and Administration or the United States Department of Agriculture pursuant to the Federal Food, Drug, and Cosmetic Act, the Poultry Products Inspection Act, or the Meat Inspection Act.

(m) *Food* includes human food, substances migrating to food from food-contact articles, pet food, and animal feed.

[41 FR 38644, Sept. 10, 1976, as amended at 42 FR 55206, Oct. 14, 1977]

§570.6 Opinion letters on food additive status.

(a) Over the years the Food and Drug Administration has given informal written opinions to inquirers as to the safety of articles intended for use as components of, or in contact with, food. Prior to the enactment of the Food Additives Amendment of 1958 (Pub. L. 85-929, Sept. 6, 1958), these opinions were given pursuant to section 402(a)(1) of the Federal Food, Drug, and Cosmetic Act, which reads in part: "A food shall be deemed to be adulterated if it bears or contains any

poisonous or deleterious substance which may render it injurious to health".

(b) Since enactment of the Food Additives Amendment, the Food and Drug Administration has advised such inquirers that an article:

(1) Is a food additive within the meaning of section 201(s) of the act; or

(2) Is generally recognized as safe (GRAS); or

(3) Has prior sanction or approval under that amendment; or

(4) Is not a food additive under the conditions of intended use.

(c) In the interest of the public health, such articles which have been considered in the past by the Food and Drug Administration to be safe under the provisions of section 402(a)(1), or to be generally recognized as safe for their intended use, or to have prior sanction or approval, or not to be food additives under the conditions of intended use, must be reexamined in the light of current scientific information and current principles for evaluating the safety of food additives if their use is to be continued.

(d) Because of the time span involved, copies of many of the letters in which the Food and Drug Administration has expressed an informal opinion concerning the status of such articles may no longer be in the file of the Food and Drug Administration. In the absence of information concerning the names and uses made of all the articles referred to in such letters, their safety of use cannot be reexamined. For this reason all food additive status opinions of the kind described in paragraph (c) of this section given by the Food and Drug Administration are hereby revoked.

(e) The prior opinions of the kind described in paragraph (c) of this section will be replaced by qualified and current opinions if the recipient of each such letter forwards a copy of each to the Department of Health and Human Services, Food and Drug Administration, Center for Veterinary Medicine, Office of Surveillance and Compliance (HFV-200), 7500 Standish Pl., Rockville, MD 20855, along with a copy of his letter of inquiry, on or before July 23, 1970.

Food and Drug Administration, HHS

§ 570.18

(f) This section does not apply to food additive status opinion letters pertaining to articles that were considered by the Food and Drug Administration to be food additives nor to articles included in regulations in this Subchapter E if the articles are used in accordance with the requirements of such regulations.

[41 FR 38644, Sept. 10, 1976, as amended at 54 FR 18281, Apr. 28, 1989; 57 FR 6476, Feb. 25, 1992]

§ 570.13 Indirect food additives resulting from packaging materials prior sanctioned for animal feed and pet food.

Regulations providing for the use of food packaging materials as prior sanctioned in part 181 of this chapter are incorporated in Subchapter E as applicable to packaging materials used for animal feed and pet food.

[42 FR 14091, Mar. 15, 1977]

§ 570.14 Indirect food additives resulting from packaging materials for animal feed and pet food.

Regulations providing for the use of food packaging materials in parts 174 through 179 of this chapter are incorporated in Subchapter E as applicable to packaging materials used for animal feed and pet food.

[42 FR 14091, Mar. 15, 1977]

§ 570.15 Adoption of regulation on initiative of Commissioner.

(a) The Commissioner upon his own initiative may propose the issuance of a regulation prescribing, with respect to any particular use of a food additive, the conditions under which such additive may be safely used. Notice of such proposal shall be published in the FEDERAL REGISTER and shall state the reasons for the proposal.

(b) Action upon a proposal made by the Commissioner shall proceed as provided in part 10 of this chapter.

[41 FR 38644, Sept. 10, 1976, as amended at 42 FR 4717, Jan. 25, 1977; 42 FR 15675, Mar. 22, 1977]

§ 570.17 Exemption for investigational use and procedure for obtaining authorization to market edible products from experimental animals.

A food additive or food containing a food additive intended for investigational use by qualified experts shall be exempt from the requirements of section 409 of the act under the following conditions:

(a) If intended for investigational use in vitro or in laboratory research animals, it bears a label which states prominently, in addition to the other information required by the act, the warning:

Caution. Contains a new food additive for investigational use only in laboratory research animals or for tests in vitro. Not for use in humans.

(b) If intended for use in animals other than laboratory research animals and if the edible products of the animals are to be marketed as food, permission for the marketing of the edible products as food has been requested by the sponsor, and authorization has been granted by the Food and Drug Administration in accordance with § 511.1 of this chapter or by the Department of Agriculture in accordance with 9 CFR 309.17, and it bears a label which states prominently, in addition to the other information required by the act, the warning:

Caution. Contains a new food additive for use only in investigational animals. Not for use in humans.

Edible products of investigational animals are not to be used for food unless authorization has been granted by the U.S. Food and Drug Administration or by the U.S. Department of Agriculture.

(c) If intended for nonclinical laboratory studies in food-producing animals, the study is conducted in compliance with the regulations set forth in part 58 of this chapter.

[41 FR 38644, Sept. 10, 1976, as amended at 43 FR 60023, Dec. 22, 1978]

§ 570.18 Tolerances for related food additives.

(a) Food additives that cause similar or related pharmacological effects will be regarded as a class, and in the absence of evidence to the contrary, as

§ 570.19

having additive toxic effects and will be considered as related food additives.

(b) Tolerances established for such related food additives may limit the amount of a common component that may be present, or may limit the amount of biological activity (such as cholinesterase inhibition) that may be present or may limit the total amount of related food additives that may be present.

(c) Where food additives from two or more chemicals in the same class are present in or on a food, the tolerance for the total of such additives shall be the same as that for the additive having the lowest numerical tolerance in this class, unless there are available methods that permit quantitative determination of the amount of each food additive present or unless it is shown that a higher tolerance is reasonably required for the combined additives to accomplish the physical or technical effect for which such combined additives are intended and that the higher tolerance will be safe.

(d) Where residues from two or more additives in the same class are present in or on a food and there are available methods that permit quantitative determination of each residue, the quantity of combined residues that are within the tolerance may be determined as follows:

(1) Determine the quantity of each residue present.

(2) Divide the quantity of each residue by the tolerance that would apply if it occurred alone, and multiply by 100 to determine the percentage of the permitted amount of residue present.

(3) Add the percentages so obtained for all residues present.

(4) The sum of the percentages shall not exceed 100 percent.

§ 570.19 Pesticide chemicals in processed foods.

When pesticide chemical residues occur in processed foods due to the use of raw agricultural commodities that bore or contained a pesticide chemical in conformity with an exemption granted or a tolerance prescribed under section 408 of the act, the processed food will not be regarded as adulterated so long as good manufacturing practice has been followed in removing

21 CFR Ch. I (4–1–99 Edition)

any residue from the raw agricultural commodity in the processing (such as by peeling or washing) and so long as the concentration of the residue in the processed food when ready to eat is not greater than the tolerance prescribed for the raw agricultural commodity. But when the concentration of residue in the processed food when ready to eat is higher than the tolerance prescribed for the raw agricultural commodity, the processed food is adulterated unless the higher concentration is permitted by a tolerance obtained under section 409 of the act. For example, if fruit bearing a residue of 7 parts per million of DDT permitted on the raw agricultural commodity is dried and a residue in excess of 7 parts per million of DDT results on the dried fruit, the dehydrated fruit is adulterated unless the higher tolerance for DDT is authorized by the regulations in this part. Food that is itself ready to eat, and which contains a higher residue than allowed for the raw agricultural commodity, may not be legalized by blending or mixing with other foods to reduce the residue in the mixed food below the tolerance prescribed for the raw agricultural commodity.

Subpart B—Food Additive Safety

§ 570.20 General principles for evaluating the safety of food additives.

(a) In reaching a decision on any petition filed under section 409 of the act, the Commissioner will give full consideration to the specific biological properties of the compound and the adequacy of the methods employed to demonstrate safety for the proposed use, and the Commissioner will be guided by the principles and procedures for establishing the safety of food additives stated in current publications of the National Academy of Sciences-National Research Council. A petition will not be denied, however, by reason of the petitioner's having followed procedures other than those outlined in the publications of the National Academy of Sciences-National Research Council if, from available evidence, the Commissioner finds that the procedures used give results as reliable as, or more reliable than, those reasonably

to be expected from the use of the outlined procedures. In reaching a decision, the Commissioner will give due weight to the anticipated levels and patterns of consumption of the additive specified or reasonably inferable. For the purposes of this section, the principles for evaluating safety of additives set forth in the above-mentioned publications will apply to any substance that may properly be classified as a food additive as defined in section 201(s) of the act.

(b) Upon written request describing the proposed use of an additive and the proposed experiments to determine its safety, the Commissioner will advise a person who wishes to establish the safety of a food additive whether he believes the experiments planned will yield data adequate for an evaluation of the safety of the additive.

§ 570.30 Eligibility for classification as generally recognized as safe (GRAS).

(a) General recognition of safety may be based only on the views of experts qualified by scientific training and experience to evaluate the safety of substances directly or indirectly added to food. The basis of such views may be either (1) scientific procedures or (2) in the case of a substance used in food prior to January 1, 1958, through experience based on common use in food. General recognition of safety requires common knowledge about the substance throughout the scientific community knowledgeable about the safety of substances directly or indirectly added to food.

(b) General recognition of safety based upon scientific procedures shall require the same quantity and quality of scientific evidence as is required to obtain approval of a food additive regulation for the ingredient. General recognition of safety through scientific procedures shall ordinarily be based upon published studies which may be corroborated by unpublished studies and other data and information.

(c) General recognition of safety through experience based on common use in food prior to January 1, 1958, may be determined without the quantity or quality of scientific procedures required for approval of a food additive

regulation. General recognition of safety through experience based on common use in food prior to January 1, 1958, shall ordinarily be based upon generally available data and information. An ingredient not in common use in food prior to January 1, 1958, may achieve general recognition of safety only through scientific procedures.

(d) The food ingredients listed as GRAS in part 582 of this chapter do not include all substances that are generally recognized as safe for their intended use in food. Because of the large number of substances the intended use of which results or may reasonably be expected to result, directly or indirectly, in their becoming a component or otherwise affecting the characteristics of food, it is impracticable to list all such substances that are GRAS. A food ingredient of natural biological origin that has been widely consumed for its nutrient properties in the United States prior to January 1, 1958, without known detrimental effects, which is subject only to conventional processing as practiced prior to January 1, 1958, and for which no known safety hazard exists, will ordinarily be regarded as GRAS without specific inclusion in part 582 of this chapter.

(e) A food ingredient that is not GRAS or subject to a prior sanction requires a food additive regulation promulgated under section 409 of the act before it may be directly or indirectly added to food.

(f) A food ingredient that is listed as GRAS in part 582 of this chapter shall be regarded as GRAS only if, in addition to all the requirements in the applicable regulation, it also meets all of the following requirements:

(1) It complies with any applicable specifications, or in the absence of such specifications, shall be of a purity suitable for its intended use.

(2) It performs an appropriate function in the food or food-contact article in which it is used.

(3) It is used at a level no higher than necessary to achieve its intended purpose in that food or, if used as a component of a food-contact article, at a level no higher than necessary to achieve its intended purpose in that article.

§ 570.35

21 CFR Ch. I (4–1–99 Edition)

(g) New information may at an time require reconsideration of the GRAS status of a food ingredient. Any change in status shall be accomplished pursuant to § 570.38.

(h) If a substance is affirmed as GRAS pursuant to § 570.35 and listed in a regulation with no limitation other than good manufacturing practice, it shall be regarded as GRAS if its conditions of use are not significantly different from those reported in the regulation as the basis on which the GRAS status of the substance was affirmed. If the conditions of use are significantly different, such use of the substance may not be GRAS. In such case a manufacturer may not rely on the regulation as authorizing the use but must independently establish that the use is GRAS or must use the substance in accordance with a food additive regulation.

(i) If an ingredient is affirmed as GRAS pursuant to § 570.35 and listed in a regulation with specific limitation(s), it may be used in food only within such limitation(s) (including the category of food(s), the functional use(s) of the ingredient, and the level(s) of use). Any use of such an ingredient not in full compliance with each such established limitation shall require a food additive regulation.

(j) Pursuant to § 570.35, a food ingredient may be affirmed as GRAS and listed in a regulation for a specific use(s) without a general evaluation of use of the ingredient. In addition to the use(s) specified in the regulation, other uses of such an ingredient may also be GRAS. Any affirmation of GRAS status for a specific use(s), without a general evaluation of use of the ingredient, is subject to reconsideration upon such evaluation.

[42 FR 55206, Oct. 14, 1977]

§ 570.35 Affirmation of generally recognized as safe (GRAS) status.

(a) The Commissioner, either on his initiative or on the petition of an interested person, may affirm the GRAS status of substances that directly or indirectly become components of food.

(b)(1) If the Commissioner proposes on his own initiative that a substance is entitled to affirmation as GRAS, he will place all of the data and information

on which he relies on public file in the office of the Dockets Management Branch and will publish in the FEDERAL REGISTER a notice giving the name of the substance, its proposed uses, and any limitations proposed for purposes other than safety.

(2) The FEDERAL REGISTER notice will allow a period of 60 days during which any interested person may review the data and information and/or file comments with the Dockets Management Branch. Copies of all comments received shall be made available for examination in the Dockets Management Branch's office.

(3) The Commissioner will evaluate all comments received. If he concludes that there is convincing evidence that the substance is GRAS as defined in § 570.3(k), he will publish a notice in the FEDERAL REGISTER listing the substance in this subchapter E as GRAS.

(4) If, after evaluation of the comments, the Commissioner concludes that there is a lack of convincing evidence that the substance is GRAS and that it should be considered a food additive subject to section 409 of the act, he shall publish a notice thereof in the FEDERAL REGISTER in accordance with § 570.38.

(c)(1) Persons seeking the affirmation of GRAS status of substances as provided for in § 570.30(e), except those subject to the NAS-NRC GRAS list survey (36 FR 20546), shall submit a petition for GRAS affirmation pursuant to part 10 of this chapter. Such petition shall contain information to establish that the GRAS criteria as set forth in § 570.30(b) have been met, in the following form:

(i) Description of the substance, including:

(a) Common or usual name.

(b) Chemical name.

(c) Chemical Abstract Service (CAS) registry number.

(d) Empirical formula.

(e) Structural formula.

(f) Specifications for food grade material, including arsenic and heavy metals. (Recommendation for any change in the Food Chemicals Codex monograph should be included where applicable.)

(g) Quantitative compositions.

(h) Manufacturing process (excluding any trade secrets).

(ii) Use of the substance, including:

(a) Date when use began.

(b) Information and reports or other data on past uses in food.

(c) Foods in which used, and levels of use in such foods, and for what purposes.

(iii) Methods for detecting the substance in food, including:

(a) References to qualitative and quantitative methods for determining the substance(s) in food, including the type of analytical procedures used.

(b) Sensitivity and reproducibility of such method(s).

(iv) Information to establish the safety and functionality of the substance in food. Published scientific literature, evidence that the substance is identical to a GRAS counterpart of natural biological origin, and other data may be submitted to support safety. Any adverse information or consumer complaints shall be included. Complete bibliographic references shall be provided where a copy of the article is not provided.

(v) A statement signed by the person responsible for the petition that to the best of his knowledge it is a representative and balanced submission that includes unfavorable information, as well as favorable information, known to him pertinent to the evaluation of the safety and functionality of the substance.

(vi) If nonclinical laboratory studies are involved, additional information and data submitted in support of filed petitions shall include, with respect to each nonclinical study, either a statement that the study was conducted in compliance with the requirements set forth in part 58 of this chapter, or, if the study was not conducted in compliance with such regulations, a brief statement of the reason for the non-compliance.

(vii) [Reserved]

(viii) A claim for categorical exclusion under § 25.30 or 25.32 of this chapter or an environmental assessment under § 25.40 of this chapter.

(2) Within 30 days after the date of filing the petition, the Commissioner will place the petition on public file in the Dockets Management Branch and

will publish a notice of filing in the FEDERAL REGISTER giving the name of the petitioner and a brief description of the petition including the name of the substance, its proposed use, and any limitations proposed for reasons other than safety. A copy of the notice will be mailed to the petitioner at the time the original is sent to the FEDERAL REGISTER.

(3) The notice of filing in the FEDERAL REGISTER will allow a period of 60 days during which any interested person may review the petition and/or file comments with the Dockets Management Branch. Copies of all comments received shall be made available for examination in the Dockets Management Branch.

(4) The Commissioner will evaluate the petition and all available information including all comments received. If the petition and such information provide convincing evidence that the substance is GRAS as defined in § 570.3, he will publish an order in the FEDERAL REGISTER listing the substance in this subchapter E as GRAS.

(5) If, after evaluation of the petition and all available information, the Commissioner concludes that there is a lack of convincing evidence that the substance is GRAS and that it should be considered a food additive subject to section 409 of the act, he shall publish a notice thereof in the FEDERAL REGISTER in accordance with § 570.38.

(6) The notice of filing in the FEDERAL REGISTER will request submission of proof of any applicable prior sanction for use of the ingredient under conditions different from those proposed to be determined to be GRAS. The failure of any person to come forward with proof of such an applicable prior sanction in response to the notice of filing will constitute a waiver of the right to assert or rely on such sanction at any later time. The notice of filing will also constitute a proposal to establish a regulation under this subchapter E, incorporating the same provisions, in the event that such a regulation is determined to be appropriate as a result of submission of proof of such an

§ 570.38

applicable prior sanction in response to the notice of filing.

[41 FR 38644, Sept. 10, 1976, as amended at 42 FR 4717, Jan. 25, 1977; 42 FR 15675, Mar. 22, 1977; 42 FR 55207, Oct. 10, 1977; 50 FR 7517, Feb. 22, 1985; 50 FR 16668, Apr. 26, 1985; 54 FR 18281, Apr. 28, 1989; 62 FR 40600, July 29, 1997]

§ 570.38 Determination of food additive status.

(a) The Commissioner may, in accordance with § 570.35 (b)(4) or (c)(5), publish a notice in the FEDERAL REGISTER determining that a substance is not GRAS and is a food additive subject to section 409 of the act.

(b)(1) The Commissioner, on his own initiative or on the petition of any interested person, pursuant to part 10 of this chapter, may issue a notice in the FEDERAL REGISTER proposing to determine that a substance is not GRAS and is a food additive subject to section 409 of the act. Any petition shall include all relevant data and information of the type described in § 571.130(b) of this chapter. The Commissioner will place all of the data and information on which he relies on public file in the Dockets Management Branch and will include in the FEDERAL REGISTER notice the name of the substance, its known uses, and a summary of the basis for the determination.

(2) The FEDERAL REGISTER notice will allow a period of 60 days during which any interested person may review the data and information and/or file comments with the Dockets Management Branch. Copies of all comments shall be made available for examination in the Dockets Management Branch.

(3) The Commissioner will evaluate all comments received. If he concludes that there is a lack of convincing evidence that the substance is GRAS or is otherwise exempt from the definition of a food additive in section 201(s) of the act, he will publish a notice thereof in the FEDERAL REGISTER. If he concludes that there is convincing evidence that the substance is GRAS, he will publish an order in the FEDERAL REGISTER listing the substance in this subchapter E as GRAS.

(c) A FEDERAL REGISTER notice determining that a substance is a food additive shall provide for the use of the ad-

21 CFR Ch. I (4–1–99 Edition)

ditive in food or food-contact surfaces as follows:

(1) It may promulgate a food additive regulation governing use of the additive.

(2) It may promulgate an interim food additive regulation governing use of the additive.

(3) It may require discontinuation of the use of the additive.

(4) It may adopt any combination of the above three approaches for different uses or levels of use of the additive.

(d) If the Commissioner of Food and Drugs is aware of any prior sanction for use of the substance, he will concurrently propose a separate regulation covering such use of the ingredient under this subchapter E. If the Commissioner is unaware of any such applicable prior sanction, the proposed regulation will so state and will require any person who intends to assert or rely on such sanction to submit proof of its existence. Any regulation promulgated pursuant to this section constitutes a determination that excluded uses would result in adulteration of the food in violation of section 402 of the act, and the failure of any person to come forward with proof of such an applicable prior sanction in response to the proposal will constitute a waiver of the right to assert or rely on such sanction at any later time. The notice will also constitute a proposal to establish a regulation under this subchapter E., incorporating the same provisions, in the event that such a regulation is determined to be appropriate as a result of submission of proof of such an applicable prior sanction in response to the proposal.

[41 FR 38644, Sept. 10, 1976, as amended at 42 FR 4717, Jan. 25, 1977; 42 FR 15675, Mar. 22, 1977; 42 FR 55207, Oct. 14, 1977; 54 FR 18281, Apr. 28, 1989]

PART 571—FOOD ADDITIVE PETITIONS

Subpart A—General Provisions

Sec.

571.1 Petitions.

571.6 Amendment of petition.

571.7 Withdrawal of petition without prejudice.

Food and Drug Administration, HHS

§ 571.1

Subpart B—Administrative Actions on Applications

- 571.100 Regulation based on petition.
571.102 Effective date of regulation.
571.110 Procedure for objections and hearings.
571.115 Application of the cancer clause of section 409 of the act.
571.130 Procedure for amending and repealing tolerances or exemptions from tolerances.

AUTHORITY: 21 U.S.C. 321, 342, 348, 371; 42 U.S.C. 241.

SOURCE: 41 FR 38647, Sept. 10, 1976, unless otherwise noted.

Subpart A—General Provisions

§ 571.1 Petitions.

(a) Petitions to be filed with the Commissioner under the provisions of section 409(b) of the act shall be submitted in triplicate. If any part of the material submitted is in a foreign language, it shall be accompanied by an accurate and complete English translation. The petition shall state petitioner's post office address to which published notices or orders issued or objections filed pursuant to section 409 of the act may be sent.

(b) Pertinent information may be incorporated in, and will be considered as part of, a petition on the basis of specific reference to such information submitted to and retained in the files of the Food and Drug Administration. However, any reference to unpublished information furnished by a person other than the applicant will not be considered unless use of such information is authorized in a written statement signed by the person who submitted it. Any reference to published information offered in support of a food-additive petition should be accompanied by reprints or photostatic copies of such references.

(c) Petitions shall include the following data and be submitted in the following form:

(Date) _____
Name of petitioner _____
Post office address _____
Date _____
Name of food additive and proposed use _____

Food and Drug Administration
CENTER FOR VETERINARY MEDICINE,
Director, Division of Animal Feeds (HFV-220),
7500 Standish Pl., Rockville, MD 20855.

DEAR SIRs: The undersigned, _____

submits this petition pursuant to section 409(b)(1) of the Federal Food, Drug, and

Cosmetic Act with respect to _____

(Name of the food additive and proposed use)

Attached hereto, in triplicate, and constituting a part of this petition, are the following:

A. The name and all pertinent information concerning the food additive, including chemical identity and composition of the food additive, its physical, chemical, and biological properties, and specifications prescribing the minimum content of the desired component(s) and identifying and limiting the reaction byproducts and other impurities. Where such information is not available, a statement as to the reasons why it is not should be submitted.

When the chemical identity and composition of the food additive is not known, the petition shall contain information in sufficient detail to permit evaluation regarding the method of manufacture and the analytical controls used during the various stages of manufacturing, processing, or packing of the food additive which are relied upon to establish that it is a substance of reproducible composition. Alternative methods and controls and variations in methods and controls within reasonable limits that do not affect the characteristics of the substance or the reliability of the controls may be specified.

If the food additive is a mixture of chemicals, the petition shall supply a list of all substances used in the synthesis, extraction, or other method of preparation, regardless of whether they undergo chemical change in the process. Each substance should be identified by its common English name and complete chemical name, using structural formulas when necessary for specific identification. If any proprietary preparation is used as a component, the proprietary name should be followed by a complete quantitative statement of composition. Reasonable alternatives for any listed substance may be specified.

If the petitioner does not himself perform all the manufacturing, processing, and packing operations for a food additive, the petition shall identify each person who will perform a part of such operations and designate the part.

The petition shall include stability data, and, if the data indicate that it is needed to ensure the identity, strength, quality, or purity of the additive, the expiration date that will be employed.

B. The amount of the food additive proposed for use and the purposes for which it is

§ 571.1

21 CFR Ch. I (4–1–99 Edition)

proposed, together with all directions, recommendations, and suggestions regarding the proposed use, as well as specimens of the labeling proposed for the food additive and any labeling that will be required by applicable provisions of the Federal Food, Drug, and Cosmetic Act on the finished food by reason of the use of the food additive. If the additive results or may reasonably be expected to result from the use of packaging material, the petitioner shall show how this may occur and what residues may reasonably be anticipated.

(Typewritten or other draft-labeling copy will be accepted for consideration of the petition, provided a statement is made that final printed labeling identical in content to the draft copy will be submitted as soon as available and prior to the marketing of the food additive.)

If the food additive is one for which a tolerance limitation is required to assure its safety, the level of use proposed should be no higher than the amount reasonably required to accomplish the intended physical or other technical effect, even though the safety data may support a higher tolerance.)

C. Data establishing that the food additive will have the intended physical or other technical effect or that it may reasonably be expected to become a component, or to affect the characteristics, directly or indirectly, of food and the amount necessary to accomplish this. These data should include information in sufficient detail to permit evaluation with control data.

D. A description of practicable methods to determine the amount of the food additive in the raw, processed, and/or finished food and of any substance formed in or on such food because of its use. The test proposed shall be one that can be used for food-control purposes and that can be applied with consistent results by any properly equipped and trained laboratory personnel.

E. Full reports of investigations made with respect to the safety of the food additive.

(A petition may be regarded as incomplete unless it includes full reports of adequate tests reasonably applicable to show whether or not the food additive will be safe for its intended use. The reports ordinarily should include detailed data derived from appropriate animal and other biological experiments in which the methods used and the results obtained are clearly set forth. The petition shall not omit without explanation any reports of investigations that would bias an evaluation of the safety of the food additive.)

F. Proposed tolerances for the food additive, if tolerances are required in order to ensure its safety. A petitioner may include a proposed regulation.

G. If submitting petition to modify an existing regulation issued pursuant to section 409(c)(1)(A) of the act, full information on each proposed change that is to be made in

the original regulation must be submitted. The petition may omit statements made in the original petition concerning which no change is proposed. A supplemental petition must be submitted for any change beyond the variations provided for in the original petition and the regulation issued on the basis of the original petition.

H. The petitioner is required to submit either a claim for categorical exclusion under § 25.30 or § 25.32 of this chapter or an environmental assessment under § 25.40 of this chapter.

Yours very truly,

Petitioner _____

By _____

(Indicate authority)

(d) The petitioner will be notified of the date on which his petition is filed, and an incomplete petition, or one that has not been submitted in triplicate, will usually be retained but not filed as a petition under section 409 of the act. The petitioner will be notified in what respects his petition is incomplete.

(e) The petition must be signed by the petitioner or by his attorney or agent, or (if a corporation) by an authorized official.

(f) The data specified under the several lettered headings should be submitted on separate sheets or sets of sheets, suitably identified. If such data have already been submitted with an earlier application, the present petition may incorporate it by specific reference to the earlier. If part of the data have been submitted by the manufacturer of the food additive as a master file, the petitioner may refer to the master file if and to the extent he obtains the manufacturer's written permission to do so. The manufacturer may authorize specific reference to the data without disclosure to the petitioner. Nothing herein shall prevent reference to published data.

(g) A petition shall be retained but shall not be filed if any of the data prescribed by section 409(b) of the act are lacking or are not set forth so as to be readily understood.

(h)(1) The following data and information in a food additive petition are available for public disclosure, unless extraordinary circumstances are shown, after the notice of filing of the petition is published in the FEDERAL REGISTER or, if the petition is not promptly filed because of deficiencies in it, after the petitioner is informed

that it will not be filed because of the deficiencies involved:

(i) All safety and functionality data and information submitted with or incorporated by reference in the petition.

(ii) A protocol for a test or study, unless it is shown to fall within the exemption established for trade secrets and confidential commercial information in § 20.61 of this chapter.

(iii) Adverse reaction reports, product experience reports, consumer complaints, and other similar data and information, after deletion of:

(a) Names and any information that would identify the person using the product.

(b) Names and any information that would identify any third party involved with the report, such as a physician or hospital or other institution.

(iv) A list of all ingredients contained in a food additive, whether or not it is in descending order of predominance. A particular ingredient or group of ingredients shall be deleted from any such list prior to public disclosure if it is shown to fall within the exemption established in § 20.61 of this chapter, and a notation shall be made that any such ingredient list is incomplete.

(v) An assay method or other analytical method, unless it serves no regulatory or compliance purpose and is shown to fall within the exemption established in § 20.61 of this chapter.

(2) The following data and information in a food additive petition are not available for public disclosure unless they have been previously disclosed to the public as defined in § 20.81 of this chapter or they relate to a product or ingredient that has been abandoned and they no longer represent a trade secret or confidential commercial or financial information as defined in § 20.61 of this chapter:

(i) Manufacturing methods or processes, including quality control procedures.

(ii) Production, sales, distribution, and similar data and information, except that any compilation of such data and information aggregated and prepared in a way that does not reveal data or information which is not available for public disclosure under this provision is available for public disclosure.

(iii) Quantitative or semiquantitative formulas.

(3) All correspondence and written summaries of oral discussions relating to a food additive petition are available for public disclosure in accordance with the provisions of part 20 of this chapter when the food additive regulation is published in the FEDERAL REGISTER.

(4) For purposes of this regulation, safety and functionality data include all studies and tests of a food additive on animals and humans and all studies and tests on a food additive for identity, stability, purity, potency, performance, and usefulness.

(i)(1) Within 15 days after receipt, the Commissioner will notify the petitioner of acceptance or nonacceptance of a petition, and if not accepted the reasons therefor. If accepted, the date of the notification letter sent to petitioner becomes the date of filing for the purposes of section 409(b)(5) of the act. If the petitioner desires, he may supplement a deficient petition after being notified regarding deficiencies. If the supplementary material or explanation of the petition is deemed acceptable, petitioner shall be notified. The date of such notification becomes the date of filing. If the petitioner does not wish to supplement or explain the petition and requests in writing that it be filed as submitted, the petition shall be filed and the petitioner so notified. The date of such notification becomes the date of filing.

(2) The Commissioner will publish in the FEDERAL REGISTER within 30 days from the date of filing of such petition, a notice of the filing, the name of the petitioner, and a brief description of the proposal in general terms. In the case of a food additive which becomes a component of food by migration from packaging material, the notice shall include the name of the migratory substance, and where it is different from that of one of the original components, the name of the parent component, the maximum quantity of the migratory substance that is proposed for use in food, and the physical or other technical effect which the migratory substance or its parent component is intended to have in the packaging material. A copy of the notice will be

§ 571.6

mailed to the petitioner when the original is forwarded to the FEDERAL REGISTER for publication.

(j) The Commissioner may request a full description of the methods used in, and the facilities and controls used for, the production of the food additive, or a sample of the food additive, articles used as components thereof, or of the food in which the additive is proposed to be used, at any time while a petition is under consideration. The Commissioner shall specify in the request for a sample of the food additive, or articles used as components thereof, or of the food in or on which the additive is proposed to be used, a quantity deemed adequate to permit tests of analytical methods to determine quantities of the food additive present in foods for which it is intended to be used or adequate for any study or investigation reasonably required with respect to the safety of the food additive or the physical or technical effect it produces. The date used for computing the 90-day limit for the purposes of section 409(c)(2) of the act shall be moved forward 1 day for each day after the mailing date of the request taken by the petitioner to submit the sample. If the information or sample is requested a reasonable time in advance of the 180 days, but is not submitted within such 180 days after filing of the petition, the petition will be considered withdrawn without prejudice.

(k) If nonclinical laboratory studies are involved, petitions filed with the Commissioner under section 409(b) of the act shall include, with respect to each study, either a statement that the study was conducted in compliance with the requirements set forth in part 58 of this chapter, or, if the study was not conducted in compliance with such regulations, a brief statement of the reason for the noncompliance.

[41 FR 38647, Sept. 10, 1976, as amended at 42 FR 15675, Mar. 22, 1977; 50 FR 7518, Feb. 22, 1985; 50 FR 16668, Apr. 26, 1985; 52 FR 8583, Mar. 19, 1987; 57 FR 6476, Feb. 25, 1992; 62 FR 40600, July 29, 1997]

§ 571.6 Amendment of petition.

After a petition has been filed, the petitioner may submit additional information or data in support thereof. In such cases, if the Commissioner de-

21 CFR Ch. I (4–1–99 Edition)

termines that the additional information or data amounts to a substantive amendment, the petition as amended will be given a new filing date, and the time limitation will begin to run anew. If nonclinical laboratory studies are involved, additional information and data submitted in support of filed petitions shall include, with respect to each such study, either a statement that the study was conducted in compliance with the requirements set forth in part 58 of this chapter, or, if the study was not conducted in compliance with such regulations, a brief statement of the reason or the noncompliance.

[41 FR 38647, Sept. 10, 1976, as amended at 50 FR 7518, Feb. 22, 1985; 50 FR 16668, Apr. 26, 1985]

§ 571.7 Withdrawal of petition without prejudice.

(a) In some cases the Commissioner will notify the petitioner that the petition, while technically complete, is inadequate to justify the establishment of a regulation or the regulation requested by petitioner. This may be due to the fact that the data are not sufficiently clear or complete. In such cases, the petitioner may withdraw the petition pending its clarification or the obtaining of additional data. This withdrawal will be without prejudice to a future filing. Upon refiling, the time limitation will begin to run anew from the date of refiling.

(b) At any time before the order provided for in § 571.100(a) has been forwarded to the FEDERAL REGISTER for publication, the petitioner may withdraw the petition without prejudice to a future filing. Upon refiling the time limitation will begin to run anew.

Subpart B—Administrative Actions on Applications

§ 571.100 Regulation based on petition.

(a) The Commissioner will forward for publication in the FEDERAL REGISTER, within 90 days after filing of the petition (or within 180 days if the time is extended as provided for in section 409(c)(2) of the act), a regulation prescribing the conditions under which the food additive may be safely used

(including, but not limited to, specifications as to the particular food or classes of food in or on which such additive may be used, the maximum quantity that may be used or permitted to remain in or on such food, the manner in which such additive may be added to or used in or on such food, and any directions or other labeling or packaging requirements for such additive deemed necessary by him to assure the safety of such use), and prior to the forwarding of the order to the FEDERAL REGISTER for publication shall notify the petitioner of such order and the reasons for such action; or by order deny the petition, and shall notify the petitioner of such order and of the reasons for such action.

(b) If the Commissioner determines that additional time is needed to study and investigate the petition, he shall by written notice to the petitioner extend the 90-day period for not more than 180 days after the filing of the petition.

§ 571.102 Effective date of regulation.

A regulation published in accordance with § 571.100(a) shall become effective upon publication in the FEDERAL REGISTER.

§ 571.110 Procedure for objections and hearings.

Objections and hearings relating to food additive regulations under section 409(c), (d), or (h) of the act shall be governed by part 12 of this chapter.

[42 FR 4717, Jan. 25, 1977, as amended at 42 FR 15676, Mar. 22, 1977]

§ 571.115 Application of the cancer clause of section 409 of the act.

Food additives intended for use as an ingredient in food for animals that are raised for food production and that have the potential to contaminate human food with residues whose consumption could present a risk of cancer to people must satisfy the requirements of subpart E of part 500 of this chapter.

[52 FR 49588, Dec. 31, 1987]

§ 571.130 Procedure for amending and repealing tolerances or exemptions from tolerances.

(a) The Commissioner, on his own initiative or on the petition of any interested person, pursuant to part 10 of this chapter, may propose the issuance of a regulation amending or repealing a regulation pertaining to a food additive or granting or repealing an exception for such additive.

(b) Any such petition shall include an assertion of facts, supported by data, showing that new information exists with respect to the food additive or that new uses have been developed or old uses abandoned, that new data are available as to toxicity of the chemical, or that experience with the existing regulation or exemption may justify its amendment or repeal. New data shall be furnished in the form specified in § 571.1 for submitting petitions.

[42 FR 4717, Jan. 25, 1977; 42 FR 15676, Mar. 22, 1977]

PART 573—FOOD ADDITIVES PERMITTED IN FEED AND DRINKING WATER OF ANIMALS

Subpart A [Reserved]

Subpart B—Food Additive Listing

Sec.

- 573.120 Acrylamide-acrylic acid resin.
- 573.130 Aminoglycoside 3'-phospho- transferase II.
- 573.140 Ammoniated cottonseed meal.
- 573.160 Ammoniated rice hulls.
- 573.180 Anhydrous ammonia.
- 573.200 Condensed animal protein hydrolysate.
- 573.220 Feed-grade biuret.
- 573.225 1,3-Butylene glycol.
- 573.240 Calcium periodate.
- 573.260 Calcium silicate.
- 573.280 Feed-grade calcium stearate and sodium stearate.
- 573.300 Choline xanthate.
- 573.310 Crambe meal, heat toasted.
- 573.320 Diammonium phosphate.
- 573.340 Diatomaceous earth.
- 573.360 Disodium EDTA.
- 573.380 Ethoxyquin in animal feeds.
- 573.400 Ethoxyquin in certain dehydrated forage crops.
- 573.420 Ethyl cellulose.
- 573.440 Ethylene dichloride.
- 573.450 Fermented ammoniated condensed whey.

§ 573.120

573.460 Formaldehyde.
573.480 Formic acid.
573.500 Condensed, extracted glutamic acid fermentation product.
573.520 Hemicellulose extract.
573.530 Hydrogenated corn syrup.
573.540 Hydrolyzed leather meal.
573.560 Iron ammonium citrate.
573.580 Iron-choline citrate complex.
573.600 Lignin sulfonates.
573.620 Menadione dimethylpyrimidinol bisulfite.
573.625 Menadione nicotinamide bisulfite.
573.640 Methyl esters of higher fatty acids.
573.660 Methyl glucoside-coconut oil ester.
573.680 Mineral oil.
573.700 Sodium nitrite.
573.720 Petrolatum.
573.740 Odorless light petroleum hydrocarbons.
573.750 *Pichia pastoris* dried yeast.
573.760 Poloxalene.
573.780 Polyethylene.
573.800 Polyethylene glycol (400) mono- and dioleate.
573.820 Polyoxyethylene glycol (400) mono- and dioleates.
573.840 Polysorbate 60.
573.860 Polysorbate 80.
573.870 Poly(2-vinylpyridine-co-styrene).
573.880 Normal propyl alcohol.
573.900 Pyrophyllite.
573.914 Salts of volatile fatty acids.
573.920 Selenium.
573.940 Silicon dioxide.
573.960 Sorbitan monostearate.
573.980 Taurine.
573.1000 Verxite.
573.1010 Xanthan gum.
573.1020 Yellow prussiate of soda.

AUTHORITY: 21 U.S.C. 321, 342, 348.

SOURCE: 41 FR 38652, Sept. 10, 1976, unless otherwise noted.

Subpart A [Reserved]

Subpart B—Food Additive Listing

§ 573.120 Acrylamide-acrylic acid resin.

Acrylamide-acrylic acid resin (hydrolyzed polyacrylamide), only for the purposes of this section as described below, may be safely used in accordance with the following prescribed conditions:

(a) The additive is produced by polymerization of acrylamide with partial hydrolysis, or by copolymerization of acrylamide and acrylic acid with the greater part of the polymer being composed of acrylamide units.

21 CFR Ch. I (4–1–99 Edition)

(b) The additive meets the following specifications:

(1) A minimum molecular weight of 3 million.

(2) Viscosity range: 3,000 to 6,000 centipoises at 77° F in a 1 percent aqueous solution as determined by LVF Brookfield Viscometer or equivalent using a number 6 spindle at 20 r.p.m.

(3) Residual acrylamide: Not more than 0.05 percent.

(c) It is used as a thickener and suspending agent in nonmedicated aqueous suspensions intended for addition to animal feeds.

[41 FR 38652, Sept. 10, 1976, as amended at 45 FR 38058, June 6, 1980]

§ 573.130 Aminoglycoside 3'-phosphotransferase II.

The food additive aminoglycoside 3'-phosphotransferase II may be safely used in the development of genetically modified cotton, oilseed rape, and tomatoes in accordance with the following prescribed conditions:

(a) The food additive is the enzyme aminoglycoside 3'-phosphotransferase II (CAS Reg. No. 58943–39–8) which catalyzes the phosphorylation of certain aminoglycoside antibiotics, including kanamycin, neomycin, and gentamicin.

(b) Aminoglycoside 3'-phosphotransferase II is encoded by the *kan^r* gene originally isolated from transposon Tn5 of the bacterium *Escherichia coli*.

(c) The level of the additive does not exceed the amount reasonably required for selection of plant cells carrying the *kan^r* gene along with the genetic material of interest.

[59 FR 26711, May 23, 1994]

§ 573.140 Ammoniated cottonseed meal.

The food additive ammoniated cottonseed meal may be safely used in accordance with the following conditions:

(a) The food additive is the product obtained by the treatment of cottonseed meal with anhydrous ammonia until a pressure of 50 pounds per square inch gauge is reached.

(b) It is used or intended for use in the feed of ruminants as a source of protein and/or as a source of non-protein nitrogen in an amount not to exceed 20 percent of the total ration.

(c) To assure safe use, the label and labeling of the additive and of any feed additive supplement, concentrate, or premix prepared therefrom shall bear, in addition to the other information required by the act, the following:

- (1) The name of the additive.
- (2) The maximum percentage of equivalent crude protein from the nonprotein nitrogen.
- (3) Directions for use to provide not more than 20 percent of the additive in the total ration.
- (4) A statement:
 - (i) That not more than one-third of the total protein in the feed should come from nonprotein nitrogen sources.
 - (ii) That the additive is not to be given to debilitated or starved animals.
 - (iii) "Warning—This feed should be used only in accordance with directions furnished on the label."

[41 FR 38652, Sept. 10, 1976, as amended at 42 FR 52397, Sept. 30, 1977]

§ 573.160 Ammoniated rice hulls.

The food additive ammoniated rice hulls may be safely used in accordance with the following prescribed conditions:

- (a) The food additive is the product obtained by the treatment of ground rice hulls with monocalcium phosphate and anhydrous ammonia at a temperature of 350° F and a pressure of 175 pounds per square inch.
- (b) It is used or intended for use in the feed of beef cattle as a source of crude fiber and as the sole source of nonprotein nitrogen in an amount not to exceed 20 percent of the total ration.
- (c) To assure safe use of the additive, the label and labeling of the additive and of any feed additive supplement, feed additive concentrate, or feed additive premix prepared therefrom, shall contain, in addition to other information required by the act, the following:
 - (1) The name of the additive.
 - (2) The maximum percentage of equivalent crude protein from the nonprotein nitrogen.
 - (3) Directions for use to provide not more than 20 percent of the additive in the total ration, and a prominent statement: "Warning—This feed should be used only in accordance with the directions furnished on the label."

§ 573.180 Anhydrous ammonia.

(a) The food additive anhydrous ammonia is applied directly to corn plant material and thoroughly blended prior to ensiling. It is used or intended for use as a source of nonprotein nitrogen in cattle feed in accordance with paragraphs (a)(1), (2), or (3) as follows:

(1)(i) The food additive anhydrous ammonia is applied as a component of an aqueous premix containing 16 to 17 percent ammonia, with molasses, minerals, and not less than 83 percent crude protein. The premix is a source of nonprotein nitrogen and minerals.

(ii) In addition to the requirements of paragraph (b) of this section, the labeling shall bear an expiration date of not more than 10 weeks after date of manufacture; a statement that additional protein should not be fed to lactating dairy cows producing less than 32 pounds of milk per day nor beef cattle consuming less than 1 percent of body weight daily in shelled corn; and a warning not to use additional trace mineral supplementation with treated silage.

(2)(i) The food additive anhydrous ammonia is applied directly to corn plant material for use in dairy or beef cattle rations.

(ii) The anhydrous ammonia is applied at a rate not to exceed the equivalent of 0.35 percent of the corn plant material.

(iii) It is applied to corn plant material containing 30 to 35 percent dry matter.

(iv) It is applied so that 75 to 85 percent of the additive is liquid at ambient pressure.

(3)(i) The food additive anhydrous ammonia is applied after being diluted to a 15 to 30 percent aqueous ammonia solution (by weight).

(ii) The anhydrous ammonia solution is applied at a rate not to exceed anhydrous ammonia equivalent to 0.3 percent of the corn plant material.

(iii) It is applied to corn plant material containing 28 to 38 percent dry matter.

(iv) The silage treated with aqueous ammonia is to be fed to dairy cattle only.

(b) Its labeling shall bear, in addition to the other requirements of the act,

§ 573.200

the name of the additive, the concentration of ammonia, the maximum percentage of equivalent crude protein from nonprotein nitrogen, and directions for use consistent with this section.

[44 FR 40284, July 10, 1979]

§ 573.200 Condensed animal protein hydrolysate.

(a) *Identity.* The condensed animal protein hydrolysate is produced from the meat byproducts scraped from cured (salted) hides taken from cattle slaughtered for food consumption. The meat byproduct is hydrolyzed with heat and phosphoric acid.

(b) *Specifications.* The additive shall conform to the following percent-by-weight specifications:

Moisture, not less than 45 percent nor more than 50 percent.

Protein, not less than 24 percent.

Salt (NaCl), not more than 15 percent.

Phosphorus, not less than 2.25 percent.

(c) *Uses.* It is used or intended for use as a source of animal protein, phosphorus, and salt (NaCl) as follows:

(1) In poultry and swine feed in an amount not to exceed 5 percent by weight of the feed.

(2) In feed concentrates for cattle in an amount not to exceed 10 percent by weight of the concentrate.

(d) *Labeling.* The label and labeling shall bear, in addition to the other information required by the act:

(1) The name of the additive, condensed animal protein hydrolysate.

(2) Adequate directions for use including maximum quantities permitted for each species and a guaranteed analysis of the additive.

§ 573.220 Feed-grade biuret.

The food additive feed grade biuret may be safely used in ruminant feed in accordance with the following prescribed conditions:

(a) The food additive is the product resulting from the controlled pyrolysis of urea conforming to the following specifications:

	Percent
Biuret	55 minimum.
Urea	15 maximum.
Cyanuric acid and triuret	30 maximum.
Mineral oil	0.5 maximum.

21 CFR Ch. I (4–1–99 Edition)

	Percent
Total nitrogen (equivalent to 218.75 pct crude protein).	35 minimum.

(b) It is used in ruminant feeds as a source of nonprotein nitrogen.

(c) To assure safe use of the additive:

(1) The label and labeling of the additive and that of any feed additive supplement, feed additive concentrate, feed additive premix, or complete feed prepared therefrom shall contain, in addition to other information required by the act, the following:

(i) The name of the additive.

(ii) The maximum percentage of equivalent crude protein from nonprotein nitrogen.

(iii) The statement “Do not feed to animals producing milk for human consumption.”

(2) The label shall recommend that the diet be balanced to provide adequate nutrients when equivalent crude protein from all forms of nonprotein nitrogen exceed one-third of the total crude protein in the total daily ration.

§ 573.225 1,3-Butylene glycol.

The food additive 1,3-butyleneglycol (1,3-butanediol) may be safely used in accordance with the following prescribed conditions:

(a) It complies with the specifications in § 173.220(a) of this chapter.

(b) It is intended for use in swine feed as a source of energy.

(c) It is to be thoroughly mixed into feed at levels not to exceed 9 percent of the dry matter of the total ration.

(d) 1,3-Butylene glycol should be mixed in feed with equipment adapted for the addition of liquids, and the feed should be mixed not less than 5 minutes after its addition.

[53 FR 40061, Oct. 13, 1988]

§ 573.240 Calcium periodate.

The food additive calcium periodate may be safely used in accordance with the following prescribed conditions:

(a) The additive is produced by reacting calcium iodate with calcium hydroxide or calcium oxide to form a substance consisting of not less than 60 percent by weight of penta calcium orthoperiodate containing 28 to 31 percent by weight of iodine.

(b) It is used or intended for use in salt for livestock as a source of iodine.

§ 573.260 Calcium silicate.

Calcium silicate, including synthetic calcium silicate, may be safely used as an anticaking agent in animal feed, provided that the amount of calcium silicate does not exceed 2 percent.

§ 573.280 Feed-grade calcium stearate and sodium stearate.

Feed-grade calcium stearate and sodium stearate may be safely used in an animal feed in accordance with the following prescribed conditions:

(a) Feed-grade calcium stearate and sodium stearate are the calcium or sodium salts of a fatty acid mixture that is predominately stearic acid. Associated fatty acids, including palmitic acid and minor amounts of lauric, myristic, pentadecanoic, margaric, arachidic, and other fatty acids may be contained in the mixture, but such associated fatty acids in aggregate do not exceed 35 percent by weight of the mixture. The fatty acids may be derived from feed-grade fats or oils.

(b) The additives meet the following specifications:

(1) Unsaponifiable matter does not exceed 2 percent.

(2) They are free of chick-edema factor.

(c) The additives are manufactured so that in aqueous solution they are exposed for 1 hour or longer to temperature in excess of 180 °F.

(d) They are used as anticaking agents in animal feeds in accordance with current good manufacturing practices.

[63 FR 8573, Feb. 20, 1998]

§ 573.300 Choline xanthate.

Choline xanthate may be safely used as a component of animal feed as an added source of choline to supplement the diets of poultry, ruminants, and swine in accordance with good feeding practice.

§ 573.310 Crambe meal, heat toasted.

(a) The additive is the seed meal of *Crambe abyssinica* obtained after the removal of oil from the seed and hull. The oil may be removed by pre-press

solvent extraction or by solvent extraction alone. The resulting seed meal is heat toasted.

(b) The additive conforms to the following percent-by-weight specifications: moisture, not more than 11 percent; oil, not more than 4 percent; crude protein, not less than 24 percent; crude fiber, not more than 26 percent; glucosinolate calculated as epiprogoitrin, not more than 4 percent; goitrin, not more than 0.1 percent; nitrile calculated as 1-cyano-2-hydroxy-3-butene, not more than 1.4 percent. At least 50 percent of the nitrogen shall be soluble in 0.5 M sodium chloride. Myrosinase enzyme activity shall be absent.

(c) The additive is used or intended for use in the feed of feedlot cattle as a source of protein in an amount not to exceed 4.2 percent of the total ration.

[46 FR 30082, June 5, 1981]

§ 573.320 Diammonium phosphate.

The food additive diammonium phosphate may be safely used in ruminant feed in accordance with the following prescribed conditions:

(a) The food additive is the product resulting from the neutralization of feeding-phosphoric-acid or defluorinated wet-process phosphoric acid with anhydrous ammonia. It contains not less than 106.25 percent equivalent crude protein (nitrogen X 6.25) and 20 percent phosphorus. It contains not more than the following:

1 part fluorine to 100 parts phosphorus.

75 parts per million or arsenic (as As).

30 parts per million of heavy metals, as lead (Pb).

(b) It is used in ruminant feeds as a source of phosphorus and nitrogen in an amount that supplies not more than 2 percent of equivalent crude protein in the total daily ration.

(c) To assure safe use of the additive, the label and labeling of the additive and that of any feed additive supplement, feed additive concentrate, feed additive premix, or complete feed prepared therefrom shall contain, in addition to other information required by the act, the following:

(1) The name of the additive.

§ 573.340

21 CFR Ch. I (4–1–99 Edition)

(2) The maximum percentage of equivalent crude protein from the non-protein nitrogen.

(3) If the feed additive premix, feed additive concentrate, or feed additive supplement contains more than 2 percent equivalent crude protein from diammonium phosphate, adequate directions for use and a prominent statement, "Warning—This feed should be used only in accordance with directions furnished on the label."

§ 573.340 Diatomaceous earth.

(a) *Identity.* The additive consists of siliceous skeletal material derived from various species of diatoms.

(b) *Specifications.* The additive shall conform to the following specifications:

Lead, not more than 15 parts per million.

Arsenic (as As), not more than 20 parts per million

Fluorine, not more than 600 parts per million.

(c) *Uses.* It is used or intended for use as an inert carrier or anticaking agent in animal feeds in an amount not to exceed 2 percent by weight of the total ration.

§ 573.360 Disodium EDTA.

The food additive disodium EDTA (disodium ethylenediaminetetraacetate) may be safely used in animal feeds, in accordance with the following prescribed conditions:

(a) The food additive contains a minimum of 99 percent disodium ethylenediaminetetraacetate dihydrate ($C_{10}H_{14}O_8N_2Na_2 \cdot 2H_2O$).

(b) It is used to solubilize trace minerals in aqueous solutions, which are then added to animal feeds.

(c) It is used or intended for use in an amount not to exceed 240 parts per million of the additive in finished feed.

(d) To assure safe use of the additive the label and labeling shall bear:

(1) The name of the additive; and

(2) Adequate mixing directions to ensure that the chelated trace-mineral mix is uniformly blended throughout the feed.

§ 573.380 Ethoxyquin in animal feeds.

Ethoxyquin (1,2-dihydro-6-ethoxy-2,2,4-trimethylquinoline) may be safely used in animal feeds, when incor-

porated therein in accordance with the following prescribed conditions.

(a) It is intended for use only: (1) As a chemical preservative for retarding oxidation of carotene, xanthophylls, and vitamins A and E in animal feed and fish food and, (2) as an aid in preventing the development of organic peroxides in canned pet food.

(b) The maximum quantity of the additive permitted to be used and to remain in or on the treated article shall not exceed 150 parts per million.

(c) To assure safe use of the additive, the label and labeling of the food additive container and that of any intermediate premixes prepared therefrom shall contain, in addition to other information required by the act:

(1) The name of the additive, ethoxyquin.

(2) A statement of the concentration or strength contained therein.

(3) Adequate use directions to provide for a finished article with the proper concentration of the additive as provided in paragraph (b) of this section, whether or not intermediate premixes are to be used.

(d) The label of any animal feed containing the additive shall, in addition to the other information required by the act, bear the statement "Ethoxyquin, a preservative" or "Ethoxyquin added to retard the oxidative destruction of carotene, xanthophylls, and vitamins A and E."

§ 573.400 Ethoxyquin in certain dehydrated forage crops.

Ethoxyquin (1,2-dihydro-6-ethoxy-2,2,4-trimethylquinoline) may be safely used in the dehydrated forage crops listed in paragraph (a) of this section when incorporated therein in accordance with the conditions prescribed in this section:

(a) It may be added to dehydrated forage prepared from:

Alfalfa	Medicago sativa.
Barley	Hordeum vulgare.
Clovers:	
Alsike clover	Trifolium hybridum.
Crimson clover	Trifolium incarnatum.
Red clover	Trifolium pratense.
White clover (including Ladino)	Trifolium repens.
White sweetclover	Melilotus alba.
Yellow sweetclover	Melilotus officinalis.
Coastal Bermudagrass	Cynodon dactylon.
Corn	Zea mays.
Fescue	Festuca sp.

Food and Drug Administration, HHS

§ 573.450

Oats	Avena sativa.
Orchardgrass	Dactylis glomerata.
Reed canarygrass	Phalaris arundinacea.
Ryegrass (annual and peren- nial)	Elymus sp. and Lolium perenne.
Sorghums	Sorghum vulgare vars, fete- rita, shallu, kaoliang, broomcorn.
Sudan grass	Sorghum vulgare sudanense.
Wheat	Triticum aestivum.

or any mixture of such forage crops, for use only as an animal feed.

(b) Such additive is used only as a chemical preservative for the purpose of retarding oxidative destruction of naturally occurring carotenes and vitamin E in the forage crops.

(c) It is added to the dehydrated forage crops in an oil mixture containing only suitable animal or suitable vegetable oil, prior to grinding and mixing.

(d) The maximum quantity of the additive permitted to be used and to remain in or on the dehydrated forage crop shall not exceed 150 parts per million.

(e) To assure the safe use of the additive, the label of the market package shall contain, in addition to other information required by the act:

(1) The name of the additive as specified in this section.

(2) Directions for the incorporation of the additive in the forage crops, as specified in paragraph (c) of this section, with the directive that only suitable animal or suitable vegetable oils are to be used in the oil mix.

(f) The label of any dehydrated forage crops treated with the additive or the label of an animal-feed supplement containing such treated forage crops, shall, in addition to other information required by the act, bear the following statements:

(1) "Ethoxyquin, a preservative," or "Ethoxyquin added to retard the oxidative destruction of carotene and vitamin E."

(2) The statement "For use in animal feed only."

§ 573.420 Ethyl cellulose.

The food additive ethyl cellulose may be safely used in animal feed in accordance with the following prescribed conditions:

(a) The food additive is a cellulose ether containing ethoxy (OC₂H₅) groups attached by an ether linkage and containing on an anhydrous basis not more

than 2.6 ethoxy groups per anhydroglucose unit.

(b) It is used or intended for use as a binder or filler in dry vitamin preparations to be incorporated into animal feed.

§ 573.440 Ethylene dichloride.

The food additive ethylene dichloride may be safely used in the manufacture of animal feeds in accordance with the following prescribed conditions:

(a) It is used as a solvent in the extraction processing of animal byproducts for use in animal feeds.

(b) The maximum quantity of the additive permitted to remain in or on the extracted byproducts shall not exceed 300 parts per million.

(c) The extracted animal byproduct is added as a source of protein to a total ration at levels consistent with good feeding practices, but in no event at levels exceeding 13 percent of the total ration.

§ 573.450 Fermented ammoniated condensed whey.

(a) *Identity.* The product is produced by the *Lactobacillus bulgaricus* fermentation of whey with the addition of ammonia.

(b) *Specifications.* The product contains 35 to 55 percent crude protein and not more than 42 percent equivalent crude protein from nonprotein nitrogen sources.

(c) *Uses.* The product is used as a source of protein and nonprotein nitrogen for cattle.

(d) *Limitations.* (1) Store in a closed vented tank equipped for agitation. Agitate 5 minutes before using. Do not store at temperature above 110° F (43° C).

(2) The maximum level of use of fermented ammoniated condensed whey and equivalent crude protein from all other added forms of nonprotein nitrogen shall not exceed 30 percent of the dietary crude protein.

(3) The additive may be used as follows:

(i) Mixed with grain, roughage, or grain and roughage prior to feeding.

(ii) As a component of free-choice liquid feeds, used to supplement the diets of cattle fed other sources of nutrients, fermented ammoniated condensed

whey shall not exceed 80 percent of the free-choice liquid feed.

(e) *Labeling*. The label shall bear, in addition to other information required by the act:

- (1) The name of the additive.
- (2) The maximum percentage of equivalent crude protein from non-protein nitrogen.

(3) Adequate directions for use in accordance with the provisions in paragraph (d) of this section.

[43 FR 33708, July 1, 1978, as amended at 46 FR 49115, Oct. 6, 1981]

§ 573.460 Formaldehyde.

The food additive formaldehyde may be safely used in the manufacture of animal feeds in accordance with the following conditions:

(a)(1) The additive is used, or intended for use, to improve the handling characteristics of animal fat in combination with certain oilseed meals by producing therefrom a dry, free-flowing product as follows:

(i) An aqueous blend of soybean and sunflower meals in a ratio of 3:1, respectively, is mixed with animal fat such that the oilseed meals and animal fat are in a ratio of 3:2. The feed ingredients are those defined by the "Official Publication" of the Association of American Feed Control Officials, Inc., 1976 ed., pages 86, 103, and 109.¹

(ii) Formaldehyde (37 percent solution) is added to the mixture at a level of 4 percent of the dry matter weight of the oilseed meals and animal fat. This mixture, upon drying, contains not more than 1 percent formaldehyde and not more than 12 percent moisture.

(2) The dried mixture described in paragraph (a) of this section is used, or intended for use, as a component of dry, nonpelleted feeds for beef and non-lactating dairy cattle.

(3) To assure safe use of the additive, in addition to the other information required by the Act, the label and labeling of the dried mixture described in paragraph (a) of this section shall bear:

- (i) The name of the additive.

(ii) Adequate directions for use providing that feed as consumed is not to contain more than 25 percent of the mixture.

(b)(1) The food additive is formaldehyde (CAS No. 50-00-0; 37 percent aqueous solution). It is used at a rate of 5.4 pounds (2.5 kilograms) per ton of animal feed or feed ingredient. It is an antimicrobial agent used to maintain complete animal feeds or feed ingredients *Salmonella* negative for up to 21 days.

(2) To assure safe use of the additive, in addition to the other information required by the Act, the label and labeling shall contain:

- (i) The name of the additive.
- (ii) A statement that formaldehyde solution which has been stored below 40 °F or allowed to freeze should not be applied to complete animal feeds or feed ingredients.

(iii) Adequate directions for use including a statement that formaldehyde should be uniformly sprayed on and thoroughly mixed into the complete animal feeds or feed ingredients and that the complete animal feeds or feed ingredients so treated shall be labeled as containing formaldehyde. The label must prominently display the statement: "Treated with formaldehyde to maintain feed *Salmonella* negative. Use within 21 days."

(iv) The labeling for feed or feed ingredients to which formaldehyde has been added under the provisions of paragraph (b)(1) of this section is required to carry the following statement: "Treated with formaldehyde to maintain feed *Salmonella* negative. Use within 21 days."

(3) To assure safe use of the additive, in addition to the other information required by the Act, the label and labeling shall contain:

- (i) Appropriate warnings and safety precautions concerning formaldehyde.
- (ii) Statements identifying formaldehyde as a poison with potentials for adverse respiratory effects.

(iii) Information about emergency aid in case of accidental inhalation.

(iv) Statements reflecting requirements of applicable sections of the

¹Copies of the current edition may be obtained from the Treasurer, Donald H. James, West Virginia Department of Agriculture, Capital Complex/Guthrie Center, Charleston, WV 25305.

Food and Drug Administration, HHS

§ 573.540

Superfund Amendments and Reauthorization Act (SARA), and the Occupational Safety and Health Administration's (OSHA) human safety guidance regulations.

(v) Contact address and phone number for reporting adverse reactions or to request a copy of the Materials Safety Data Sheet (MSDS).

[41 FR 38652, Sept. 10, 1976, as amended at 54 FR 18281, Apr. 28, 1989; 61 FR 15704, Apr. 9, 1996; 63 FR 53580, Oct. 6, 1998]

§ 573.480 Formic acid.

Formic acid may be safely used as a preservative in hay crop silage in an amount not to exceed 2.25 percent of the silage on a dry weight basis or 0.45 percent when direct-cut. The top foot of silage stored should not contain formic acid and silage should not be fed to livestock within 4 weeks of treatment.

§ 573.500 Condensed, extracted glutamic acid fermentation product.

Condensed, extracted glutamic acid fermentation product may be safely used in animal feed under the following conditions:

(a) The additive is a concentrated mixture of the liquor remaining from the extraction of glutamic acid, combined with the cells of *Corynebacterium lilium* used to produce the glutamic acid.

(b) It is used or intended for use as follows:

(1) In poultry feed as a source of protein in an amount not to exceed 5 percent of the total ration.

(2) In cattle feed as a source of protein in an amount not to exceed 10 percent of the feed.

(c) In order to assure safe use, the label and labeling of the additive shall bear, in addition to the other information required by the Act, the following:

(1) The name of the additive.

(2) A statement of the concentration of the additive contained in any mixture.

(3) Adequate directions for use.

§ 573.520 Hemicellulose extract.

Hemicellulose extract may be safely used in animal feed when incorporated therein in accordance with the following conditions:

(a) The additive is produced from the aqueous extract obtained by the treatment of wood with water at elevated temperatures (325 degrees–535 degrees F) and pressure (80 to 900 pounds per square inch) and contains primarily pentose and hexose sugars.

(b) The additive may be used in a liquid or dry state with the liquid product containing not less than 55 percent carbohydrate and the dry product containing not less than 84 percent carbohydrate.

(c) The additive is used as a source of metabolizable energy in animal feed in accordance with good manufacturing and feeding practices.

[41 FR 38652, Sept. 10, 1976, as amended at 43 FR 11181, Mar. 17, 1978]

§ 573.530 Hydrogenated corn syrup.

(a) *Identity*. The product is produced by hydrogenation of corn syrup over a nickel catalyst.

(b) *Specifications*. The product contains 70 percent hydrogenated corn syrup and a maximum of 0.5 percent reducing sugars.

(c) *Uses*. The product is used as a humectant and plasticizer in preparation of soft-moist dog and cat foods.

(d) *Limitations*. The product is preferably stored in a closed, stainless steel or aluminum container. The level of use of the product shall not exceed 15 percent of the total weight of the pet food formulation.

(e) *Labeling*. The labeling shall bear, in addition to other information required by the Act:

(1) The name of the additive.

(2) Adequate directions for use in accordance with the provisions in paragraph (d) of this section.

[45 FR 22920, Apr. 4, 1980]

§ 573.540 Hydrolyzed leather meal.

(a) *Identity*. Hydrolyzed leather meal is produced from leather scraps that are treated with steam for not less than 33 minutes at a pressure of not less than 125 pounds per square inch.

(b) *Specifications*. The additive shall conform to the following percent-by-weight specifications:

Moisture, not less than 5 percent nor more than 10 percent.

Crude protein, not less than 60 percent.

§ 573.560

Crude fat, not less than 5 percent.
Crude fiber, not more than 6 percent.
Chromium, not more than 2.75 percent.

(c) *Use.* It is used or intended for use as a source of protein in swine feeds in an amount not to exceed 1.0 percent by weight of the finished feed.

(d) *Labeling.* The labels and labeling shall bear, in addition to the other information required by the Act:

(1) The name of the additive, hydrolyzed leather meal.

(2) Adequate directions to provide finished feeds complying with paragraph (c) of this section.

§ 573.560 Iron ammonium citrate.

Iron ammonium citrate may be safely used in animal feed in accordance with the following prescribed conditions:

(a) The additive is the chemical green ferric ammonium citrate.

(b) The additive is used or intended for use as an anticaking agent in salt for animal consumption so that the level of iron ammonium citrate does not exceed 25 parts per million (0.0025 percent) in the finished salt.

(c) To assure safe use of the additive the label or labeling of the additive shall bear, in addition to the other information required by the Act:

(1) The name of the additive.

(2) Adequate directions to provide a final product that complies with the limitations prescribed in paragraph (b) of this section.

§ 573.580 Iron-choline citrate complex.

Iron-choline citrate complex made by reacting approximately equimolecular quantities of ferric hydroxide, choline, and citric acid may be safely used as a source of iron in animal feed.

§ 573.600 Lignin sulfonates.

Lignin sulfonates may be safely used in animal feeds in accordance with the following prescribed conditions:

(a) For the purpose of this section, the food additive is either one, or a combination of, the ammonium, calcium, magnesium, or sodium salts of the extract of spent sulfite liquor derived from the sulfite digestion of wood or of abaca (*Musa textilis*) or of sisal (*Agave sisalana*) in either a liquid form (moisture not to exceed 50 percent by

21 CFR Ch. I (4–1–99 Edition)

weight) or dry form (moisture not to exceed 6 percent by weight).

(b) It is used or intended for use in an amount calculated on a dry weight basis, as follows:

(1) As a pelleting aid in the liquid or dry form in an amount not to exceed 4 percent of the finished pellets.

(2) As a binding aid in the liquid form in the flaking of feed grains in an amount not to exceed 4 percent of the flaked grain.

(3) As a surfactant in molasses used in feeds, as liquid lignin sulfonate, in an amount not to exceed 11 percent of the molasses.

(4) As a source of metabolizable energy, in the liquid or dry form, in an amount not to exceed 4 percent of the finished feed.

§ 573.620 Menadione dimethylpyrimidinol bisulfite.

The food additive, menadione dimethylpyrimidinol bisulfite, may be safely used in accordance with the following conditions:

(a) The additive is the 2-hydroxy-4,6-dimethylpyrimidinol salt of menadione ($C_{17}H_{18}O_6N_2S$).

(b) The additive is used or intended for use as a nutritional supplement for the prevention of vitamin K deficiency as follows:

(1) In chicken and turkey feed at a level not to exceed 2 grams per ton of complete feed.

(2) In the feed of growing and finishing swine at a level not to exceed 10 grams per ton of feed.

(c) To assure safe use, the label and labeling of the additive shall bear adequate directions for use.

§ 573.625 Menadione nicotinamide bisulfite.

The food additive may be safely used as follows:

(a) *Product.* The additive is 1,2,3,4-tetrahydro-2-methyl-1,4-dioxo-2-naphthalene sulfonic acid with 3-pyridine carboxylic acid amine (CAS No. 73581-79-0).

(b) *Conditions of use.* As a nutritional supplement in chicken and turkey feeds for both the prevention of vitamin K deficiency and as a source of supplemental niacin.

(c) *Limitations.* Not to exceed 2 grams per ton of complete feed. To assure safe use, the label and labeling shall bear adequate directions for use.

[61 FR 5, Jan. 2, 1996]

§ 573.640 Methyl esters of higher fatty acids.

The food additive methyl esters of higher fatty acids may be safely used in animal feeds in accordance with the following prescribed conditions:

(a) The food additive is manufactured by reaction of methyl alcohol with feed-grade fats or oils and consists of not less than 70 percent methyl esters of the following straight-chain monocarboxylic acids: Docosahexanoic acid, eicosapentanoic acid, linoleic acid, myristic acid, oleic acid, palmitic acid, palmitoleic acid, and stearic acid, and lesser amounts of the associated acid esters.

(b) The food additive meets the following specifications:

(1) Free methyl alcohol not to exceed 150 parts per million.

(2) Unsaponifiable matter not to exceed 2 percent.

(3) It is free of chick-edema factor or other factors toxic to chicks, as evidenced during the bioassay method for determining the chick-edema factor as prescribed in paragraph (b)(4)(ii) of this section.

(4) For the purposes of this section:

(i) Unsaponifiable matter shall be determined by the method described in Section 28.081, "Unsaponifiable Residue (20)—Official Final Action" of the "Official Methods of Analysis of the Association of Official Analytical Chemists," 13th Ed., 1980, p. 451, which is incorporated by reference. Copies are available from the Association of Official Analytical Chemists, 2200 Wilson Blvd., Suite 400, Arlington, VA 22201-3301, or available for inspection at the Office of the Federal Register, 800 North Capitol Street, NW., suite 700, Washington, DC 20001.

(ii) The chick-edema factor bioassay method described under "26. Oils, Fats, and Waxes" in the *Journal of the Association of Official Agricultural Chemists*, Vol. 44, Page 146 (1961), or the method described under "Chick-Edema Factor—Bioassay Method (34)—Official Final Action" in §§ 28.113–28.117, "Offi-

cial Methods of Analysis of the Association of Official Analytical Chemists," 12th Ed., 1975, pp. 509–511, which is incorporated by reference, shall be employed. (Copies of the methods are available from the Association of Official Analytical Chemists, 2200 Wilson Blvd., Suite 400, Arlington, VA 22201-3301, or available for inspection at the Office of the Federal Register, 800 North Capitol Street, NW., suite 700, Washington, DC 20001.) The presence of chick-edema factor shall be determined by a comparison between the mean log of the pericardial fluid volumes of a test group and of a concurrent negative control group. The significance of the difference in pericardial fluid volumes between the test group and the negative control group is determined by calculating a "t" value according to the formula:

$$t = \frac{\bar{x}_t - \bar{x}_c}{\sqrt{(s_t^2/n_t) + (s_c^2/n_c)}}$$

where:

$\bar{x}_t$ and $\bar{x}_c$ are the means of the logs of the pericardial fluid volumes of the test and control groups, respectively;

n_t and n_c are the number of chicks in the respective groups;

s_t^2 and s_c^2 are the variances of the test and control groups, respectively.

The variances are calculated as follows:

$$s^2 = \frac{n(\sum x^2) - (\sum x)^2}{n(n-1)}$$

where:

$\sum x$ is the sum of the logs of the pericardial fluid volumes;

$\sum x^2$ is the sum of the squares of the logs of the pericardial fluid volumes for either the test t or control c group data.

The test sample is judged to contain chick-edema factor if the calculated "t" exceeds +1.3 and the mean log of the pericardial fluid volume obtained from the negative control group multiplied by 100 is less than 1.1461.

(iii) "Other factors toxic to chicks" referred to in paragraph (b)(3) of this section shall be determined during the course of the bioassay test described in paragraph (b)(4)(ii) of this section, on the basis of chick deaths or other abnormalities not attributable to chick-

§ 573.660

edema factor or to the experimental conditions of the test.

(c) It is used or intended for use as a supplementary source of fat for animal feed.

(d) To assure safe use of the additive, in addition to the other information required by the act:

(1) The label and labeling of the additive, and any feed additive supplement, feed additive concentrate, feed additive premix, or complete feed prepared therefrom shall bear:

(i) The name of the additive.

(ii) The designation "feed grade" in juxtaposition with the name and equally as prominent.

(2) The label or labeling of the additive and any feed additive supplement, feed additive concentrate, feed additive premix, or complete feed prepared therefrom shall bear adequate directions for use.

[41 FR 38652, Sept. 10, 1976, as amended at 47 FR 9397, Mar. 5, 1982; 54 FR 18281, Apr. 28, 1989]

§ 573.660 Methyl glucoside-coconut oil ester.

Methyl glucoside-coconut oil ester may be safely used in accordance with the following conditions:

(a) The additive meets the specifications prescribed in § 172.816 of this chapter.

(b) It is used as a surfactant in molasses intended for use in animal feed at a level not to exceed 320 parts per million.

§ 573.680 Mineral oil.

Mineral oil may be safely used in animal feed, subject to the provisions of this section.

(a) Mineral oil, for the purpose of this section, is that complying with the definition and specifications contained in § 172.878 (a) and (b) or in § 178.3620(b)(1) (i) and (ii) of this chapter.

(b) It is used in animal feeds for the following purposes:

(1) To reduce dustiness of feeds or mineral supplements.

(2) To serve as a lubricant in the preparation of pellets, cubes, or blocks and to improve resistance to moisture of such pellets, cubes, or blocks.

(3) To prevent the segregation of trace minerals in mineralized salt.

21 CFR Ch. I (4-1-99 Edition)

(4) To serve as a diluent carrier in the manufacture of feed grade biuret in accordance with good manufacturing practice.

(5) For the removal of water from substances intended as ingredients of animal feed.

(c) The quantity of mineral oil used in animal feed shall not exceed 3.0 percent in mineral supplements, nor shall it exceed 0.06 percent of the total ration when present in feed or feed concentrates.

[41 FR 38652, Sept. 10, 1976, as amended at 47 FR 41106, Sept. 17, 1982]

§ 573.700 Sodium nitrite.

Sodium nitrite may be safely used in canned pet food containing meat and fish in accordance with the following prescribed conditions:

(a) It is used or intended for use alone as a preservative and color fixative in canned pet food containing fish, meat, and fish and meat byproducts so that the level of sodium nitrite does not exceed 20 parts per million.

(b) To assure safe use of the additive, in addition to the other information required by the act:

(1) The label of the additive shall bear:

(i) The name of the additive.

(ii) A statement of the concentration of the additive in any mixture.

(2) The label or labeling shall bear adequate directions to provide a final product that complies with the limitations prescribed in paragraph (a) of this section.

§ 573.720 Petrolatum.

Petrolatum may be safely used in or on animal feed, subject to the following prescribed conditions:

(a) Petrolatum complies with the specifications set forth in the U.S. Pharmacopeia XVI for white petrolatum or in The National Formulary XII for yellow petrolatum.

(b) Petrolatum meets the following ultraviolet absorbance limits when subjected to the analytical procedure described in § 172.886(b) of this chapter.

Ultraviolet absorbance per centimeter path length:

Food and Drug Administration, HHS

§ 573.780

Millimicrons	Maximum
280 to 289	0.25
290 to 299	.20
300 to 359	.14
360 to 400	.04

(c) It is used in animal feed for the following purposes:

(1) To reduce dustiness of feeds or mineral supplements.

(2) To serve as a lubricant in the preparation of pellets, cubes, or blocks, and to improve resistance to moisture of such pellets, cubes, or blocks.

(d) The quantity of petrolatum present in animal feeds from the uses specified in paragraph (c) of this section shall not exceed 3 percent in mineral supplements nor shall it exceed 0.06 percent of the total ration when present in feed or feed concentrates.

(e) When used in combination with technical white mineral oil for the uses described in paragraph (c) of this section, the total quantity of combined petrolatum and technical white mineral oil shall not exceed the limits prescribed in paragraph (d) of this section.

(f) Petrolatum may contain any antioxidant permitted in food by regulations issued in accordance with section 409 of the act, in an amount not greater than that required to produce its intended effect.

§ 573.740 Odorless light petroleum hydrocarbons.

Odorless light petroleum hydrocarbons complying with §172.884(a) and (b) of this chapter may be safely used in an amount not in excess of that required as a component of insecticide formulations used in compliance with regulations issued in this part.

§ 573.750 *Pichia pastoris* dried yeast.

(a) *Identity*. The food additive *Pichia pastoris* dried yeast may be used in feed formulations of broiler chickens as a source of protein not to exceed 10 percent by weight of the total formulation.

(b) *Specifications*. The additive shall conform to the following percent-by-weight specifications:

(1) Crude protein, not less than 60 percent.

(2) Crude fat, not less than 2 percent.

(3) Crude fiber, not more than 2 percent.

(4) Ash, not more than 13 percent.

(5) Moisture, not more than 6 percent.

(c) *Use*. To ensure safe use, the labeling of the additive and any feed additive supplement, concentrate, or premix prepared therefrom shall bear, in addition to other required information, the name of the additive, directions for use to provide not more than 10 percent by weight of the total ration, and the statement "Caution: Not to be used in layers or other poultry intended for breeding."

[58 FR 59170, Nov. 8, 1993]

§ 573.760 Poloxalene.

The food additive poloxalene may be safely used in accordance with the following prescribed conditions:


(a) The additive consists of polyoxypropylene-polyoxyethylene glycol non-ionic block polymer meeting the following specifications:

(1) Molecular weight range: 2,850–3,150.

(2) Hydroxyl number: 35.7–39.4.

(3) Cloud point (10 percent solution): 42° C–46° C.

(4) Structural formula:



(b) In feed as a surfactant for the flaking of feed grains when added to liquid grain conditioner in an amount not to exceed 1.0 percent of the conditioner. The conditioner is added to the feed at a rate of 1 quart per ton of feed.

(c) The label and labeling shall bear, in addition to the other information required by the Act:

(1) The name of the additive.

(2) Adequate directions and warnings for use.

§ 573.780 Polyethylene.

(a) *Identity*. Polyethylene consists of basic polymers manufactured by the catalytic polymerization of ethylene.

(b) *Specifications*. (1) For the purposes of this section, polyethylene shall meet the specifications in item 2.1 of §177.1520(c) of this chapter.

§ 573.800

(2) The polyethylene is designed in a pellet form in a configuration presenting maximum angular surface having the following dimensions in centimeters:

$$0.9 \pm 0.1 \times 0.8 \pm 0.1 \times 1.2 \pm 0.1$$

(c) *Use.* It is used as a replacement for roughage in feedlot rations for finishing slaughter cattle.

(d) *Labeling.* The labels and labeling shall bear in addition to the other information required by the Act:

(1) The name of the additive “polyethylene roughage replacement.”

(2) Adequate directions for use which shall provide for the administration of one-half pound of polyethylene pellets per head per day for 6 successive days. All natural roughage should be removed for a minimum of 12 hours prior to administration of polyethylene roughage replacement. Roughage replacement must be adequately mixed in the ration for uniform distribution.

[41 FR 38652, Sept. 10, 1976, as amended at 54 FR 18282, Apr. 28, 1989]

§ 573.800 Polyethylene glycol (400) mono- and dioleate.

(a) The food additive polyethylene glycol (400) mono- and dioleate meets the following specifications: Saponification number, 80–88; acid number, 5.0 maximum; and average molecular weight range, 640–680.

(b) It is used as a processing aid in the production of animal feeds when present as a result of its addition to molasses in an amount not to exceed 250 parts per million of the molasses.

§ 573.820 Polyoxyethylene glycol (400) mono- and dioleates.

The food additive polyoxyethylene glycol (400) mono- and dioleates may be safely used as an emulsifier in calf-milk replacer formulations.

§ 573.840 Polysorbate 60.

The food additive polysorbate 60 (polyoxyethylene (20) sorbitan monostearate) may be safely used in animal feeds in accordance with the following prescribed conditions:

(a) It is used alone or in combination with sorbitan monostearate as an

21 CFR Ch. I (4–1–99 Edition)

emulsifier in mineral premixes and dietary supplements for animal feeds.

(b) It is used as an emulsifier in milk-replacer formulations for calves.

§ 573.860 Polysorbate 80.

The food additive polysorbate 80 (polyoxyethylene (20) sorbitan monooleate) may be safely used as an emulsifier in milk-replacer formulations for calves.

§ 573.870 Poly(2-vinylpyridine-co-styrene).

The food additive poly(2-vinylpyridine-co-styrene) may be safely used as nutrient protectant in feed for beef cattle and dairy cattle and replacement dairy heifers when used in accordance with the following conditions:

(a) The additive meets the following specifications:

Component/property	Limitation
Inherent viscosity	1.0–1.6 deciliter per gram. ¹
Styrene moiety	40 percent maximum.
2-Vinylpyridine moiety.	90 percent maximum.
Residual styrene	200 parts per billion maximum.
Residual 2-vinylstyrene	200 parts per billion maximum.
Heavy metals such as lead	>10 parts per million.
Arsenic	>3 parts per million.

¹ Inherent viscosity of a 0.25 percent (weight/volume) solution in dimethylformamide.

(b) The additive is used in the manufacture of rumen-stable, abomasum-dispersible nutrient(s) for beef cattle and dairy cattle and replacement dairy heifers such that the maximum use of the additive from all sources does not exceed 5.1 grams per head per day. The additive may be used to protect the following nutrients:

(1) *Methionine.* The resulting product must contain a maximum of 10 percent poly(2-vinylpyridine-co-styrene) by weight and a minimum of 55 percent methionine by weight. The coated methionine must be established through in vitro tests to be at least 90 percent rumen-stable, of which at least 90 percent is subsequently dispersible under abomasal conditions.

(2) *Methionine and lysine.* The resulting product must contain a maximum of 10 percent poly(2-vinylpyridine-co-styrene) by weight and a minimum of a combined total of 55 percent methionine and lysine by weight. The coated

Food and Drug Administration, HHS

§ 573.920

methionine and lysine must be established through in vitro tests to be at least 90 percent rumen-stable, of which at least 90 percent is subsequently dispersible under abomasal conditions.

(c) *Label and labeling.* To ensure safe use of the additive, the label and labeling of the additive and of any feed additive supplement, feed additive concentrate, feed additive premix, or liquid feed supplement prepared therefrom, shall bear, in addition to the other information required by the Federal Food, Drug, and Cosmetic Act, the following:

(1) The name of the additive.

(2) A statement of the concentration of poly(2-vinylpyridine-co-styrene) in any product or mixture.

(3) Adequate directions for the use of the rumen-stable, abomasum-dispersible nutrient(s) products.

(4) The following statement: "Warning: Maximum use of poly(2-vinylpyridine-co-styrene) from all sources is not to exceed 5.1 grams per head per day."

[57 FR 7875, Mar. 5, 1992, as amended at 57 FR 24187, June 8, 1992; 61 FR 11547, Mar. 21, 1996]

§ 573.880 Normal propyl alcohol.

Normal propyl alcohol may be safely used in feeds and feed supplements for cattle as a source of metabolizable energy. It is incorporated in the feed or feed supplement in an amount which provides not more than 54.5 grams of the additive per head per day.

§ 573.900 Pyrophyllite.

Pyrophyllite (aluminum silicate monohydrate) may be safely used as the sole anticaking aid, blending agent, pelleting aid, or carrier in animal feed when incorporated therein in an amount not to exceed 2 percent in complete animal feed.

§ 573.914 Salts of volatile fatty acids.

(a) *Identity.* The food additive is a blend containing the ammonium or calcium salt of isobutyric acid and the ammonium or calcium salts of a mixture of 5-carbon acids—isovaleric, 2-methylbutyric, and *n*-valeric.

(b) *Specifications.* The additive contains ammonium or calcium salts of volatile fatty acids and shall conform to the following specifications:

(1) Ammonium salts:

Components	Amount
Ammonium salts of mixed 5-carbon acids (as identified in paragraph (a) of this section).	48 to 54 percent.
Ammonium salt of isobutyric acid	22 to 26 percent.
Water	28 percent maximum.
Ammonia	0.3 percent maximum.
Arsenic	>3 parts per million.
Heavy metals as lead	>10 parts per million.

(2) Calcium salts:

Components	Amount
Calcium salts of mixed 5-carbon acids (as identified in paragraph (a) of this section).	58 to 72 percent.
Calcium salt of isobutyric acid	26 to 34 percent.
Calcium hydroxide	3 percent maximum.
Water	14 percent maximum.
Arsenic	>3 parts per million.
Heavy metals as lead	>10 parts per million.

(c) *Use.* The additive is used or intended for use as a source of energy in dairy cattle feed.

(d) *Labeling.* The label and labeling of the additive in any feed, feed supplement, feed concentrate, feed premix, or liquid feed supplement prepared therefrom shall bear, in addition to other information required by the act, the following:

(1) The name of the additive.

(2) Adequate directions for use, including statements expressing maximum use levels. For ammonium salts of volatile fatty acids, the statements: "Not to exceed 160 grams per head per day thoroughly mixed in dairy cattle feed as a source of energy." For calcium salts of volatile fatty acids, the statement: "Not to exceed 135 grams per head per day thoroughly mixed in dairy cattle feed as a source of energy."

[49 FR 45125, Nov. 15, 1984; 50 FR 8606, Mar. 4, 1985]

§ 573.920 Selenium.

(a) Public Law 103-354 enacted October 13, 1994 (the 1994 Act), states that FDA shall not implement or enforce the final rule issued on September 13, 1993 (58 FR 47962), in which FDA stayed the 1987 amendments and any modification of such rule issued after enactment of the 1994 Act; unless the Commissioner of Food and Drugs makes a determination that:

(1) Selenium additives are not essential at levels authorized in the absence of such final rule, to maintain animal nutrition and protect animal health;

(2) selenium at such levels is not safe to the animals consuming the additive;

(3) selenium at such levels is not safe to individuals consuming edible portions of animals that receive the additive;

(4) selenium at such levels does not achieve its intended effect of promoting normal growth and reproduction of livestock and poultry; and

(5) the manufacture and use of selenium at such levels cannot reasonably be controlled by adherence to current good manufacturing practice requirements.

(6) Paragraphs (b) through (g) of this section provide the currently acceptable levels of selenium supplementation.

(b) The food additive selenium is a nutrient administered in animal feed as sodium selenite or sodium selenate or in a controlled-release sodium selenite bolus, as provided in paragraph (f) of this section.

(c) It is added to feed as follows:

(1) In complete feed for chickens, swine, turkeys, sheep, cattle, and ducks at a level not to exceed 0.3 part per million.

(2) In feed supplements for limit feeding as follows:

(i) *Sheep*: At a level not to exceed an intake of 0.7 milligram per head per day.

(ii) *Beef cattle*: At a level not to exceed an intake of 3 milligrams per head per day.

(3) In salt-mineral mixtures for free-choice feeding as follows:

(i) *Sheep*: Up to 90 parts per million in a mixture for free-choice feeding at a rate not to exceed an intake of 0.7 milligram per head per day.

(ii) *Beef cattle*: Up to 120 parts per million in a mixture for free-choice feeding at a rate not to exceed an intake of 3 milligrams per head per day.

(d) The additive shall be incorporated into feed as follows:

(1) It shall be incorporated into each ton of complete feed by adding no less than 1 pound of a premix containing no more than 272.4 milligrams of added selenium per pound.

(2) It shall be incorporated into each ton of salt-mineral mixture for sheep or beef cattle from a premix containing no more than 4.5 grams of added selenium per pound.

(e) The premix manufacturer shall follow good manufacturing practices in the production of selenium premixes. Inventory, production, and distribution records must provide a complete and accurate history of product production. Production controls must assure products to be what they are purported and labeled. Production controls shall include analysis sufficient to adequately monitor quality.

(f) The label or labeling of any selenium premix shall bear adequate directions and cautions for use including this statement: "Caution: Follow label directions. The addition to feed of higher levels of this premix containing selenium is not permitted."

(g) The additive is orally administered to beef and dairy cattle as an osmotically controlled, constant release bolus containing sodium selenite. Each bolus contains 360 milligrams of selenium as sodium selenite, and delivers 3 milligrams of selenium per day for 120 days. To ensure safe use of the additive:

(1) The osmotically controlled, constant release bolus is for use only in beef and dairy cattle more than 3 months of age or over 200 pounds body weight.

(2) Only one bolus containing 360 milligrams of selenium as sodium selenite is administered orally to each animal in 120 days.

(3) The labeling shall bear the following: "This bolus delivers the maximum daily allowable amount of selenium and shall be the sole source of supplementation. Do not use in areas containing excess selenium. Do not rebolus within 4 months."

[52 FR 10888, Apr. 6, 1987; 52 FR 21001, June 4, 1987, as amended at 54 FR 14215, Apr. 10, 1989; 54 FR 15874, Apr. 19, 1989; 60 FR 53703, Oct. 17, 1995]

EFFECTIVE DATE NOTE: At 58 FR 47973, Sept. 13, 1993, the amendments to § 573.920 that were published at 52 FR 10887, Apr. 6, 1987; 52 FR 21001, June 4, 1987; and 54 FR 14214, Apr. 10, 1989 were stayed until further notice. At 59 FR 45973, Sept. 6, 1994, the stay was confirmed.

§ 573.940 Silicon dioxide.

The food additive silicon dioxide may be safely used in animal feed in accordance with the following conditions:

(a) The food additive is manufactured by vapor phase hydrolysis or by other means whereby the particle size is such as to accomplish the intended effect.

(b) It is used or intended for use in feed components as an anticaking agent, and/or grinding aid, as follows:

Feed component	Limitations (percent)
BHT (butylated hydroxytoluene)	2
Methionine hydroxy analog and its calcium salts	1
Piperazine, piperazine salts	0.8
Sodium propionate	1
Urea	1
Vitamins	3

(c) It is used in feed as an anticaking agent in an amount not to exceed that reasonably required to accomplish its intended effect and in no case in an amount to exceed 2 percent by weight of the finished feed.

§ 573.960 Sorbitan monostearate.

The food additive sorbitan monostearate may be safely used alone or in combination with polysorbate 60 as an emulsifier in mineral premixes and dietary supplements for animal feeds.

§ 573.980 Taurine.

The food additive taurine (2-aminoethanesulfonic acid) may be safely used in feed in accordance with the following prescribed conditions:

(a) It is used as a nutritional supplement in the feed of growing chickens.

(b) It is added to complete feeds so that the total taurine content does not exceed 0.054 percent of the feed.

(c) To assure safe use of the additive, the label and labeling shall bear in addition to the other information required by the Act:

(1) The name of the additive.

(2) The quantity of the additive contained therein.

(3) Adequate directions for use.

§ 573.1000 Verxite.

The food additive verxite may be safely used in animal feed in accordance with the following prescribed conditions:

(a) The additive is a magnesium-aluminum-iron silicate conforming to one of the following:

(1)(i) Verxite granules: The additive contains a minimum of 98 percent of hydrobiotite; it is thermally expanded and has a bulk density of from 5 to 9 pounds per cubic foot.

(ii) It is used or intended for use:

(a) In poultry feed at a level not to exceed 5 percent of the weight of the finished feed as a nonnutritive bulking agent for restricting calorie intake in pullet replacement feeds.

(b) As an anticaking or blending agent, pelleting aid, or nonnutritive carrier for the incorporation of nutrients in poultry, swine, dog, or ruminant feeds, in an amount not to exceed that necessary to accomplish its intended effect and in no case to exceed 1.5 percent of the dog feed or 5 percent of the final feed for other animals.

(2)(i) Verxite flakes: The additive contains a minimum of 98 percent of hydrobiotite; it has a bulk density of from 20 to 30 pounds per cubic foot.

(ii) It is used or intended for use as an anticaking or blending agent in ruminant feeds in an amount not to exceed that necessary to accomplish its intended effect and in no case to exceed 1 percent by weight of the final feed for ruminants.

(3)(i) Verxite grits: The additive contains a minimum of 80 percent of hydrobiotite; it has a bulk density of from 40 to 50 pounds per cubic foot.

(ii) It is used or intended for use as a partial roughage replacement in ruminant feeds in an amount not to exceed that necessary to accomplish its intended effect and in no case to exceed 1 percent by weight of the final feed.

(b) To assure safe use of the additive, the label of any feed additive supplement, feed additive concentrate, feed additive premix, or complete feed prepared therefrom shall bear, in addition to the other information required by the Act, the name of the additive (verxite granules, verxite flakes, or verxite grits), adequate directions for use, and, when the additive is present in excess of 1 percent, a statement of the quantity of the additive contained therein and the term "nonnutritive" in juxtaposition therewith.

§ 573.1010

§ 573.1010 Xanthan gum.

The food additive xanthan gum may be safely used in animal feed as follows:

(a) The food additive is xanthan gum as defined in §172.695 of this chapter and meets all of the specifications thereof.

(b) It is used or intended for use as a stabilizer, emulsifier, thickener, suspending agent, or bodying agent in animal feed as follows:

(1) In calf milk replacers at a maximum use level of 0.1 percent, as fed.

(2) In liquid feed supplements for ruminant animals at a maximum use level of 0.25 percent (5 pounds per ton).

(c) To assure safe use of the additive:

(1) The label of its container shall bear, in addition to other information required by the act, the name of the additive.

(2) The label or labeling of the additive container shall bear adequate directions for use.

[49 FR 44630, Nov. 8, 1984]

§ 573.1020 Yellow prussiate of soda.

Yellow prussiate of soda (sodium ferrocyanide decahydrate: $\text{Na}_4\text{Fe}(\text{Cn})_6 \cdot 10\text{H}_2\text{O}$) may be safely used as an anticaking agent in salt for animal consumption at a level not to exceed 13 parts per million. The additive contains a minimum of 99.0 percent by weight of sodium ferrocyanide decahydrate.

[41 FR 38657, Sept. 10, 1976; 41 FR 48100, Nov. 2, 1976]

PART 579—IRRADIATION IN THE PRODUCTION, PROCESSING, AND HANDLING OF ANIMAL FEED AND PET FOOD

Subpart A—General Provisions

Sec.

579.12 Incorporation of regulations in part 179.

Subpart B—Radiation and Radiation Sources

579.22 Ionizing radiation for treatment of laboratory animal diets.

579.40 Ionizing radiation for the treatment of poultry feed and poultry feed ingredients.

21 CFR Ch. I (4–1–99 Edition)

AUTHORITY: 21 U.S.C. 321, 342, 343, 348, 371.

Subpart A—General Provisions

§ 579.12 Incorporation of regulations in part 179.

Regulations providing for irradiation in the production, processing, and handling of food in part 179 of this chapter are incorporated in subchapter E as applicable to use in the production, processing, handling, and labeling of animal feed and pet food, except where specifically provided for in this part.

[51 FR 5993, Feb. 19, 1986]

Subpart B—Radiation and Radiation Sources

§ 579.22 Ionizing radiation for treatment of laboratory animal diets.

Ionizing radiation for treatment of complete diets for laboratory animals (mice, rats, and hamsters) may be safely used under the following conditions:

(a) *Energy sources.* Ionizing radiation is limited to:

(1) Gamma rays for sealed units of the radionuclides cobalt-60 or cesium-137.

(2) Electrons generated from machine sources at energy levels not to exceed 10 million electron volts.

(b) *Uses.* The ionizing radiation is used or intended for use in single treatment as follows:

Food for irradiation	Limitations	Use
Bagged complete diets for laboratory animals (mice, rats, hamsters, rabbits, and guinea pigs).	Absorbed dose: Not to exceed 50 kiloGrays (5 megarads) Feeds treated by irradiation should be formulated to account for nutritional loss..	Microbial disinfection.

[51 FR 5993, Feb 19, 1986; 51 FR 8315, Mar. 11, 1986, as amended at 58 FR 18148, Apr. 8, 1993]

§ 579.40 Ionizing radiation for the treatment of poultry feed and poultry feed ingredients.

Ionizing radiation for the treatment of complete poultry diets and poultry feed ingredients may be safely used as follows:

(a) *Energy sources.* Ionizing radiation is limited to gamma rays from sealed units of cobalt-60.

(b) *Limitation.* The ionizing radiation is used for feed or feed ingredients that do not contain drugs.

(c) *Use.* Ionizing radiation is used as a single treatment for rendering complete poultry diets or poultry feed ingredients salmonella negative as follows:

(1) Minimum dose 2.0 kiloGrays (kGy) (0.2 megarad (Mrad)); maximum dose 25 kGy (2.5 megarads Mrad). The absorbed dose of irradiation is to be based on initial concentration of salmonella using the relationship that 1.0 kGy (0.1 Mrad) reduces salmonella concentration by one log cycle (one decimal reduction).

(2) Feeds treated by irradiation should be formulated to account for nutritional loss.

(3) If an irradiated feed ingredient is less than 5 percent of the final product, the final product can be irradiated without being considered to be re-irradiated.

[60 FR 50099, Sept. 28, 1995]

PART 582—SUBSTANCES GENERALLY RECOGNIZED AS SAFE

Subpart A—General Provisions

Sec.

582.1 Substances that are generally recognized as safe.

582.10 Spices and other natural seasonings and flavorings.

582.20 Essential oils, oleoresins (solvent-free), and natural extractives (including distillates).

582.30 Natural substances used in conjunction with spices and other natural seasonings and flavorings.

582.40 Natural extractives (solvent-free) used in conjunction with spices, seasonings, and flavorings.

582.50 Certain other spices, seasonings, essential oils, oleoresins, and natural extracts.

582.60 Synthetic flavoring substances and adjuvants.

582.80 Trace minerals added to animal feeds.

582.99 Adjuvants for pesticide chemicals.

Subpart B—General Purpose Food Additives

582.1005 Acetic acid.

582.1009 Adipic acid.

582.1033 Citric acid.

582.1057 Hydrochloric acid.

582.1061 Lactic acid.

582.1069 Malic acid.

582.1073 Phosphoric acid.

582.1077 Potassium acid tartrate.

582.1087 Sodium acid pyrophosphate.

582.1091 Succinic acid.

582.1095 Sulfuric acid.

582.1099 Tartaric acid.

582.1125 Aluminum sulfate.

582.1127 Aluminum ammonium sulfate.

582.1129 Aluminum potassium sulfate.

582.1131 Aluminum sodium sulfate.

582.1135 Ammonium bicarbonate.

582.1137 Ammonium carbonate.

582.1139 Ammonium hydroxide.

582.1141 Ammonium phosphate.

582.1143 Ammonium sulfate.

582.1155 Bentonite.

582.1165 Butane.

582.1191 Calcium carbonate.

582.1193 Calcium chloride.

582.1195 Calcium citrate.

582.1199 Calcium gluconate.

582.1205 Calcium hydroxide.

582.1207 Calcium lactate.

582.1210 Calcium oxide.

582.1217 Calcium phosphate.

582.1235 Caramel.

582.1240 Carbon dioxide.

582.1275 Dextrans.

582.1320 Glycerin.

582.1324 Glyceryl monostearate.

582.1355 Helium.

582.1366 Hydrogen peroxide.

582.1400 Lecithin.

582.1425 Magnesium carbonate.

582.1428 Magnesium hydroxide.

582.1431 Magnesium oxide.

582.1480 Methylcellulose.

582.1500 Monoammonium glutamate.

582.1516 Monopotassium glutamate.

582.1540 Nitrogen.

582.1585 Papain.

582.1613 Potassium bicarbonate.

582.1619 Potassium carbonate.

582.1625 Potassium citrate.

582.1631 Potassium hydroxide.

582.1643 Potassium sulfate.

582.1655 Propane.

582.1666 Propylene glycol.

582.1685 Rennet.

582.1711 Silica aerogel.

582.1721 Sodium acetate.

582.1736 Sodium bicarbonate.

582.1742 Sodium carbonate.

582.1745 Sodium carboxymethylcellulose.

582.1748 Sodium caseinate.

582.1751 Sodium citrate.

582.1763 Sodium hydroxide.

582.1775 Sodium pectinate.

582.1778 Sodium phosphate.

582.1781 Sodium aluminum phosphate.

582.1792 Sodium sesquicarbonate.

582.1804 Sodium potassium tartrate.

582.1810 Sodium tripolyphosphate.

582.1901 Triacetin.

Pt. 582

582.1973 Beeswax.
 582.1975 Bleached beeswax.
 582.1978 Carnauba wax.

Subpart C—Anticaking Agents

582.2122 Aluminum calcium silicate.
 582.2227 Calcium silicate.
 582.2437 Magnesium silicate.
 582.2727 Sodium aluminosilicate.
 582.2729 Hydrated sodium calcium aluminosilicate.
 582.2906 Tricalcium silicate.

Subpart D—Chemical Preservatives

582.3013 Ascorbic acid.
 582.3021 Benzoic acid.
 582.3041 Erythorbic acid.
 582.3081 Propionic acid.
 582.3089 Sorbic acid.
 582.3109 Thiodipropionic acid.
 582.3149 Ascorbyl palmitate.
 582.3169 Butylated hydroxyanisole.
 582.3173 Butylated hydroxytoluene.
 582.3189 Calcium ascorbate.
 582.3221 Calcium propionate.
 582.3225 Calcium sorbate.
 582.3280 Dilauryl thiodipropionate.
 582.3336 Gum guaiac.
 582.3490 Methylparaben.
 582.3616 Potassium bisulfite.
 582.3637 Potassium metabisulfite.
 582.3640 Potassium sorbate.
 582.3660 Propyl gallate.
 582.3670 Propylparaben.
 582.3731 Sodium ascorbate.
 582.3733 Sodium benzoate.
 582.3739 Sodium bisulfite.
 582.3766 Sodium metabisulfite.
 582.3784 Sodium propionate.
 582.3795 Sodium sorbate.
 582.3798 Sodium sulfite.
 582.3845 Stannous chloride.
 582.3862 Sulfur dioxide.
 582.3890 Tocopherols.

Subpart E—Emulsifying Agents

582.4101 Diacetyl tartaric acid esters of mono- and diglycerides of edible fats or oils, or edible fat-forming fatty acids.
 582.4505 Mono- and diglycerides of edible fats or oils, or edible fat-forming fatty acids.
 582.4521 Monosodium phosphate derivatives of mono- and diglycerides of edible fats or oils, or edible fat-forming fatty acids.
 582.4666 Propylene glycol.

Subpart F—Nutrients and/or Dietary Supplements

582.5013 Ascorbic acid.
 582.5017 Aspartic acid.
 582.5049 Aminoacetic acid.
 582.5065 Linoleic acid.
 582.5118 Alanine.
 582.5145 Arginine.

21 CFR Ch. I (4–1–99 Edition)

582.5159 Biotin.
 582.5191 Calcium carbonate.
 582.5195 Calcium citrate.
 582.5201 Calcium glycerophosphate.
 582.5210 Calcium oxide.
 582.5212 Calcium pantothenate.
 582.5217 Calcium phosphate.
 582.5223 Calcium pyrophosphate.
 582.5230 Calcium sulfate.
 582.5245 Carotene.
 582.5250 Choline bitartrate.
 582.5252 Choline chloride.
 582.5260 Copper gluconate.
 582.5271 Cysteine.
 582.5273 Cystine.
 582.5301 Ferric phosphate.
 582.5304 Ferric pyrophosphate.
 582.5306 Ferric sodium pyrophosphate.
 582.5308 Ferrous gluconate.
 582.5311 Ferrous lactate.
 582.5315 Ferrous sulfate.
 582.5361 Histidine.
 582.5370 Inositol.
 582.5375 Iron reduced.
 582.5381 Isoleucine.
 582.5406 Leucine.
 582.5411 Lysine.
 582.5431 Magnesium oxide.
 582.5434 Magnesium phosphate.
 582.5443 Magnesium sulfate.
 582.5446 Manganese chloride.
 582.5449 Manganese citrate.
 582.5452 Manganese gluconate.
 582.5455 Manganese glycerophosphate.
 582.5458 Manganese hypophosphite.
 582.5461 Manganese sulfate.
 582.5464 Manganous oxide.
 582.5470 Mannitol.
 582.5475 Methionine.
 582.5477 Methionine hydroxy analog and its calcium salts.
 582.5530 Niacin.
 582.5535 Niacinamide.
 582.5580 D-Pantothenyl alcohol.
 582.5590 Phenylalanine.
 582.5622 Potassium chloride.
 582.5628 Potassium glycerophosphate.
 582.5634 Potassium iodide.
 582.5650 Proline.
 582.5676 Pyridoxine hydrochloride.
 582.5695 Riboflavin.
 582.5697 Riboflavin-5-phosphate.
 582.5701 Serine.
 582.5772 Sodium pantothenate.
 582.5778 Sodium phosphate.
 582.5835 Sorbitol.
 582.5875 Thiamine hydrochloride.
 582.5878 Thiamine mononitrate.
 582.5881 Threonine.
 582.5890 Tocopherols.
 582.5892 α -Tocopherol acetate.
 582.5915 Tryptophane.
 582.5920 Tyrosine.
 582.5925 Valine.
 582.5930 Vitamin A.
 582.5933 Vitamin A acetate.
 582.5936 Vitamin A palmitate.

Food and Drug Administration, HHS

§ 582.1

582.5945	Vitamin B ₁₂ .
582.5950	Vitamin D ₂ .
582.5953	Vitamin D ₃ .
582.5985	Zinc chloride.
582.5988	Zinc gluconate.
582.5991	Zinc oxide.
582.5994	Zinc stearate.
582.5997	Zinc sulfate.

Subpart G—Sequestrants

582.6033	Citric acid.
582.6085	Sodium acid phosphate.
582.6099	Tartaric acid.
582.6185	Calcium acetate.
582.6193	Calcium chloride.
582.6195	Calcium citrate.
582.6197	Calcium diacetate.
582.6199	Calcium gluconate.
582.6203	Calcium hexametaphosphate.
582.6215	Monobasic calcium phosphate.
582.6219	Calcium phytate.
582.6285	Dipotassium phosphate.
582.6290	Disodium phosphate.
582.6386	Isopropyl citrate.
582.6511	Monoisopropyl citrate.
582.6625	Potassium citrate.
582.6751	Sodium citrate.
582.6754	Sodium diacetate.
582.6757	Sodium gluconate.
582.6760	Sodium hexametaphosphate.
582.6769	Sodium metaphosphate.
582.6778	Sodium phosphate.
582.6787	Sodium pyrophosphate.
582.6789	Tetra sodium pyrophosphate.
582.6801	Sodium tartrate.
582.6804	Sodium potassium tartrate.
582.6807	Sodium thiosulfate.
582.6810	Sodium tripolyphosphate.
582.6851	Stearyl citrate.

Subpart H—Stabilizers

582.7115	Agar-agar.
582.7133	Ammonium alginate.
582.7187	Calcium alginate.
582.7255	Chondrus extract.
582.7330	Gum arabic.
582.7333	Gum ghatti.
582.7339	Guar gum.
582.7343	Locust bean gum.
582.7349	Sterculia gum.
582.7351	Gum tragacanth.
582.7610	Potassium alginate.
582.7724	Sodium alginate.

AUTHORITY: 21 U.S.C. 321, 342, 348, 371.

SOURCE: 41 FR 38657, Sept. 10, 1976, unless otherwise noted.

Subpart A—General Provisions

§ 582.1 Substances that are generally recognized as safe.

(a) It is impracticable to list all substances that are generally recognized

as safe for their intended use. However, by way of illustration, the Commissioner regards such common food ingredients as salt, pepper, sugar, vinegar, baking powder, and monosodium glutamate as safe for their intended use. The lists in subparts B through H of this part include additional substances that, when used for the purposes indicated, in accordance with good manufacturing or feeding practice, are regarded by the Commissioner as generally recognized as safe for such uses.

(b) For the purposes of this section, good manufacturing or feeding practice shall be defined to include the following restrictions:

(1) The quantity of a substance added to animal food does not exceed the amount reasonably required to accomplish its intended physical, nutritional, or other technical effect in food; and

(2) The quantity of a substance that becomes a component of animal food as a result of its use in the manufacturing, processing, or packaging of food, and which is not intended to accomplish any physical or other technical effect in the food itself, shall be reduced to the extent reasonably possible.

(3) The substance is of appropriate grade and is prepared and handled as a food ingredient. Upon request the Commissioner will offer an opinion, based on specifications and intended use, as to whether or not a particular grade or lot of the substance is of suitable purity for use in food and would generally be regarded as safe for the purpose intended, by experts qualified to evaluate its safety.

(c) The inclusion of substances in the list of nutrients does not constitute a finding on the part of the Department that the substance is useful as a supplement to the diet for animals.

(d) Substances that are generally recognized as safe for their intended use within the meaning of section 409 of the Act are listed in subparts B through H of this part. When the status of a substance has been reevaluated and affirmed as GRAS or deleted from subparts B through H of this part, an appropriate explanation will be noted,

§ 582.10

e.g., “affirmed as GRAS,” “food additive regulation,” “interim food additive regulation,” or “prohibited from use in food,” with a reference to the appropriate new regulation. Such notation will apply only to the specific use covered by the review, e.g., direct animal food use and/or indirect animal food use and/or animal feed use and will not affect its status for other uses not specified in the referenced regula-

21 CFR Ch. I (4–1–99 Edition)

tion, pending a specific review of such other uses.

§ 582.10 Spices and other natural seasonings and flavorings.

Spices and other natural seasonings and flavorings that are generally recognized as safe for their intended use, within the meaning of section 409 of the act, are as follows:

Common name	Botanical name of plant source
Alfalfa herb and seed	<i>Medicago sativa</i> L.
Allspice	<i>Pimenta officinalis</i> Lindl.
Ambrette seed	<i>Hibiscus abelmoschus</i> L.
Angelica	<i>Angelica archangelica</i> L. or other spp. of <i>Angelica</i> .
Angelica root	Do.
Angelica seed	Do.
Angostura (cusparia bark)	<i>Galipea officinalis</i> Hancock.
Anise	<i>Pimpinella anisum</i> L.
Anise, star	<i>Illicium verum</i> Hook. f.
Balm (lemon balm)	<i>Melissa officinalis</i> L.
Basil, bush	<i>Ocimum minimum</i> L.
Basil, sweet	<i>Ocimum basilicum</i> L.
Bay	<i>Laurus nobilis</i> L.
Calendula	<i>Calendula officinalis</i> L.
Camomile (chamomile), English or Roman	<i>Anthemis nobilis</i> L.
Camomile (chamomile), German or Hungarian	<i>Matricaria chamomilla</i> L.
Capers	<i>Capparis spinosa</i> L.
Capsicum	<i>Capsicum frutescens</i> L. or <i>Capsicum annuum</i> L.
Caraway	<i>Carum carvi</i> L.
Caraway, black (black cumin)	<i>Nigella sativa</i> L.
Cardamom (cardamon)	<i>Elettaria cardamomum</i> Maton.
Cassia, Chinese	<i>Cinnamomum cassia</i> Blume.
Cassia, Padang or Batavia	<i>Cinnamomum burmanni</i> Blume.
Cassia, Saigon	<i>Cinnamomum loureirii</i> Nees.
Cayenne pepper	<i>Capsicum frutescens</i> L. or <i>Capsicum annuum</i> L.
Celery seed	<i>Apium graveolens</i> L.
Chervil	<i>Anthriscus cerefolium</i> (L.) Hoffm.
Chives	<i>Allium schoenoprasum</i> L.
Cinnamon, Ceylon	<i>Cinnamomum zeylanicum</i> Nees.
Cinnamon, Chinese	<i>Cinnamomum cassia</i> Blume.
Cinnamon, Saigon	<i>Cinnamomum loureirii</i> Nees.
Clary (clary sage)	<i>Salvia sclarea</i> L.
Clover	<i>Trifolium</i> spp.
Cloves	<i>Eugenia caryophyllata</i> Thunb.
Coriander	<i>Coriandrum sativum</i> L.
Cumin (cummin)	<i>Cuminum cyminum</i> L.
Cumin, black (black caraway)	<i>Nigella sativa</i> L.
Dill	<i>Anethum graveolens</i> L.
Elder flowers	<i>Sambucus canadensis</i> L.
Fennel, common	<i>Foeniculum vulgare</i> Mill.
Fennel, sweet (finocchio, Florence fennel)	<i>Foeniculum vulgare</i> Mill. var. <i>duice</i> (DC.) Alex.
Fenugreek	<i>Trigonella foenum-graecum</i> L.
Galanga (galangal)	<i>Alpinia officinarum</i> Hance.
Garlic	<i>Allium sativum</i> L.
Geranium	<i>Pelargonium</i> spp.
Ginger	<i>Zingiber officinale</i> Rosc.
Glycyrrhiza	<i>Glycyrrhiza glabra</i> L. and other spp. of <i>Glycyrrhiza</i> .
Grains of paradise	<i>Amomum melegueta</i> Rosc.
Horehound (hoarhound)	<i>Marrubium vulgare</i> L.
Horseradish	<i>Armoracia lapathifolia</i> Gilib.
Hyssop	<i>Hyssopus officinalis</i> L.
Lavender	<i>Lavandula officinalis</i> Chaix.
Licorice	<i>Glycyrrhiza glabra</i> L. and other spp. of <i>Glycyrrhiza</i> .
Linden flowers	<i>Tilia</i> spp.
Mace	<i>Myristica fragrans</i> Houtt.
Marigold, pot	<i>Calendula officinalis</i> L.
Marjoram, pot	<i>Majorana onites</i> (L.) Benth.
Marjoram, sweet	<i>Majorana hortensis</i> Moench.
Mustard, black or brown	<i>Brassica nigra</i> (L.) Koch.

Food and Drug Administration, HHS

§ 582.20

Common name	Botanical name of plant source
Mustard, brown	<i>Brassica juncea</i> (L.) Coss.
Mustard, white or yellow	<i>Brassica hirta</i> Moench.
Nutmeg	<i>Myristica fragrans</i> Houtt.
Oregano (oreganum, Mexican oregano, Mexican sage, origan)	<i>Lippia</i> spp.
Paprika	<i>Capsicum annuum</i> L.
Parsley	<i>Petroselinum crispum</i> (Mill.) Mansf.
Pepper, black	<i>Piper nigrum</i> L.
Pepper, cayenne	<i>Capsicum frutescens</i> L. or <i>Capsicum annuum</i> L.
Pepper, red	Do.
Pepper, white	<i>Piper nigrum</i> L.
Peppermint	<i>Mentha piperita</i> L.
Poppy seed	<i>Papaver somniferum</i> L.
Pot marigold	<i>Calendula officinalis</i> L.
Pot marjoram	<i>Majorana onites</i> (L.) Benth.
Rosemary	<i>Rosmarinus officinalis</i> L.
Rue	<i>Ruta graveolens</i> L.
Saffron	<i>Crocus sativus</i> L.
Sage	<i>Salvia officinalis</i> L.
Sage, Greek	<i>Salvia triloba</i> L.
Savory, summer	<i>Satureia hortensis</i> L. (Satureja).
Savory, winter	<i>Satureia montana</i> L. (Satureja).
Sesame	<i>Sesamum indicum</i> L.
Spearmint	<i>Mentha spicata</i> L.
Star anise	<i>Illicium verum</i> Hook. f.
Tarragon	<i>Artemisia dracunculus</i> L.
Thyme	<i>Thymus vulgaris</i> L.
Thyme, wild or creeping	<i>Thymus serpyllum</i> L.
Turmeric	<i>Curcuma longa</i> L.
Vanilla	<i>Vanilla planifolia</i> Andr. or <i>Vanilla tahitensis</i> J. W. Moore.
Zedoary	<i>Curcuma zedoaria</i> Rosc.

§ 582.20 Essential oils, oleoresins (solvent-free), and natural extractives (including distillates).

Essential oils, oleoresins (solvent-free), and natural extractives (includ-

ing distillates) that are generally recognized as safe for their intended use, within the meaning of section 409 of the act, are as follows:

Common name	Botanical name of plant source
Alfalfa	<i>Medicago sativa</i> L.
Allspice	<i>Pimenta officinalis</i> Lindl.
Almond, bitter (free from prussic acid)	<i>Prunus amygdalus</i> Batsch, <i>Prunus armeniaca</i> L. or <i>Prunus persica</i> (L.) Batsch.
Ambrette (seed)	<i>Hibiscus moschatus</i> Moench.
Angelica root	<i>Angelica archangelica</i> L.
Angelica seed	Do.
Angelica stem	Do.
Angostura (cusparia bark)	<i>Galipea officinalis</i> Hancock.
Anise	<i>Pimpinella anisum</i> L.
Asafetida	<i>Ferula assa-foetida</i> L. and related spp. of <i>Ferula</i> .
Balm (lemon balm)	<i>Melissa officinalis</i> L.
Balsam of Peru	<i>Myroxylon pereirae</i> Klotzsch.
Basil	<i>Ocimum basilicum</i> L.
Bay leaves	<i>Laurus nobilis</i> L.
Bay (myrcia oil)	<i>Pimenta racemosa</i> (Mill.) J. W. Moore.
Bergamot (bergamot orange)	<i>Citrus aurantium</i> L. subsp. <i>bergamia</i> Wright et Arn.
Bitter almond (free from prussic acid)	<i>Prunus amygdalus</i> Batsch, <i>Prunus armeniaca</i> L., or <i>Prunus persica</i> (L.) Batsch.
Bois de rose	<i>Aniba roseodora</i> Ducke.
Cacao	<i>Theobroma cacao</i> L.
Camomile (chamomile) flowers, Hungarian	<i>Matricaria chamomilla</i> L.
Camomile (chamomile) flowers, Roman or English	<i>Anthemis nobilis</i> L.
Cananga	<i>Cananga odorata</i> Hook. f. and Thoms.
Capsicum	<i>Capsicum frutescens</i> L. and <i>Capsicum annuum</i> L.
Caraway	<i>Carum carvi</i> L.
Cardamom seed (cardamon)	<i>Elettaria cardamomum</i> Maton.
Carob bean	<i>Ceratonia siliqua</i> L.
Carrot	<i>Daucus carota</i> L.
Cascarilla bark	<i>Croton eluteria</i> Benn.
Cassia bark, Chinese	<i>Cinnamomum cassia</i> Blume.

Common name	Botanical name of plant source
Cassia bark, Padang or Batavia	<i>Cinnamomum burmanni</i> Blume.
Cassia bark, Saigon	<i>Cinnamomum loureirii</i> Nees.
Celery seed	<i>Apium graveolens</i> L.
Cherry, wild, bark	<i>Prunus serotina</i> Ehrh.
Chervil	<i>Anthriscus cerefolium</i> (L.) Hoffm.
Chicory	<i>Cichorium intybus</i> L.
Cinnamon bark, Ceylon	<i>Cinnamomum zeylanicum</i> Nees.
Cinnamon bark, Chinese	<i>Cinnamomum cassia</i> Blume.
Cinnamon bark, Saigon	<i>Cinnamomum loureirii</i> Nees.
Cinnamon leaf, Ceylon	<i>Cinnamomum zeylanicum</i> Nees.
Cinnamon leaf, Chinese	<i>Cinnamomum cassia</i> Blume.
Cinnamon leaf, Saigon	<i>Cinnamomum loureirii</i> Nees.
Citronella	<i>Cymbopogon nardus</i> Rendle.
Citrus peels	<i>Citrus</i> spp.
Clary (clary sage)	<i>Salvia sclarea</i> L.
Clove bud	<i>Eugenia caryophyllata</i> Thunb.
Clove leaf	Do.
Clove stem	Do.
Clover	<i>Trifolium</i> spp.
Coca (decocainized)	<i>Erythroxylum coca</i> Lam. and other spp. of <i>Erythroxylum</i> .
Coffee	<i>Coffea</i> spp.
Cola nut	<i>Cola acuminata</i> Schott and Endl., and other spp. of <i>Cola</i> .
Coriander	<i>Coriandrum sativum</i> L.
Corn silk	<i>Zea mays</i> L.
Cumin (cumin)	<i>Cuminum cyminum</i> L.
Curacao orange peel (orange, bitter peel)	<i>Citrus aurantium</i> L.
Cusparia bark	<i>Galipea officinalis</i> Hancock.
Dandelion	<i>Taraxacum officinale</i> Weber and <i>T. laevigatum</i> DC.
Dandelion root	Do.
Dill	<i>Anethum graveolens</i> L.
Dog grass (quackgrass, triticum)	<i>Agropyron repens</i> (L.) Beauv.
Elder flowers	<i>Sambucus canadensis</i> L. and <i>S. nigra</i> L.
Estragole (esdragol, esdragon, tarragon)	<i>Artemisia dracunculus</i> L.
Estragon (tarragon)	Do.
Fennel, sweet	<i>Foeniculum vulgare</i> Mill.
Fenugreek	<i>Trigonella foenum-graecum</i> L.
Galanga (galangal)	<i>Alpinia officinarum</i> Hance.
Garlic	<i>Allium sativum</i> L.
Geranium	<i>Pelargonium</i> spp.
Geranium, East Indian	<i>Cymbopogon martini</i> Stapf.
Geranium, rose	<i>Pelargonium graveolens</i> L'Her.
Ginger	<i>Zingiber officinale</i> Rosc.
Glycyrrhiza	<i>Glycyrrhiza glabra</i> L. and other spp. of <i>Glycyrrhiza</i> .
Glycyrrhizin, ammoniated	Do.
Grapefruit	<i>Citrus paradisi</i> Macf.
Guava	<i>Psidium</i> spp.
Hickory bark	<i>Carya</i> spp.
Horehound (hoarhound)	<i>Marrubium vulgare</i> L.
Hops	<i>Humulus lupulus</i> L.
Horsemint	<i>Monarda punctata</i> L.
Hyssop	<i>Hyssopus officinalis</i> L.
Immortelle	<i>Helichrysum angustifolium</i> DC.
Jasmine	<i>Jasminum officinale</i> L. and other spp. of <i>Jasminum</i> .
Juniper (berries)	<i>Juniperus communis</i> L.
Kola nut	<i>Cola acuminata</i> Schott and Endl., and other spp. of <i>Cola</i> .
Laurel berries	<i>Laurus nobilis</i> L.
Laurel leaves	<i>Laurus</i> spp.
Lavender	<i>Lavandula officinalis</i> Chaix.
Lavender, spike	<i>Lavandula latifolia</i> Vill.
Lavandin	Hybrids between <i>Lavandula officinalis</i> Chaix and <i>Lavandula latifolia</i> Vill.
Lemon	<i>Citrus limon</i> (L.) Burm. f.
Lemon balm (see balm).	
Lemon grass	<i>Cymbopogon citratus</i> DC. and <i>Cymbopogon flexuosus</i> Stapf.
Lemon peel	<i>Citrus limon</i> (L.) Burm. f.
Licorice	<i>Glycyrrhiza glabra</i> L. and other spp. of <i>Glycyrrhiza</i> .
Lime	<i>Citrus aurantifolia</i> Swingle.
Linden flowers	<i>Tilia</i> spp.
Locust bean	<i>Ceratonia siliqua</i> L.
Lupulin	<i>Humulus lupulus</i> L.
Mace	<i>Myristica fragrans</i> Houtt.
Malt (extract)	<i>Hordeum vulgare</i> L., or other grains.
Mandarin	<i>Citrus reticulata</i> Blanco.
Marjoram, sweet	<i>Majorana hortensis</i> Moench.

Food and Drug Administration, HHS

§ 582.20

Common name	Botanical name of plant source
Mate 1	<i>Ilex paraguariensis</i> St. Hil.
Melissa (see balm)	
Menthol	<i>Mentha</i> spp.
Menthyl acetate	Do.
Molasses (extract)	<i>Saccharum officinarum</i> L.
Mustard	<i>Brassica</i> spp.
Naringin	<i>Citrus paradisi</i> Macf.
Neroli, bigarade	<i>Citrus aurantium</i> L.
Nutmeg	<i>Myristica fragrans</i> Houtt.
Onion	<i>Allium cepa</i> L.
Orange, bitter, flowers	<i>Citrus aurantium</i> L.
Orange, bitter, peel	Do.
Orange leaf	<i>Citrus sinensis</i> (L.) Osbeck.
Orange, sweet	Do.
Orange, sweet, flowers	Do.
Orange, sweet, peel	Do.
Origanum	<i>Origanum</i> spp.
Palmarosa	<i>Cymbopogon martini</i> Stapf.
Paprika	<i>Capsicum annum</i> L.
Parsley	<i>Petroselinum crispum</i> (Mill.) Mansf.
Pepper, black	<i>Piper nigrum</i> L.
Pepper, white	<i>Piper nigrum</i> L.
Peppermint	<i>Mentha piperita</i> L.
Peruvian balsam	<i>Myroxylon pereirae</i> Klotzsch.
Petitgrain	<i>Citrus aurantium</i> L.
Petitgrain lemon	<i>Citrus limon</i> (L.) Burm. f.
Petitgrain mandarin or tangerine	<i>Citrus reticulata</i> Blanco.
Pimenta	<i>Pimenta officinalis</i> Lindl.
Pimenta leaf	<i>Primenta officinalis</i> Lindl.
Pipsissewa leaves	<i>Chimaphila umbellata</i> Nutt.
Pomegranate	<i>Punica granatum</i> L.
Prickly ash bark	<i>Xanthoxylum</i> (or <i>Zanthoxylum</i>) <i>Americanum</i> Mill. or <i>Xanthoxylum clava-herculis</i> L.
Rose absolute	<i>Rosa alba</i> L., <i>Rosa centifolia</i> L., <i>Rosa damascena</i> Mill., <i>Rosa gallica</i> L., and vars. of these spp.
Rose (otto of roses, attar of roses)	Do.
Rose buds	Do.
Rose flowers	Do.
Rose fruit (hips)	Do.
Rose geranium	<i>Pelargonium graveolens</i> L'Her.
Rose leaves	<i>Rosa</i> spp.
Rosemary	<i>Rosmarinus officinalis</i> L.
Rue	<i>Ruta graveolens</i> L.
Saffron	<i>Crocus sativus</i> L.
Sage	<i>Salvia officinalis</i> L.
Sage, Greek	<i>Salvia triloba</i> L.
Sage, Spanish	<i>Salvia lavandulaefolia</i> Vahl.
St. John's bread	<i>Ceratonja siliqua</i> L.
Savory, summer	<i>Satureia hortensis</i> L.
Savory, winter	<i>Satureia montana</i> L.
Schinus molle	<i>Schinus molle</i> L.
Sloe berries (blackthorn berries)	<i>Prunus spinosa</i> L.
Spearmint	<i>Mentha spicata</i> L.
Spike lavender	<i>Lavandula latifolia</i> Vill.
Tamarind	<i>Tamarindus indica</i> L.
Tangerine	<i>Citrus reticulata</i> Blanco.
Tannic acid	Nutgalls of <i>Quercus infectoria</i> Oliver and related spp. of <i>Quercus</i> . Also in many other plants.
Tarragon	<i>Artemisia dracunculus</i> L.
Tea	<i>Thea sinensis</i> L.
Thyme	<i>Thymus vulgaris</i> L. and <i>Thymus zygis</i> var. <i>gracilis</i> Boiss.
Thyme, white	Do.
Thyme, wild or creeping	<i>Thymus serpyllum</i> L.
Triticum (see dog grass)	
Tuberose	<i>Polianthes tuberosa</i> L.
Turmeric	<i>Curcuma longa</i> L.
Vanilla	<i>Vanilla planifolia</i> Andr. or <i>Vanilla tahitensis</i> J. W. Moore.
Violet flowers	<i>Viola odorata</i> L.
Violet leaves	Do.
Violet leaves absolute	Do.
Wild cherry bark	<i>Prunus serotina</i> Ehrh.
Ylang-ylang	<i>Cananga odorata</i> Hook. f. and Thoms.
Zedoary bark	<i>Curcuma zedoaria</i> Rosc.

§ 582.30

21 CFR Ch. I (4–1–99 Edition)

§ 582.30 Natural substances used in conjunction with spices and other natural seasonings and flavorings.

Natural substances used in conjunction with spices and other natural

seasonings and flavorings that are generally recognized as safe for their intended use, within the meaning of section 409 of the act, are as follows:

Common name	Botanical name of plant source
Algae, brown (kelp)	<i>Laminaria</i> spp. and <i>Nereocystis</i> spp.
Algae, red	<i>Porphyra</i> spp. and <i>Rhodomenia palmata</i> (L.) Grev.
Dulse	<i>Rhodomenia palmata</i> (L.)

§ 582.40 Natural extractives (solvent-free) used in conjunction with spices, seasonings, and flavorings.

Natural extractives (solvent-free) used in conjunction with spices,

seasonings, and flavorings that are generally recognized as safe for their intended use, within the meaning of section 409 of the act, are as follows:

Common name	Botanical name of plant source
Algae, brown	<i>Laminaria</i> spp. and <i>Nereocystis</i> spp.
Algae, red	<i>Porphyra</i> spp. and <i>Rhodomenia palmata</i> (L.) Grev.
Apricot kernel (persic oil)	<i>Prunus armeniaca</i> L.
Dulse	<i>Rhodomenia palmata</i> (L.) Grev.
Kelp (see algae, brown).	
Peach kernel (persic oil)	<i>Prunus persica</i> Sieb. et Zucc.
Peanut stearine	<i>Arachis hypogaea</i> L.
Persic oil (see apricot kernel and peach kernel).	
Quince seed	<i>Cydonia oblonga</i> Miller.

§ 582.50 Certain other spices, seasonings, essential oils, oleoresins, and natural extracts.

Certain other spices, seasonings, essential oils, oleoresins, and natural ex-

tracts that are generally recognized as safe for their intended use, within the meaning of section 409 of the act, are as follows:

Common name	Derivation
Ambergris	<i>Physeter macrocephalus</i> L.
Castoreum	Castor fiber L. and <i>C. canadensis</i> Kuhl.
Civet (zibeth, zibet, zibetum)	Civet cats, <i>Viverra civetta</i> Schreber and <i>Viverra zibetha</i> Schreber.
Cognac oil, white and green	<i>Ethyl oenanthate</i> , so-called.
Musk (Tonquin musk)	Musk deer, <i>Moschus moschiferus</i> L.

§ 582.60 Synthetic flavoring substances and adjuvants.

Synthetic flavoring substances and adjuvants that are generally recognized as safe for their intended use, within the meaning of section 409 of the act, are as follows:

Acetaldehyde (ethanal).
 Acetoin (acetyl methylcarbinol).
 Aconitic acid (equisetic acid, citridic acid, achilleic acid).
 Anethole (parapropenyl anisole).
 Benzaldehyde (benzoic aldehyde).
 N-Butyric acid (butanoic acid).
 d- or l-Carvone (carvol).

Cinnamaldehyde (cinnamic aldehyde).
 Citral (2,6-dimethyloctadien-2,6-al-8, geranial, neral).
 Decanal (N-decylaldehyde, capraldehyde, capric aldehyde, caprinaldehyde, aldehyde C-10).
 Diacetyl (2,3-butanedione). Ethyl acetate. Ethyl butyrate.
 3-Methyl-3-phenyl glycidic acid ethyl ester (ethyl-methyl-phenyl-glycidate, so-called strawberry aldehyde, C-16 aldehyde).
 Ethyl vanillin.
 Eugenol.
 Geraniol (3,7-dimethyl-2,6 and 3,6-octadien-1-ol).
 Geranyl acetate (geraniol acetate).

Food and Drug Administration, HHS

§ 582.1061

Glycerol (glyceryl) tributyrate (tributyryl, butyrin).
 Limonene (*d*-, *l*-, and *dl*-).
 Linalool (linalol, 3,7-dimethyl-1,6-octadien-3-ol).
 Linalyl acetate (bergamol).
 1-Malic acid.
 Methyl anthranilate (methyl-2-aminobenzoate).
 Piperonal (3,4-methylenedioxy-benzaldehyde, heliotropin).
 Vanillin.

§ 582.80 Trace minerals added to animal feeds.

These substances added to animal feeds as nutritional dietary supplements are generally recognized as safe when added at levels consistent with good feeding practice.¹

Element	Source compounds
Cobalt	Cobalt acetate. Cobalt carbonate. Cobalt chloride. Cobalt oxide. Cobalt sulfate.
Copper	Copper carbonate. Copper chloride. Copper gluconate. Copper hydroxide. Copper orthophosphate. Copper oxide. Copper pyrophosphate. Copper sulfate.
Iodine	Calcium iodate. Calcium iodobenzenate. Cuprous iodide. 3,5-Diiodosalicylic acid. Ethylenediamine dihydroiodide. Potassium iodate. Potassium iodide. Sodium iodate. Sodium iodide. Thymol iodide.
Iron	Iron ammonium citrate. Iron carbonate. Iron chloride. Iron gluconate. Iron oxide. Iron phosphate. Iron pyrophosphate. Iron sulfate. Reduced iron.
Manganese	Manganese acetate. Manganese carbonate. Manganese citrate (soluble). Manganese chloride. Manganese gluconate. Manganese orthophosphate. Manganese phosphate (dibasic). Manganese sulfate. Manganous oxide.
Zinc	Zinc acetate. Zinc carbonate. Zinc chloride. Zinc oxide.

¹ All substances listed may be in anhydrous or hydrated form.

Element	Source compounds
	Zinc sulfate.

§ 582.99 Adjuvants for pesticide chemicals.

Adjuvants, identified and used in accordance with 40 CFR 180.1001(c) and (d), which are added to pesticide use dilutions by a grower or applicator prior to application to the raw agricultural commodity, are exempt from the requirement of tolerances under section 409 of the act.

Subpart B—General Purpose Food Additives

§ 582.1005 Acetic acid.

- (a) *Product*. Acetic acid.
 (b) *Conditions of use*. This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1009 Adipic acid.

- (a) *Product*. Adipic acid.
 (b) [Reserved]
 (c) *Limitations, restrictions, or explanation*. This substance is generally recognized as safe when used as a buffer and neutralizing agent in accordance with good manufacturing or feeding practice.

§ 582.1033 Citric acid.

- (a) *Product*. Citric acid.
 (b) *Conditions of use*. This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1057 Hydrochloric acid.

- (a) *Product*. Hydrochloric acid.
 (b) [Reserved]
 (c) *Limitations, restrictions, or explanation*. This substance is generally recognized as safe when used as a buffer and neutralizing agent in accordance with good manufacturing or feeding practice.

§ 582.1061 Lactic acid.

- (a) *Product*. Lactic acid.
 (b) *Conditions of use*. This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1069

§ 582.1069 Malic acid.

- (a) *Product.* Malic acid.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1073 Phosphoric acid.

- (a) *Product.* Phosphoric acid.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1077 Potassium acid tartrate.

- (a) *Product.* Potassium acid tartrate.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1087 Sodium acid pyrophosphate.

- (a) *Product.* Sodium acid pyrophosphate.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1091 Succinic acid.

- (a) *Product.* Succinic acid.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1095 Sulfuric acid.

- (a) *Product.* Sulfuric acid.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1099 Tartaric acid.

- (a) *Product.* Tartaric acid.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1125 Aluminum sulfate.

- (a) *Product.* Aluminum sulfate.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

21 CFR Ch. I (4–1–99 Edition)

§ 582.1127 Aluminum ammonium sulfate.

- (a) *Product.* Aluminum ammonium sulfate.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1129 Aluminum potassium sulfate.

- (a) *Product.* Aluminum potassium sulfate.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1131 Aluminum sodium sulfate.

- (a) *Product.* Aluminum sodium sulfate.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1135 Ammonium bicarbonate.

- (a) *Product.* Ammonium bicarbonate.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1137 Ammonium carbonate.

- (a) *Product.* Ammonium carbonate.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1139 Ammonium hydroxide.

- (a) *Product.* Ammonium hydroxide.
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1141 Ammonium phosphate.

- (a) *Product.* Ammonium phosphate (mono- and dibasic).
(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1143 Ammonium sulfate.

- (a) *Product.* Ammonium sulfate.

Food and Drug Administration, HHS

§ 582.1355

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1155 Bentonite.

(a) *Product.* Bentonite.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1165 Butane.

(a) *Product.* Butane.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1191 Calcium carbonate.

(a) *Product.* Calcium carbonate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1193 Calcium chloride.

(a) *Product.* Calcium chloride.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1195 Calcium citrate.

(a) *Product.* Calcium citrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1199 Calcium gluconate.

(a) *Product.* Calcium gluconate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1205 Calcium hydroxide.

(a) *Product.* Calcium hydroxide.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1207 Calcium lactate.

(a) *Product.* Calcium lactate.

(b) *Conditions of use.* This substance is generally recognized as safe when

used in accordance with good manufacturing or feeding practice.

§ 582.1210 Calcium oxide.

(a) *Product.* Calcium oxide.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1217 Calcium phosphate.

(a) *Product.* Calcium phosphate (mono-, di-, and tribasic).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1235 Caramel.

(a) *Product.* Caramel.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1240 Carbon dioxide.

(a) *Product.* Carbon dioxide.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1275 Dextrans.

(a) *Product.* Dextrans of average molecular weight below 100,000.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1320 Glycerin.

(a) *Product.* Glycerin.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1324 Glyceryl monostearate.

(a) *Product.* Glyceryl monostearate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1355 Helium.

(a) *Product.* Helium.

(b) *Conditions of use.* This substance is generally recognized as safe when

§ 582.1366

used in accordance with good manufacturing or feeding practice.

§ 582.1366 Hydrogen peroxide.

(a) *Product.* Hydrogen peroxide.

(b) [Reserved]

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used as a bleaching agent in accordance with good manufacturing or feeding practice.

§ 582.1400 Lecithin.

(a) *Product.* Lecithin.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1425 Magnesium carbonate.

(a) *Product.* Magnesium carbonate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1428 Magnesium hydroxide.

(a) *Product.* Magnesium hydroxide.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1431 Magnesium oxide.

(a) *Product.* Magnesium oxide.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1480 Methylcellulose.

(a) *Product.* U.S.P. methylcellulose, except that the methoxy content shall not be less than 27.5 percent and not more than 31.5 percent on a dry-weight basis.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1500 Monoammonium glutamate.

(a) *Product.* Monoammonium glutamate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

21 CFR Ch. I (4–1–99 Edition)

§ 582.1516 Monopotassium glutamate.

(a) *Product.* Monopotassium glutamate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1540 Nitrogen.

(a) *Product.* Nitrogen.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1585 Papain.

(a) *Product.* Papain.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1613 Potassium bicarbonate.

(a) *Product.* Potassium bicarbonate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1619 Potassium carbonate.

(a) *Product.* Potassium carbonate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1625 Potassium citrate.

(a) *Product.* Potassium citrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1631 Potassium hydroxide.

(a) *Product.* Potassium hydroxide.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1643 Potassium sulfate.

(a) *Product.* Potassium sulfate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1655 Propane.

(a) *Product.* Propane.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1666 Propylene glycol.

(a) *Product.* Propylene glycol.

(b) *Conditions of use.* This substance is generally recognized as safe (except in cat food) when used in accordance with good manufacturing or feeding practice.

[41 FR 38657, Sept. 10, 1976, as amended at 61 FR 19544, May 2, 1996]

§ 582.1685 Rennet.

(a) *Product.* Rennet (rennin).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1711 Silica aerogel.

(a) *Product.* Silica aerogel as a finely powdered microcellular silica foam having a minimum silica content of 89.5 percent.

(b) [Reserved]

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used as a component of antifoaming agents in accordance with good manufacturing or feeding practice.

§ 582.1721 Sodium acetate.

(a) *Product.* Sodium acetate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1736 Sodium bicarbonate.

(a) *Product.* Sodium bicarbonate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1742 Sodium carbonate.

(a) *Product.* Sodium carbonate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1745 Sodium carboxymethylcellulose.

(a) *Product.* Sodium carboxymethylcellulose is the sodium salt of carboxymethylcellulose not less than 99.5 percent on a dry-weight basis, with maximum substitution of 0.95 carboxymethyl groups per anhydroglucose unit, and with a minimum viscosity of 25 centipoises for 2 percent by weight aqueous solution at 25° C.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1748 Sodium caseinate.

(a) *Product.* Sodium caseinate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1751 Sodium citrate.

(a) *Product.* Sodium citrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1763 Sodium hydroxide.

(a) *Product.* Sodium hydroxide.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1775 Sodium pectinate.

(a) *Product.* Sodium pectinate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1778 Sodium phosphate.

(a) *Product.* Sodium phosphate (mono-, di-, and tribasic).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1781 Sodium aluminum phosphate.

(a) *Product.* Sodium aluminum phosphate.

§ 582.1792

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1792 Sodium sesquicarbonate.

(a) *Product.* Sodium sesquicarbonate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1804 Sodium potassium tartrate.

(a) *Product.* Sodium potassium tartrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1810 Sodium tripolyphosphate.

(a) *Product.* Sodium tripolyphosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1901 Triacetin.

(a) *Product.* Triacetin (glyceryl triacetate).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1973 Beeswax.

(a) *Product.* Beeswax (yellow wax).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1975 Bleached beeswax.

(a) *Product.* Bleached beeswax (white wax).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.1978 Carnauba wax.

(a) *Product.* Carnauba wax.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

21 CFR Ch. I (4–1–99 Edition)

Subpart C—Anticaking Agents

§ 582.2122 Aluminum calcium silicate.

(a) *Product.* Aluminum calcium silicate.

(b) *Tolerance.* 2 percent.

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in table salt in accordance with good manufacturing or feeding practice.

§ 582.2227 Calcium silicate.

(a) *Product.* Calcium silicate.

(b) *Tolerance.* 2 percent and 5 percent.

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used at levels not exceeding 2 percent in table salt and 5 percent in baking powder in accordance with good manufacturing or feeding practice.

§ 582.2437 Magnesium silicate.

(a) *Product.* Magnesium silicate.

(b) *Tolerance.* 2 percent.

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in table salt in accordance with good manufacturing or feeding practice.

§ 582.2727 Sodium aluminosilicate.

(a) *Product.* Sodium aluminosilicate (sodium silicoaluminate).

(b) *Tolerance.* This substance is generally recognized as safe for use at a level not exceeding 2 percent in accordance with good manufacturing or feeding practice.

§ 582.2729 Hydrated sodium calcium aluminosilicate.

(a) *Product.* Hydrated sodium calcium aluminosilicate (sodium calcium silicoaluminate).

(b) *Tolerance.* This substance is generally recognized as safe for use at a level not exceeding 2 percent in accordance with good manufacturing or feeding practice.

§ 582.2906 Tricalcium silicate.

(a) *Product.* Tricalcium silicate.

(b) *Tolerance.* 2 percent.

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in table salt

Food and Drug Administration, HHS

§ 582.3280

in accordance with good manufacturing or feeding practice.

Subpart D—Chemical Preservatives

§ 582.3013 Ascorbic acid.

(a) *Product.* Ascorbic acid.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.3021 Benzoic acid.

(a) *Product.* Benzoic acid.

(b) *Tolerance.* This substance is generally recognized as safe for use at a level not exceeding 0.1 percent in accordance with good manufacturing or feeding practice.

§ 582.3041 Erythorbic acid.

(a) *Product.* Erythorbic acid.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.3081 Propionic acid.

(a) *Product.* Propionic acid.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.3089 Sorbic acid.

(a) *Product.* Sorbic acid.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.3109 Thiodipropionic acid.

(a) *Product.* Thiodipropionic acid.

(b) *Tolerance.* This substance is generally recognized as safe for use in food when the total content of antioxidants is not over 0.02 percent of fat or oil content including essential (volatile) oil content of the food, provided the substance is used in accordance with good manufacturing or feeding practice.

§ 582.3149 Ascorbyl palmitate.

(a) *Product.* Ascorbyl palmitate.

(b) *Conditions of use.* This substance is generally recognized as safe when

used in accordance with good manufacturing or feeding practice.

§ 582.3169 Butylated hydroxyanisole.

(a) *Product.* Butylated hydroxyanisole.

(b) *Tolerance.* This substance is generally recognized as safe for use in food when the total content of antioxidants is not over 0.02 percent of fat or oil content, including essential (volatile) oil content of food provided the substance is used in accordance with good manufacturing or feeding practice.

§ 582.3173 Butylated hydroxytoluene.

(a) *Product.* Butylated hydroxytoluene.

(b) *Tolerance.* This substance is generally recognized as safe for use in food when the total content of antioxidants is not over 0.02 percent of fat or oil content, including essential (volatile) oil content of food provided the substance is used in accordance with good manufacturing or feeding practice.

§ 582.3189 Calcium ascorbate.

(a) *Product.* Calcium ascorbate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.3221 Calcium propionate.

(a) *Product.* Calcium propionate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.3225 Calcium sorbate.

(a) *Product.* Calcium sorbate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.3280 Dilauryl thiodipropionate.

(a) *Product.* Dilauryl thiodipropionate.

(b) *Tolerance.* This substance is generally recognized as safe for use in food when the total content of antioxidants is not over 0.02 percent of fat or oil content, including essential (volatile) oil content of the food, provided the substance is used in accordance with

§ 582.3336

good manufacturing or feeding practice.

§ 582.3336 Gum guaiac.

- (a) *Product.* Gum guaiac.
- (b) *Tolerance.* 0.1 percent (equivalent antioxidant activity 0.01 percent).
- (c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in edible fats or oils in accordance with good manufacturing or feeding practice.

§ 582.3490 Methylparaben.

- (a) *Product.* Methylparaben (methyl *p*-hydroxybenzoate).
- (b) *Tolerance.* This substance is generally recognized as safe for use at a level not exceeding 0.1 percent in accordance with good manufacturing or feeding practice.

§ 582.3616 Potassium bisulfite.

- (a) *Product.* Potassium bisulfite.
- (b) [Reserved]
- (c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice, except that it is not used in meats or in food recognized as source of vitamin B₁.

§ 582.3637 Potassium metabisulfite.

- (a) *Product.* Potassium metabisulfite.
- (b) [Reserved]
- (c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice, except that it is not used in meats or in food recognized as source of vitamin B₁.

§ 582.3640 Potassium sorbate.

- (a) *Product.* Potassium sorbate.
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.3660 Propyl gallate.

- (a) *Product.* Propyl gallate.
- (b) *Tolerance.* This substance is generally recognized as safe for use in food when the total content of antioxidants is not over 0.02 percent of fat or oil content, including essential (volatile) oil content of the food, provided the

21 CFR Ch. I (4–1–99 Edition)

substance is used in accordance with good manufacturing or feeding practice.

§ 582.3670 Propylparaben.

- (a) *Product.* Propylparaben (propyl *p*-hydroxybenzoate).
- (b) *Tolerance.* This substance is generally recognized as safe for use at a level not exceeding 0.1 percent in accordance with good manufacturing or feeding practice.

§ 582.3731 Sodium ascorbate.

- (a) *Product.* Sodium ascorbate.
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.3733 Sodium benzoate.

- (a) *Product.* Sodium benzoate.
- (b) *Tolerance.* This substance is generally recognized as safe for use at a level not exceeding 0.1 percent in accordance with good manufacturing or feeding practice.

§ 582.3739 Sodium bisulfite.

- (a) *Product.* Sodium bisulfite.
- (b) [Reserved]
- (c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice, except that it is not used in meats or in food recognized as source of vitamin B₁.

§ 582.3766 Sodium metabisulfite.

- (a) *Product.* Sodium metabisulfite.
- (b) [Reserved]
- (c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice, except that it is not used in meats or in food recognized as source of vitamin B₁.

§ 582.3784 Sodium propionate.

- (a) *Product.* Sodium propionate.
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.3795 Sodium sorbate.

- (a) *Product.* Sodium sorbate.

Food and Drug Administration, HHS

§ 582.5049

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.3798 Sodium sulfite.

(a) *Product.* Sodium sulfite.

(b) [Reserved]

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice, except that it is not used in meats or in food recognized as source of vitamin B₁.

§ 582.3845 Stannous chloride.

(a) *Product.* Stannous chloride.

(b) *Tolerance.* This substance is generally recognized as safe for use at a level not exceeding 0.0015 percent calculated as tin in accordance with good manufacturing or feeding practice.

§ 582.3862 Sulfur dioxide.

(a) *Product.* Sulfur dioxide.

(b) [Reserved]

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice, except that it is not used in meats or in food recognized as source of vitamin B₁.

§ 582.3890 Tocopherols.

(a) *Product.* Tocopherols.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

Subpart E—Emulsifying Agents

§ 582.4101 Diacetyl tartaric acid esters of mono- and diglycerides of edible fats or oils, or edible fat-forming fatty acids.

(a) *Product.* Diacetyl tartaric acid esters of mono- and diglycerides of edible fats or oils, or edible fat-forming fatty acids.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.4505 Mono- and diglycerides of edible fats or oils, or edible fat-forming acids.

(a) *Product.* Mono- and diglycerides of edible fats or oils, or edible fat-forming acids.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.4521 Monosodium phosphate derivatives of mono- and diglycerides of edible fats or oils, or edible fat-forming fatty acids.

(a) *Product.* Monosodium phosphate derivatives of mono- and diglycerides of edible fats or oils, or edible fat-forming fatty acids.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.4666 Propylene glycol.

(a) *Product.* Propylene glycol.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

Subpart F—Nutrients and/or Dietary Supplements¹

§ 582.5013 Ascorbic acid.

(a) *Product.* Ascorbic acid.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5017 Aspartic acid.

(a) *Product.* Aspartic acid (L- and DL-forms).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5049 Aminoacetic acid.

(a) *Product.* Glycine (aminoacetic acid).

(b) [Reserved]

¹Amino acids listed in this subpart may be free hydrochloride salt, hydrated, or anhydrous form, where applicable.

§ 582.5065

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in animal feeds in accordance with good manufacturing or feeding practice.

§ 582.5065 Linoleic acid.

(a) *Product.* Linoleic acid prepared from edible fats and oils and free from chick-edema factor.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5118 Alanine.

(a) *Product.* Alanine (L- and DL-forms).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5145 Arginine.

(a) *Product.* Arginine (L- and DL-forms).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5159 Biotin.

(a) *Product.* Biotin.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5191 Calcium carbonate.

(a) *Product.* Calcium carbonate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5195 Calcium citrate.

(a) *Product.* Calcium citrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5201 Calcium glycerophosphate.

(a) *Product.* Calcium glycerophosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

21 CFR Ch. I (4–1–99 Edition)

§ 582.5210 Calcium oxide.

(a) *Product.* Calcium oxide.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5212 Calcium pantothenate.

(a) *Product.* Calcium pantothenate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5217 Calcium phosphate.

(a) *Product.* Calcium phosphate (mono-, di-, and tribasic).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5223 Calcium pyrophosphate.

(a) *Product.* Calcium pyrophosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5230 Calcium sulfate.

(a) *Product.* Calcium sulfate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5245 Carotene.

(a) *Product.* Carotene.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5250 Choline bitartrate.

(a) *Product.* Choline bitartrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5252 Choline chloride.

(a) *Product.* Choline chloride.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

Food and Drug Administration, HHS

§ 582.5431

§ 582.5260 Copper gluconate.

- (a) *Product.* Copper gluconate.
- (b) *Tolerance.* This substance is generally recognized as safe for use at a level not exceeding 0.005 percent in accordance with good manufacturing or feeding practice.

§ 582.5271 Cysteine.

- (a) *Product.* Cysteine (L-forms).
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5273 Cystine.

- (a) *Product.* Cystine (L- and DL-forms).
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5301 Ferric phosphate.

- (a) *Product.* Ferric phosphate.
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5304 Ferric pyrophosphate.

- (a) *Product.* Ferric pyrophosphate.
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5306 Ferric sodium pyrophosphate.

- (a) *Product.* Ferric sodium pyrophosphate.
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5308 Ferrous gluconate.

- (a) *Product.* Ferrous gluconate.
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5311 Ferrous lactate.

- (a) *Product.* Ferrous lactate.
- (b) *Conditions of use.* This substance is generally recognized as safe when

used in accordance with good manufacturing or feeding practice.

§ 582.5315 Ferrous sulfate.

- (a) *Product.* Ferrous sulfate.
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5361 Histidine.

- (a) *Product.* Histidine (L- and DL-forms).
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5370 Inositol.

- (a) *Product.* Inositol.
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5375 Iron reduced.

- (a) *Product.* Iron reduced.
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5381 Isoleucine.

- (a) *Product.* Isoleucine (L- and DL-forms).
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5406 Leucine.

- (a) *Product.* Leucine (L- and DL-forms).
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5411 Lysine.

- (a) *Product.* Lysine (L- and DL-forms).
- (b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5431 Magnesium oxide.

- (a) *Product.* Magnesium oxide.

§ 582.5434

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5434 Magnesium phosphate.

(a) *Product.* Magnesium phosphate (di- and tribasic).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5443 Magnesium sulfate.

(a) *Product.* Magnesium sulfate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5446 Manganese chloride.

(a) *Product.* Manganese chloride.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5449 Manganese citrate.

(a) *Product.* Manganese citrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5452 Manganese gluconate.

(a) *Product.* Manganese gluconate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5455 Manganese glycerophosphate.

(a) *Product.* Manganese glycerophosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5458 Manganese hypophosphite.

(a) *Product.* Manganese hypophosphite.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

21 CFR Ch. I (4–1–99 Edition)

§ 582.5461 Manganese sulfate.

(a) *Product.* Manganese sulfate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5464 Manganous oxide.

(a) *Product.* Manganous oxide.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5470 Mannitol.

(a) *Product.* Mannitol.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5475 Methionine.

(a) *Product.* Methionine.

(b) [Reserved]

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in animal feeds in accordance with good manufacturing or feeding practice.

§ 582.5477 Methionine hydroxy analog and its calcium salts.

(a) *Product.* Methionine hydroxy analog and its calcium salts.

(b) [Reserved]

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in animal feeds in accordance with good manufacturing or feeding practice.

§ 582.5530 Niacin.

(a) *Product.* Niacin.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5535 Niacinamide.

(a) *Product.* Niacinamide.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5580 D-Pantothenyl alcohol.

(a) *Product.* D-Pantothenyl alcohol.

Food and Drug Administration, HHS

§ 582.5881

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5590 Phenylalanine.

(a) *Product.* Phenylalanine (L- and DL-forms).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5622 Potassium chloride.

(a) *Product.* Potassium chloride.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5628 Potassium glycerophosphate.

(a) *Product.* Potassium glycerophosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5634 Potassium iodide.

(a) *Product.* Potassium iodide.

(b) *Tolerance.* 0.01 percent.

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in table salt as a source of dietary iodine in accordance with good manufacturing or feeding practice.

§ 582.5650 Proline.

(a) *Product.* Proline (L- and DL-forms).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5676 Pyridoxine hydrochloride.

(a) *Product.* Pyridoxine hydrochloride.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5695 Riboflavin.

(a) *Product.* Riboflavin.

(b) *Conditions of use.* This substance is generally recognized as safe when

used in accordance with good manufacturing or feeding practice.

§ 582.5697 Riboflavin-5-phosphate.

(a) *Product.* Riboflavin-5-phosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5701 Serine.

(a) *Product.* Serine (L- and DL-forms).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5772 Sodium pantothenate.

(a) *Product.* Sodium pantothenate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5778 Sodium phosphate.

(a) *Product.* Sodium phosphate (mono-, di-, and tribasic).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5835 Sorbitol.

(a) *Product.* Sorbitol.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5875 Thiamine hydrochloride.

(a) *Product.* Thiamine hydrochloride.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5878 Thiamine mononitrate.

(a) *Product.* Thiamine mononitrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5881 Threonine.

(a) *Product.* Threonine (L- and DL-forms).

(b) *Conditions of use.* This substance is generally recognized as safe when

§ 582.5890

used in accordance with good manufacturing or feeding practice.

§ 582.5890 Tocopherols.

(a) *Product.* Tocopherols.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5892 *a*-Tocopherol acetate.

(a) *Product.* *a*-Tocopherol acetate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5915 Tryptophane.

(a) *Product.* Tryptophane (L- and DL-forms).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5920 Tyrosine.

(a) *Product.* Tyrosine (L- and DL-forms).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5925 Valine.

(a) *Product.* Valine (L- and DL-forms).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5930 Vitamin A.

(a) *Product.* Vitamin A.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5933 Vitamin A acetate.

(a) *Product.* Vitamin A acetate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5936 Vitamin A palmitate.

(a) *Product.* Vitamin A palmitate.

21 CFR Ch. I (4–1–99 Edition)

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5945 Vitamin B₁₂.

(a) *Product.* Vitamin B₁₂.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5950 Vitamin D₂.

(a) *Product.* Vitamin D₂.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5953 Vitamin D₃.

(a) *Product.* Vitamin D₃.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5985 Zinc chloride.

(a) *Product.* Zinc chloride.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5988 Zinc gluconate.

(a) *Product.* Zinc gluconate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5991 Zinc oxide.

(a) *Product.* Zinc oxide.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5994 Zinc stearate.

(a) *Product.* Zinc stearate prepared from stearic acid free from chik-edema factor.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.5997 Zinc sulfate.

(a) *Product.* Zinc sulfate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

Subpart G—Sequestrants²

§ 582.6033 Citric acid.

(a) *Product.* Citric acid.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6085 Sodium acid phosphate.

(a) *Product.* Sodium acid phosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6099 Tartaric acid.

(a) *Product.* Tartaric acid.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6185 Calcium acetate.

(a) *Product.* Calcium acetate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6193 Calcium chloride.

(a) *Product.* Calcium chloride.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6195 Calcium citrate.

(a) *Product.* Calcium citrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6197 Calcium diacetate.

(a) *Product.* Calcium diacetate.

(b) *Conditions of use.* This substance is generally recognized as safe when

used in accordance with good manufacturing or feeding practice.

§ 582.6199 Calcium gluconate.

(a) *Product.* Calcium gluconate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6203 Calcium hexametaphosphate.

(a) *Product.* Calcium hexametaphosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6215 Monobasic calcium phosphate.

(a) *Product.* Monobasic calcium phosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6219 Calcium phytate.

(a) *Product.* Calcium phytate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6285 Dipotassium phosphate.

(a) *Product.* Dipotassium phosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6290 Disodium phosphate.

(a) *Product.* Disodium phosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6386 Isopropyl citrate.

(a) *Product.* Isopropyl citrate.

(b) *Tolerance.* This substance is generally recognized as safe for use at a level not exceeding 0.02 percent in accordance with good manufacturing or feeding practice.

§ 582.6511 Monoisopropyl citrate.

(a) *Product.* Monoisopropyl citrate.

²For the purpose of this subpart, no attempt has been made to designate those sequestrants that may also function as chemical preservatives.

§ 582.6625

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6625 Potassium citrate.

(a) *Product.* Potassium citrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

[41 FR 38657, Sept. 10, 1976. Redesignated at 42 FR 14091, Mar. 15, 1977]

§ 582.6751 Sodium citrate.

(a) *Product.* Sodium citrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

[41 FR 38657, Sept. 10, 1976. Redesignated at 42 FR 14091, Mar. 15, 1977]

§ 582.6754 Sodium diacetate.

(a) *Product.* Sodium diacetate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6757 Sodium gluconate.

(a) *Product.* Sodium gluconate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6760 Sodium hexametaphosphate.

(a) *Product.* Sodium hexametaphosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6769 Sodium metaphosphate.

(a) *Product.* Sodium metaphosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6778 Sodium phosphate.

(a) *Product.* Sodium phosphate (mono-, di-, and tribasic).

(b) *Conditions of use.* This substance is generally recognized as safe when

21 CFR Ch. I (4–1–99 Edition)

used in accordance with good manufacturing or feeding practice.

§ 582.6787 Sodium pyrophosphate.

(a) *Product.* Sodium pyrophosphate.

(b) *Condition of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6789 Tetra sodium pyrophosphate.

(a) *Product.* Tetra sodium pyrophosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6801 Sodium tartrate.

(a) *Product.* Sodium tartrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6804 Sodium potassium tartrate.

(a) *Product.* Sodium potassium tartrate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6807 Sodium thiosulfate.

(a) *Product.* Sodium thiosulfate.

(b) *Tolerance.* 0.1 percent.

(c) *Limitations, restrictions, or explanation.* This substance is generally recognized as safe when used in salt in accordance with good manufacturing or feeding practice.

§ 582.6810 Sodium tripolyphosphate.

(a) *Product.* Sodium tripolyphosphate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.6851 Stearyl citrate.

(a) *Product.* Stearyl citrate.

(b) *Tolerance.* This substance is generally recognized as safe for use at a level not exceeding 0.15 percent in accordance with good manufacturing or feeding practice.

Subpart H—Stabilizers**§ 582.7115 Agar-agar.**

(a) *Product.* Agar-agar.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.7133 Ammonium alginate.

(a) *Product.* Ammonium alginate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.7187 Calcium alginate.

(a) *Product.* Calcium alginate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.7255 Chondrus extract.

(a) *Product.* Chondrus extract (carrageenin).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.7330 Gum arabic.

(a) *Product.* Acacia (gum arabic).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.7333 Gum ghatti.

(a) *Product.* Gum ghatti.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.7339 Guar gum.

(a) *Product.* Guar gum.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.7343 Locust bean gum.

(a) *Product.* Locust (carob) bean gum.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.7349 Sterculia gum.

(a) *Product.* Sterculia gum (karaya gum).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.7351 Gum tragacanth.

(a) *Product.* Tragacanth (gum tragacanth).

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.7610 Potassium alginate.

(a) *Product.* Potassium alginate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

§ 582.7724 Sodium alginate.

(a) *Product.* Sodium alginate.

(b) *Conditions of use.* This substance is generally recognized as safe when used in accordance with good manufacturing or feeding practice.

PART 584—FOOD SUBSTANCES AFFIRMED AS GENERALLY RECOGNIZED AS SAFE IN FEED AND DRINKING WATER OF ANIMALS

Subpart A [Reserved]**Subpart B—Listing of Specific Substances Affirmed as GRAS**

Sec.

584.200 Ethyl alcohol containing ethyl acetate.

584.700 Hydrophobic silicas.

AUTHORITY: 21 U.S.C. 321, 342, 348, 371.

Subpart A [Reserved]**Subpart B—Listing of Specific Substances Affirmed as GRAS****§ 584.200 Ethyl alcohol containing ethyl acetate.**

The feed additive ethyl alcohol containing ethyl acetate meets the requirement of 27 CFR 212.45, being not less than 92.5 percent ethyl alcohol, each 100 gallons having had added the

§ 584.700

equivalent of 4.25 gallons of 100 percent ethyl acetate. It is used in accordance with good feeding practices in ruminant feed supplements as a source of added energy.

[46 FR 52333, Oct. 27, 1981]

§ 584.700 Hydrophobic silicas.

(a) *Product.* Amorphous fumed hydrophobic silica or precipitated hydrophobic silica (CAS Reg. No. 68611-0944-099, silane, dichlorodimethyl-, reaction products with silica).

(b) *Conditions of use.* An anticaking/free-flow agent in vitamin preparations for animal feed.

(c) *Limitations.* Not to exceed 5 percent in the vitamin preparation. It shall be used in accordance with good manufacturing or feeding practices. It must be of purity suitable for intended use, and it must comply with the following specifications:

(i) Amorphous fumed hydrophobic silica: Not less than 99.0 percent silicon dioxide after ignition. Not more than 3 ppm arsenic. Not more than 0.003 percent heavy metals (as lead). Not more than 10 ppm lead. Not more than 2.5 percent loss on drying. Not more than 2 percent loss on ignition after drying. Not more than 1 percent insoluble substances. Not more than 50 parts per million dichlorodimethylsilane.

(ii) Precipitated hydrophobic silica: Not less than 94.0 percent silicon dioxide after ignition. Not more than 3 ppm arsenic. Not more than 0.003 percent heavy metals (as lead). Not more than 10 ppm lead. Not more than 7 percent loss on drying. Not more than 8.5 percent loss on ignition after drying. Not more than 5 percent soluble ionizable salts (as sodium sulfate). Not more than 1 percent insoluble substances. Not more than 50 parts per million dichlorodimethylsilane.

[61 FR 43453, Aug. 23, 1996]

PART 589—SUBSTANCES PROHIBITED FROM USE IN ANIMAL FOOD OR FEED

Subpart A—General Provisions

Sec.

589.1 Substances prohibited from use in animal food or feed.

21 CFR Ch. I (4–1–99 Edition)

Subpart B—Listing of Specific Substances Prohibited From Use in Animal Food or Feed

589.1000 Gentian violet.

589.1001 Propylene glycol in or on cat food.

589.2000 Animal proteins prohibited in ruminant feed.

AUTHORITY: 21 U.S.C. 321, 342, 343, 348, 371.

Subpart A—General Provisions

§ 589.1 Substances prohibited from use in animal food or feed.

(a) The substances listed in this part have been prohibited from use in animal food or feed by the Food and Drug Administration because of a determination that they present a potential risk to the public health or have not been shown by adequate scientific data to be safe for use in such food or feed. Use of any of these substances in violation of this part causes the animal food or feed involved to be adulterated and in violation of the Act.

(b) This part includes only a partial list of substances prohibited from use in animal food or feed; it is for easy reference purposes and is not a complete list of substances that may not lawfully be used in such animal food or feed. No substance may be used in animal food or feed unless it meets all applicable requirements of the Act.

(c) The Food and Drug Administration either on its own initiative or on behalf of any interested person who has submitted a petition, may publish a proposal to establish, amend, or repeal a regulation under this part on the basis of new scientific evaluation or information. Any such petition shall include an adequate scientific basis to support the petition, shall be the form set forth in § 571.1 of this chapter, and will be published in the FEDERAL REGISTER for comment if it contains reasonable ground.

[45 FR 28319, Apr. 29, 1980]

Subpart B—Listing of Specific Substances Prohibited From Use in Animal Food or Feed

§ 589.1000 Gentian violet.

The Food and Drug Administration has determined that gentian violet has

not been shown by adequate scientific data to be safe for use in animal feed. Use of gentian violet in animal feed causes the feed to be adulterated and in violation of the Federal Food, Drug, and Cosmetic Act (the act), in the absence of a regulation providing for its safe use as a food additive under section 409 of the act, unless it is subject to an effective notice of claimed investigational exemption for a food additive under § 570.17 of this chapter, or unless the substance is intended for use as a new animal drug and is subject to an approved application under section 512 of the act or an effective notice of claimed investigational exemption for a new animal drug under part 511 of this chapter.

[56 FR 40507, Aug. 15, 1991]

§ 589.1001 Propylene glycol in or on cat food.

The Food and Drug Administration has determined that propylene glycol in or on cat food has not been shown by adequate scientific data to be safe for use. Use of propylene glycol in or on cat food causes the feed to be adulterated and in violation of the Federal Food, Drug, and Cosmetic Act (the act), in the absence of a regulation providing for its safe use as a food additive under section 409 of the act, unless it is subject to an effective notice of claimed investigational exemption for a food additive under § 570.17 of this chapter, or unless the substance is intended for use as a new animal drug and is subject to an approved application under section 512 of the act or an effective notice of claimed investigational exemption for a new animal drug under part 511 of this chapter.

[61 FR 19544, May 2, 1996]

§ 589.2000 Animal proteins prohibited in ruminant feed.

(a) *Definitions*—(1) *Protein derived from mammalian tissues* means any protein-containing portion of mammalian animals, excluding: Blood and blood products; gelatin; inspected meat products which have been cooked and offered for human food and further heat processed for feed (such as plate waste and used cellulosic food casings); milk products (milk and milk proteins); and any prod-

uct whose only mammalian protein consists entirely of porcine or equine protein.

(2) *Renderer* means any firm or individual that processes slaughter byproducts, animals unfit for human consumption, or meat scraps. The term includes persons who collect such materials and subject them to minimal processing, or distribute them to firms other than renderers (as defined here) whose intended use for the products may include animal feed. The term includes renderers that also blend animal protein products.

(3) *Blender* means any firm or individual which obtains processed animal protein from more than one source or from more than one species, and subsequently mixes (blends) or redistributes an animal protein product.

(4) *Feed manufacturer* includes manufacturers of complete and intermediate feeds intended for animals, and includes on-farm in addition to off-farm feed manufacturing and mixing operations.

(5) *Nonmammalian protein* includes proteins from nonmammalian animals.

(6) *Distributor* includes persons who distribute or transport feeds or feed ingredients intended for animals.

(7) *Ruminant* includes any member of the order of animals which has a stomach with four chambers (rumen, reticulum, omasum, and abomasum) through which feed passes in digestion. The order includes, but is not limited to, cattle, buffalo, sheep, goats, deer, elk, and antelopes.

(b) *Food additive status*. The Food and Drug Administration has determined that protein derived from mammalian tissues for use in ruminant feed is a food additive subject to section 409 of the Federal Food, Drug, and Cosmetic Act (the act). The use or intended use in ruminant feed of any material that contains protein derived from mammalian tissues causes the feed to be adulterated and in violation of the act, unless it is the subject of an effective notice of claimed investigational exemption for a food additive under § 570.17 of this chapter.

(c) *Requirements for renderers that are not included in paragraph (e) of this section*. (1) Renderers that manufacture products that contain or may contain

protein derived from mammalian tissues and that are intended for use in animal feed shall take the following measures to ensure that materials identified in paragraph (b) of this section are not used in the feed of ruminants:

(i) Label the materials as follows: “Do not feed to cattle or other ruminants”; and

(ii) Maintain records sufficient to track the materials throughout their receipt, processing, and distribution, and make the copies available for inspection and copying by the Food and Drug Administration.

(2) Renderers described in paragraph (c)(1) of this section will be exempted from the requirements of paragraphs (c)(1)(i) and (c)(1)(ii) of this section if they:

(i) Use exclusively a manufacturing method that has been validated by the Food and Drug Administration to deactivate the agent that causes transmissible spongiform encephalopathy (TSE) and whose design has been made available to the public;

(ii) Use routinely a test method that has been validated by the Food and Drug Administration to detect the presence of the agent that causes TSE's and whose design has been made available to the public. Renderers whose products test positive for agents that cause TSE's must comply with paragraphs (c)(1)(i) and (c)(1)(ii) of this section. Records of the test results shall be made available for inspection by the Food and Drug Administration; or

(iii) Use exclusively a method for controlling the manufacturing process that minimizes the risk of the TSE agent entering the product and whose design has been made available to the public and validated by the Food and Drug Administration.

(3) Renderers described in paragraph (c)(1) of this section will be exempted from the requirements of paragraph (c)(1)(ii) of this section if they use a permanent method, approved by FDA, to make a mark indicating that the product contains or may contain protein derived from mammalian tissue. If the marking is by the use of an agent that cannot be detected on visual inspection, the renderer must use an

agent whose presence can be detected by a method that has been validated by the Food and Drug Administration and whose design has been made available to the public.

(d) *Requirements for protein blenders, feed manufacturers, and distributors that are not included in paragraph (e) of this section.* (1) Protein blenders, feed manufacturers, and distributors that manufacture, blend, process, and distribute products that contain or may contain protein derived from mammalian tissues shall comply with paragraph (c)(1) of this section.

(2) Protein blenders, feed manufacturers, and distributors, shall be exempt from paragraphs (d)(1) of this section if they:

(i) Purchase animal products from renderers that certified compliance with paragraph (c)(2) of this section or purchase such materials from parties that certify that the materials were purchased from renderers that certified compliance with paragraph (c)(2) of this section; or

(ii) Comply with the requirements of paragraph (c)(2) of this section where appropriate.

(3) Protein blenders, feed manufacturers, and distributors, shall be exempt from paragraph (c)(1)(ii) of this section if they:

(i) Purchase animal protein products that are marked in accordance with paragraph (c)(3) of this section or purchase such materials from renderers that certified compliance with paragraph (c)(3) of this section, or purchase such materials from parties that certify that the materials were purchased from renderers that certified compliance with paragraph (c)(3) of this section; or

(ii) Comply with the requirements of paragraph (c)(3) of this section where appropriate.

(4) Pet food products that are sold or are intended for sale at retail and feeds for nonruminant laboratory animals are exempt from the labeling requirements in paragraphs (c) and (d) of this section. However, if the pet food products or feeds for nonruminant laboratory animals are sold or are intended for sale as distressed or salvage items, then such products shall be labeled in

accordance with paragraph (c) or (d) of this section, as appropriate.

(5) Copies of certifications as described in paragraphs (d)(2) and (d)(3) of this section, shall be made available for inspection and copying by the Food and Drug Administration.

(e) *Requirements for persons that intend to separate mammalian and nonmammalian materials.* (1) Renderers, protein blenders, feed manufacturers, distributors, and others that manufacture, process, blend and distribute both products that contain or may contain protein derived from mammalian tissues or feeds containing such products, and protein products from other animal tissues or feeds containing such products, and that intend to keep those products separate shall:

(i) Comply with paragraphs (c)(1) or (d)(1) of this section as appropriate except that the labeling requirement shall apply only to products that contain or may contain protein derived from mammalian tissues or feeds containing such products;

(ii) In the case of a renderer, obtain nonmammalian or pure porcine or pure equine materials only from single-species slaughter facilities;

(iii) Provide for measures to avoid commingling or cross-contamination;

(A) Maintain separate equipment or facilities for the manufacture, processing, or blending of such materials; or

(B) Use clean-out procedures or other means adequate to prevent carry-over of products that contain or may contain protein derived from mammalian tissues into animal protein or feeds that may be used for ruminants; and

(iv) Maintain written procedures specifying the clean-out procedures or other means, and specifying the procedures for separating products that contain or may contain protein derived from mammalian tissue from all other protein products from the time of receipt until the time of shipment.

(2) Renderers, blenders, feed manufacturers, and distributors will be ex-

empted from applicable requirements of paragraph (e)(1) of this section, if they meet the criteria for exemption under paragraphs (c)(2) or (c)(3) of this section, and (d)(2) or (d)(3) of this section.

(f) *Requirements for establishments and individuals that are responsible for feeding ruminant animals.* Establishments and individuals that are responsible for feeding ruminant animals shall maintain copies of purchase invoices and labeling for all feeds containing animal protein products received, and make the copies available for inspection and copying by the Food and Drug Administration.

(g) *Adulteration and misbranding.* (1) Animal protein products, and feeds containing such products, that are not in compliance with paragraphs (c) through (f) of this section, excluding labeling requirements, will be deemed adulterated under section 402(a)(2)(C) or 402(a)(4) of the act.

(2) Animal protein products, and feeds containing such products, that are not in compliance with the labeling requirements of paragraphs (c) through (f) of this section will be deemed misbranded under section 403(a)(1) or 403(f) of the act.

(h) *Inspection; records retention.* (1) Records that are to be made available for inspection and copying, as required by this section, shall be kept for a minimum of 1 year.

(2) Written procedures required by this section shall be made available for inspection and copying by the Food and Drug Administration.

[62 FR 30976, June 5, 1997]

EFFECTIVE DATE NOTE: At 62 FR 30976, June 5, 1997, §589.2000 was added. Paragraph (e)(1)(iv) of this section contains information collection and recordkeeping requirements and will not become effective until approval has been given by the Office of Management and Budget.

PARTS 590–599 [RESERVED]

FINDING AIDS

A list of CFR titles, subtitles, chapters, subchapters and parts and an alphabetical list of agencies publishing in the CFR are included in the CFR Index and Finding Aids volume to the Code of Federal Regulations which is published separately and revised annually.

Material Approved for Incorporation by Reference
Table of CFR Titles and Chapters
Alphabetical List of Agencies Appearing in the CFR
Redesignation Table
List of CFR Sections Affected

Material Approved for Incorporation by Reference

(Revised as of April 1, 1999)

The Director of the Federal Register has approved under 5 U.S.C. 552(a) and 1 CFR Part 51 the incorporation by reference of the following publications. This list contains only those incorporations by reference effective as of the revision date of this volume. Incorporations by reference found within a regulation are effective upon the effective date of that regulation. For more information on incorporation by reference, see the preliminary pages of this volume.

21 CFR (PARTS 500 TO 599)

FOOD AND DRUG ADMINISTRATION, DEPARTMENT OF HEALTH AND HUMAN SERVICES

21 CFR

Academic Press, Inc.

111 Fifth Ave., New York, NY 10003

Method published in "Methods in Hormone Research," New York, 556.710(b)
Academic Press, Volume II, p. 286 (1962).

AOAC International (Association of Official Analytical Chemists)

481 N. Frederick Ave., Suite 500, Gaithersburg, MD 20877-2417

Telephone: (301) 924-7077

Official Methods of Analysis, 13th Ed., 1980 520.2320;
573.640(b)(4)(i)
and (ii)

NOTE: The following materials are available through the Food
and Drug Administration at the addresses indicated.

Center for Food Safety and Applied Nutrition, Food and Drug Administration

200 C St. SW., Washington, DC 20204

Journal of the Association of Official Analytical Chemists:

"Oils, Fats, and Waxes" Vol. 44(1) 146 (1961) 573.640(b)(4)(ii)

Weight Method of E. J. Emberg, G. H. Gass and J. M. Curtis, published 556.250(b)
in "Endocrinology", Vol. 63, p. 806 (1958).

Dockets Management Branch (HFA-305), Food and Drug Administration

5600 Fishers Lane, Room 4-62, Rockville, MD 20857

Determination of Factor Content in Tylosin by High Performance 520.2640; 522.2640a;
Liquid Chromatography. 522.2640b;
524.2640; 558.625;
558.630

Table of CFR Titles and Chapters

(Revised as of March 31, 1999)

Title 1—General Provisions

- I Administrative Committee of the Federal Register (Parts 1—49)
- II Office of the Federal Register (Parts 50—299)
- IV Miscellaneous Agencies (Parts 400—500)

Title 2—[Reserved]

Title 3—The President

- I Executive Office of the President (Parts 100—199)

Title 4—Accounts

- I General Accounting Office (Parts 1—99)
- II Federal Claims Collection Standards (General Accounting Office—Department of Justice) (Parts 100—299)

Title 5—Administrative Personnel

- I Office of Personnel Management (Parts 1—1199)
- II Merit Systems Protection Board (Parts 1200—1299)
- III Office of Management and Budget (Parts 1300—1399)
- IV Advisory Committee on Federal Pay (Parts 1400—1499)
- V The International Organizations Employees Loyalty Board (Parts 1500—1599)
- VI Federal Retirement Thrift Investment Board (Parts 1600—1699)
- VII Advisory Commission on Intergovernmental Relations (Parts 1700—1799)
- VIII Office of Special Counsel (Parts 1800—1899)
- IX Appalachian Regional Commission (Parts 1900—1999)
- XI Armed Forces Retirement Home (Part 2100)
- XIV Federal Labor Relations Authority, General Counsel of the Federal Labor Relations Authority and Federal Service Impasses Panel (Parts 2400—2499)
- XV Office of Administration, Executive Office of the President (Parts 2500—2599)
- XVI Office of Government Ethics (Parts 2600—2699)
- XXI Department of the Treasury (Parts 3100—3199)

Title 5—Administrative Personnel—Continued

Chap.	
XXII	Federal Deposit Insurance Corporation (Part 3201)
XXIII	Department of Energy (Part 3301)
XXIV	Federal Energy Regulatory Commission (Part 3401)
XXV	Department of the Interior (Part 3501)
XXVI	Department of Defense (Part 3601)
XXVIII	Department of Justice (Part 3801)
XXIX	Federal Communications Commission (Parts 3900—3999)
XXX	Farm Credit System Insurance Corporation (Parts 4000—4099)
XXXI	Farm Credit Administration (Parts 4100—4199)
XXXIII	Overseas Private Investment Corporation (Part 4301)
XXXV	Office of Personnel Management (Part 4501)
XL	Interstate Commerce Commission (Part 5001)
XLI	Commodity Futures Trading Commission (Part 5101)
XLII	Department of Labor (Part 5201)
XLIII	National Science Foundation (Part 5301)
XLV	Department of Health and Human Services (Part 5501)
XLVI	Postal Rate Commission (Part 5601)
XLVII	Federal Trade Commission (Part 5701)
XLVIII	Nuclear Regulatory Commission (Part 5801)
L	Department of Transportation (Part 6001)
LII	Export-Import Bank of the United States (Part 6201)
LIII	Department of Education (Parts 6300—6399)
LIV	Environmental Protection Agency (Part 6401)
LVII	General Services Administration (Part 6701)
LVIII	Board of Governors of the Federal Reserve System (Part 6801)
LIX	National Aeronautics and Space Administration (Part 6901)
LX	United States Postal Service (Part 7001)
LXI	National Labor Relations Board (Part 7101)
LXII	Equal Employment Opportunity Commission (Part 7201)
LXIII	Inter-American Foundation (Part 7301)
LXV	Department of Housing and Urban Development (Part 7501)
LXVI	National Archives and Records Administration (Part 7601)
LXIX	Tennessee Valley Authority (Part 7901)
LXXI	Consumer Product Safety Commission (Part 8101)
LXXIV	Federal Mine Safety and Health Review Commission (Part 8401)
LXXVI	Federal Retirement Thrift Investment Board (Part 8601)
LXXVII	Office of Management and Budget (Part 8701)

Title 6—[Reserved]

Title 7—Agriculture

SUBTITLE A—OFFICE OF THE SECRETARY OF AGRICULTURE (PARTS 0—26)

SUBTITLE B—REGULATIONS OF THE DEPARTMENT OF AGRICULTURE

Title 7—Agriculture—Continued

Chap.	
I	Agricultural Marketing Service (Standards, Inspections, Marketing Practices), Department of Agriculture (Parts 27—209)
II	Food and Nutrition Service, Department of Agriculture (Parts 210—299)
III	Animal and Plant Health Inspection Service, Department of Agriculture (Parts 300—399)
IV	Federal Crop Insurance Corporation, Department of Agriculture (Parts 400—499)
V	Agricultural Research Service, Department of Agriculture (Parts 500—599)
VI	Natural Resources Conservation Service, Department of Agriculture (Parts 600—699)
VII	Farm Service Agency, Department of Agriculture (Parts 700—799)
VIII	Grain Inspection, Packers and Stockyards Administration (Federal Grain Inspection Service), Department of Agriculture (Parts 800—899)
IX	Agricultural Marketing Service (Marketing Agreements and Orders; Fruits, Vegetables, Nuts), Department of Agriculture (Parts 900—999)
X	Agricultural Marketing Service (Marketing Agreements and Orders; Milk), Department of Agriculture (Parts 1000—1199)
XI	Agricultural Marketing Service (Marketing Agreements and Orders; Miscellaneous Commodities), Department of Agriculture (Parts 1200—1299)
XIII	Northeast Dairy Compact Commission (Parts 1300—1399)
XIV	Commodity Credit Corporation, Department of Agriculture (Parts 1400—1499)
XV	Foreign Agricultural Service, Department of Agriculture (Parts 1500—1599)
XVI	Rural Telephone Bank, Department of Agriculture (Parts 1600—1699)
XVII	Rural Utilities Service, Department of Agriculture (Parts 1700—1799)
XVIII	Rural Housing Service, Rural Business-Cooperative Service, Rural Utilities Service, and Farm Service Agency, Department of Agriculture (Parts 1800—2099)
XXVI	Office of Inspector General, Department of Agriculture (Parts 2600—2699)
XXVII	Office of Information Resources Management, Department of Agriculture (Parts 2700—2799)
XXVIII	Office of Operations, Department of Agriculture (Parts 2800—2899)
XXIX	Office of Energy, Department of Agriculture (Parts 2900—2999)
XXX	Office of the Chief Financial Officer, Department of Agriculture (Parts 3000—3099)
XXXI	Office of Environmental Quality, Department of Agriculture (Parts 3100—3199)
XXXII	Office of Procurement and Property Management, Department of Agriculture (Parts 3200—3299)

Title 7—Agriculture—Continued	
Chap.	
XXXIII	Office of Transportation, Department of Agriculture (Parts 3300—3399)
XXXIV	Cooperative State Research, Education, and Extension Service, Department of Agriculture (Parts 3400—3499)
XXXV	Rural Housing Service, Department of Agriculture (Parts 3500—3599)
XXXVI	National Agricultural Statistics Service, Department of Agriculture (Parts 3600—3699)
XXXVII	Economic Research Service, Department of Agriculture (Parts 3700—3799)
XXXVIII	World Agricultural Outlook Board, Department of Agriculture (Parts 3800—3899)
XLI	[Reserved]
XLII	Rural Business-Cooperative Service and Rural Utilities Service, Department of Agriculture (Parts 4200—4299)

Title 8—Aliens and Nationality

I	Immigration and Naturalization Service, Department of Justice (Parts 1—499)

Title 9—Animals and Animal Products

I	Animal and Plant Health Inspection Service, Department of Agriculture (Parts 1—199)
II	Grain Inspection, Packers and Stockyards Administration (Packers and Stockyards Programs), Department of Agriculture (Parts 200—299)
III	Food Safety and Inspection Service, Department of Agriculture (Parts 300—599)

Title 10—Energy

I	Nuclear Regulatory Commission (Parts 0—199)
II	Department of Energy (Parts 200—699)
III	Department of Energy (Parts 700—999)
X	Department of Energy (General Provisions) (Parts 1000—1099)
XVII	Defense Nuclear Facilities Safety Board (Parts 1700—1799)

Title 11—Federal Elections

I	Federal Election Commission (Parts 1—9099)

Title 12—Banks and Banking

I	Comptroller of the Currency, Department of the Treasury (Parts 1—199)
II	Federal Reserve System (Parts 200—299)
III	Federal Deposit Insurance Corporation (Parts 300—399)

Chap. **Title 12—Banks and Banking—Continued**

- IV Export-Import Bank of the United States (Parts 400—499)
- V Office of Thrift Supervision, Department of the Treasury (Parts 500—599)
- VI Farm Credit Administration (Parts 600—699)
- VII National Credit Union Administration (Parts 700—799)
- VIII Federal Financing Bank (Parts 800—899)
- IX Federal Housing Finance Board (Parts 900—999)
- XI Federal Financial Institutions Examination Council (Parts 1100—1199)
- XIV Farm Credit System Insurance Corporation (Parts 1400—1499)
- XV Department of the Treasury (Parts 1500—1599)
- XVII Office of Federal Housing Enterprise Oversight, Department of Housing and Urban Development (Parts 1700—1799)
- XVIII Community Development Financial Institutions Fund, Department of the Treasury (Parts 1800—1899)

Title 13—Business Credit and Assistance

- I Small Business Administration (Parts 1—199)
- III Department of Commerce, Economic Development Administration, (Parts 300—399)

Title 14—Aeronautics and Space

- I Federal Aviation Administration, Department of Transportation (Parts 1—199)
- II Office of the Secretary, Department of Transportation (Aviation Proceedings) (Parts 200—399)
- III Commercial Space Transportation, Federal Aviation Administration, Department of Transportation (Parts 400—499)
- V National Aeronautics and Space Administration (Parts 1200—1299)

Title 15—Commerce and Foreign Trade

SUBTITLE A—OFFICE OF THE SECRETARY OF COMMERCE (PARTS 0—29)

SUBTITLE B—REGULATIONS RELATING TO COMMERCE AND FOREIGN TRADE

- I Bureau of the Census, Department of Commerce (Parts 30—199)
- II National Institute of Standards and Technology, Department of Commerce (Parts 200—299)
- III International Trade Administration, Department of Commerce (Parts 300—399)
- IV Foreign-Trade Zones Board, Department of Commerce (Parts 400—499)
- VII Bureau of Export Administration, Department of Commerce (Parts 700—799)

Title 15—Commerce and Foreign Trade—Continued

Chap.

- VIII Bureau of Economic Analysis, Department of Commerce (Parts 800—899)
- IX National Oceanic and Atmospheric Administration, Department of Commerce (Parts 900—999)
- XI Technology Administration, Department of Commerce (Parts 1100—1199)
- XIII East-West Foreign Trade Board (Parts 1300—1399)
- XIV Minority Business Development Agency (Parts 1400—1499)
SUBTITLE C—REGULATIONS RELATING TO FOREIGN TRADE AGREEMENTS
- XX Office of the United States Trade Representative (Parts 2000—2099)
SUBTITLE D—REGULATIONS RELATING TO TELECOMMUNICATIONS AND INFORMATION
- XXIII National Telecommunications and Information Administration, Department of Commerce (Parts 2300—2399)

Title 16—Commercial Practices

- I Federal Trade Commission (Parts 0—999)
- II Consumer Product Safety Commission (Parts 1000—1799)

Title 17—Commodity and Securities Exchanges

- I Commodity Futures Trading Commission (Parts 1—199)
- II Securities and Exchange Commission (Parts 200—399)
- IV Department of the Treasury (Parts 400—499)

Title 18—Conservation of Power and Water Resources

- I Federal Energy Regulatory Commission, Department of Energy (Parts 1—399)
- III Delaware River Basin Commission (Parts 400—499)
- VI Water Resources Council (Parts 700—799)
- VIII Susquehanna River Basin Commission (Parts 800—899)
- XIII Tennessee Valley Authority (Parts 1300—1399)

Title 19—Customs Duties

- I United States Customs Service, Department of the Treasury (Parts 1—199)
- II United States International Trade Commission (Parts 200—299)
- III International Trade Administration, Department of Commerce (Parts 300—399)

	Title 20—Employees' Benefits
Chap.	
I	Office of Workers' Compensation Programs, Department of Labor (Parts 1—199)
II	Railroad Retirement Board (Parts 200—399)
III	Social Security Administration (Parts 400—499)
IV	Employees' Compensation Appeals Board, Department of Labor (Parts 500—599)
V	Employment and Training Administration, Department of Labor (Parts 600—699)
VI	Employment Standards Administration, Department of Labor (Parts 700—799)
VII	Benefits Review Board, Department of Labor (Parts 800—899)
VIII	Joint Board for the Enrollment of Actuaries (Parts 900—999)
IX	Office of the Assistant Secretary for Veterans' Employment and Training, Department of Labor (Parts 1000—1099)

Title 21—Food and Drugs

I	Food and Drug Administration, Department of Health and Human Services (Parts 1—1299)
II	Drug Enforcement Administration, Department of Justice (Parts 1300—1399)
III	Office of National Drug Control Policy (Parts 1400—1499)

Title 22—Foreign Relations

I	Department of State (Parts 1—199)
II	Agency for International Development (Parts 200—299)
III	Peace Corps (Parts 300—399)
IV	International Joint Commission, United States and Canada (Parts 400—499)
V	United States Information Agency (Parts 500—599)
VII	Overseas Private Investment Corporation, International Development Cooperation Agency (Parts 700—799)
IX	Foreign Service Grievance Board Regulations (Parts 900—999)
X	Inter-American Foundation (Parts 1000—1099)
XI	International Boundary and Water Commission, United States and Mexico, United States Section (Parts 1100—1199)
XII	United States International Development Cooperation Agency (Parts 1200—1299)
XIII	Board for International Broadcasting (Parts 1300—1399)
XIV	Foreign Service Labor Relations Board; Federal Labor Relations Authority; General Counsel of the Federal Labor Relations Authority; and the Foreign Service Impasse Disputes Panel (Parts 1400—1499)
XV	African Development Foundation (Parts 1500—1599)
XVI	Japan-United States Friendship Commission (Parts 1600—1699)
XVII	United States Institute of Peace (Parts 1700—1799)

Chap.

Title 23—Highways

- I Federal Highway Administration, Department of Transportation (Parts 1—999)
- II National Highway Traffic Safety Administration and Federal Highway Administration, Department of Transportation (Parts 1200—1299)
- III National Highway Traffic Safety Administration, Department of Transportation (Parts 1300—1399)

Title 24—Housing and Urban Development

SUBTITLE A—OFFICE OF THE SECRETARY, DEPARTMENT OF HOUSING AND URBAN DEVELOPMENT (PARTS 0—99)

SUBTITLE B—REGULATIONS RELATING TO HOUSING AND URBAN DEVELOPMENT

- I Office of Assistant Secretary for Equal Opportunity, Department of Housing and Urban Development (Parts 100—199)
- II Office of Assistant Secretary for Housing-Federal Housing Commissioner, Department of Housing and Urban Development (Parts 200—299)
- III Government National Mortgage Association, Department of Housing and Urban Development (Parts 300—399)
- IV Office of Multifamily Housing Assistance Restructuring, Department of Housing and Urban Development (Parts 400—499)
- V Office of Assistant Secretary for Community Planning and Development, Department of Housing and Urban Development (Parts 500—599)
- VI Office of Assistant Secretary for Community Planning and Development, Department of Housing and Urban Development (Parts 600—699) [Reserved]
- VII Office of the Secretary, Department of Housing and Urban Development (Housing Assistance Programs and Public and Indian Housing Programs) (Parts 700—799)
- VIII Office of the Assistant Secretary for Housing—Federal Housing Commissioner, Department of Housing and Urban Development (Section 8 Housing Assistance Programs, Section 202 Direct Loan Program, Section 202 Supportive Housing for the Elderly Program and Section 811 Supportive Housing for Persons With Disabilities Program) (Parts 800—899)
- IX Office of Assistant Secretary for Public and Indian Housing, Department of Housing and Urban Development (Parts 900—999)
- X Office of Assistant Secretary for Housing—Federal Housing Commissioner, Department of Housing and Urban Development (Interstate Land Sales Registration Program) (Parts 1700—1799)
- XII Office of Inspector General, Department of Housing and Urban Development (Parts 2000—2099)
- XX Office of Assistant Secretary for Housing—Federal Housing Commissioner, Department of Housing and Urban Development (Parts 3200—3899)
- XXV Neighborhood Reinvestment Corporation (Parts 4100—4199)

Chap.

Title 25—Indians

- I Bureau of Indian Affairs, Department of the Interior (Parts 1—299)
- II Indian Arts and Crafts Board, Department of the Interior (Parts 300—399)
- III National Indian Gaming Commission, Department of the Interior (Parts 500—599)
- IV Office of Navajo and Hopi Indian Relocation (Parts 700—799)
- V Bureau of Indian Affairs, Department of the Interior, and Indian Health Service, Department of Health and Human Services (Part 900)
- VI Office of the Assistant Secretary-Indian Affairs, Department of the Interior (Part 1001)
- VII Office of the Special Trustee for American Indians, Department of the Interior (Part 1200)

Title 26—Internal Revenue

- I Internal Revenue Service, Department of the Treasury (Parts 1—799)

Title 27—Alcohol, Tobacco Products and Firearms

- I Bureau of Alcohol, Tobacco and Firearms, Department of the Treasury (Parts 1—299)

Title 28—Judicial Administration

- I Department of Justice (Parts 0—199)
- III Federal Prison Industries, Inc., Department of Justice (Parts 300—399)
- V Bureau of Prisons, Department of Justice (Parts 500—599)
- VI Offices of Independent Counsel, Department of Justice (Parts 600—699)
- VII Office of Independent Counsel (Parts 700—799)

Title 29—Labor

SUBTITLE A—OFFICE OF THE SECRETARY OF LABOR (PARTS 0—99)

SUBTITLE B—REGULATIONS RELATING TO LABOR

- I National Labor Relations Board (Parts 100—199)
- II Office of Labor-Management Standards, Department of Labor (Parts 200—299)
- III National Railroad Adjustment Board (Parts 300—399)
- IV Office of Labor-Management Standards, Department of Labor (Parts 400—499)
- V Wage and Hour Division, Department of Labor (Parts 500—899)
- IX Construction Industry Collective Bargaining Commission (Parts 900—999)
- X National Mediation Board (Parts 1200—1299)

Title 29—Labor—Continued

Chap.	
XII	Federal Mediation and Conciliation Service (Parts 1400—1499)
XIV	Equal Employment Opportunity Commission (Parts 1600—1699)
XVII	Occupational Safety and Health Administration, Department of Labor (Parts 1900—1999)
XX	Occupational Safety and Health Review Commission (Parts 2200—2499)
XXV	Pension and Welfare Benefits Administration, Department of Labor (Parts 2500—2599)
XXVII	Federal Mine Safety and Health Review Commission (Parts 2700—2799)
XL	Pension Benefit Guaranty Corporation (Parts 4000—4999)

Title 30—Mineral Resources

I	Mine Safety and Health Administration, Department of Labor (Parts 1—199)
II	Minerals Management Service, Department of the Interior (Parts 200—299)
III	Board of Surface Mining and Reclamation Appeals, Department of the Interior (Parts 300—399)
IV	Geological Survey, Department of the Interior (Parts 400—499)
VI	Bureau of Mines, Department of the Interior (Parts 600—699)
VII	Office of Surface Mining Reclamation and Enforcement, Department of the Interior (Parts 700—999)

Title 31—Money and Finance: Treasury

	SUBTITLE A—OFFICE OF THE SECRETARY OF THE TREASURY (PARTS 0—50)
	SUBTITLE B—REGULATIONS RELATING TO MONEY AND FINANCE
I	Monetary Offices, Department of the Treasury (Parts 51—199)
II	Fiscal Service, Department of the Treasury (Parts 200—399)
IV	Secret Service, Department of the Treasury (Parts 400—499)
V	Office of Foreign Assets Control, Department of the Treasury (Parts 500—599)
VI	Bureau of Engraving and Printing, Department of the Treasury (Parts 600—699)
VII	Federal Law Enforcement Training Center, Department of the Treasury (Parts 700—799)
VIII	Office of International Investment, Department of the Treasury (Parts 800—899)

Title 32—National Defense

	SUBTITLE A—DEPARTMENT OF DEFENSE
I	Office of the Secretary of Defense (Parts 1—399)
V	Department of the Army (Parts 400—699)
VI	Department of the Navy (Parts 700—799)

Chap. **Title 32—National Defense—Continued**

- VII Department of the Air Force (Parts 800—1099)
- SUBTITLE B—OTHER REGULATIONS RELATING TO NATIONAL DEFENSE
- XII Defense Logistics Agency (Parts 1200—1299)
- XVI Selective Service System (Parts 1600—1699)
- XIX Central Intelligence Agency (Parts 1900—1999)
- XX Information Security Oversight Office, National Archives and Records Administration (Parts 2000—2099)
- XXI National Security Council (Parts 2100—2199)
- XXIV Office of Science and Technology Policy (Parts 2400—2499)
- XXVII Office for Micronesian Status Negotiations (Parts 2700—2799)
- XXVIII Office of the Vice President of the United States (Parts 2800—2899)
- XXIX Presidential Commission on the Assignment of Women in the Armed Forces (Part 2900)

Title 33—Navigation and Navigable Waters

- I Coast Guard, Department of Transportation (Parts 1—199)
- II Corps of Engineers, Department of the Army (Parts 200—399)
- IV Saint Lawrence Seaway Development Corporation, Department of Transportation (Parts 400—499)

Title 34—Education

- SUBTITLE A—OFFICE OF THE SECRETARY, DEPARTMENT OF EDUCATION (PARTS 1—99)
- SUBTITLE B—REGULATIONS OF THE OFFICES OF THE DEPARTMENT OF EDUCATION
- I Office for Civil Rights, Department of Education (Parts 100—199)
- II Office of Elementary and Secondary Education, Department of Education (Parts 200—299)
- III Office of Special Education and Rehabilitative Services, Department of Education (Parts 300—399)
- IV Office of Vocational and Adult Education, Department of Education (Parts 400—499)
- V Office of Bilingual Education and Minority Languages Affairs, Department of Education (Parts 500—599)
- VI Office of Postsecondary Education, Department of Education (Parts 600—699)
- VII Office of Educational Research and Improvement, Department of Education (Parts 700—799)
- XI National Institute for Literacy (Parts 1100—1199)
- SUBTITLE C—REGULATIONS RELATING TO EDUCATION
- XII National Council on Disability (Parts 1200—1299)

Chap.

Title 35—Panama Canal

- I Panama Canal Regulations (Parts 1—299)

Title 36—Parks, Forests, and Public Property

- I National Park Service, Department of the Interior (Parts 1—199)
- II Forest Service, Department of Agriculture (Parts 200—299)
- III Corps of Engineers, Department of the Army (Parts 300—399)
- IV American Battle Monuments Commission (Parts 400—499)
- V Smithsonian Institution (Parts 500—599)
- VII Library of Congress (Parts 700—799)
- VIII Advisory Council on Historic Preservation (Parts 800—899)
- IX Pennsylvania Avenue Development Corporation (Parts 900—999)
- X Presidio Trust (Parts 1000—1099)
- XI Architectural and Transportation Barriers Compliance Board (Parts 1100—1199)
- XII National Archives and Records Administration (Parts 1200—1299)
- XIV Assassination Records Review Board (Parts 1400—1499)

Title 37—Patents, Trademarks, and Copyrights

- I Patent and Trademark Office, Department of Commerce (Parts 1—199)
- II Copyright Office, Library of Congress (Parts 200—299)
- IV Assistant Secretary for Technology Policy, Department of Commerce (Parts 400—499)
- V Under Secretary for Technology, Department of Commerce (Parts 500—599)

Title 38—Pensions, Bonuses, and Veterans' Relief

- I Department of Veterans Affairs (Parts 0—99)

Title 39—Postal Service

- I United States Postal Service (Parts 1—999)
- III Postal Rate Commission (Parts 3000—3099)

Title 40—Protection of Environment

- I Environmental Protection Agency (Parts 1—799)
- V Council on Environmental Quality (Parts 1500—1599)

Title 41—Public Contracts and Property Management

SUBTITLE B—OTHER PROVISIONS RELATING TO PUBLIC CONTRACTS

- 50 Public Contracts, Department of Labor (Parts 50—1—50—999)

Title 41—Public Contracts and Property Management—Continued

Chap.

- 51 Committee for Purchase From People Who Are Blind or Severely Disabled (Parts 51–1—51–99)
- 60 Office of Federal Contract Compliance Programs, Equal Employment Opportunity, Department of Labor (Parts 60–1—60–999)
- 61 Office of the Assistant Secretary for Veterans Employment and Training, Department of Labor (Parts 61–1—61–999)
- SUBTITLE C—FEDERAL PROPERTY MANAGEMENT REGULATIONS SYSTEM
- 101 Federal Property Management Regulations (Parts 101–1—101–99)
- 105 General Services Administration (Parts 105–1—105–999)
- 109 Department of Energy Property Management Regulations (Parts 109–1—109–99)
- 114 Department of the Interior (Parts 114–1—114–99)
- 115 Environmental Protection Agency (Parts 115–1—115–99)
- 128 Department of Justice (Parts 128–1—128–99)
- SUBTITLE D—OTHER PROVISIONS RELATING TO PROPERTY MANAGEMENT [RESERVED]
- SUBTITLE E—FEDERAL INFORMATION RESOURCES MANAGEMENT REGULATIONS SYSTEM
- 201 Federal Information Resources Management Regulation (Parts 201–1—201–99) [Reserved]
- SUBTITLE F—FEDERAL TRAVEL REGULATION SYSTEM
- 300 General (Parts 300–1—300–99)
- 301 Temporary Duty (TDY) Travel Allowances (Parts 301–1—301–99)
- 302 Relocation Allowances (Parts 302–1—302–99)
- 303 Payment of Expenses Connected with the Death of Certain Employees (Parts 303–1—303–2)
- 304 Payment from a Non-Federal Source for Travel Expenses (Parts 304–1—304–99)

Title 42—Public Health

- I Public Health Service, Department of Health and Human Services (Parts 1—199)
- IV Health Care Financing Administration, Department of Health and Human Services (Parts 400—499)
- V Office of Inspector General-Health Care, Department of Health and Human Services (Parts 1000—1999)

Title 43—Public Lands: Interior

- SUBTITLE A—OFFICE OF THE SECRETARY OF THE INTERIOR (PARTS 1—199)
- SUBTITLE B—REGULATIONS RELATING TO PUBLIC LANDS
- I Bureau of Reclamation, Department of the Interior (Parts 200—499)
- II Bureau of Land Management, Department of the Interior (Parts 1000—9999)

- Chap. **Title 43—Public Lands: Interior—Continued**
- III Utah Reclamation Mitigation and Conservation Commission (Parts 10000—10005)

Title 44—Emergency Management and Assistance

- I Federal Emergency Management Agency (Parts 0—399)
- IV Department of Commerce and Department of Transportation (Parts 400—499)

Title 45—Public Welfare

- SUBTITLE A—DEPARTMENT OF HEALTH AND HUMAN SERVICES (PARTS 1—199)
- SUBTITLE B—REGULATIONS RELATING TO PUBLIC WELFARE
- II Office of Family Assistance (Assistance Programs), Administration for Children and Families, Department of Health and Human Services (Parts 200—299)
- III Office of Child Support Enforcement (Child Support Enforcement Program), Administration for Children and Families, Department of Health and Human Services (Parts 300—399)
- IV Office of Refugee Resettlement, Administration for Children and Families Department of Health and Human Services (Parts 400—499)
- V Foreign Claims Settlement Commission of the United States, Department of Justice (Parts 500—599)
- VI National Science Foundation (Parts 600—699)
- VII Commission on Civil Rights (Parts 700—799)
- VIII Office of Personnel Management (Parts 800—899)
- X Office of Community Services, Administration for Children and Families, Department of Health and Human Services (Parts 1000—1099)
- XI National Foundation on the Arts and the Humanities (Parts 1100—1199)
- XII Corporation for National and Community Service (Parts 1200—1299)
- XIII Office of Human Development Services, Department of Health and Human Services (Parts 1300—1399)
- XVI Legal Services Corporation (Parts 1600—1699)
- XVII National Commission on Libraries and Information Science (Parts 1700—1799)
- XXIII Harry S. Truman Scholarship Foundation (Parts 1800—1899)
- XXI Commission on Fine Arts (Parts 2100—2199)
- XXII Christopher Columbus Quincentenary Jubilee Commission (Parts 2200—2299)
- XXIII Arctic Research Commission (Part 2301)
- XXIV James Madison Memorial Fellowship Foundation (Parts 2400—2499)
- XXV Corporation for National and Community Service (Parts 2500—2599)

Chap.

Title 46—Shipping

- I Coast Guard, Department of Transportation (Parts 1—199)
- II Maritime Administration, Department of Transportation (Parts 200—399)
- III Coast Guard (Great Lakes Pilotage), Department of Transportation (Parts 400—499)
- IV Federal Maritime Commission (Parts 500—599)

Title 47—Telecommunication

- I Federal Communications Commission (Parts 0—199)
- II Office of Science and Technology Policy and National Security Council (Parts 200—299)
- III National Telecommunications and Information Administration, Department of Commerce (Parts 300—399)

Title 48—Federal Acquisition Regulations System

- 1 Federal Acquisition Regulation (Parts 1—99)
- 2 Department of Defense (Parts 200—299)
- 3 Department of Health and Human Services (Parts 300—399)
- 4 Department of Agriculture (Parts 400—499)
- 5 General Services Administration (Parts 500—599)
- 6 Department of State (Parts 600—699)
- 7 United States Agency for International Development (Parts 700—799)
- 8 Department of Veterans Affairs (Parts 800—899)
- 9 Department of Energy (Parts 900—999)
- 10 Department of the Treasury (Parts 1000—1099)
- 12 Department of Transportation (Parts 1200—1299)
- 13 Department of Commerce (Parts 1300—1399)
- 14 Department of the Interior (Parts 1400—1499)
- 15 Environmental Protection Agency (Parts 1500—1599)
- 16 Office of Personnel Management Federal Employees Health Benefits Acquisition Regulation (Parts 1600—1699)
- 17 Office of Personnel Management (Parts 1700—1799)
- 18 National Aeronautics and Space Administration (Parts 1800—1899)
- 19 United States Information Agency (Parts 1900—1999)
- 20 Nuclear Regulatory Commission (Parts 2000—2099)
- 21 Office of Personnel Management, Federal Employees Group Life Insurance Federal Acquisition Regulation (Parts 2100—2199)
- 23 Social Security Administration (Parts 2300—2399)
- 24 Department of Housing and Urban Development (Parts 2400—2499)
- 25 National Science Foundation (Parts 2500—2599)
- 28 Department of Justice (Parts 2800—2899)
- 29 Department of Labor (Parts 2900—2999)

Title 48—Federal Acquisition Regulations System—Continued

Chap.

- 34 Department of Education Acquisition Regulation (Parts 3400—3499)
- 35 Panama Canal Commission (Parts 3500—3599)
- 44 Federal Emergency Management Agency (Parts 4400—4499)
- 51 Department of the Army Acquisition Regulations (Parts 5100—5199)
- 52 Department of the Navy Acquisition Regulations (Parts 5200—5299)
- 53 Department of the Air Force Federal Acquisition Regulation Supplement (Parts 5300—5399)
- 54 Defense Logistics Agency, Department of Defense (Part 5452)
- 57 African Development Foundation (Parts 5700—5799)
- 61 General Services Administration Board of Contract Appeals (Parts 6100—6199)
- 63 Department of Transportation Board of Contract Appeals (Parts 6300—6399)
- 99 Cost Accounting Standards Board, Office of Federal Procurement Policy, Office of Management and Budget (Parts 9900—9999)

Title 49—Transportation

SUBTITLE A—OFFICE OF THE SECRETARY OF TRANSPORTATION
(PARTS 1—99)

SUBTITLE B—OTHER REGULATIONS RELATING TO TRANSPORTATION

- I Research and Special Programs Administration, Department of Transportation (Parts 100—199)
- II Federal Railroad Administration, Department of Transportation (Parts 200—299)
- III Federal Highway Administration, Department of Transportation (Parts 300—399)
- IV Coast Guard, Department of Transportation (Parts 400—499)
- V National Highway Traffic Safety Administration, Department of Transportation (Parts 500—599)
- VI Federal Transit Administration, Department of Transportation (Parts 600—699)
- VII National Railroad Passenger Corporation (AMTRAK) (Parts 700—799)
- VIII National Transportation Safety Board (Parts 800—999)
- X Surface Transportation Board, Department of Transportation (Parts 1000—1399)
- XI Bureau of Transportation Statistics, Department of Transportation (Parts 1400—1499)

Title 50—Wildlife and Fisheries

- I United States Fish and Wildlife Service, Department of the Interior (Parts 1—199)

Title 50—Wildlife and Fisheries—Continued	
Chap.	
II	National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Department of Commerce (Parts 200—299)
III	International Fishing and Related Activities (Parts 300—399)
IV	Joint Regulations (United States Fish and Wildlife Service, Department of the Interior and National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Department of Commerce); Endangered Species Committee Regulations (Parts 400—499)
V	Marine Mammal Commission (Parts 500—599)
VI	Fishery Conservation and Management, National Oceanic and Atmospheric Administration, Department of Commerce (Parts 600—699)

CFR Index and Finding Aids

Subject/Agency Index
 List of Agency Prepared Indexes
 Parallel Tables of Statutory Authorities and Rules
 List of CFR Titles, Chapters, Subchapters, and Parts
 Alphabetical List of Agencies Appearing in the CFR

Alphabetical List of Agencies Appearing in the CFR

(Revised as of March 31, 1999)

Agency	CFR Title, Subtitle or Chapter
Administrative Committee of the Federal Register	1, I
Advanced Research Projects Agency	32, I
Advisory Commission on Intergovernmental Relations	5, VII
Advisory Committee on Federal Pay	5, IV
Advisory Council on Historic Preservation	36, VIII
African Development Foundation	22, XV
Federal Acquisition Regulation	48, 57
Agency for International Development	22, II
Federal Acquisition Regulation	48, 7
Agricultural Marketing Service	7, I, IX, X, XI
Agricultural Research Service	7, V
Agriculture Department	
Agricultural Marketing Service	7, I, IX, X, XI
Agricultural Research Service	7, V
Animal and Plant Health Inspection Service	7, III; 9, I
Chief Financial Officer, Office of	7, XXX
Commodity Credit Corporation	7, XIV
Cooperative State Research, Education, and Extension Service	7, XXXIV
Economic Research Service	7, XXXVII
Energy, Office of	7, XXIX
Environmental Quality, Office of	7, XXXI
Farm Service Agency	7, VII, XVIII
Federal Acquisition Regulation	48, 4
Federal Crop Insurance Corporation	7, IV
Food and Nutrition Service	7, II
Food Safety and Inspection Service	9, III
Foreign Agricultural Service	7, XV
Forest Service	36, II
Grain Inspection, Packers and Stockyards Administration	7, VIII; 9, II
Information Resources Management, Office of	7, XXVII
Inspector General, Office of	7, XXVI
National Agricultural Library	7, XLI
National Agricultural Statistics Service	7, XXXVI
Natural Resources Conservation Service	7, VI
Operations, Office of	7, XXVIII
Procurement and Property Management, Office of	7, XXXII
Rural Business-Cooperative Service	7, XVIII, XLII
Rural Development Administration	7, XLII
Rural Housing Service	7, XVIII, XXXV
Rural Telephone Bank	7, XVI
Rural Utilities Service	7, XVII, XVIII, XLII
Secretary of Agriculture, Office of	7, Subtitle A
Transportation, Office of	7, XXXIII
World Agricultural Outlook Board	7, XXXVIII
Air Force Department	32, VII
Federal Acquisition Regulation Supplement	48, 53
Alcohol, Tobacco and Firearms, Bureau of	27, I
AMTRAK	49, VII
American Battle Monuments Commission	36, IV
American Indians, Office of the Special Trustee	25, VII
Animal and Plant Health Inspection Service	7, III; 9, I
Appalachian Regional Commission	5, IX

Agency	CFR Title, Subtitle or Chapter
Architectural and Transportation Barriers Compliance Board	36, XI
Arctic Research Commission	45, XXIII
Armed Forces Retirement Home	5, XI
Army Department	32, V
Engineers, Corps of	33, II; 36, III
Federal Acquisition Regulation	48, 51
Assassination Records Review Board	36, XIV
Benefits Review Board	20, VII
Bilingual Education and Minority Languages Affairs, Office of	34, V
Blind or Severely Disabled, Committee for Purchase From People Who Are	41, 51
Board for International Broadcasting	22, XIII
Census Bureau	15, I
Central Intelligence Agency	32, XIX
Chief Financial Officer, Office of	7, XXX
Child Support Enforcement, Office of	45, III
Children and Families, Administration for	45, II, III, IV, X
Christopher Columbus Quincentenary Jubilee Commission	45, XXII
Civil Rights, Commission on	45, VII
Civil Rights, Office for	34, I
Coast Guard	33, I; 46, I; 49, IV
Coast Guard (Great Lakes Pilotage)	46, III
Commerce Department	44, IV
Census Bureau	15, I'
Economic Affairs, Under Secretary	37, V
Economic Analysis, Bureau of	15, VIII
Economic Development Administration	13, III
Emergency Management and Assistance	44, IV
Export Administration, Bureau of	15, VII
Federal Acquisition Regulation	48, 13
Fishery Conservation and Management	50, VI
Foreign-Trade Zones Board	15, IV
International Trade Administration	15, III; 19, III
National Institute of Standards and Technology	15, II
National Marine Fisheries Service	50, II, IV, VI
National Oceanic and Atmospheric Administration	15, IX; 50, II, III, IV, VI
National Telecommunications and Information Administration	15, XXIII; 47, III
National Weather Service	15, IX
Patent and Trademark Office	37, I
Productivity, Technology and Innovation, Assistant Secretary for	37, IV
Secretary of Commerce, Office of	15, Subtitle A
Technology, Under Secretary for	37, V
Technology Administration	15, XI
Technology Policy, Assistant Secretary for	37, IV
Commercial Space Transportation	14, III
Commodity Credit Corporation	7, XIV
Commodity Futures Trading Commission	5, XLI; 17, I
Community Planning and Development, Office of Assistant Secretary for	24, V, VI
Community Services, Office of	45, X
Comptroller of the Currency	12, I
Construction Industry Collective Bargaining Commission	29, IX
Consumer Product Safety Commission	5, LXXI; 16, II
Cooperative State Research, Education, and Extension Service	7, XXXIV
Copyright Office	37, II
Corporation for National and Community Service	45, XII, XXV
Cost Accounting Standards Board	48, 99
Council on Environmental Quality	40, V
Customs Service, United States	19, I
Defense Contract Audit Agency	32, I
Defense Department	5, XXVI; 32, Subtitle A
Advanced Research Projects Agency	32, I
Air Force Department	32, VII

Agency	CFR Title, Subtitle or Chapter
Army Department	32, V; 33, II; 36, III, 48, 51
Defense Intelligence Agency	32, I
Defense Logistics Agency	32, I, XII; 48, 54
Engineers, Corps of	33, II; 36, III
Federal Acquisition Regulation	48, 2
National Imagery and Mapping Agency	32, I
Navy Department	32, VI; 48, 52
Secretary of Defense, Office of	32, I
Defense Contract Audit Agency	32, I
Defense Intelligence Agency	32, I
Defense Logistics Agency	32, XII; 48, 54
Defense Nuclear Facilities Safety Board	10, XVII
Delaware River Basin Commission	18, III
Drug Enforcement Administration	21, II
East-West Foreign Trade Board	15, XIII
Economic Affairs, Under Secretary	37, V
Economic Analysis, Bureau of	15, VIII
Economic Development Administration	13, III
Economic Research Service	7, XXXVII
Education, Department of	5, LIII
Bilingual Education and Minority Languages Affairs, Office of	34, V
Civil Rights, Office for	34, I
Educational Research and Improvement, Office of	34, VII
Elementary and Secondary Education, Office of	34, II
Federal Acquisition Regulation	48, 34
Postsecondary Education, Office of	34, VI
Secretary of Education, Office of	34, Subtitle A
Special Education and Rehabilitative Services, Office of	34, III
Vocational and Adult Education, Office of	34, IV
Educational Research and Improvement, Office of	34, VII
Elementary and Secondary Education, Office of	34, II
Employees' Compensation Appeals Board	20, IV
Employees Loyalty Board	5, V
Employment and Training Administration	20, V
Employment Standards Administration	20, VI
Endangered Species Committee	50, IV
Energy, Department of	5, XXIII; 10, II, III, X
Federal Acquisition Regulation	48, 9
Federal Energy Regulatory Commission	5, XXIV; 18, I
Property Management Regulations	41, 109
Energy, Office of	7, XXIX
Engineers, Corps of	33, II; 36, III
Engraving and Printing, Bureau of	31, VI
Environmental Protection Agency	5, LIV; 40, I
Federal Acquisition Regulation	48, 15
Property Management Regulations	41, 115
Environmental Quality, Office of	7, XXXI
Equal Employment Opportunity Commission	5, LXII; 29, XIV
Equal Opportunity, Office of Assistant Secretary for	24, I
Executive Office of the President	3, I
Administration, Office of	5, XV
Environmental Quality, Council on	40, V
Management and Budget, Office of	25, III, LXXVII; 48, 99
National Drug Control Policy, Office of	21, III
National Security Council	32, XXI; 47, 2
Presidential Documents	3
Science and Technology Policy, Office of	32, XXIV; 47, II
Trade Representative, Office of the United States	15, XX
Export Administration, Bureau of	15, VII
Export-Import Bank of the United States	5, LII; 12, IV
Family Assistance, Office of	45, II
Farm Credit Administration	5, XXXI; 12, VI
Farm Credit System Insurance Corporation	5, XXX; 12, XIV
Farm Service Agency	7, VII, XVIII
Federal Acquisition Regulation	48, 1

Agency	CFR Title, Subtitle or Chapter
Federal Aviation Administration	14, I
Commercial Space Transportation	14, III
Federal Claims Collection Standards	4, II
Federal Communications Commission	5, XXIX; 47, I
Federal Contract Compliance Programs, Office of	41, 60
Federal Crop Insurance Corporation	7, IV
Federal Deposit Insurance Corporation	5, XXII; 12, III
Federal Election Commission	11, I
Federal Emergency Management Agency	44, I
Federal Acquisition Regulation	48, 44
Federal Employees Group Life Insurance Federal Acquisition Regulation	48, 21
Federal Employees Health Benefits Acquisition Regulation	48, 16
Federal Energy Regulatory Commission	5, XXIV; 18, I
Federal Financial Institutions Examination Council	12, XI
Federal Financing Bank	12, VIII
Federal Highway Administration	23, I, II; 49, III
Federal Home Loan Mortgage Corporation	1, IV
Federal Housing Enterprise Oversight Office	12, XVII
Federal Housing Finance Board	12, IX
Federal Labor Relations Authority, and General Counsel of the Federal Labor Relations Authority	5, XIV; 22, XIV
Federal Law Enforcement Training Center	31, VII
Federal Maritime Commission	46, IV
Federal Mediation and Conciliation Service	29, XII
Federal Mine Safety and Health Review Commission	5, LXXIV; 29, XXVII
Federal Pay, Advisory Committee on	5, IV
Federal Prison Industries, Inc.	28, III
Federal Procurement Policy Office	48, 99
Federal Property Management Regulations	41, 101
Federal Property Management Regulations System	41, Subtitle C
Federal Railroad Administration	49, II
Federal Register, Administrative Committee of	1, I
Federal Register, Office of	1, II
Federal Reserve System	12, II
Board of Governors	5, LVIII
Federal Retirement Thrift Investment Board	5, VI, LXXVI
Federal Service Impasses Panel	5, XIV
Federal Trade Commission	5, XLVII; 16, I
Federal Transit Administration	49, VI
Federal Travel Regulation System	41, Subtitle F
Fine Arts, Commission on	45, XXI
Fiscal Service	31, II
Fish and Wildlife Service, United States	50, I, IV
Fishery Conservation and Management	50, VI
Food and Drug Administration	21, I
Food and Nutrition Service	7, II
Food Safety and Inspection Service	9, III
Foreign Agricultural Service	7, XV
Foreign Assets Control, Office of	31, V
Foreign Claims Settlement Commission of the United States	45, V
Foreign Service Grievance Board	22, IX
Foreign Service Impasse Disputes Panel	22, XIV
Foreign Service Labor Relations Board	22, XIV
Foreign-Trade Zones Board	15, IV
Forest Service	36, II
General Accounting Office	4, I, II
General Services Administration	5, LVII
Contract Appeals, Board of	48, 61
Federal Acquisition Regulation	48, 5
Federal Property Management Regulations System	41, 101, 105
Federal Travel Regulation System	41, Subtitle F
General	41, 300
Payment From a Non-Federal Source for Travel Expenses	41, 304
Payment of Expenses Connected With the Death of Certain Employees	41, 303
Relocation Allowances	41, 302

Agency	CFR Title, Subtitle or Chapter
Temporary Duty (TDY) Travel Allowances	41, 301
Geological Survey	30, IV
Government Ethics, Office of	5, XVI
Government National Mortgage Association	24, III
Grain Inspection, Packers and Stockyards Administration	7, VIII; 9, II
Harry S. Truman Scholarship Foundation	45, XVIII
Health and Human Services, Department of	5, XLV; 45, Subtitle A
Child Support Enforcement, Office of	45, III
Children and Families, Administration for	45, II, III, IV, X
Community Services, Office of	45, X
Family Assistance, Office of	45, II
Federal Acquisition Regulation	48, 3
Food and Drug Administration	21, I
Health Care Financing Administration	42, IV
Human Development Services, Office of	45, XIII
Indian Health Service	25, V
Inspector General (Health Care), Office of	42, V
Public Health Service	42, I
Refugee Resettlement, Office of	45, IV
Health Care Financing Administration	42, IV
Housing and Urban Development, Department of	5, LXV; 24, Subtitle B
Community Planning and Development, Office of Assistant Secretary for	24, V, VI
Equal Opportunity, Office of Assistant Secretary for	24, I
Federal Acquisition Regulation	48, 24
Federal Housing Enterprise Oversight, Office of	12, XVII
Government National Mortgage Association	24, III
Housing—Federal Housing Commissioner, Office of Assistant Secretary for	24, II, VIII, X, XX
Inspector General, Office of	24, XII
Multifamily Housing Assistance Restructuring, Office of	24, IV
Public and Indian Housing, Office of Assistant Secretary for	24, IX
Secretary, Office of	24, Subtitle A, VII
Housing—Federal Housing Commissioner, Office of Assistant Secretary for	24, II, VIII, X, XX
Human Development Services, Office of	45, XIII
Immigration and Naturalization Service	8, I
Independent Counsel, Office of	28, VII
Indian Affairs, Bureau of	25, I, V
Indian Affairs, Office of the Assistant Secretary	25, VI
Indian Arts and Crafts Board	25, II
Indian Health Service	25, V
Information Agency, United States	22, V
Federal Acquisition Regulation	48, 19
Information Resources Management, Office of	7, XXVII
Information Security Oversight Office, National Archives and Records Administration	32, XX
Inspector General	
Agriculture Department	7, XXVI
Health and Human Services Department	42, V
Housing and Urban Development Department	24, XII
Institute of Peace, United States	22, XVII
Inter-American Foundation	5, LXIII; 22, X
Intergovernmental Relations, Advisory Commission on	5, VII
Interior Department	
American Indians, Office of the Special Trustee	25, VII
Endangered Species Committee	50, IV
Federal Acquisition Regulation	48, 14
Federal Property Management Regulations System	41, 114
Fish and Wildlife Service, United States	50, I, IV
Geological Survey	30, IV
Indian Affairs, Bureau of	25, I, V
Indian Affairs, Office of the Assistant Secretary	25, VI
Indian Arts and Crafts Board	25, II
Land Management, Bureau of	43, II
Minerals Management Service	30, II
Mines, Bureau of	30, VI

Agency	CFR Title, Subtitle or Chapter
National Indian Gaming Commission	25, III
National Park Service	36, I
Reclamation, Bureau of	43, I
Secretary of the Interior, Office of	43, Subtitle A
Surface Mining and Reclamation Appeals, Board of	30, III
Surface Mining Reclamation and Enforcement, Office of	30, VII
Internal Revenue Service	26, I
International Boundary and Water Commission, United States and Mexico, United States Section	22, XI
International Development, Agency for	22, II
Federal Acquisition Regulation	48, 7
International Development Cooperation Agency, United States	22, XII
International Development, Agency for	22, II; 48, 7
Overseas Private Investment Corporation	5, XXXIII; 22, VII
International Fishing and Related Activities	50, III
International Investment, Office of	31, VIII
International Joint Commission, United States and Canada	22, IV
International Organizations Employees Loyalty Board	5, V
International Trade Administration	15, III; 19, III
International Trade Commission, United States	19, II
Interstate Commerce Commission	5, XL
James Madison Memorial Fellowship Foundation	45, XXIV
Japan–United States Friendship Commission	22, XVI
Joint Board for the Enrollment of Actuaries	20, VIII
Justice Department	5, XXVIII; 28, I
Drug Enforcement Administration	21, II
Federal Acquisition Regulation	48, 28
Federal Claims Collection Standards	4, II
Federal Prison Industries, Inc.	28, III
Foreign Claims Settlement Commission of the United States	45, V
Immigration and Naturalization Service	8, I
Offices of Independent Counsel	28, VI
Prisons, Bureau of	28, V
Property Management Regulations	41, 128
Labor Department	5, XLII
Benefits Review Board	20, VII
Employees' Compensation Appeals Board	20, IV
Employment and Training Administration	20, V
Employment Standards Administration	20, VI
Federal Acquisition Regulation	48, 29
Federal Contract Compliance Programs, Office of	41, 60
Federal Procurement Regulations System	41, 50
Labor-Management Standards, Office of	29, II, IV
Mine Safety and Health Administration	30, I
Occupational Safety and Health Administration	29, XVII
Pension and Welfare Benefits Administration	29, XXV
Public Contracts	41, 50
Secretary of Labor, Office of	29, Subtitle A
Veterans' Employment and Training, Office of the Assistant Secretary for	41, 61; 20, IX
Wage and Hour Division	29, V
Workers' Compensation Programs, Office of	20, I
Labor-Management Standards, Office of	29, II, IV
Land Management, Bureau of	43, II
Legal Services Corporation	45, XVI
Library of Congress	36, VII
Copyright Office	37, II
Management and Budget, Office of	5, III, LXXVII; 48, 99
Marine Mammal Commission	50, V
Maritime Administration	46, II
Merit Systems Protection Board	5, II
Micronesia Status Negotiations, Office for	32, XXVII
Mine Safety and Health Administration	30, I
Minerals Management Service	30, II
Mines, Bureau of	30, VI

Agency	CFR Title, Subtitle or Chapter
Minority Business Development Agency	15, XIV
Miscellaneous Agencies	1, IV
Monetary Offices	31, I
Multifamily Housing Assistance Restructuring, Office of	24, IV
National Aeronautics and Space Administration	5, LIX; 14, V
Federal Acquisition Regulation	48, 18
National Agricultural Library	7, XLI
National Agricultural Statistics Service	7, XXXVI
National Archives and Records Administration	5, LXVI; 36, XII
Information Security Oversight Office	32, XX
National Bureau of Standards	15, II
National Capital Planning Commission	1, IV
National Commission for Employment Policy	1, IV
National Commission on Libraries and Information Science	45, XVII
National and Community Service, Corporation for	45, XII, XXV
National Council on Disability	34, XII
National Credit Union Administration	12, VII
National Drug Control Policy, Office of	21, III
National Foundation on the Arts and the Humanities	45, XI
National Highway Traffic Safety Administration	23, II, III; 49, V
National Imagery and Mapping Agency	32, I
National Indian Gaming Commission	25, III
National Institute for Literacy	34, XI
National Institute of Standards and Technology	15, II
National Labor Relations Board	5, LXI; 29, I
National Marine Fisheries Service	50, II, IV, VI
National Mediation Board	29, X
National Oceanic and Atmospheric Administration	15, IX; 50, II, III, IV, VI
National Park Service	36, I
National Railroad Adjustment Board	29, III
National Railroad Passenger Corporation (AMTRAK)	49, VII
National Science Foundation	5, XLIII; 45, VI
Federal Acquisition Regulation	48, 25
National Security Council	32, XXI
National Security Council and Office of Science and Technology Policy	47, II
National Telecommunications and Information Administration	15, XXIII; 47, III
National Transportation Safety Board	49, VIII
National Weather Service	15, IX
Natural Resources Conservation Service	7, VI
Navajo and Hopi Indian Relocation, Office of	25, IV
Navy Department	32, VI
Federal Acquisition Regulation	48, 52
Neighborhood Reinvestment Corporation	24, XXV
Northeast Dairy Compact Commission	7, XIII
Nuclear Regulatory Commission	5, XLVIII; 10, I
Federal Acquisition Regulation	48, 20
Occupational Safety and Health Administration	29, XVII
Occupational Safety and Health Review Commission	29, XX
Offices of Independent Counsel	28, VI
Operations Office	7, XXVIII
Overseas Private Investment Corporation	5, XXXIII; 22, VII
Panama Canal Commission	48, 35
Panama Canal Regulations	35, I
Patent and Trademark Office	37, I
Payment From a Non-Federal Source for Travel Expenses	41, 304
Payment of Expenses Connected With the Death of Certain Employees	41, 303
Peace Corps	22, III
Pennsylvania Avenue Development Corporation	36, IX
Pension and Welfare Benefits Administration	29, XXV
Pension Benefit Guaranty Corporation	29, XL
Personnel Management, Office of	5, I, XXXV; 45, VIII
Federal Acquisition Regulation	48, 17
Federal Employees Group Life Insurance Federal Acquisition Regulation	48, 21

Agency	CFR Title, Subtitle or Chapter
Federal Employees Health Benefits Acquisition Regulation	48, 16
Postal Rate Commission	5, XLVI; 39, III
Postal Service, United States	5, LX; 39, I
Postsecondary Education, Office of	34, VI
President's Commission on White House Fellowships	1, IV
Presidential Commission on the Assignment of Women in the Armed Forces	32, XXIX
Presidential Documents	3
Presidio Trust	36, X
Prisons, Bureau of	28, V
Procurement and Property Management, Office of	7, XXXII
Productivity, Technology and Innovation, Assistant Secretary	37, IV
Public Contracts, Department of Labor	41, 50
Public and Indian Housing, Office of Assistant Secretary for	24, IX
Public Health Service	42, I
Railroad Retirement Board	20, II
Reclamation, Bureau of	43, I
Refugee Resettlement, Office of	45, IV
Regional Action Planning Commissions	13, V
Relocation Allowances	41, 302
Research and Special Programs Administration	49, I
Rural Business-Cooperative Service	7, XVIII, XLII
Rural Development Administration	7, XLII
Rural Housing Service	7, XVIII, XXXV
Rural Telephone Bank	7, XVI
Rural Utilities Service	7, XVII, XVIII, XLII
Saint Lawrence Seaway Development Corporation	33, IV
Science and Technology Policy, Office of	32, XXIV
Science and Technology Policy, Office of, and National Security Council	47, II
Secret Service	31, IV
Securities and Exchange Commission	17, II
Selective Service System	32, XVI
Small Business Administration	13, I
Smithsonian Institution	36, V
Social Security Administration	20, III; 48, 23
Soldiers' and Airmen's Home, United States	5, XI
Special Counsel, Office of	5, VIII
Special Education and Rehabilitative Services, Office of	34, III
State Department	22, I
Federal Acquisition Regulation	48, 6
Surface Mining and Reclamation Appeals, Board of	30, III
Surface Mining Reclamation and Enforcement, Office of	30, VII
Surface Transportation Board	49, X
Susquehanna River Basin Commission	18, VIII
Technology Administration	15, XI
Technology Policy, Assistant Secretary for	37, IV
Technology, Under Secretary for	37, V
Tennessee Valley Authority	5, LXIX; 18, XIII
Thrift Supervision Office, Department of the Treasury	12, V
Trade Representative, United States, Office of	15, XX
Transportation, Department of	5, L
Coast Guard	33, I; 46, I; 49, IV
Coast Guard (Great Lakes Pilotage)	46, III
Commercial Space Transportation	14, III
Contract Appeals, Board of	48, 63
Emergency Management and Assistance	44, IV
Federal Acquisition Regulation	48, 12
Federal Aviation Administration	14, I
Federal Highway Administration	23, I, II; 49, III
Federal Railroad Administration	49, II
Federal Transit Administration	49, VI
Maritime Administration	46, II
National Highway Traffic Safety Administration	23, II, III; 49, V
Research and Special Programs Administration	49, I
Saint Lawrence Seaway Development Corporation	33, IV

Agency	CFR Title, Subtitle or Chapter
Secretary of Transportation, Office of	14, II; 49, Subtitle A
Surface Transportation Board	49, X
Transportation Statistics Bureau	49, XI
Transportation, Office of	7, XXXIII
Transportation Statistics Bureau	49, XI
Travel Allowances, Temporary Duty (TDY)	41, 301
Treasury Department	5, XXI; 12, XV; 17, IV
Alcohol, Tobacco and Firearms, Bureau of	27, I
Community Development Financial Institutions Fund	12, XVIII
Comptroller of the Currency	12, I
Customs Service, United States	19, I
Engraving and Printing, Bureau of	31, VI
Federal Acquisition Regulation	48, 10
Federal Law Enforcement Training Center	31, VII
Fiscal Service	31, II
Foreign Assets Control, Office of	31, V
Internal Revenue Service	26, I
International Investment, Office of	31, VIII
Monetary Offices	31, I
Secret Service	31, IV
Secretary of the Treasury, Office of	31, Subtitle A
Thrift Supervision, Office of	12, V
Truman, Harry S. Scholarship Foundation	45, XVIII
United States and Canada, International Joint Commission	22, IV
United States and Mexico, International Boundary and Water Commission, United States Section	22, XI
Utah Reclamation Mitigation and Conservation Commission	43, III
Veterans Affairs Department	38, I
Federal Acquisition Regulation	48, 8
Veterans' Employment and Training, Office of the Assistant Secretary for	41, 61; 20, IX
Vice President of the United States, Office of	32, XXVIII
Vocational and Adult Education, Office of	34, IV
Wage and Hour Division	29, V
Water Resources Council	18, VI
Workers' Compensation Programs, Office of	20, I
World Agricultural Outlook Board	7, XXXVIII

Redesignation Table

At 53 FR 24666, June 29, 1988, 21 CFR part 561 was redesignated as 40 CFR part 186. The following redesignation table shows the relationship of the CFR section numbers formerly in 21 CFR part 561 and those sections being redesignated as 40 CFR part 186.

Old section	New section	Old section	New section
561.20	186.100	561.260	186.3700
561.30	186.150	561.263	186.3750
561.40	186.200	561.268	186.3800
561.41	186.250	561.270	186.3850
561.50	186.350	561.273	186.4000
561.51	186.375	561.280	186.4050
561.53	186.4975	561.282	186.4150
561.60	186.450	561.283	186.4450
561.65	186.500	561.285	186.4575
561.66	186.550	561.289	186.4700
561.67	186.600	561.290	186.4750
561.70	186.700	561.300	186.4800
561.80	186.750	561.305	186.4850
561.91	186.950	561.310	186.4900
561.92	186.400	561.330	186.5000
561.93	186.800	561.340	186.5200
561.95	186.850	561.350	186.5350
561.96	186.1250	561.360	186.1550
561.97	186.1300	561.365	186.5400
561.98	186.1000	561.380	186.5550
561.99	186.1400	561.385	186.5600
561.100	186.1450	561.386	186.5650
561.110	186.1500	561.387	186.5700
561.130	186.1600	561.390	186.5800
561.140	186.1650	561.400	186.1350
561.145	186.1700	561.410	186.6300
561.150	186.1875	561.415	186.1750
561.160	186.1950	561.420	186.2000
561.170	186.2100	561.425	186.2050
561.180	186.2225	561.427	186.1800
561.190	186.2325	561.428	186.3250
561.191	186.2150	561.429	186.3650
561.197	186.2275	561.430	186.2800
561.200	186.2400	561.432	186.4950
561.210	186.2450	561.434	186.5100
561.215	186.2500	561.435	186.3300
561.220	186.2550	561.436	186.3400
561.225	186.2700	561.437	186.1050
561.230	186.2750	561.438	186.3200
561.232	186.2950	561.439	186.1100
561.233	186.3000	561.440	186.1850
561.234	186.3050	561.441	186.300
561.235	186.2775	561.442	186.5950
561.240	186.3350	561.443	186.4350
561.250	186.3450	561.444	186.5850
561.253	186.3500	561.445	186.5250
561.255	186.3550		

List of CFR Sections Affected

All changes in this volume of the Code of Federal Regulations which were made by documents published in the FEDERAL REGISTER since January 1, 1986, are enumerated in the following list. Entries indicate the nature of the changes effected. Page numbers refer to FEDERAL REGISTER pages. The user should consult the entries for chapters and parts as well as sections for revisions.

For the period before January 1, 1986, see the "List of CFR Sections Affected, 1949–1963, 1964–1972, and 1973–1985," published in seven separate volumes.

21 CFR	1986	51 FR Page
Chapter I		
Mandatory compliance date 1–1–		
89.....		34085
510.6 (d)(3), (e), and (f) amended;		
eff. 5–2–86.....		7391
510.305 Introductory text and (a)		
amended; eff. 5–2–86.....		7391
510.455 Revised.....		19827
510.505 Removed.....		30211
510.520 (a) introductory text		
amended.....		30211
510.600 (c)(1) and (2) tables amend-		
ed.....		594,
		5990, 5991, 11438, 11439, 18883,
		19829, 22799, 28546, 29916, 31100,
		33897, 47013
Effective date corrected to 12–		
13–85.....		8315
514 Authority citation revised.....		7391
514.1 (e) removed; (b)(3)(v)(a) and		
(b), (5)(vii)(a) and (x) revised;		
OMB number; eff. 5–2–86.....		7391
514.2 Revised; eff. 5–2–86.....		7391
514.9 (c) amended; eff. 5–2–86.....		7391
514.55 Amended; eff. 5–2–86.....		7391
514.105 (b) amended; eff. 5–2–86.....		7392
514.112 Added; eff. 5–2–86.....		7392
520 Authority citation cor-		
rected.....		42163
520.314 (c) introductory text, (1),		
(2), and (3) redesignated as		
(c)(1) introductory text, (i),		
(ii), and (iii); new (c)(2)		
added.....		4165
520.446 (c) revised.....		34960
520.447 (c) revised.....		34960

21 CFR—Continued	51 FR Page
Chapter I—Continued	
520.580 Effective date corrected to	
12–13–85.....	8315
520.600 (c) amended.....	28546
520.905e Added.....	41783
520.1192 (a) revised; (c)(1) redesi-	
gnated as (c)(1)(i) and revised;	
(c)(2) and (3) redesignated as	
(c)(1)(ii) and (iii); new (c)(2)	
added.....	44449
520.1236 (a) revised.....	42834
520.1242f (d)(1), (2), and (3) redesi-	
gnated as (d)(1) (i), (ii), and (iii);	
new (d)(2) added; (d) introduc-	
tory text redesignated as new	
(d)(1) heading and revised.....	29215
520.1326 Redesignated as 520.1326a	
and heading revised; new	
520.1326 added.....	13212
520.1326a Redesignated from	
520.1326 and heading revised.....	13212
520.1326b Added.....	13212
520.1450a (d)(3) amended.....	9005
520.1450b (d) revised.....	23415
(d)(3) amended.....	41081
520.1468 (c)(1) revised.....	24525
520.2043 (b)(4) added.....	11440
520.2260b (f) and (g) added.....	30212
520.2455 (b) amended.....	28546
520.2473a (c)(1)(ii) corrected.....	2693
520.2480 (a) and footnote 1 re-	
moved; (b) and (c) redesignated	
as (a) and (b); new (c) added;	
(d)(1) and (2) amended; (d)(3)	
and (4) revised.....	26002

21 CFR (4–1–99 Edition)

21 CFR—Continued

51 FR
Page

Chapter I—Continued

520.2481 (a) removed; (b) and (c) re-designated as (a) and (b); (c) and (d)(4) added; new (b) and (d)(3) revised.....	26002
520.2482 (c) redesignated as (d); new (c) added; new (d)(3) re-vised.....	26002
522 Heading corrected.....	19545
Authority citation corrected.....	20646
Authority citation revised.....	25032
522.234 (b) amended.....	19329
522.840 Revised.....	22276
522.850 (c)(2) revised.....	33592
522.940 (d) removed.....	14989
(c)(1) amended.....	18314
522.1044 (b)(3) and (d)(5) added.....	15606
522.1145 Heading, (a), and (b) re-vised.....	11438
(a), (b), and (c) (1), (2), and (3) re-designated as (a) (1), (2), and (3) (i), (ii), and (iii); new (b) added.....	25032
522.1192 (d)(2) (ii) and (iii) re-vised.....	25686
(a)(2) heading revised; (d)(4) added.....	27021
(d)(4)(ii) corrected.....	28932, 29463
522.1410 (b) revised.....	741
522.1468 Removed; new 522.1468 re-designated from 522.1468b and heading and (c)(2) revised.....	24525
522.1468a Removed.....	24525
522.1468b Redesignated as 522.1468 and heading and (c)(2) re-vised.....	24525
522.1662a (a)(2) and (g)(2) amend-ed.....	28547
522.1881 Removed.....	35632
522.1940 (d)(1) (ii) and (iii) amend-ed.....	21746
Technical correction.....	25198
522.2063 (b) revised; (c)(2) redesign-ated as (c)(2)(i); new (c)(2)(ii) added.....	41477
522.2470 (b) amended.....	24142
522.2615 Added.....	44450
529 Authority citation revised.....	594
529.1030 Added.....	11441
529.1186 Added.....	594
529.2503 (b) amended.....	11439
540 Authority citation revised.....	4484
540.103g (c)(3)(i)(b) and (c) re-vised.....	4484
(a)(1) amended; (c)(3)(ii) added.....	5318
540.103h Added.....	4483
(c)(3)(ii) added.....	5317

21 CFR—Continued

51 FR
Page

Chapter I—Continued

(a)(1) corrected.....	6218
546 Authority citation revised.....	28547
546.110c (c)(2) amended.....	28547
546.180d (c)(5) revised; (c)(6)(iv)(c) added.....	44451
548.314a (a)(1)(i) and (ii) and (c)(2)(i) and (ii) amended.....	19546
548.314b (a)(1) and (2) amended.....	19546
555 Authority citation revised.....	1367
555.110b (c)(2)(i) amended.....	11441
555.110c Removed.....	1367
556.344 Revised.....	27021
556.425 Revised.....	9005
556.428 Added.....	29097
556.490 Revised.....	18884
556.640 (a) revised.....	18884
(d) table amended.....	29097
558 Authority citation revised.....	6910
Authority citation corrected.....	7064
558.3 Revised; eff. 5–2–86.....	7392
558.4 Revised; eff. 5–2–86 and 3–3–87.....	7392
(d) table corrected.....	9005
(d) table amended; eff. 5–2–86.....	9190
558.15 (g) introductory text and (2) introductory text revised; eff. 5–2–86.....	7394
Revised; eff. 4–14–86.....	8811
(g)(1) table corrected.....	11014
(g)(1) and (2) tables amended.....	28547
558.20 Added; eff. 5–2–86.....	7394
558.35 (a), (b), (d), and (e) removed; (c), (f), and (g) redesignated as (a), (b), and (c); new (a) intro-ductory text revised; eff. 5–2–86.....	7395
Technical correction.....	9005
558.45 (b), (c), (d), and (e) removed; (f) redesignated as (b); (a) re-vised; eff. 5–2–86.....	7395
558.55 (a) revised; (b) removed; (c), (d), and (e) redesignated as (b), (c), and (d); new (b) revised; eff. 5–2–86.....	7395
558.58 (b) removed; (c), (d), and (e) redesignated as (b), (c), and (d); (a), new (b), and (d) introduc-tory text revised; eff. 5–2–86.....	7395
558.59 Added.....	9190
558.60 (a) added; (b) and (c) re-moved; (d) and (e) redesignated as (b) and (c); new (c)(1) intro-ductory text revised; eff. 5–2–86.....	7395
(a) amended.....	33897

List of CFR Sections Affected

21 CFR—Continued

51 FR
Page

Chapter I—Continued

558.62 (a) added; (b) and (c) removed; (d) and (e) redesignated as (b) and (c); new (c)(1) introductory text revised; eff. 5-2-86	7395
(a) amended	33897
558.76 (b) removed; (c), (d), and (e) redesignated as (b), (c), and (d); (a), new (b), and (d)(1) introductory text revised; eff. 5-2-86	7396
558.78 (b) removed; (c), (d) and (e) redesignated as (b), (c) and (d); (a), new (b), and (d)(1) introductory text revised; eff. 5-2-86	7396
(d)(3)(x) added	18314
(d)(3)(x) revised	29463
558.95 (b), (c), and (d) removed; (e) redesignated as (b); (a), and new (b)(1) introductory text and (2) introductory text revised; eff. 5-2-86	7396
558.105 (b) and (c) removed; (d), (e), and (f) redesignated as (b), (c), and (d); (a), new (b), and (d) introductory text revised; eff. 5-2-86	7396
558.115 (b) and (c) removed; (d), (e) and (f) redesignated as (b), (c) and (d); (a), new (c), and (d) introductory text revised; eff. 5-2-86	7396
558.120 (a) added; (b) and (c) removed; (d) and (e) redesignated as (b) and (c); new (c)(1) introductory text revised; eff. 5-2-86	7396
558.128 (b) and (c) removed; (d) and (e) redesignated as (b) and (c); (a) revised; eff. 5-2-86	7396
558.145 (c) and (d) removed; (e) and (f) redesignated as (c) and (d); (a) and (b) revised; eff. 5-2-86	7396
558.155 (c) and (d) removed; (e) and (f) redesignated as (c) and (d); (a), (b), and new (d) introductory text revised; eff. 5-2-86	7397
(a) (1) and (2) amended	28547
558.175 (b) and (c) removed; (d) and (e) redesignated as (b) and (c); (a) revised; eff. 5-2-86	7397
558.185 (b) and (c) removed; (d), (e) and (f) redesignated as (b), (c) and (d); (a) and new (b) revised; eff. 5-2-86	7397

21 CFR—Continued

51 FR
Page

Chapter I—Continued

558.195 (b), (c), and (d) removed; (e), (f), and (g) redesignated as (b), (c), and (d); (a) revised; eff. 5-2-86	7397
(a) and (d) table amended	34589
558.205 (b) and (c) removed; (d), (e) and (f) redesignated as (b), (c) and (d); (a) revised; eff. 5-2-86	7397
(a) amended	28547
558.240 (b) and (c) removed; (d) and (e) redesignated as (b) and (c); (a) added	7397
558.248 (b) removed; (c), (d), and (e) redesignated as (b), (c), and (d); (a), new (b), and (d)(1) introductory text revised; eff. 5-2-86	7397
558.254 (b) removed; (c), (d), and (e) redesignated as (b), (c), and (d); (a) added; eff. 5-2-86	7397
558.258 (b) removed; (c) and (d) redesignated as (b) and (c); (a) revised; eff. 5-2-86	7397
558.262 (b) and (c) removed; (d) and (e) redesignated as (b) and (c); (a) added; eff. 5-2-86	7397
558.265 (b) removed; (d) and (e) redesignated as (b) and (c); (a) revised; eff. 5-2-86	7397
(a) amended	11439
(c) (3) and (4) added	14989
(c) (2), (3), and (4) introductory texts revised	23737
558.274 (b) and (c) removed; (d) and (e) redesignated as (b) and (c); (a) revised; eff. 5-2-86	7398
(c)(1) table amended; (a)(8) added	19547
558.305 (b), (c), and (e) removed; (d) and (f) redesignated as (b) and (c); (a) and new (c) introductory text revised; eff. 5-2-86	7398
558.311 (b)(3) amended; (f)(11) added	5163
(f) table corrected	7064
(b) and (c) removed; (d), (e), and (f) redesignated as (b), (c), and (d); (a) revised; eff. 5-2-86	7398
(a) (1) and (3) and (d) table amended	37271

21 CFR (4-1-99 Edition)

21 CFR—Continued

51 FR
Page

Chapter I—Continued

558.315 (b), (c), (d), and (f) removed; (e) and (g) redesignated as (b) and (c); (a) and new (c) introductory text revised; eff. 5-2-86	7398
558.325 (b), (c) and (e) removed; (d) and (f) redesignated as (b) and (c); (a) revised; eff. 5-2-86	7398
(a) revised; (c)(2)(v) added	12137
(a) and (c)(2)(v) correctly designated; (a) introductory text, (1)(i), (ii), and (iii), and (2), (4), (5), (6), and (8) through (15) corrected	28547
558.342 (b) and (c) removed; (d) and (e) redesignated as (b) and (c); (a) revised; eff. 5-2-86	7398
(c)(2)(ii) revised	36222
558.355 (a), (d) (3) and (4) removed; (b) introductory text and (c) revised; eff. 5-2-86	7398
(b)(12) and (f)(1)(xxii) added	16676
(b)(11) revised; (f)(1)(xxiii) added	32631
(f)(3)(iv)(b) revised	36222
(f)(1)(xii)(b) amended	36392
(b)(11) revised; (f)(1)(xxiv) added	45105
558.360 (b) removed; (c), (d), and (e) redesignated as (b), (c), and (d); (a) and new (c)(1) revised; eff. 5-2-86	7399
(d)(3) amended	9005
558.363 Added	29098
558.365 (b) and (c) removed; (d), (e), and (f) redesignated as (b), (c), and (d); (a) and new (c) revised; eff. 5-2-86	7399
558.366 (b) and (d) removed; (c) and (e) redesignated as (b) and (c); (a) revised; eff. 5-2-86	7399
558.367 (b) removed; (c) redesignated as (b); (a) revised; eff. 5-2-86	7399
558.369 (b) and (c) removed; (d), (e), and (f) redesignated as (b), (c), and (d); (a) revised; eff. 5-2-86	7399
558.370 (a) added; (b) and (c) removed; (d) redesignated as (b); eff. 5-2-86	7399
558.376 (b) and (c) removed; (d) and (e) redesignated as (b) and (c); (a) added; eff. 5-2-86	7399

21 CFR—Continued

51 FR
Page

Chapter I—Continued

558.415 (b), (c), and (d) removed; (e) and (f) redesignated as (b) and (c); (a) revised; eff. 5-2-86	7399
558.430 (b) and (c) removed; (d) and (e) redesignated as (b) and (c); (a) added; eff. 5-2-86	7399
558.435 (b) and (c) removed; (d), (e) and (f) redesignated as (b), (c) and (d); (a) and new (c) revised; eff. 5-2-86	7399
558.450 (b) removed; (c), (d), and (e) redesignated as (b), (c), and (d); (a) and new (b) revised; eff. 5-2-86	7399
(d)(1) Table 1 amended	16676
558.460 (b), (c), and (d) removed; (e) and (f) redesignated as (b) and (c); (a) revised; eff. 5-2-86	7399
558.464 (b) removed; (c) redesignated as (b); (a) revised	7399
558.465 (b) and (c) removed; (d) redesignated as (b); heading and (a) revised; eff. 5-2-86	7399
558.485 (b) removed; (a) introductory text and (d) revised; eff. 5-2-86	7400
(a)(27) added	33898
Technical correction	43121
558.515 (b) and (c) removed; (d), (e), and (f) redesignated as (b), (c), and (d); (a) revised; new (b) amended; eff. 5-2-86	7400
558.525 (b) and (c) removed; (d), (e), and (f) redesignated as (b), (c), and (d); (a) revised; eff. 5-2-86	7400
558.530 (b) and (c) removed; (d), (e), and (f) redesignated as (b), (c), and (d); (a) revised; eff. 5-2-86	7400
(d)(4) (iii) and (iv) added	29463
(a)(1) amended	44055
558.550 (b) revised; (c)(2) redesignated as (c)(3); new (c)(2) added	595
(c)(1)(vi) added	6911
(b) removed; (c) redesignated as (b); (a) revised; eff. 5-2-86	7400
(b)(1)(vii) added	18314
(b)(1)(viii) added	29463
(b)(1)(ix) added	31764
558.575 (b) and (c) removed; (d) and (e) redesignated as (b) and (c); (a) revised; eff. 5-2-86	7400

List of CFR Sections Affected

21 CFR—Continued

51 FR
Page

Chapter I—Continued	
(c)(6) added	18884
558.579 (a) added; (b) and (c) removed; (d) and (e) redesignated as (b) and (c); eff. 5-2-86	7400
558.582 (a) added; (b) and (c) removed; (d) and (e) redesignated as (b) and (c); eff. 5-2-86	7400
558.586 (b) and (c) removed; heading revised; (a) added; eff. 5-2-86	7400
558.615 (b) removed; (c), (d), and (e) redesignated as (b), (c), and (d); (a) and new (b) revised; eff. 5-2-86	7400
558.625 (b)(86) added	5991
(c) and (d) removed and reserved; (b) introductory text revised; eff. 5-2-86	7400
(b)(25) amended	7785
(b)(6) removed	11439
(b)(87) added	22799
(b)(88) added	26378
558.630 (b)(3) and (10) amended	3333
(b)(10) amended	5992
(a), (c), and (d) removed; (b) introductory text revised; eff. 5-2-86	7400
(b)(3) amended	7785
(b)(10) amended	19829
(b)(8) and (10) amended	34961
558.635 (f)(2)(ii) revised	7262
(a), (c), and (e) (3) and (4) removed; (b) revised; eff. 5-2-86	7400
(f)(3)(v) added	14990
558.680 (a) revised; (b) and (c) removed; (d) and (e) redesignated as (b) and (c); eff. 5-2-86	7401
561.92 Revised	9439
Corrected	12138, 17174
561.120 Removed	46617
561.260 (c) removed	1785
561.370 Removed	46618
561.432 (c) added	28224
561.436 Added	15317
Correctly designated	19169
561.438 Added	39655
570 Authority citation revised	5992
570.12 Removed	5993
579 Added	5993
579.22 Correctly designated	8315

1987

21 CFR

52 FR
Page

Chapter I	
Uniform compliance date 1-1-91	44861
500.80—500.92 (Subpart E)	
Added	49586
510 Authority citation revised	25976
510.600 (c)(1) and (2) tables amended	5455,
7831, 7833, 11041, 11988, 12153,	
18691, 19501, 20385, 20597, 25212,	
25976, 27801, 32540, 32917, 36022,	
36023, 39912, 41295, 41988	
(c) (1) and (2) tables amended; eff. 4-3-87	9295
Technical correction	10668, 18495,
	23397
511.1 (c)(2) revised; (b)(4)(vi) and (f) added; eff. 6-17-87	8847
514.1 (b)(8)(viii) and (ix) added; eff. 6-17-87	8847
(b)(7) introductory text and (ii) revised; OMB number	49588
514.111 (a)(10) added	49588
520 Technical correction	10668, 18495
Heading corrected	28067
520.23 (a)(2) revised	666
520.314 (a) revised	11989
520.540a (b) amended	7832
520.540b (a)(2) and (b)(2) amended	7832
520.580 (b)(1) revised	39512
520.680 Removed	25213
520.680a Removed	25213
520.680b Removed	25213
Correctly designated	27108
520.970a (b) amended	7832
520.970b (b) amended	7832
520.1044a (b) amended	7832
520.1044b (b) amended	7832
520.1044c (b) amended	7832
(a) revised	48675
520.1100 (c)(1) amended	7832
520.1193 Added	11042
520.1195 Added	34637
520.1242a (f)(2)(i) amended	20598
520.1341 (b) amended	7832
520.1448a (b) removed	27197
(d)(4)(iii) amended	30151
520.1540 Removed	20597
520.1660d (e) revised	48094
520.1720a (b)(5) removed	36023
520.2041 Added	37937
520.2100 (b) amended	7832
(b) corrected	9756
520.2455 Revised	15718
520.2473a (b) amended	7832

21 CFR (4-1-99 Edition)

21 CFR—Continued

52 FR
Page

Chapter I—Continued	
520.2473b	Added.....43059
522	Technical correction.....10668, 18495
522.56	Added.....11816
	(a) corrected.....15412
522.161	(c) amended.....7832
522.163	(b) amended.....7832
522.518	(b) amended.....7832
522.540	(a)(2) amended.....7832
	(b)(2)(ii) and (c)(2) amended.....25212
	(e)(2) amended.....36023
522.900	(b)(2) amended.....7832
522.970	(b) amended.....7832
522.1044	(b)(1) amended.....7832
522.1060b	(b) amended.....7832
522.1081	(a)(2)(i) amended.....25212
522.1182	(a)(4)(i) amended.....7832
522.1183	(e)(1) revised.....18691
	(e)(1) amended.....36023
522.1258	(b) amended.....25212
522.1660	(c)(1)(i) and (ii) amend- ed.....19502
522.1662a	(e)(1) amended.....7834
	(i)(2) amended.....36023
522.1680	(b) revised.....18691
	(b) amended.....25212, 36023
522.1704	(b)(2) amended.....25212
522.1720	(b)(2) amended.....25212, 36023
522.1820	(b) amended.....7832
522.1822	(b) amended.....7832
522.1881	Added.....23032
522.1884	(c) amended.....25212
522.1890	(b)(2) amended.....7832
522.1940	(d)(2)(iii) amended; (e) added.....45312
522.2063	(b) amended.....7832
522.2100	(a)(2) and (b)(2) amend- ed.....7832
522.2476	Added.....24995
522.2483	(b) amended.....1903
524	Technical correction.....10668, 18495
524.1044a	(b) amended.....7832
524.1044b	(b) amended.....7832
524.1044c	(b) amended.....7832
524.1044d	(b) amended.....7832
524.1044e	(b) amended.....7832
524.1044f	(b) amended.....7833
524.1240	Added.....10887
524.1580b	(b) amended.....18691
524.1580d	(b) and (d)(1)(i) re- vised.....4897
	(b) revised.....18691
	(b) amended.....36023
524.2350	(b) amended.....7833
529.1044a	(b) amended.....7833
	Technical correction.....10668, 18495
529.1044b	(b) amended.....7833

21 CFR—Continued

52 FR
Page

Chapter I—Continued	
	Technical correction.....10668, 18495
540.274b	(c)(3)(ii) amended; (c)(5) added.....32917
540.874a	(c)(3)(i) and (4)(i) amend- ed.....36023
544.110	Removed.....22438
546.180h	(a)(1) and (c)(4)(i) amend- ed.....43060
546.180i	(a)(1) and (c)(4)(i) amend- ed.....43061
556.170	Revised.....43061
556.210	Removed.....25213
556.420	(b) revised.....15718
556.500	(d) revised.....24293
556.710	Revised.....27683
556.739	Added.....24995
558.3	(b)(3) amended.....2682
558.4	(d) table revised.....2682
	(d) table amended.....4284
	(d) correctly designated.....4992
	(d) table corrected.....7834
	(d) table amended.....25213, 26956, 39912
558.5	Heading revised; (a), (b), and (c) introductory text and (2) amended.....2684
558.45	Removed.....39912
558.55	(d)(1)(i)(b), (2) introductory text and table and (3) introduc- tory text amended.....2684
	(d)(1)(ii)(b) corrected.....6649
558.58	(a)(2) amended.....2684
558.76	(a) amended.....2684
558.78	(d)(1) table amended.....2684
	(d)(3)(xi) added.....2687
558.120	(c)(1)(iii) added.....2687
558.128	(c)(3) Tables 1 and 2 head- ings revised.....2684
558.145	(d)(1) amended.....2684
558.155	(d)(3) table heading re- vised.....2684
558.175	(c) introductory text, (1) (ii) and (iii), and (4)(ii) amend- ed.....2684
558.185	(d) introductory text, (1)(i)(b), (2)(iii), and (3)(iii) amended.....2684
558.195	(c) revised; (b) (1) and (2) tables combined and amended; (d)(2) introductory text re- moved; (d)(1) redesignated as (d) and introductory text re- vised.....2684
	(c) corrected.....4992
	(d) table amended.....38924, 43061, 43062
558.205	(b)(3) amended; (b)(4) re- moved.....2684

List of CFR Sections Affected

21 CFR—Continued

52 FR
Page

Chapter I—Continued	
558.240 (c) introductory text amended	2684
Removed.....	25213
558.248 (a)(2) revised; (a) (3), (4), and (5) added; (b) amended	2684
558.274 (c)(1) introductory text amended	2684
(a)(4), (c)(1)(i) and (ii) amended; eff. 4–3–87.....	9295
(c)(1) table amended; (a)(7) revised.....	29010
558.311 (a) through (d) redesignated as (b) through (e); new (a) added; new (e) table amended; new (b), (d), and (e) introductory text revised.....	2685
(d)(5) (i) and (ii) and (e) table amended	16240
(b) revised; (e) introductory text and (1) through (11) redesignated as (e)(1) introductory text and (i) through (xi); (e)(2) added	24996
(b)(5) added; (e)(1) table amended.....	26299
(b)(3) amended; (e)(1) table amended	33804
(e)(2)(ii) correctly revised	33930
Technical correction	35518
(b)(3) and (e)(1) table amended	41988
(e)(1) table amended	48095
558.325 (c)(1) introductory text and (2) introductory text revised; (c)(3) (i) and (ix) amended.....	2685
(a)(1)(iv) added.....	28470
558.342 (a), (c) introductory text, (1)(ii) and (2)(ii) amended.....	2685
(c)(3) added	33804
Technical correction	35518
558.355 (a) added; (b)(3), (c), (d) (1), (2), (5), (7), and (9), and (f)(1) (ii), (iv), (vi), (vii), (x) introductory text and (b), (xi) introductory text and (b), (xvii), (3)(i)(b)(I) and (2), and (iv)(b) amended.....	2685
(f)(1)(xiii)(b) and (xxi)(b) revised	13641
(b)(4) and (f)(2) added	15718
(b)(13) and (f)(3)(v) added.....	27197
558.360 (d)(3) amended	11642
558.365 (d)(1)(ii) amended	2685
558.369 (d) introductory text amended	2686
558.370 (b) introductory text amended	2686

21 CFR—Continued

52 FR
Page

Chapter I—Continued	
558.376 (c) introductory text, (1)(ii) and (2)(ii) amended.....	2686
558.415 (a) revised.....	36402
558.430 (c) introductory text amended	2686
558.435 (c) revised	2686
558.450 (b) and (d)(1) table 1 heading amended.....	2686
(d)(1) Table 3 amended	24293
(d)(1) Table 3 corrected	26401
(d)(2)(ii) added	48096
558.460 (c)(1) introductory text amended	2686
558.464 (b)(2) amended.....	2686
558.465 Heading revised; (b) (1), (2), (3), and (4) amended	2686
558.485 (a)(26) revised	781
(e)(1)(ii), (2)(ii), (3)(ii), (4)(ii), (10)(ii) and (11)(ii) amended	2686
558.515 (b) and (d)(1)(ii) amended.....	2686
558.525 (b)(1), (2) (i) and (ii), (d)(1)(ii), (2)(ii), (3)(ii), (4)(ii) and (5)(ii) amended	2686
(a)(1) removed.....	12521
558.526 Heading revised; (a), (d) (1) and (2) amended; (b) removed	2686
Removed.....	12521
558.530 (d)(1)(iii) and (2)(i)(b) amended	2686
558.550 (b)(2) removed	1626
Correctly designated.....	7261
558.575 (a) revised; (c)(1)(ii) amended	2686
558.579 (c)(1)(iii) amended.....	2686
558.586 (e)(2) removed; (f) amended.....	2686
558.600 Added	26956
558.615 (a), (b), and (d)(5)(iii) amended	2686
558.625 (b)(42) revised	530
(b)(81) removed.....	7833
(b)(18) removed; eff. 4–3–87	9295
(b) (37) and (64) removed.....	11041
(b)(7) removed.....	11988
(b)(70) removed	12153
(b)(89) added.....	27801
558.630 (b)(10) amended	1626
(b)(10) amended.....	28470
558.635 (e)(2), (f)(1) introductory text and (2) introductory text amended	2686
558.680 (c)(1) introductory text revised; (c)(1) table amended.....	2686
561.96 Added	29009

21 CFR (4-1-99 Edition)

21 CFR—Continued

52 FR
Page

Chapter I—Continued	
561.99 Revised	27545
561.273 (a) table amended	12154
(d) added	41418
561.360 Revised	27543
561.400 Heading and text nomen- clature change	32293
561.415 Heading and text nomen- clature change	32293
561.425 Heading and text nomen- clature change	32293
561.430 Table amended	1447
Table amended	27544
561.439 Added	10563
561.440 Added	17941
561.441 Added	17942
Technical correction	23137, 23916, 34903
561.442 Added	39222
571.1 (c) amended	8583
571.115 Added	49588
573 Authority citation revised	10888
573.920 Revised	10888
(c)(1) corrected	21001

1988

21 CFR

53 FR
Page

Chapter I	
510 Technical correction	49823
510.600 (c) (1) and (2) tables amended	11493, 20843, 21993, 22297, 25151, 32610, 39256, 40056, 40057, 40727, 40728, 40729, 50514, 52682
Effective date corrected	39839
(c)(1) table corrected	49823
520 Technical correction	49823
Heading corrected	53120
520.23 (a)(2) amended	40727
520.62 (b) amended	27851
520.110 (a) revised	37753
520.246 (b) amended	27851
520.314 (b) amended	27851
520.315 Added	27344
520.580 (b)(2) amended	21993, 40727
520.622a (a)(1) amended	40056
(a)(6) amended	40727
520.622b (c)(2) amended	40727
520.622c (b)(1) amended	40056
(b)(6) amended	40727
520.623 (a) revised	45759
520.763a (c) amended	40727
520.763b (c) amended	40727
520.763c (b) amended	40727
520.903d (c)(2)(i) revised	48533

21 CFR—Continued

53 FR
Page

Chapter I—Continued	
520.905a (d)(2) (ii) and (iii) redesign- ated as (d)(2)(i) (A) and (B); new (d)(2)(i)(A) revised; new (d)(2)(ii) added	40058
520.1010a (b) amended	40727
520.1192 (c)(1) (ii) and (iii) amend- ed	51273
520.1194 Added	27958
520.1195 (c) (2) and (3) amended	51273
520.1204 Heading revised; (b) amended	27851
520.1242g Added	23757
520.1330 (c) amended	23390
520.1331 (b) amended	23390
520.1408 (b) amended	40727
520.1801b Removed	48634
520.1900 (b) amended	40727
520.2260b (f)(1) amended	40727
520.2481 (b) amended	40727
520.2611 (b) and (c)(1) revised	11063
522 Heading correctly revised	26559
Technical correction	49823
522.23 (c) introductory text amended	40727
522.46 (b) amended	40057
522.56 (b) amended	27851
522.62 (c) amended	27851
522.246 (b) amended	27851
522.311 Added	40057
522.313 Added	5369
522.480 (a), (b), (c) and (d) heading and (1) through (3) redesign- ated as (a) (1), (2), (3) and (4) heading and (i) through (iii); new (a)(4)(i) amended; new (b) and (c) added	45760
522.844 Removed	15812
522.1010 (b) amended	40727
522.1044 (b)(3) amended	40727
522.1145 (c) added	19773
(d) added	22297
522.1182 (b)(2)(i) amended	40727
522.1183 (e)(1) amended	40728, 40729
522.1192 (a)(3) added; (d)(4)(i) re- vised	11064
(d)(4)(ii) revised	27006
522.1204 (b) amended	27851
522.1222a (c)(2) amended	23390
(c)(1) amended	27851
522.1222b (c) amended	27851
522.1410 (b) amended	40728
522.1662a (k) added	11494
(h)(2) amended	40728
522.1680 (b) revised	32610
Effective date corrected	39839

List of CFR Sections Affected

21 CFR—Continued

53 FR
Page

Chapter I—Continued	
(b) amended	40728
522.2220 (c)(2) amended	40728
522.2424 (b) amended	23390, 40728
522.2483 (b) amended	40728
522.2640a (b)(2) amended	40728
522.2662 (b) revised	4848, 23608
(b) amended	40728
522.2680 (c) amended	20843
Correctly designated; heading	
correctly revised	23340
524 Authority citation revised	4615,
	7504, 12512
Technical correction	49823
524.1200a (b) amended	27851
524.1200b (b) amended	27851
524.1204 (b) amended	7504
(a)(2) revised	12512
Technical correction	13217
(b) amended	27851
524.1443 Heading and (a) revised;	
(c)(2) amended	26242
524.1465 Added	39085
524.1484h Removed	4615
524.1484j Removed	11065
524.1580b (b) amended	32610, 40728
Effective date corrected	39839
524.1580c (b) amended	40728
524.1580d (b) revised	32610
Effective date corrected	39839
(b) amended	40728
524.1600a (b) amended	39257
526 Authority citation revised	27851
526.363 (b) amended	27851
529.50 (b) amended	27852
529.365 (b) amended	27852
540.103g (a)(1) amended	235
540.119 (c)(2) amended	27852
540.129a (c)(2) amended	27852
540.129c (c)(2) amended	27852
540.181b (c)(2) amended	40729
540.203 (c)(2) (i), (iii), (iv) heading	
and (b) revised; (c)(2)(iv)(c)	
amended	40059
540.207b (c)(2) amended	27852
540.255c (a)(2) (i) and (ii) amend-	
ed	27852
540.274b (c)(3)(ii) amended	11493
540.814 (c)(2)(i) amended	27852
540.829 (c)(2) amended	27852
544.170b (c)(2) text added	52683
546 Technical correction	49823
546.110e Removed	235
546.180d (c)(6)(i)(c)(3), (iii)(d)(3)	
and (iv)(c)(3) amended	40728
548.114 (c)(2) amended	20843
555 Technical correction	49823

21 CFR—Continued

53 FR
Page

Chapter I—Continued	
555.110a (c)(1)(ii) amended	23390,
	40728
555.111 (c)(2) amended	23390
555.310c (c)(2) amended	23390
556.113 Added	5370
556.344 (c) added	27958
556.420 (b) revised	40060
558.4 (d) table amended	14788,
	25152, 40060
Technical correction	18022
558.15 (g)(1) table and (2) table	
amended	20843
558.58 (d)(1) table amended	20843
558.76 (d)(3)(xi) added	1018
(a) amended	4010
(d)(3)(xii) added	11065
Technical correction	14888
558.78 (a)(2) and (d)(1) table and	
(d)(ii) amended	20843
558.105 (d)(1)(xi)(b) amended	20843
558.120 (c)(1)(iii)(b) amended	20843
558.128 (a) revised; (c)(4) redesi-	
gnated as (c)(5); new (c)(4)	
added	31316
558.175 (c)(1)(iii)(b) and (iv)(b)	
amended	20843
558.195 (d) table amended	20843,
	22298
558.258 (c) introductory text and	
(1) through (3) redesignated as	
(c)(1) introductory text and (i)	
through (iii); new (c)(2)	
added	14788
Technical correction	18022
(c)(1) revised	48533
558.265 (c)(5) added	1018
(c)(6) added	11065
Technical correction	14888
558.311 Technical correction	11251
(e)(1) table amended	20843, 38708
558.342 (c)(3)(ii) revised	27959
558.355 (f)(3)(ii) introductory text	
and (b) revised	236
(b)(9), (f)(1)(iv)(b), (v)(b), (xv)(b)	
and (xvi)(b) amended	20843
(b)(10) removed	27345
(b)(6) and (f)(5) added	40060
(f)(3)(vi) added	50401
558.363 (c) revised	24260
558.430 (a) amended	40729
558.450 (a)(2) revised	27006
558.485 (a)(26) removed	236
558.515 (d)(1)(vi)(b) amended	20843
558.530 (d)(4)(v) added	1018
(d)(4)(vi) added	24260
558.550 (b)(1)(vii)(c) and (ix)(c)	
amended	20843

21 CFR (4-1-99 Edition)

21 CFR—Continued

	53 FR Page
Chapter I—Continued	
(b)(1) (x) and (xi) added	35313
(b)(1)(xii) added	44010
558.600 (c)(2) (i) and (ii) revised	39257
558.635 (f)(2)(iv) added	27852
(f)(3)(vi) added	35313
(f)(3)(vi) revised	44010
561 Redesignated as 40 CFR Part	
186	24666
Correctly redesignated as 40	
CFR Part 186	26131
561.20 (b), (c), and (d) removed	9435
561.55 Removed	9435
561.90 Removed	9435
561.96 (a) added	1917
(c) added	18837
561.98 (b), (c), and (d) removed	9435
561.175 Removed	9435
561.195 Removed	9435
561.197 Existing text designated	
as (a); (b) added; eff. to 6-30-	
89	23386
561.225 (a) table amended	5367
(a) table corrected	11938, 12640
561.231 Removed	9435
561.237 Removed	9435
561.253 (b) removed	9435
561.265 Removed	9435
561.273 (c) removed	8874
561.277 Removed	8874
561.280 (b), (c), and (d) removed	9435
561.281 Removed	9435
561.320 Removed	9435
561.330 Revised	9435
561.371 Removed	9435
561.380 (b) removed	9435
561.395 Removed	9435
561.400 Revised	23389
561.415 Introductory text re-	
vised	23389
561.425 Revised	23389
561.430 Table amended	234, 23388
561.432 Revised	8874
561.433 Removed	8874
561.440 Introductory text amend-	
ed	20308
561.441 Amended	23107
561.443 Added; exp. 2-28-88	21
Added	12943
561.444 Added	15813
561.445 Added	23387
573.225 Added	40061

1989

21 CFR

	54 FR Page
Chapter I	
500 Authority citation revised	39637
500.27 (d) amended	18279
500.35 (a) amended	18279
500.49 Authority citation re-	
moved	39637
500.51 (c) amended	18279
501 Authority citation revised;	
sectional authority citations	
removed	39637
501.3 (e)(3) revised	18279
501.105 (b)(3) amended	18279
502 Authority citation revised;	
sectional authority citations	
removed	39637
505 Authority citation revised;	
sectional authority citations	
removed	39638
505.10 Removed	18279
507 Authority citation revised	39638
508 Authority citation revised	39638
508.5 (a)(1) amended	18279
508.35 (c) (1) and (2) introductory	
text, (ii), (j)(1), and (l) amend-	
ed	18280
509 Authority citation revised	39638
509.5 Amended	18280
510 Technical correction	6232, 8880, 25447, 41713
Authority citation revised; sec-	
tional and subpart authority	
citations removed	39638
510.3 (g)(2) amended; (g)(3) re-	
moved	22741
510.5 Removed	22741
510.6 Removed	22741
510.55 Removed	22741
510.110 (f) amended	18280
510.112 (a) and (e) amended	18280
510.301 Introductory text amend-	
ed	18280
510.302 (d) amended	18280
510.310 (f) amended	18280
510.350 Removed	22741
510.510 Removed	22741
510.520 Removed	22741
510.521 Removed	22742
510.600 (c) (1) and (2) tables	
amended	1163, 1164, 6659, 7406, 17705, 22885, 24901, 25565, 26958, 27328, 32632, 33673, 38514, 38515, 38646, 40657, 41442, 43290, 47678, 50365, 51021
(c)(1) table corrected	9979

List of CFR Sections Affected

21 CFR—Continued

54 FR
Page

Chapter I—Continued	
Authority citation revised; sectional and subpart authority citations removed.....	39638
511 Authority citation revised	18280, 39638
511.1 (c)(1) and (e) amended.....	18280
Authority citation removed	39638
514 Technical correction.....	25447
Authority citation revised; sectional authority citations removed.....	39638
514.1 (b)(6) introductory text, (d)(2) introductory text, (i), and (ii) amended	18280
(d)(2) introductory text and (i) corrected.....	20235
(b)(9) removed.....	22742
514.11 (e)(2)(ii)(a) and (b) amended.....	18280
Heading corrected	20235
514.50 Removed.....	22742
514.51 Removed.....	22742
514.55 Removed.....	22742
514.60 Removed.....	22742
514.111 (a)(5)(ii)(a)(6) amended	18280
514.150 Removed	22742
514.155 Removed	22742
514.160 Removed	22742
520 Technical correction.....	6232, 8880, 22278
Effective date corrected	33814
Authority citation revised; sectional authority citations removed.....	39638
520.45 Added	25115
520.45a Added.....	25115
520.45b Added.....	51385
520.260 (a)(2) amended	38515
520.312 Added	32336
520.443 Removed	30205
520.446 (c) revised	47767
520.447 (c) revised.....	47766
520.580 (b)(1) amended.....	1164
(b)(1) revised	6659
520.623 (c)(2) revised	3776
(c)(2) corrected.....	6804
520.812 Added	3444
520.816 Added	50615
520.905e (a) and (d)(3) revised	20787
520.1100 (c) revised; (d) (3) and (4) redesignated as (d) (2) and (3).....	30205
520.1162 Removed	1686
520.1193 Section heading and (a) revised	32337
520.1236 Removed.....	29543

21 CFR—Continued

54 FR
Page

Chapter I—Continued	
520.1242a (f)(2)(ii)(a) amended	18280
520.1242g (f)(3) amended	1353
520.1660d (d) amended.....	18280
520.1720a (b)(4) amended	22885
520.1760 Removed.....	30205
520.1780 Removed.....	30205
520.1801a (d) removed.....	1163
520.1801c Removed.....	1163
520.1803 (b) amended	38515
520.1840 (c)(4) and (d)(4) added	33501
520.2080b Removed	30205
520.2184 (b) amended	12188
520.2220d Added.....	48593
520.2260a Revised	15751
(a)(3)(ii)(A) corrected.....	19283
520.2260b (h) added	14341
520.2320 (b)(1)(iii) amended.....	18280
520.2560 Removed	1164
522 Technical correction	6232
Authority citation corrected	9590
Authority citation revised; sectional authorities removed.....	39638
522.443 Removed	30205
522.514 Added	17706
522.536 Added	50365
Heading and (a) corrected.....	51551
522.540 (d)(2)(i) and (e)(2) amended	41442
522.850 (b) amended.....	1165
522.914 (e) (1), (2) and (3) redesignated as (e)(1) (i), (ii), and (iii); (a) and new (e)(1)(i) heading revised; new (e)(2) added	400
522.1077 Added	50235
522.1182 (a)(2) removed	41442
522.1235 Removed.....	29543
522.1462 (b) amended.....	32632
522.1662a (f) and (j) removed.....	41442
522.1680 (b) amended.....	41442
522.1720 (b)(2) amended.....	41442
522.1800 Removed.....	30205
522.1850 Revised	53053
522.1940 (d)(1)(iii) amended	7406
522.2063 (b) amended	1164
522.2220 (c)(2), (3)(i), and (ii) revised; (c)(3)(iii) removed; (c)(3)(iv) redesignated as (c)(3)(iii).....	30205
522.2670 Added	40657
522.2680 (f)(1)(iii) amended; (f)(3)(iii) removed	31950
524 Technical correction	6232
524 Authority citation revised	39638
524.1580b (b) revised	29544
(b) amended	37097
524.1580c (b) revised.....	30542

21 CFR (4-1-99 Edition)

21 CFR—Continued

54 FR
Page

Chapter I—Continued	
524.1600a (b) revised	32633, 48090
(b) amended.....	5431
524.1742 (b) amended	51021
524.2080 Removed.....	30205
524.2101 (c) introductory text amended; (d) removed	36962
524.2542 Removed	1164
524.2620 (b)(2) amended.....	25565
526 Authority citation revised	39638
529 Authority citation revised	39638
529.1030 (b) amended	5432
529.1186 (c) revised	23472
536 Authority citation revised	39638
539 Authority citation revised; sectional authority citations removed	39638
540 Authority citation revised; sectional authority citations removed	39638
540.181b (c)(1) amended	18280
540.255c (c)(2)(i) amended; (c)(2)(ii) revised	41442
540.274b (c)(3)(ii) and (5)(i) re- vised.....	33673
(c)(4) removed.....	41442
544 Authority citation revised; sectional authority citations removed	39638
544.275 (c)(2) and (4)(iii) amend- ed	41442
546 Technical correction	6232
546 Authority citation revised; sectional authority citations removed	39638
546.110b (b)(2) amended	18281
546.180a (c)(5)(i)(c) amended; eff. 4-3-89.....	11698
Correctly redesignated	12989
Correctly designated	13977
548 Authority citation revised	39638
548.112b Removed	30206
555 Authority citation revised; sectional authority citations removed	39638
555 Technical correction	6232
555.110b (c)(2) amended	26958
555.210 (c)(2) amended	41442
556 Authority citation cor- rected.....	20235
Authority citation revised	39638
556.34 Added	25115
556.250 (b) amended	18281
556.308 Revised	28052
556.340 Removed.....	1686
556.344 (a) amended	18281
556.375 Added	5229

21 CFR—Continued

54 FR
Page

Chapter I—Continued	
556.420 (a) revised.....	32633
556.739 Amended	12595
556.760 Introductory text re- moved; (a) and (b) introductory text added.....	31950
558 Authority citation revised	39639
Technical correction	41713
558.3 (b)(4) amended	51386
558.4 (d) table amended.....	1686, 1928, 5229, 9430, 18281, 32633, 53319
(d) table corrected	40657
558.15 (g)(1) table amended	37098
(g) introductory text amended	51386
558.62 (c)(2)(iii) amended.....	18281
558.76 (d)(1)(x) table amended	7189
(a) revised.....	12189
(d)(3)(xiii) added	29335
(d)(3)(xiv) added.....	32635
558.95 (b)(4)(ii) added	5930
558.128 (c)(5) (vii) and (viii) redes- ignated as (c)(5) (vi) and (vii); new (c)(5)(viii) added.....	24901
(c)(5)(xi) redesignated as (c)(5)(xii); new (c)(5)(xi) added	25115
(c)(5)(xiii) added	26732
558.175 (a)(1) and (2) amended.....	6518
558.195 (c)(3) added; (d) table amended	11183
558.248 (a) (3), (4), and (5) removed; (d)(1) table amended; eff. 4-3- 89.....	12189
558.258 (c)(3) added	36963
558.265 (c) revised	28052
(c)(7) added	11519
558.274 (a)(4) and (c)(1) table amended; eff. 4-7-89.....	12596
(a)(2) and (c)(1) table amended	51021
558.305 Removed.....	1686
558.311 (b)(6) added	29544
558.325 (c)(3)(xiv) added	11520
(c)(3)(xv) added	21940
(c)(3)(xv) revised	23209
558.340 Added	5229
(c)(2) and (3) redesignated as (c)(1)(i) and (ii); new (c)(2) added	26732
Technical correction	24789
(c)(3) added	32635
(c)(4) added	33885
558.355 (b)(14), (d) (3), (10), (11), and (f)(6) added.....	32633
558.363 (a) revised; (c)(1)(iii) added	1928
(a)(4) and (c)(1)(iv) added.....	5930

List of CFR Sections Affected

21 CFR—Continued

54 FR
Page

Chapter I—Continued	
(c)(1)(iii)(B) amended	9430
(c)(1)(v) added	29335
558.366 (c) table amended	1928
558.485 (a)(24) removed	3776
558.530 (d)(4)(vii) added	24901
(d)(3)(xxiii) added	29335
(d)(3)(xxiv) added	33885
558.550 (b)(1)(xiii) added	21940
(b)(1)(xiv) added	23209
Technical correction	24789
(b)(1)(xv) added	24901
(b)(1)(xvi) added	25116
(b)(1)(xv)(b) corrected	28154
(b)(1)(xvi)(c) corrected	32963
558.575 (c)(3)(ii) removed;	
(c)(3)(i)(a) and (b) redesignated	
as (c)(3)(ii) and (iii)	1686
558.625 (b)(33) revised; (f)(1)(i)(b)	
amended	10980
(b)(38) amended	38646
558.630 (b)(3) amended	109
(b)(10) amended	3776
(b)(4) removed	51021
(b)(10) amended	51021
564 Authority citation revised	39639
564.17 (k) amended	18281
570 Authority citation revised;	
sectional authority citations	
removed	39639
570.6 (e) amended	18281
570.35 (c)(2) and (3) amended	18281
570.38 (b)(1) and (2) amended	18281
571 Authority citation revised;	
sectional authority citations	
removed	39639
573 Opportunity for comment	29019
Comment time extended	33673
Authority citation revised; sec-	
tional authority citations re-	
moved	39639
573.460 (a)(1) footnote 1 amend-	
ed	18281
573.640 (b)(4) (i) and (ii) amend-	
ed	18281
573.780 (b)(1) amended	18282
573.920 (a) revised; (f) added	14215
(f)(3) corrected	15874
579 Authority citation revised	39639
582 Authority citation revised	39639
582.99 Authority citation re-	
moved	39639
582 Authority citation revised	39639
584.200 Authority citation re-	
moved	39639
589 Authority citation revised	39639

1990

21 CFR

55 FR
Page

Chapter I	
509 Subpart C heading redesign-	
ated as Subpart D; new Sub-	
part C heading added	20786
Subpart D heading redesignated	
from Subpart C	20786
509.4 (c) redesignated as (d); head-	
ing, (b), and new (d) revised;	
new (c) added	20786
509.6 (c) and (d) redesignated as	
(d) and (e); (a), new (d) and (e)	
revised; new (c) added	20786
509.7 (b) revised	20786
509.8 Removed	20787
510.450 Removed	32396
510.600 (c)(1) table and (2) table	
amended	1405,
1673, 8460, 8461, 13902, 14830, 17952,	
18330, 19874, 23703, 24226, 25301,	
25971, 26683, 29842, 30214, 32616,	
37226, 39615, 39616, 45798, 51409	
514.2 (d) added	14831
514.9 Interpretative rule	25972
514.106 Added	46052
(b)(2)(iii) corrected	49973
520.48 (c)(1) and (3) amended	26431
520.260 (b)(2) amended	24556
520.540 (b)(2) amended	8461
520.580 (b)(1) amended	39616
520.622a (a)(3) amended	8461
520.622c (b)(4) amended	8462
(b)(3) amended	32616
520.812 (c)(3) amended	24226
(c)(2) redesignated as (c)(2)(i);	
(c)(2)(ii) added; (c)(3) revised	43327
520.1263c (d) revised	3209
520.1445 Added	25301
(c)(3) corrected	49888
520.1448 Amended	3586
520.1630 Revised	46943
520.1720a (b)(3) amended	8462
520.1720d (b) amended	8462
520.1870 Nomenclature change;	
(c)(1)(i), (iii), and (2)(iii) re-	
vised	2234
520.2080a Removed	24556
520.2087 (b) amended	8460
520.2088 (a)(2) and (b)(2) amend-	
ed	8460
520.2150a (b) amended	23076
520.2150b (b) amended	23076
520.2184 (b) amended	8460
520.2260b (c)(1) and (e)(1) amend-	
ed	8462
520.2320 (c) amended	8460

21 CFR (4-1-99 Edition)

21 CFR—Continued

55 FR
Page

Chapter I—Continued	
520.2325a (c)(1) amended	29843
520.2325b (c) amended	29843
522 Technical correction	36751
522.144 (c) revised	26683
522.313 (d)(1)(i) amended	13768
522.540 (d)(2)(i) and (e)(2) amend- ed	8462
(d)(2)(ii) amended	13902
522.812 Added	26683
522.1079 Added	1405
522.1183 (e)(1) amended	8462
(f) removed	33670
522.1192 (d)(4)(ii) correctly added	3586
522.1193 Added	38984
522.1225 Added	40653
522.1228 Removed	33670
522.1290 Added	1185
522.1372 (b) amended	23076
522.1662a (i)(2) amended	8462
522.1680 (b) amended	8462
522.1698 (b) amended	23076
522.1720 (b)(1) amended	8462
(b)(1) amended	13902
522.1880 (b) amended	13902
522.1940 (a) removed; (b) and (d)(1)(iii) revised	13769
522.2150 (b) amended	23076
522.2662 (b) revised	18724
(b) amended	32616
522.2670 (b) amended	32616
524 Technical correction	797
524.1193 Added	50551
524.1580b (b) amended	8462
(b) revised	20455
524.1580d (b) amended	8462
529.2090 (b) correctly amended	31481
(c) amended	29842
540.181b (c)(2) amended	8460
540.255a Removed	25973
540.874a (c)(3)(i) and (4)(i) amend- ed	8462
544.275 (c)(2) amended	13902
555.110a (c)(2)(ii) amended	8462
555.210 (c)(2) amended	8462
556.360 Revised	3209
556.495 Added	46943
558 Technical correction	37287
558.4 (d) table amended	1406
18599, 23424, 46513	
558.15 (g)(1) table amended	8462
(g)(2) table amended	8460
558.35 (a) amended	8460
558.55 (d)(2) table amended	8460
558.58 (d)(1) table amended	8460

21 CFR—Continued

55 FR
Page

Chapter I—Continued	
558.95 (b)(1)(iii)(b), (iv)(b), (v)(b), (vii)(b), (ix)(b), (x)(b), and (xi)(b) amended	8460
558.105 (d)(1)(ix)(b) amended	8460
558.195 (d) table amended	8460
558.258 (c)(1)(i) and (ii) revised	48231
558.274 (a)(4) and (c)(1) table amended	25972
558.311 (e)(1) table amended	8460
Correctly designated	3685
Corrected	5446
Technical correction	15099
(e)(1) table amended; (e)(3) added	31828
558.325 (c)(2)(i)(b), (ii)(b), and (v)(b) amended	23424
(c)(2)(v)(b) corrected	25202
(c)(2)(ii)(b) corrected	34011
558.340 (c)(2)(ii) amended	23
(c)(4)(ii) amended	8460
(c)(3) and (4) removed	49616
558.342 (a), (c)(1)(ii), (2) introduc- tory text, (ii), (3) introductory text, and (ii) revised; (c)(4) through (6) added	31828
(c)(4)(ii) corrected	34007
558.355 (a) revised	3586
(f)(1)(x)(b), (xi)(b), (xii)(b), (xv)(b), (xvi)(b), (xviii)(b), (xix)(b), (xx)(b), and (xxiii)(b) amended	8460
(f)(2)(iii) revised	11012
(f)(3)(iv) revised	31829
(b)(4) through (9), (11), and (12) revised	42704
(f)(3)(vii) added	43328
558.363 (c)(1)(V)(B) amended	8460
558.369 (a) amended	8460
558.376 (a) amended	8460
558.430 (a) amended	8461
558.450 (d)(1) Table 1 amended	17599
558.485 (e) introductory text amended; (e)(1) introductory text through (12)(iii) redesign- ated as (e)(1)(i) through (xii)(C); new (e)(1) introduc- tory text and (2) added; (a)(1) through (23), (25), (27), new (e)(1)(x)(B) and (xi)(B) re- vised	18599
(a)(14) removed	30215
558.515 (d)(1)(ii)(b) amended	8461
558.530 (a)(2) amended	8461
558.550 (b)(1)(ii)(c), (iv)(c), (v)(c), (ix)(c), (xii)(c), (xiv)(c), (xv)(c) amended	8461

List of CFR Sections Affected

21 CFR—Continued

55 FR
Page

Chapter I—Continued

(b)(3)(iii) added	17599
558.586 (a) amended	29843
558.625 (b)(39) amended	8462
(b)(20) removed	17952
(b)(47) and (75) removed	18330
(b)(46) removed	24226
(b)(69) removed	25972
(b)(49) and (56) removed	30215
(f)(2)(ii) added	31829
(b)(26) and (86) removed	39615
(b)(82) removed	39616
558.630 (b)(10) amended	25972
(b)(8) amended	30215
(b)(10) amended	39615, 39616
558.680 (a) amended	8461

1991

21 CFR

56 FR
Page

Chapter I

500.29 Added	40506
500.30 Added	40507
510.515 (a)(4), (5), (b)(11), (15) and (17)(ii) removed; (c) table amended	41912
510.600 (c)(1) table and (2) table amended	7805, 9623, 9840, 14019, 14641, 16269, 19022, 19023, 19263, 20126, 27196, 37472—37474, 43698, 49410, 49845, 50652, 63875, 64553, 65834
(c)(1) table amended	9841
514.106 (a), (b)(1)(ii), (iii), (v), (vii), (xiii), (xiv), (2)(iii), (iv), (ix) and (xiii) corrected	12422
520 Authority citation cor- rected	14978
520.23 (a)(2) amended	37473
520.45a (b) amended	50653
Corrected	59331
520.45b (b) amended	50653
520.182 (b) amended	43699
520.540c (b) amended	50653
520.550 (b) amended	50653
520.622c (b)(2) amended; (b)(7) re- moved	50653
520.623 (b) amended	50653
520.816 (b) amended	50653
520.903d (c)(4) added	50813
520.903e Added	50655
520.1010a (b) amended	37473
520.1065 Removed	31075
520.1204 Heading revised; (a) and (c)(2) amended	8710
520.1205 Added	8710
520.1284 (b) amended	50653

21 CFR—Continued

56 FR
Page

Chapter I—Continued

520.1450c Added	13396
520.1630 (d)(2) introductory text and (iii) corrected	8710
520.1638 (b) amended	50653
520.1640 (b) amended	50653
520.1840 (c)(1) and (2) amended	50653
520.1846 (b) amended	9841
520.1900 (b) amended	37473
520.1920 (b) amended	50653
520.1921 (b) amended	50653
520.2260a (b)(1) amended	50653
520.2260b (a) amended	50653
Corrected	59331, 64288
520.2260c (b)(1) amended	50653
Corrected	59331, 64288
520.2604 (b) amended	50653
520.2605 (b) amended	50653
522.313 (d)(1)(ii) and (iii) amend- ed	12119
522.540 (d)(2)(ii) amended	19022
(d)(2)(i) amended	50653
522.562 Removed	31075
522.900 (a) removed	9623
522.1081 (a)(2)(ii) amended	14642
(a)(2)(iii) added	67175
522.1145 (e) added	50814
522.1192 (d)(4)(i) revised	14020
522.1290 (b) amended	50653
522.1301 Removed	31075
522.1680 (b) amended	14642, 16002
522.1720 (b)(2) amended	16002
(b)(1) amended	19022
522.1880 (b) amended	19022
522.1920 (b) amended	50653
522.2477 Added	67175
524.1005 (b)(1) amended	50653
524.1465 (b) amended	50653
524.1580b (b) amended	37473
(b) amended	50653
524.1580c (b) amended	50653
524.1580d (b) amended	37473
524.1600a (b) amended	50653
524.2101 (c) introductory text amended	9623
524.2620 (b)(2) amended	37474
529.1526 (b) amended	43699
540.103a (c)(2) amended	50653
540.103b (c)(2) amended	50653
540.103c (c)(2) amended	50654
540.103d (c)(2) amended	50654
540.103e (c)(2) amended	50654
540.103g (c)(2) amended	50654
540.103h (c)(2) amended	50654
540.107a (c)(2) amended	50654
540.107e (c)(4)(ii) introductory text amended	50654

21 CFR (4-1-99 Edition)

21 CFR—Continued

56 FR
Page

Chapter I—Continued

540.203 (c)(1)(ii) and (2)(ii) amend- ed	50654
540.207a (c)(2)(i) amended	50654
540.209 (c)(2) amended	50654
540.255c (c)(2)(i) and (ii) amend- ed	50654
540.680 (c)(2) amended	50654
540.803 (c)(2) amended	50654
540.814 (c)(2)(ii) amended	50654
540.814a (c)(2)(ii) amended	50654
540.815 (c)(2) amended	50654
540.815a (c)(2) amended	50654
546.110c (c)(2) amended	20126
546.180d (c)(6)(iv)(d) added	7806
555.110b (c)(2)(ii) amended	19263
556.240 Heading and introductory text revised	67175
556.250 Removed	67175
556.308 Amended	8711
558.3 (b)(1) introductory text re- vised	19268
558.4 (d) table amended	19268, 29897, 41912
558.15 (g)(1) table and (2) table amended	41912, 64702
558.20 Table amended	19269
558.62 (a) and (c)(1) revised	19268
Corrected	37839
558.76 (d)(3)(xiv) removed	9623
(d)(1) table amended	21076
558.95 (a)(5) removed	9623
(a)(1) revised; (b)(1) heading and (2) heading amended	49410
558.108 Added	19269
558.145 (a)(1) amended	14019
558.195 (d) table amended	15498
Technical correction	23105
558.258 (a) revised	9841
558.262 Removed	41912
558.274 (a)(2), (4) and (c)(1) table amended	9623
(a)(4) and (c)(1) table amended	16269
558.325 (a)(2) and (6) removed	9623
(a)(5) amended	48732
(a)(4) removed	49846
(a)(11) removed	50049
(a)(9), (10), (12) and (15) removed	50655
(a)(8) and (14) removed	66573
558.342 (c)(4)(ii), (5)(ii) and (6)(ii) amended	14021
558.370 Removed	41912
558.464 (a)(1) amended	50654
558.465 (a) amended	50654
558.485 (a)(6) removed	9623
558.530 (d)(3)(xxiv) removed	12349

21 CFR—Continued

56 FR
Page

Chapter I—Continued

558.625 (b)(3), (32), (50), (60), (65), (87), and (88) removed	9623
(b)(23) removed	16269
(b)(27) removed	50655
558.630 (b)(9) removed; (b)(10) amended	9623
(b)(10) amended	16269, 19023, 50655
558.635 (b)(1) amended	50654
589.1000 (Subpart B) Added	40507

1992

21 CFR

57 FR
Page

Chapter I

500.27 (d) amended	6475
500.51 (c) amended	6475
510 Technical correction	21854
510.112 (e) amended	6475
510.302 (d) amended	6475
510.310 (f) amended	6475
510.600 (c)(1) table and (2) table amended	2836, 5052, 7651, 8577, 11683, 12711, 26995, 30641, 31313, 32175, 35988, 38442, 45984, 48163, 54176, 61796
511.1 (e) amended	6475
514.1 (d)(2) amended	6475
520 Authority citation cor- rected	6352
520 Heading revised	37319
520.88 Added	37319
520.88a Added	37319
520.88b Added	37319
(b)(1)(ii)(B) corrected	42623
520.88c Added	37319
520.88d Added	37319
(c) corrected	42623
520.88e Added	37320
520.88f Added	37320
520.88g Added	37320
520.88h Added	37320
520.90 Added	37321
520.90a Added	37321
520.90b Added	37321
520.90c Added	37321
520.90d Added	37321
520.90e Added	37322
520.90f Added	37322
520.154 Added	37322
520.154a Added	37322
(d)(3)(iii) corrected	42623
520.154b Added	37322
520.154c Added	37322
520.390 Added	37323
520.390a Added	37323
520.390b Added	37323

List of CFR Sections Affected

21 CFR—Continued

57 FR
Page

Chapter I—Continued

520.390c Added	37323
(c)(1) corrected	42623
520.445 Added	37323
520.445a Added	37323
520.445b Added	37324
(b), (d)(1)(iv)(A) introductory text and (2)(i)(D)(1) corrected; (d)(2)(i)(C) correctly des- ignated	42623
520.445c Added	37325
520.580 (b)(1) amended	35989, 45984
520.608 Added	37325
520.1130 Added	37325
520.1130a Added	37325
520.1130b Added	37326
520.1130c Added	37326
520.1242g (c) amended	7652
520.1448b (b) amended	5052
520.1484 Added	19085
(b) revised	26604
520.1660d (b) revised	42490
520.1696 Added	37326
520.1696a Added	37326
520.1696b Added	37326
520.1696c Added	37326
(c) corrected	42623
520.1696d Added	37327
520.1720b (b) amended	2836
520.1806 (b) amended	48163
520.1880 Added	4718
520.2041 (b) amended	48163
520.2042 (b) amended	48163
520.2087 (b) amended	8577
520.2088 (a)(2) and (b)(2) amend- ed	8577
520.2158 Added	37327
520.2158a Added	37327
520.2158b Added	37327
(b) corrected	42623
520.2158c Added	37327
520.2220a (b) amended	32176
520.2345 Added	37327
520.2345a Added	37327
520.2345b Added	37327
520.2345c Added	37328
520.2345d Added	37328
(e)(1)(i)(A), (ii)(A) and (4)(iii)(A)(1) corrected	42623
520.2345e Added	37329
520.2345f Added	37329
520.2345g Added	37329
520.2345h Added	37329
522 Technical correction	5295
Heading revised	37330
522.88 Added	37330
522.90 Added	37330

21 CFR—Continued

57 FR
Page

Chapter I—Continued

522.90a Added	37330
522.90b Added	37331
(d)(3)(ii) corrected	42623
522.90c Added	37331
522.313 (d)(2) and (3) added	41862
522.390 Added	37331
522.650 Added	37331
(b) corrected	42623
522.840 (a), (c)(1) and (3) amend- ed	41861
522.1145 (d)(2) amended	2837
522.1696 Added	37332
522.1696a Added	37332
(d)(2) introductory text cor- rected	42623
522.1696b Added	37332
Heading, (d)(3)(ii) and (e)(1)(ii) corrected	42623
522.1696c Added	37333
522.2100 (a)(1) amended	21209
522.2471 Added	12712
522.2477 (b) corrected	5052
524 Heading revised	37333
524.154 Added	37333
524.155 Added	37333
524.390 Added	37333
524.390a Added	37333
524.390b Added	37333
524.390c Added	37334
524.390d Added	37334
524.900 (c) amended	7652
524.1005 (b)(1) amended	31314
526 Heading revised	37334
526.88 Added	37334
526.464 Added	37334
526.464a Added	37334
(c)(2) corrected	42623
526.464b Added	37334
(c)(2) and (3) corrected	42623
526.464c Added	37335
526.464d Added	37335
526.1130 Added	37335
526.1696 Added	37335
526.1696a Added	37335
526.1696b Added	37336
526.1696c Added	37336
(a) corrected	42623
526.1696d Added	37336
(c)(1)(iii) corrected	42623
529 Heading revised	37336
529.1030 (b) amended	42490
529.2464 Added	37336
536 Removed	37336
539 Removed	37336
540 Removed	37336
540.274b (c)(3)(iv)(c) amended	32896

21 CFR (4-1-99 Edition)

21 CFR—Continued

57 FR
Page

Chapter I—Continued

544	Removed.....	37336
546	Removed.....	37336
546.180d	(c)(6)(i)(a)(3), (b)(3), (ii)(d), (iii)(a)(3), (b)(3), (c)(3), (iv)(a)(3) and (b)(3)	7651
	Corrected	8961
	(c)(6)(iv)(d)(3) amended.....	12711
	(c)(6)(i)(a)(3), (b)(3), (ii)(d), (iii)(a)(3), (b)(3), (c)(3), (iv)(a)(3) and (b)(3) amended	26996
548	Removed.....	37336
548.114	(c)(2) amended.....	7652
555	Removed.....	37336
556.113	Revised	41862
556.308	Amended	21209
556.735	Added	12712
558	Technical correction.....	21854, 44791
558.4	(d) table amended	525
	(c) revised	1642
	Technical correction.....	5210
	(d) table amended; eff. 6-30-92	6072
	Corrected	6892
	(d) amended	38442
558.5	(c)(2) amended.....	6475
558.15	(d) and (e) amended	6476
	(g)(2) table amended.....	8577
	(g)(1) table and (2) table cor- rected	14639
558.20	Removed	1642
	Technical correction.....	5210
558.55	(d)(2) table corrected; CFR correction	12712
558.58	(d)(1) table amended	7652, 8577
558.76	(d)(3)(xiii) revised.....	32176
558.78	(a)(2), (d)(1) table and (d)(2)(ii) amended	7652
558.95	(b)(1)(iii)(b), (iv)(b), (v)(b), (vii)(b), (ix)(b), (x)(b) and (xi)(b) amended	8577
	(a)(2) amended	23059
558.105	(d)(1)(ix)(b) amended	8577
558.108	Removed	525
558.120	(c)(1)(iii)(b) amended.....	7652
558.195	(d) table amended	8577
558.235	Added	38442
558.254	(a) amended.....	7652
558.258	(a) and (c)(1)(i) revised; (c)(1)(ii) and (iii) redesignated as (c)(1)(i)(A) and (B); new (c)(1)(ii) through (v) added	34516
558.274	(a)(2), (4) and (c)(1) table amended	37410
558.311	(e)(1) table amended	8577
558.325	(c)(4)(ii) added	34517
558.342	(c)(4)(ii), (5)(ii) and (6)(ii) revised	5053

21 CFR—Continued

57 FR
Page

Chapter I—Continued

	Technical correction.....	6892
558.355	(f)(3)(viii) added	5054
	(f)(1)(xxiv) revised	6554
	Technical correction.....	6892
	(b)(9), (f)(1)(iv)(b), (v)(b), (xv)(b) and (xvi)(b) amended	7652
	(f)(1)(x)(b), (xi)(b), (xii)(b), (xv)(b), (xvi)(b), (xviii)(b), (xix)(b), (xx)(b) and (xxiii)(b) amended.....	8578
	(f)(3)(ii)(a) revised; (f)(3)(ix) added	11683
	(b)(7) and (14) revised	23953
	(b)(11) revised; (f)(1)(xxv) added	46504
558.363	(c)(1)(v)(B) amended	8578
	(c)(1)(vi) added.....	32176
558.369	(a) amended.....	8578
558.430	(a) amended	8578
558.515	(d)(1)(ii)(b) amended	8578
558.530	(a)(2) amended.....	8578
558.550	(b)(1)(vii)(c) and (ix)(c) amended	7652
558.625	(b)(51) removed	23059
	(b)(3) added	37410
	(b)(28) removed	54176
558.630	(b)(10) amended.....	37410
558.680	(c)(1) table revised	8403
	(a) amended.....	8578
570.6	(e) amended.....	6476
571.1	(c) amended.....	6476
573	Hearing.....	28606
	Hearing denied	33244
	Technical correction	40318
573.870	Added	7875
	(a) table amended	24187

1993

21 CFR

58 FR
Page

Chapter I

510	Authority citation revised	44611
510.515	Introductory text, (b)(7)(i) and (c) table amended; (b)(7)(i)(c) removed	30119
510.600	(c)(1) table and (2) table amended.....	4316, 5607, 26523, 30118, 36135, 37855, 38971, 44611, 47056, 47377, 59169, 59947, 63890
514	Authority citation revised	51011
514.210	Removed	51011
520.88d	Heading corrected.....	18304
520.90b	(d)(2)(ii) correctly des- ignated.....	18304
520.90c	(b) amended	61016

List of CFR Sections Affected

21 CFR—Continued

58 FR
Page

Chapter I—Continued	
520.90d (b) amended.....	61016
520.90e (b) amended.....	61016
520.90f (b) amended.....	61016
520.182 (b) amended.....	5608
520.240 Removed.....	41025
520.445b (b) amended; (b) heading and (d)(4) introductory text re- vised.....	61015
520.580 (b)(1) amended.....	41025
520.1448a (a)(2) amended.....	26523
520.1485 Added.....	38972
520.1660d (a) and (b) revised; (e)(1)(iii), (iv), (v) and (2) added.....	42854
(a)(5) added; (b)(3) revised; (e)(1)(ii)(A)(3), (B)(3) and (C)(3) amended.....	43794
520.1720c (b) amended.....	29777
520.1806 (b) amended.....	44611
520.1840 (c)(3) amended.....	26523
520.1870 (c)(1)(i) revised.....	42853
520.1871 Added.....	58652
520.1196 Added.....	8542
520.1445 (c)(2) revised.....	5608
520.1870 (a) and (c)(2)(ii) re- vised.....	7864
520.2041 (b) amended.....	44611
520.2042 (b) amended.....	44611
520.2088 (c) added.....	65664
520.2089 Added.....	65665
520.2158a (d)(1) and (2) revised; (d)(3) removed.....	47211
520.2220a Heading, (a), (b) and (e) introductory text revised.....	6092
520.2345d (e)(1)(iii)(A), (B), (C), (3)(iv)(A), (B) and (C) re- vised.....	33331
520.2455 Heading revised; (d)(2)(i) and (ii) amended.....	14313
520.2456 Added.....	14313
520.2613 Added.....	36135
522.90b (d)(2)(ii) correctly des- ignated.....	18304
522.575 Added.....	500
522.820 Revised.....	43795
522.1002 Redesignated from 522.1822 and revised.....	47377
522.1079 (c)(2) and (3) revised.....	52222
522.1696a (b)(1) amended.....	65285
522.1696b (d) introductory text re- vised.....	11964
522.1822 Redesignated as 522.1002.....	47377
522.1870 (c)(1)(ii) revised.....	42853
522.2100 (c), (d) and (e) added.....	57556
522.2112 Added.....	59947

21 CFR—Continued

58 FR
Page

Chapter I—Continued	
522.2220 (a)(2) revised.....	38972
522.2670 Revised.....	8543
524.1044g Added.....	38973
524.2101 (c) amended.....	41025
526.464b (c) introductory text amended.....	61016
526.1696a (f) removed.....	500
526.1810 Added.....	58486
529.1030 (c)(1)(iii) and (2)(iii) added.....	59169
529.1044a (b) amended.....	14314
529.1186 (b) amended.....	17346
556.230 (a) and (b) revised.....	43795
556.500 (e) added.....	42855
556.515 Added.....	58486
556.610 Revised.....	47211
558 Technical correction.....	17346
558.4 (d) table amended.....	30119, 58653
558.15 (g)(1) table and (2) table amended.....	17515
(g)(1) table amended.....	30119
(g)(2) table amended.....	30120
558.55 (d)(2) table amended.....	30120
558.58 (d)(1) table amended.....	30120
558.95 (b)(1)(xi)(b) and (xii)(b) amended.....	47056
(a)(1) revised; (b)(4) redesi- gnated as (b)(5); new (b)(4) added.....	54287
558.258 (c)(3)(iv)(C) amended.....	47973
558.274 (c)(1) table amended.....	30120
558.300 Added.....	58653
558.311 (b)(4) and (6) revised; (e)(3) redesignated as (e)(4); new (e)(3) added.....	43560
(e)(2)(i) table amended.....	47973
558.355 (f)(3)(ix) amended.....	4316
(d)(7)(vi) added.....	17516
(b)(13) revised.....	53883
558.460 (c)(1) table amended; (c)(2)(vii) and (viii) removed; (c)(2)(ix) and (x) redesignated as (c)(2)(vii) and (viii).....	30120
558.550 (a) amended.....	47056
558.625 (b)(54) amended.....	63890
573.750 Added.....	59170
573.920 Regulations at 52 FR 10888, 21001, and 54 FR 14215 stayed.....	47973
579.22 (b) table revised.....	18148

1994

21 CFR

59 FR
Page

Chapter I	
500.80 (a) amended.....	14365

21 CFR (4-1-99 Edition)

21 CFR—Continued

59 FR
Page

Chapter I—Continued	
509.30 (b) amended.....	14365
510 Technical correction.....	49291
510.600 (c)(1) table and (2) table amended	1919, 14367, 28768, 31139, 33197, 33198, 33908, 38119, 41664, 41976, 44315, 49576, 50829, 59364
520 Technical correction.....	50829
520.45a (a), (b), (c), (d)(1), (2) and (3) redesignated as (a)(1), (2), and (3), (a)(4)(i), (ii) and (iii); new(b) added	65711
520.445b (a) amended; (b), (d)(1), (2), (4)(iii)(B), (C), (iv)(B) and (C) revised; (d)(3) removed; (d)(4)(i)(C) added	39439
520.531 (a) removed; (b) revised	26942
520.580 (b)(1) amended	59365
520.804 Added	17694
520.903c Removed.....	33198
520.905a (d)(3) added	26943
520.905b (c)(3) added	35252
520.1660d (a)(4), (b)(2) and (e)(1)(iii)(C) revised.....	3318
(a)(6) and (b)(4) added; (e)(1)(ii)(A)(3), (B)(3), (C)(3) and (iii)(C) amended	41664
(e)(1)(ii)(A)(3), (B)(3) and (C)(3) corrected	56388
520.1696b (b) revised; (c)(2) amend- ed	42493
(b) revised.....	58775
520.1720a (b)(3) amended	53585
520.1806 (b) amended.....	28769
520.1872 Added	33908
520.1962 (b) removed.....	5705
520.2045 (b)(2) amended.....	28769
520.2155 Removed	1919
520.2220a (b) amended.....	56000
520.2220b (b)(1) amended.....	56000
520.2220c (c) amended.....	56000
520.2220d (b) amended.....	56000
520.2260a (b)(1) amended	22754
520.2260b (b)(1) amended	22754
520.2260c (a) amended.....	22754
520.2325a (c)(2) amended	28769
(c) amended	33197
520.2325b (c) amended.....	33197
520.2345a (b)(1) amended	59365
520.2345d Revised	17693
(a)(1) amended	19133
520.2520h Removed	1919
520.2640 (a) amended.....	14365
522.246 (a) amended; (c)(3) added.....	41665
522.313 (d)(4) added	41666
(d)(5) added	54518

21 CFR—Continued

59 FR
Page

Chapter I—Continued	
522.900 (b)(2) amended	14367
522.1145 (a)(2) amended.....	33198
522.1222a (c) revised.....	41664, 41976
Corrected.....	49291
522.1350 Added	37422
522.1680 (b) amended.....	31139
522.1696a (b)(1) and (2) amend- ed	38119
522.1720 (b)(1) amended.....	53585
522.1940 (e) redesignated as (d)(2)(iv); (b), (d)(2)(ii), (iii) and (iv) revised; interim	49808
522.2220 (a)(2)(i) amended	56000
522.2640a (a) amended.....	14365
522.2640b (a) amended.....	14365
522.2662 (b) amended.....	14367
522.2680 Revised	19639
524.900 (c) amended	28769
524.1484h Added	5105
524.1580d (b) amended	5705
524.1580b (b) amended.....	33197
524.1580c (b) amended.....	33197
524.1580d Removed	53586
524.2640 (a) amended.....	14365
529.1030 (b) revised; (d) redesis- gnated as (c)(3).....	60076
529.1044a (b) amended.....	31140
529.1186 (b) amended.....	44315
556.34 Revised	65711
556.200 Revised	41977
556.275 Revised	26943
556.344 (a) revised; (d) added	50830
556.380 Revised	41241
556.425 Revised	17922
556.750 (c) added	38902
558.4 (d) table amended	17477, 18297, 41241
558.95 (b)(1)(xi)(b) amended.....	14368
(b)(4)(ii) added	15624
(b)(1)(xii)(b) amended	19134
(b)(1)(x)(b) and (xi)(b) amended	33197
558.128 (c)(5)(xiii) removed	8134
558.258 (c)(4) added	35252
558.265 (c)(3) added	51498
558.274 (a)(4) and (c)(1) table amended.....	59365
558.305 Added	18297
558.311 (e)(1) table amended.....	33197
558.340 (c)(2) removed	8134
558.342 (c)(1)(ii), (2)(ii), (3)(ii), (4)(ii), (5)(ii) and (6)(ii) amend- ed	41241
558.355 (f)(1)(xii)(b) and (xx)(b) amended	33197
(f)(3)(iv)(b) and (viii) amended	41241

List of CFR Sections Affected

21 CFR—Continued

59 FR
Page

Chapter I—Continued	
(f)(2)(ii) and (iii) redesignated as (f)(2)(i)(a) and (f)(2)(i)(b); new (f)(2)(i)(b) revised; new (f)(2)(ii) added.....	62320
558.360 (d)(2) redesignated as (d)(2)(i); new (d)(2)(i) heading and (ii) added; (d)(3) amended.....	17922
558.550 (a) revised.....	12547
(a) revised; (b)(1)(ii)(c), (iv)(c), (v)(c), (ix)(c), (xii)(c), (xiv)(c) and (xv)(c) amended.....	14368
(a)(2) amended.....	15339
(a)(2) revised.....	15339
(a)(2) amended.....	19134, 21930, 26423, 26943
(a)(2), (b)(1)(x)(c), (xii)(b) and (c) amended.....	26424
(b)(1)(ii)(c) amended.....	33197
(b)(2) added.....	49577
(a)(2) corrected.....	64739
(a) revised.....	67185
558.555 Added.....	17477
558.586 (a) amended.....	33197
558.600 (c)(1)(i) amended; (c)(3) added.....	41975
558.625 (a) amended.....	14365
(b)(71) removed.....	33198
558.635 (f)(3) redesignated as (f)(4); new (f)(3) added.....	38902
(b)(2) amended.....	54518
573.130 Added.....	26711
573.920 Stay at 58 FR 47973 confirmed.....	45973

1995

21 CFR

60 FR
Page

Chapter I	
500.26 (a) amended.....	38480
500.27 (a)(3) amended.....	38480
501.4 (b)(13) amended.....	38480
510.413 (b) amended.....	38480
510.600 (c)(1) table and (2) table amended.....	7121, 11027, 14216, 16571
(c)(1) table and (2) table amended.....	29755, 32447, 33110, 33342, 34135, 35122, 35838, 39846, 40454, 40455, 47052, 47480, 54193, 55658, 63622
520.45a (b) amended.....	55658
520.45b (b) amended.....	55658
520.88a (b) amended.....	55658
520.88b (b) amended.....	55658
520.88c (b) amended.....	55658
520.88d (b) amended.....	55658
520.88e (b) amended.....	55659

21 CFR—Continued

60 FR
Page

Chapter I—Continued	
520.88g (b) amended.....	55659
520.88h (b) amended.....	55659
520.90b (b) amended.....	55659
520.90f (b) amended.....	55659
520.445b (d)(4) introductory text amended.....	26827
(b) and (d)(4)(iii)(C) amended.....	47052
520.540c (b) amended.....	55659
520.550 (b) amended.....	55659
520.622c (b)(2) amended.....	55659
(b)(3) amended.....	57832
520.623 (b) amended.....	55659
520.704 Removed.....	63622
520.816 (b) amended.....	55659
520.1263c (a) revised.....	14217
520.1284 (b) amended.....	55659
520.1288 Added.....	362
520.1289 Added.....	20402
520.1445 (c)(2) and (3) amended.....	50097
520.1484 (b) revised.....	14217
520.1485 Heading and (b) revised.....	3079
520.1560 Removed.....	39847
520.1560a Removed.....	39847
520.1560b Removed.....	39847
520.1638 (b) amended.....	55659
520.1640 (b) amended.....	55659
520.1660d (b)(2) amended; (c) removed.....	40455
520.1696b (b) revised.....	26359
520.1801 Removed.....	39847
520.1801a Removed.....	39847
520.1840 (c)(1) and (2) amended.....	55659
520.1880 (b) amended.....	57832
520.1920 (b) amended.....	55659
520.1921 (b) amended.....	55659
520.2095 Added.....	50097
520.2260a (b)(1) amended.....	55659
520.2260b (b)(1) amended.....	55659
520.2260c (a) amended.....	55659
520.2604 (b) amended.....	55659
520.2605 (b) amended.....	55659
522.88 (b) amended.....	55659
522.90a (b)(1) amended.....	55659
522.90c (b) amended.....	55659
522.313 (d)(1)(ii) amended.....	51719
522.514 Removed.....	63622
522.540 (a)(2) revised.....	29985
(d)(2)(i) amended.....	55659
(d)(2)(ii) removed; (d)(2)(iii) redesignated as (d)(2)(ii).....	63622
522.723 (c) amended.....	39847
522.883 (c) amended.....	39847
522.970 (b) revised.....	48650, 54942
522.1044 (a) and (b)(1) revised; (b)(4) added.....	29985

21 CFR (4–1–99 Edition)

21 CFR—Continued

60 FR
Page

Chapter I—Continued	
522.1060	Removed.....27223
522.1060a	Removed.....27223
522.1060b	Removed.....27223
522.1085	(b) amended.....27223
522.1086	Added.....27223
522.1192	Heading and (a)(3) heading revised; (d)(5) added.....45041
522.1222a	(a) and (d) removed; (c) revised.....49339
522.1290	(b) amended.....55659
522.1362	Added.....49340
522.1563	Removed.....39847
522.1660	(b) and (c)(2)(iii) amended.....363, 14218
	(b) and (c)(2)(iii) amended.....29755
522.1662a	(h)(2) revised.....48651
522.1720	(c) and (d) amended; (b)(2) revised.....53509
	(b)(1) amended.....63622
522.1880	Removed.....63622
522.1920	(b) amended.....55659
522.2095	Added.....57833
522.2100	(d)(2) revised.....57833
522.2477	Revised.....4376
522.2662	(b) amended.....33110, 35122, 35123
522.2670	(b) amended.....57832
522.2680	(d)(3) added.....26360
524.575	Added.....48651
524.1005	(b)(1) amended.....55659
524.1465	(b) amended.....55659
524.1580a	Removed.....39847
524.1580b	(b) amended.....55659
524.1580c	(b) amended.....55659
524.1600a	(b) amended.....55660
526.88	(b) amended.....55660
526.464a	(d) amended.....55660
526.464b	(d) amended.....55660
526.464c	(b) amended.....55660
526.464d	(b) amended.....55660
529.1044a	(b) amended.....45042, 48894
529.1186	(b) revised.....40456
529.2464	(b) amended.....55660
556.594	Added.....50098
558.4	(d) table amended.....39848, 57928, 66495
558.62	(c)(2)(v) removed; (c)(2)(vi) redesignated as (c)(2)(v).....39847
558.76	(d)(3)(v) added.....29483
558.95	(b)(1)(xi)(b) and (xii)(b) amended.....11027
	(b)(1)(x)(b) and (xi)(b) amended.....18741
	(a)(4) amended.....47052
558.105	Removed.....39847
558.128	(c)(5)(iii) removed.....39847
558.195	(c)(2) removed; (d) table amended.....34461

21 CFR—Continued

60 FR
Page

Chapter I—Continued	
	(d) table amended.....53702
558.274	(a)(7), (c)(1)(i) table and (ii) table amended.....47052
558.300	(c)(1)(i) and (iii) revised.....39848
558.305	(a) amended.....53509
558.311	(e)(1) table amended.....18741
	(b)(6) and (e)(1) table amended; (b)(7) added.....29481
	(b)(7) revised; (e)(1) table amended.....29482, 57929
	(b)(3) and (e)(1) table amended.....54194
558.325	(c)(3)(iv) removed.....39847
558.342	(c)(4)(ii)(C) amended.....11028
558.355	(f)(1)(xii)(b) and (xx)(b) amended.....18741
	(f)(3)(x) added.....63623
558.366	(c) table amended.....29483
	(a) amended.....33343
558.450	(d)(1) Table 1 amended.....3080
558.460	(c)(2)(v) removed.....39847
558.464	(a)(1) and (2) amended.....55660
558.465	(a) amended.....55660
558.485	(a)(11) amended.....47052
558.530	(d)(3)(vii) removed.....39847
558.550	(a)(2) revised; (b)(1)(ii)(a) and (xv)(a) amended.....3080
	(a)(1) amended; (a)(2) revised.....11027
	(b)(1)(ii)(c) amended.....18741
558.555	(b)(1)(iii) amended.....57928
558.625	(b)(10) revised.....7121
	(b)(11) and (15) removed.....8547
	(b)(52) amended.....47053
	(b)(83) amended.....66496
558.630	(b)(3), (8) and (10) amended.....47053
	(b)(10) amended.....66496
558.635	(b)(2) revised.....8547
	(b)(1) amended.....55660
573.920	Introductory text and (a) through (f) redesignated as (a) through (g); new (a) revised; interim.....53703
579.40	Added.....50099

1996

21 CFR

61 FR
Page

Chapter I	
500.23	Added.....37681
500.24	Added.....37681
500.49	Removed.....37682
500.50	Added.....19544
501.17	(c) revised; interim.....20101
505	Removed.....37682
507	Removed.....37682
508	Removed.....37682

List of CFR Sections Affected

21 CFR—Continued

61 FR
Page

Chapter I—Continued	
508.35 (c)(1), (2) introductory text and (ii) amended	14481
510.120 Removed	37682
510.200 (Subpart C) Removed	37682
510.310 Removed	37682
510.413 Removed	37682
510.515 (b)(7), (17), (25) and (29) removed; (b)(10) and (13) redesignated as (b)(1) and (2); (c) table amended	35950
510.600 (c)(1) table and (2) table amended	259, 515, 4735, 5506, 8872, 15703, 17565, 18672, 21075, 24441, 46547, 58630, 58631, 59003, 63710, 63711, 66580, 68146
520 Technical correction	43964
520.45a (a)(2) amended	4875
520.82a (b) amended	8873
520.82b (b) amended	8873
520.154b (b) amended	66581
520.222 (c) amended	8873
520.309 Added	66581
520.447 (c)(3) amended; (d) added	59003
520.500 Removed	34728
520.580 (b)(2) amended	34728
520.620a Removed	34728
520.620b Removed	34728
520.622a (a) amended	34728
520.622b (b)(2) revised	34728
520.622c (b)(5) amended	8873
520.784 (b) amended	8873
520.813 Added	56893
520.863 (b) amended	8873
520.905a (d)(2) heading revised; (d)(2)(i)(B) amended	29478
520.905c (d)(2) heading revised; (d)(2)(iii) amended	29478
520.960 (b) amended	5506
520.1120a (c) amended	8873
520.1120b (c) amended	8873
520.1193 (c)(2) and (3) revised	39868
520.1196 Heading revised; (c)(1)(iii) amended	15186
(c)(1)(ii) amended	59004
520.1197 Added	67452
520.1445 (c)(2) amended	43654
520.1448a (c)(4)(iii) amended	24444
(a)(4)(iii) and (d)(4)(iii) amended	53615
520.1468 (b) amended	5506
520.1484 (b) revised; (c)(3) amended	31027
(c)(3) amended	31399
520.1485 (b) revised; (d)(3) amended	31398

21 CFR—Continued

61 FR
Page

Chapter I—Continued	
520.1520 Removed	34728
520.1628 (b) amended	5506
520.1629 (a)(2) and (b)(2) amended	5506
520.1630 (b) amended	5506
520.1631 (b) amended	5506
520.1660d (a)(7) and (b)(5) added	2915
520.1660d (a)(7) revised; (e)(1)(ii)(A)(3), (B)(3), (C)(3), (iii)(C) and (iv)(C) amended	52690
520.1720a (b)(1) amended	8873
520.1720b (b) amended	8873
520.1720c (b) amended	8873
520.1805 (b) amended	8873
520.1872 (c)(1)(ii) amended	29651
520.2002 (c) amended	5506
520.2022 Removed	34728
520.2043 (b)(2) revised	29477
(b)(2) amended	63712
520.2122 (b)(2) amended	5506
520.2160a Removed	34728
520.2160b Removed	34729
520.2160c Removed	34729
520.2160d Removed	34729
520.2220a (b) amended	4875
520.2220b (b)(1) amended	4875
520.2220c (c) amended	4875
520.2220d (b) amended	4875
(c)(2) amended	46719
520.2260a (b)(1) amended	4875
520.2260b (b)(1) amended	4875
520.2260c (a) amended	4875
520.2325a (b) added; (d) removed; (c), (e) and (f) redesignated as (a), (c) and (d); new (a) revised; new (d) amended	24443
(a) revised	63711
520.2362 (c) amended	8873
520.2480 Removed	34729
520.2520a (b) amended	34729
520.2520c Removed	34729
520.2520d Removed	34729
520.2610 (b) amended	5506, 8873
520.2611 (b) amended	5506, 8873
520.2612 (b) amended	8873
522.82 (b) amended	8873
522.147 Added	48830
522.281 Removed	34729
522.313 (d)(3)(ii) revised; (d)(4) and (5) redesignated as (d)(5) and (6); new (d)(4) added	35130
(d)(7) added	66583
522.314 Added	29479
522.740 Removed	34729
522.770 Added	53321
522.784 (b) amended	8873

21 CFR (4-1-99 Edition)

21 CFR—Continued

61 FR
Page

Chapter I—Continued

522.842	(b) amended.....	5506
522.863	(b) amended.....	8873
522.883	(c) amended.....	260
522.914	(b) amended.....	5506
522.955	Added.....	42383
522.960a	(c) amended.....	5506
522.960b	(c) amended.....	5507
522.960c	(b) amended.....	5507
522.1004	Added.....	68147
522.1044	(b), (d)(2)(i) and (3)(i) re- vised.....	24441
522.1055	(b) amended.....	18672
522.1078	(b) amended.....	37682
522.1145	(a)(2) amended.....	59003
522.1155	(b) amended.....	8873
522.1182	(a)(4)(i) amended.....	18672
522.1290	(b) amended.....	66582
522.1335	Added.....	21075
522.1468	(b) amended.....	5507
522.1660	(b), (c)(1)(iii) and (2)(iii) amended.....	31028, 36291
522.1720	(b)(1) amended.....	8873
	(b)(1) and (2) revised.....	54333
522.1850	(c)(1) revised; (c)(2)(i) and (ii) amended.....	54333
522.1940	(b) and (d)(1)(iii) amend- ed.....	5507
522.2002	(c) amended.....	5507
522.2005	Added.....	66582
522.2012	(b) amended.....	5507
522.2022	Removed.....	34729
522.2120	(b) amended.....	5507
	(b) revised; (d)(4) amended.....	31028
522.2220	(a)(2)(i) amended.....	4875
522.2474	Added.....	25785
522.2477	(c)(3) added.....	29480
	(c)(3) heading and (iii) revised.....	41499
522.2478	Added.....	14482
	(a) amended.....	29479
522.2480	Removed.....	34729
522.2610	(a)(2) and (b)(2) amend- ed.....	5507, 8873
522.2615	(b) amended; (d) re- moved.....	29480
522.2662	(b) amended.....	46548
524.154	(a)(2) amended.....	8873
524.155	(a)(1) amended.....	8873
524.660a	(b) amended.....	5507
524.660b	(b) amended.....	5507
524.960	(b) amended.....	5507
524.981d	(b) amended.....	5507
524.1044b	(b) amended.....	48624
524.1484e	(b) amended.....	5507
524.1600a	(b) amended.....	63712
524.1742	(b) amended.....	8873
529.810	Removed.....	34729

21 CFR—Continued

61 FR
Page

Chapter I—Continued

529.1044a	(b) amended.....	17830
530	Added.....	57743
556.113	Amended.....	66583
556.150	Revised.....	67453
556.225	Added.....	53321
556.228	Added.....	56893
556.275	(c) added.....	29478
556.283	Added.....	42383
556.300	(a) amended.....	24441
556.430	Revised.....	31399
556.500	Revised.....	67453
556.600	Revised.....	31028
556.685	Added.....	24443
556.720	Revised.....	67453
556.735	Revised.....	68148
558.4	(d) table amended.....	68148
558.15	(g)(1) table amended.....	51589
558.55	(d)(2) table amended.....	515, 35950
558.58	(d)(1) table amended....	4875, 18082, 35951
558.76	(d)(3)(xiv) added.....	43451
	(d)(3)(xiv) revised.....	66584
558.78	(a)(2), (d)(1) table and (2)(ii) amended.....	18082
558.95	(b)(3)(iv)(b) amended.....	515
	(b)(1)(xiii)(b) amended.....	18082
	(b)(4)(iii) added.....	43655
558.120	(a)(1) and (2) amended.....	515
	(c)(1)(iii)(b) amended.....	18082
558.128	(a) amended.....	18082
	(a), (b) and (c)(1) revised; (c)(2) and (3) removed; (c)(4) and (5) redesignated as (c)(2) and (3).....	35952
	(c)(4) amended.....	36291
	(c)(2) corrected.....	53615
558.140	Added.....	35954
558.145	(a)(1) and (2) amended.....	18082
558.155	Heading, (d)(2) and (3) re- vised.....	2415
558.175	(c)(2)(ii) revised.....	35954
558.185	(a)(1) removed.....	34729
558.195	(d) table amended.....	35954
558.258	(c)(2) introductory text and (3) introductory text re- vised; (c)(2)(iii) and (3)(iii) amended.....	29478
558.265	(c)(2)(ii) added.....	21076
	(c)(2)(iii) added.....	24694
558.274	(c)(1) table amended.....	35955
558.300	(a) revised; (c)(2) through (5) added.....	32652
558.311	(b)(7)(iii) and (e)(1) table corrected.....	4349
	(b)(2) amended.....	4875
558.325	(c)(4)(iii) added.....	32653
558.340	(a) amended.....	18082

List of CFR Sections Affected

21 CFR—Continued

61 FR
Page

Chapter I—Continued	
558.355 (b)(5) amended.....	4875
(b)(8), (9), (f)(1)(iv)(b), (v)(b), (xiv)(b), (xv)(b), (xvi)(b) amended; (b)(10) added.....	18082
(f)(3)(iii)(b) and (v)(b) amended	24444
(f)(3)(x)(c) amended.....	51003
558.366 (a) revised; (c) table amended	1832
(c) table amended	14021, 14483
558.367 Removed	34729
558.450 Revised	51590
558.485 (a)(16) removed	34729
558.515 (a), (d)(1)(iii)(b), (iv)(b), (v)(b) and (vi)(b) amended.....	18082
(d)(1)(iii), (iv) and (v)(b) revised	35956
(d)(2) revised	51593
558.530 (d)(2), (3) and (4) redesignated as (d)(4), (5) and (6); (a) and new (d)(4) revised; new (d)(2) and new (3) added; new (d)(6)(i)(a) through (d) redesignated as (d)(6)(i)(A) through (D).....	35956
(d)(5)(xxiv) added.....	43451
558.550 (b)(1)(vii)(c), (ix)(c), (xv)(c) and (xvi)(c) amended	18082
558.555 (b)(2) added	29481
(b)(2)(ii) amended	43451
(b)(3) added	66584
558.565 Removed	34729
558.582 (a) amended	18082
558.618 Added	68148
558.625 (b)(25) amended	4875
(b)(79) amended.....	17566
(b)(72) removed	34729
(b)(76) removed	58631
(f)(1)(vi)(e) added	67713
558.630 (b)(10) amended.....	34729
558.635 (b)(1) amended	30133
558.680 (c)(1) table amended.....	35957
570.22 Removed.....	37682
573.460 (a) introductory text, (1), (2), (b), (c) introductory text, (1) and (2) redesignated as (a)(1) introductory text, (i), (ii), (2), (3) introductory text, (i) and (ii); new (b) added.....	15704
573.625 Added	5
573.870 Introductory text and (b) amended.....	11547
582.1666 (b) revised	19544
584.700 Added	43453
589.1001 Added	19544

1997

21 CFR

62 FR
Page

Chapter I	
500 Authority citation revised	51518
500.80 (b) amended.....	66983
501 Authority citation revised	51518
502 Authority citation revised	51518
509 Authority citation revised	51518
510 Technical correction.....	23534
Authority citation revised	51518
510.515 (c) table corrected; CFR correction	52659
510.600 (c)(1) table and (2) table amended	6724, 22887, 27691, 29010, 34631, 35075, 38906, 39442, 40931, 48939, 55159, 60781, 61624, 61626, 62241, 63270
(c)(1) table and (2) table amended; eff. 4-7-97.....	14300
Regulation at 62 FR 14300 eff. date corrected to 4-7-97	15751
511 Technical correction.....	47760
Authority citation revised	51518
511.1 (b)(10) revised	40599
514 Technical correction.....	47760
Authority citation revised	51518
514.1 (b)(14) revised	40599
514.8 (a)(1) amended	40600
514.110 (b)(10) revised	40600
514.111 (a)(9) revised.....	40600
520 Technical correction	8370
Authority citation revised	51518
520.23 (a)(2) amended.....	35075
520.82a (b) amended.....	61624
520.82b (b) amended.....	61624
520.88b (c) amended.....	13302
520.88d (d) heading and (2) revised; (d)(3) amended.....	5525
520.88e (d) heading and (2) revised; (d)(3) amended.....	5526
520.88f (b) amended	13302
520.100a (b) amended	63270
520.100b (b) amended	63270
520.100c (b) amended	63270
520.222 (c) amended	61624
520.300a (b) amended	63270
520.300b (b) amended	63270
520.300c (b) amended	63270
520.390a (a)(2) amended	35076
520.420 (a)(2) amended	63270
520.445b (b) amended ...	27691, 35076, 60656
520.447 (b) revised.....	46669
520.462 (b) amended	63270
520.580 (b)(2) amended	35076, 61624
520.600 (c) amended	35076
520.622a (a)(6) amended	35076
520.622b (c)(2) amended	35076
(a)(2) amended	38906

21 CFR (4-1-99 Edition)

21 CFR—Continued

62 FR
Page

Chapter I—Continued

520.622c (b)(6) amended	35076
(b)(6) corrected	40932
(b)(5) amended	61624
520.763a (c) amended	35076
520.763b (c) amended	35076
520.763c (b) amended	35076
520.784 (b) amended	61624
520.804 (b) amended	63270
520.812 (c) redesignated as (d); new (d) revised	38906
520.863 (b) amended	61624
520.1010a (b) amended	6724, 35076
520.1044a (b) amended	34169
520.1044b (b) amended	29011
520.1044c (c) amended	29013
520.1120a (c) amended	61624
520.1120b (c) amended	61625
520.1192 (b) amended	63270
520.1193 Heading and (a) revised; (d) added	5319
(b) amended	63270
520.1194 (b) amended	63270
520.1195 (b) amended	63270
520.1196 (b) amended	63270
520.1197 (b) amended	63270
520.1242a (c)(2) amended	61625
520.1242b (c) amended	61625
520.1242g (e) amended	55160
(c) amended	61625
520.1263c (b) amended	65020
520.1288 (c)(3) amended	8371
(c) heading revised; (d) added	12085
(a), (d)(1) and (3) revised	14301
520.1289 (c)(3) amended	8371
520.1320 (c) amended	61625
520.1326a (b) amended	61625
520.1326b (b) amended	61625
520.1408 (b) amended	35076
520.1446 Added	28629
520.1451 Added	37713
520.1452 Added	42902
520.1484 (b) and (c)(3) amended	55160
520.1485 (d)(3) amended	60657
520.1616 Added	29012
(d)(1) heading and (iii) amended	61627
520.1660d (a)(8) and (b)(6) added	23356
(b)(3), (e)(1)(ii)(A)(3), (B)(3) and (C)(3) amended	35076
520.1696b (b) amended	55160
520.1720a (b)(1) amended	61625
520.1720b (b) amended	61625
520.1720c (b) amended	61625
520.1805 (b) amended	61625
520.1900 (b) amended	35076
520.2043 (b)(2) revised	37712

21 CFR—Continued

62 FR
Page

Chapter I—Continued

520.2080 Removed	15111
520.2098 Added	34632
(d) heading and (2) corrected	55159
520.2170 (c) amended	63270
520.2220a (b) amended ... 8371, 23357, 35076 (b) corrected	40932
520.2220b (b)(3) amended	61625
520.2220c (c) amended	61625
520.2260b (f)(1) and (h)(1) amend- ed	35076
520.2325a (a)(4) added	37712
520.2345d (a)(1) amended; (c) re- moved	5319
(a)(1), (d)(1)(iii) and (2)(iii) amended	35076
(a)(1) amended	46668
(a)(1), (4), (d)(1)(iii) and (2)(iii) amended	55160
520.2362 (c) amended	61625
520.2380a (c)(2) amended	63271
520.2380b (c) amended	63271
520.2380c (c) amended	63271
520.2380d (b) amended	63271
520.2380f (b) amended	63271
520.2455 (b) amended	35076
520.2456 (b) amended	35076
520.2481 (b) amended	35076
520.2610 (b) amended	61625
520.2611 (b) amended	61625
520.2612 (b) amended	61625
520.2640 (e)(1)(iii), (2)(iii) and (3)(iii) amended	39443
522 Authority citation revised	51518
522.23 (c) amended	35076
522.56 (b) amended	23357
522.82 (b) amended	61625
522.150 (b) amended	61625
522.518 (b) amended	28630
522.540 (a)(2) amended	37713
522.770 (d)(2) revised	44410
(d) heading revised; (d)(1), (2) and (3) redesignated as (d)(1)(i), (ii) and (iii); new (d)(1) heading and (2) added	62242
522.784 (b) amended	61625
522.812 (c) redesignated as (d); new (d)(2) revised	38907
522.863 (b) amended	61625
522.970 (b) revised	22888
522.1002 (b)(2) amended	62242
522.1010 (b) amended	35076
522.1044 (b)(3) amended	35076
(b)(3) and (4) added	45157
522.1150 (b) amended	63271
522.1155 (b) amended	61625
522.1156 Added	66984

List of CFR Sections Affected

21 CFR—Continued

62 FR
Page

Chapter I—Continued

522.1182 (b)(2)(i) amended	35076
522.1183 (e)(1) amended	35076
522.1192 (d)(2)(ii) amended	14634
(b) amended	63271
522.1193 (d)(2) amended	14302
(b) amended	63271
522.1222a (c) amended	22888, 35076
522.1244 (b) amended	61625
522.1410 (b) amended	35076
522.1452 (b) amended	63271
522.1465 Added	5320
522.1503 (b) amended	61625
522.1660 (c)(1)(iii) corrected	13825
(c) redesignated as (d); new	
(d)(1) heading revised; new	
(d)(1)(ii), new (iii) and new	
(2)(ii) amended	27692
522.1662a (g)(3)(i)(c) revised	14303
(a)(2), (g)(2) and (h)(2) amended	
.....	35076
522.1680 (b) amended	35076, 38906
522.1704 (a)(2) amended	61625
522.1720 (b)(2) amended	38906
(b)(1) amended	61625
522.1850 (d) added	45158
522.1885 (b) amended	63271
522.1940 (d)(1)(iii) amended;	
(d)(2)(iv) removed	8372
522.2005 (b) amended	61625
522.2095 (d) revised	10220
522.2220 (a)(2)(iii) added	23128
(c)(2) amended	35076
522.2424 (b) amended	35076
522.2471 (d)(1)(ii) revised;	
(d)(1)(iii) amended	5526
522.2476 (b) amended	29013
522.2477 (a) revised	28629
522.2483 (b) amended	35077
522.2610 (b)(2) corrected; CFR cor-	
rection	41272
(a)(2) amended	61625
522.2615 (c) redesignated as (d);	
new (c) added; new (d)(3) re-	
vised	4164
522.2640a (b)(2) amended	35077
522.2662 (b) amended	35077
522.2680 (b) amended	61625
524 Authority citation revised	51518
524.154 (a)(2) amended	61625
524.155 (a)(1) amended	61626
524.575 (c)(2) revised	48940
524.770 Added	65753
524.814 Added	33997
524.900 (e) amended	55161
(c) amended	61626
524.981a (b) amended	40932

21 CFR—Continued

62 FR
Page

Chapter I—Continued

524.981b (b) amended	40932
524.981c (b) amended	40932
524.1044f (b) revised	10220
524.1193 (d)(2) amended	38908
(b) amended	63271
524.1240 (b) amended	61626
524.1443 (b) revised	55161
(b) amended	61626
524.1484g (b) amended	63271
524.1580b (b) amended	35077
524.1580c (b) amended	35077
524.1742 (b) amended	61626
524.1883 (b) amended	63271
526 Authority citation revised	51518
529 Authority citation revised	51518
529.50 (c)(2) amended	15110
(b) amended	23358
529.1044a (b) amended	611, 35077
529.1044b (b) amended	22889
529.1115 (b) amended	29014
529.2090 (a)(1) amended	8372
530 Authority citation revised	51518
530.41 Revised	27947
556 Authority citation revised	51518
556.52 Revised	40933
556.225 Revised	62243
556.227 Added	33998
556.273 Added	55161
556.735 Corrected	15391
556.738 Revised	12086
556.741 Added	4164
558 Technical correction	23534, 44892
Authority citation revised	51518
558.4 (d) table amended	25477, 60657
558.55 (a) amended	63271
558.58 (d)(1)(iii) table amended	60658,
61627, 66987	
(d)(1) table amended	63271
558.76 (d)(1)(iv) table amended	10304
(d)(1) table amended	12952
558.78 (a) revised; (d)(1) table	
amended	55162
558.95 (b) redesignated as (d);	
(a)(5) and new (b) added;	
(d)(4)(i)(b) amended	8373
(d)(1)(ii)(b), (iii)(b), (iv)(b), (v)(b)	
and (3)(iii)(b) amended	63271
558.120 (c) redesignated as (d); new	
(d)(1)(iii)(b) amended	61011
558.128 (c)(3)(xiii) added	12086
(c)(3)(xiv) added	12952
(c) redesignated as (d); new	
(d)(1)(vi), new (xii) and new	
(xvii) amended	14303
(a)(4) and (d)(1) table amended	27692
(d)(1) table and (2) amended	61628

21 CFR (4-1-99 Edition)

21 CFR—Continued

	62 FR Page
Chapter I—Continued	
558.145 (a)(1) amended	14300
558.155 (a)(1) and (2) amended.....	35077
(a)(1) amended	67725
558.175 (c)(1)(iii)(b) and (iv)(b) amended.....	61626
(c) redesignated as (d); new (d)(1)(iv)(b) amended.....	61912
(d)(1)(iii)(b) amended	62243
558.195 (c)(2) added; (d) table amended.....	23129
(d) table amended.....	61626, 66523, 67273
(a) and (d) table amended.....	67725
558.205 (a) amended	35077
558.235 (a) amended	63271
558.254 (c) amended	55161
(a) amended	61626
558.274 (a)(4), (c)(1)(i) and (ii) amended; eff. 4-7-97.....	14301
Regulation at 62 FR 14301 eff. date corrected to 4-7-97	15751
(a)(7) and (c)(1) table amended	27692
(a)(8), (c)(1)(i) table and (ii) table amended	55160
558.300 (a)(1) and (2) amended.....	63271
558.305 (b) redesignated as (d); new (b) and (c) added; new (d)(3) heading revised.....	9929
558.311 (e)(3)(i) table amended	28630
(e)(1)(ii) table amended.....	61626
558.342 (c)(5)(ii)(C) amended	8373
(c) redesignated as (d); new (c) and (d)(7) added.....	14304
558.355 (b)(7) and (14) amended	14304
(f)(3)(ii)(a), (ix)(a) and (x)(b) table amended	14306
(f)(1)(xv)(b) and (xvi)(b) amended	61912
558.366 (a) and (c) table amend- ed.....	29011
558.485 (a)(4) removed; eff. 4-7- 97.....	14301
Regulation at 62 FR 14301 eff. date corrected to 4-7-97	15751
(a)(11) amended.....	27692
(a)(28) added.....	46444
558.515 (d)(1)(vi)(b) amended.....	61626, 66986
558.525 Removed	15111
558.530 (a) amended	67725
558.550 (b)(3) redesignated as (b)(4); new (b)(3) added; eff. 5- 27-97	14305
(b) redesignated as (d); (d)(1)(xvii) added.....	29014
(b)(1)(vii)(c) amended.....	60781
(b)(1)(ix)(c) amended.....	66985

21 CFR—Continued

	62 FR Page
Chapter I—Continued	
558.553 (b)(3) added	66584
558.555 (b)(4) added	66985
558.600 (c)(4) added	12086
(a) amended	35077
558.615 (a) amended	63271
558.618 (d)(1) corrected	15391
558.625 (b)(53) removed; eff. 4-7- 97.....	14301
(f)(1)(i)(b) amended	14306
(f)(1)(vi)(e) correctly revised.....	15111
Regulation at 62 FR 14301 eff. date corrected to 4-7-97	15751
(b)(10) and (52) amended	27692
558.630 (b)(3) and (10) amended; eff. 4-7-97	14301
Regulation at 62 FR 14301 eff. date corrected to 4-7-97	15751
(b)(3), (8) and (10) amended.....	27692
564 Authority citation revised	51518
570 Technical correction.....	47760
Authority citation revised	51518
570.35 (c)(1)(viii) revised.....	40600
571 Technical correction.....	47760
Authority citation revised	51518
571.1 (c) amended	40600
573 Authority citation revised	51518
573.920 Regulation at 60 FR 53703 confirmed.....	44894
579 Denial of request for hear- ing.....	611
Authority citation revised	51518
582 Authority citation revised	51518
584 Authority citation revised	51518
589 Authority citation revised	30976, 51519
589.2000 Added (OMB number pending)	30976

1998

21 CFR

	63 FR Page
Chapter I	
510 Technical correction.....	408
510.105 (c)(2) revised.....	32980
510.106 Revised	32980
510.600 (c)(1) table and (2) table amended	5254, 7700, 8122, 8347, 8349, 11598, 14612, 24106, 25164, 26981, 29551, 31624, 31931, 36178, 41188, 44381, 44383, 51822, 59216, 65553
514.111 (a)(5) revised; eff. 4-6-98.....	10770
514.117 Added; eff. 4-6-98.....	10770
520 Technical correction.....	408, 46652
520.88g (c)(1)(ii) amended	13121
520.88h (c)(1)(ii) amended	13121

List of CFR Sections Affected

21 CFR—Continued

63 FR
Page

Chapter I—Continued

520.154a (a) amended; (d)(4) added.....	38474
520.390b (b)(1) amended.....	5255
520.452 Added.....	41419
520.580 (b)(1) amended.....	26981
520.645 Added.....	8123
520.870 Added.....	51300
520.905a (a) removed; (b), (c) and (d)(3) redesignated as (a), (b) and (d)(4); (d)(2) heading revised; (d)(2)(i)(A), (B), (ii) introductory text, (A) and (B) redesignated as (d)(2)(ii), (iii), (3)(i), (ii) and (iii); new (c) and new (d)(3) heading added.....	63983
520.905c (d)(1)(iii) added.....	31624
520.1195 (b) amended.....	38474
520.1242a (a) and (d) removed; (b), (c), (e) and (f) redesignated as (a) through (d); new (a), (b), (c), (d)(1)(i), (iii) and (2)(i) revised; new (d)(2)(iii) added.....	59713
520.1288 Revised.....	52968
520.1445 (a) and (c) heading revised; (d) added.....	29352
(c) removed; (d) revised.....	41189
520.1446 (c) redesignated as (d); new (d) revised.....	41190
520.1448b Removed.....	6862
520.1484 (b) revised; (c)(3) amended.....	17329
520.1485 (d)(3) amended.....	45944
520.1660c Heading, (a) and (b) revised; (c) removed; (d) and (e) redesignated as (c) and (d); new (d)(3) amended.....	70334
520.1660d (a)(3) revised.....	44384
520.1720a (b)(3) amended.....	36178
520.1806 (b) amended.....	8348
520.1880 (a) amended.....	148
520.2043 (a)(2) revised.....	39727
(b)(2) amended.....	45945
520.2098 (b) amended.....	29551
520.2158a (b) amended.....	51821
520.2220a (a) revised; (b) removed; (c), (d) and (e) redesignated as (b), (c) and (d).....	59714
520.2345a (b)(4) removed.....	5255
522 Technical correction.....	46652
522.84 Added.....	44382
522.90b (b) revised.....	41420
522.246 (b) revised.....	66432
522.314 (d)(2) added.....	53578
522.468 Added.....	13123
522.533 Added.....	44383
522.535 Added.....	13122

21 CFR—Continued

63 FR
Page

Chapter I—Continued

522.770 (d)(1)(ii) revised.....	68183
522.778 Added.....	8349
522.812 (a) revised; (d)(1), (2) and (3) redesignated as (d)(1)(i), (ii) and (iii); (c), new (d)(1) heading and new (2) added.....	49003
522.955 (d)(1)(iii) amended.....	26981
(d)(1)(i) revised; (d)(1)(iii) amended.....	41191
522.970 (c) redesignated as (d); new (c) added; (b) and new (d) revised.....	38749
522.1044 (b)(3) amended.....	59714
(b)(4) revised.....	68182
522.1081 (a)(2)(ii) amended.....	51822
522.1086 (b) amended; (c) redesignated as (d).....	29352
522.1125 Added.....	11598
522.1145 (a)(2) revised; (f) added.....	59216
522.1182 (b)(2)(i) amended.....	53578
522.1183 (b) removed; (e)(1) amended.....	44384
522.1192 (a)(2) heading revised; (d)(6) added.....	7702
522.1222a (c) amended.....	51822, 65553
522.1289 Added.....	29552
522.1462 (b) amended.....	7701
522.1642 (b) amended.....	7701
522.1660 (c) added; (d)(1) heading revised; (d)(1)(iii) amended.....	52157
522.1662a (h)(2) amended.....	6644
522.1940 (d)(3) added.....	45945
522.2005 (b) amended.....	24420
522.2120 Heading revised.....	24107
522.2121 Added.....	24107
522.2471 (d)(1)(iii) amended.....	7701
(d)(1)(iii) corrected.....	14818
522.2478 (c)(2) added.....	63789
524.770 (d)(2) revised.....	68183
524.814 (d)(2) revised.....	59715
524.1044g (b) amended.....	31932
524.1193 (b) and (d)(2) revised.....	44385
524.1451 Added.....	14036
524.1742 (b) amended.....	5255
524.2101 (c) amended.....	26981
526.365 Redesignated from 529.365.....	8349
529.365 Redesignated to 526.365.....	8349
529.469 Added.....	25164
529.1030 (c) redesignated as (d); (b)(1), (2), new (d) introductory text, (1) and (2)(i) revised; (d)(2)(iv) and (v) added.....	38304
529.1044a (b) amended.....	6644
529.1186 (b) revised.....	8122, 24106
529.2503 (b) amended.....	7702

21 CFR (4-1-99 Edition)

21 CFR—Continued

63 FR
Page

Chapter I—Continued	
556.100 Revised	13337
556.113 Revised	53579
556.150 Revised	52158
(b) revised	57246
556.167 Added	13123
556.225 Revised	68184
556.227 Revised	59715
556.228 Undesignated text des- ignated as (a); new introduc- tory text, (a) heading and (b) added	49003
556.283 Revised	41191
556.286 Added	38750
556.344 Revised	7702, 54352
556.426 Added	14036
556.500 Revised	52158, 57246
556.600 Revised	24107
(c) corrected	38304
556.720 Revised	52158, 57246
558 Technical correction	408, 46389
Determination	15292
558.4 (d) table amended	11599
558.58 (d)(1)(iii) table amended	27845
558.76 (d)(1)(iv) table amended	10304
(d)(1) table amended	40825, 44385
(d)(3)(xv) added	38475
(d)(3)(xv) added	38750
(d)(3)(xvi) added	39028
(d)(3)(xvii) correctly designated	48576
(d)(3)(xviii) added	51824
558.78 Regulation at 62 FR 55162 eff. date corrected to 10-23- 97	9310
(a)(3) corrected	14613
(d)(3) revised	19185
(a)(2), (d)(1) table and (2)(ii) amended	27845
558.95 (d)(5)(iii) added	13124
(a) revised	14356
(d)(4)(iv) added	18836
(d)(1)(xi)(b) and (xii)(b) amended	27845
(d)(4)(i)(b), (ii)(b), (iii) introduc- tory text, (d), (iv) introduc- tory text and (c) amended	44386
558.115 (d)(1)(ii) and (2)(ii) re- vised	59216
558.120 (d)(1)(iii)(b) amended	27845
558.128 (d)(3)(xiv) redesignated as (d)(3)(iii)	10304
(a)(1) and (d)(1) table amended	27845
(d)(1) table amended	57246, 65554
558.145 (a)(1) and (2) amended	27845
558.155 (a)(2) amended	2307
(a)(2) corrected	11599

21 CFR—Continued

63 FR
Page

Chapter I—Continued	
558.175 (d)(1)(iii)(b) and (iv)(b) amended	17948, 27845
558.195 (d) table amended	17948, 27845, 38475, 51824
558.274 (a)(6) removed; (c)(1)(i) table amended	5255
558.300 (c) redesignated as (d); new (d) introductory text added; new (d)(1) revised	54353
558.305 (a) amended	27845
558.311 (e)(1)(ii) table amended	17948
(b)(2) through (7), (e)(1)(v) table, (2)(v) and (3)(v) amended	27845
558.325 (c)(3)(xii) revised	57248
558.340 (a) amended	27845
558.342 (d)(3)(ii) and (6)(ii) amend- ed	27845
(d)(8) added	41191
(c) and (d)(3)(ii) amended	66018
558.355 (c) removed	14035
(a) amended	24420
(b)(8), (9), (f)(1)(iv)(b), (v)(b), (xiv)(b), (xv)(b) and (xvi)(b) amended	27845
(f)(2)(iii) added	52969
(b)(11) and (f)(1)(xiv)(b) amended	70335
558.363 (a) revised; (c) redesign- nated as (d); new (d)(1)(vii) added	13124
(a)(6), (d)(1)(viii) and (ix) added	51823
(d)(2) revised	57248
558.366 (c) table amended	13124, 57248
558.369 (c) removed; (d) introduc- tory text revised; (d)(1), (2) and (3) redesignated as (d)(1)(i), (ii) and (iii); (d)(1) heading and (2) added	39028
558.450 (d)(3)(iii) added	41192
558.460 (b) and (c) redesignated as (c) and (d); new (b) added; (d)(1) table revised	36179
558.485 (a)(21) and (25) removed	5255
558.515 (d)(1)(iv)(b) amended	17948
(a) and (d)(1)(vi)(b) amended	27845
558.530 (c) removed; (d)(5)(x) re- vised	38475
558.550 (a)(1) amended	6644
(a)(1), (d)(1)(vii)(c), (ix)(c), (xv)(c) and (xvi)(c) amended	27846
(a) revised; (d)(1)(xv)(c) amended	65554
(a)(3) revised	66432
558.575 (a)(1) and (2) amended	27846
558.582 (a) amended	27846

List of CFR Sections Affected

21 CFR—Continued

63 FR
Page

Chapter I—Continued	
558.600 (c)(4)(ii) amended.....	27846, 52969
558.625 (b)(84) removed.....	5255
558.680 (c)(1) table amended.....	38750
573.280 Revised	8573
573.460 (b)(1), (2)(ii), (iii) and (3)(iv) revised; (b)(2)(iv) added.....	53580

1999

(Regulations published January 1, 1999
through April 1, 1999)

21 CFR

64 FR
Page

Chapter I	
510.45 Removed; eff. 5–20–99	403
510.110 (e) and (f) amended; eff. 5– 20–99	403
510.600 (c)(1) table and (2) table amended.....	15683
514.10 Removed; eff. 5–20–99	403
520.45a (a)(4)(i)(I) designation and (2) removed; (b)(1) revised	1504
520.154a (a) and (d)(2) amended	13068
520.260 (b) amended	15684
520.455 Added	1762
520.534 Added	10103
520.1204 (a) amended; eff. 5–20– 99.....	403
520.1263a (a) revised; eff. 5–20– 99.....	403
520.1263b (a) amended; eff. 5–20– 99.....	403
520.1263c (d)(1)(i)(C) amended	13341
(a) amended; (b) revised	13508
520.1660d (a)(9) and (b)(7) added; (b)(2) revised.....	10389
(a)(7) revised; (c) removed; (d) and (e) redesignated as (c) and (d).....	13341
(a)(8), (d)(1)(ii)(A)(3), (B)(3) and (C)(3) amended	13678
520.2098 (d)(2) and (d)(3) redesi- gnated as (d)(1)(i) and (ii); new (d)(2) added.....	2122
520.2158b (d) correctly added; CFR correction.....	4540

21 CFR—Continued

64 FR
Page

Chapter I—Continued	
520.2184 (b) amended.....	15684
520.2220b (a) removed; (b), (d) and (e) redesignated as (a), (b) and (d); new (a) and new (d)(2) re- vised.....	15684
522.690 (b) amended	15685
522.723 (c) amended	15684
522.770 (d)(1)(ii) amended	13509
522.800 (b) amended	15684
522.955 (d)(1)(i) revised; (d)(1)(ii) redesignated as (d)(1)(i)(B); (d)(1)(ii)(A) and (B) added	5596
(d)(1)(i)(B) revised	9435
522.1204 (a) revised; eff. 5–20–99.....	403
522.1484 (a) amended; eff. 5–20– 99.....	403
522.2005 (b) revised; (c)(2) added.....	13510
524.1200a (a)(1) and (2) designation removed; eff. 5–20–99.....	404
524.1200b (a) revised; eff. 5–20– 99.....	404
524.1204 (a)(1) removed; (a)(2)(i), (ii), (iii) and (3) redesignated as (a)(1)(i), (ii), (iii) and (2); eff. 5– 20–99	404
529.360 (a) amended; eff. 5–20– 99.....	404
556.34 Revised	1504
556.170 Revised	10103
556.347 Revised	13069
556.360 Revised	13342
556.420 Revised	5159
556.735 Revised	13679
558.15 (g)(1) table and (2) table amended	992
558.76 (d)(3)(xix) added.....	4966
(d)(3)(xx) added	13342
558.140 (a) amended	15684
558.300 (d)(1)(ii), (2)(ii), (3)(ii), (4)(ii) and (5)(ii) amended; (d)(6) added.....	13342
558.311 (b)(4) revised.....	13069
558.325 (c) redesignated as (d); (d)(2)(ii)(b), (iii)(b) and (iv)(b) amended.....	13342

21 CFR (4-1-99 Edition)

21 CFR—Continued

64 FR
Page

Chapter I—Continued

558.355	(f)(3)(ii)(b) amended	2855
	(d)(7)(ii), (f)(3)(iii)(a), (b), (vi)(a),	
	(b), (vii)(a) and (b) revised;	
	(f)(3)(xi) added	5159
	(b)(11) amended; (f)(1)(xxvi) and	
	(xxvii) added	5597
	(f)(2)(iv) added	13069

21 CFR—Continued

64 FR
Page

Chapter I—Continued

558.363	(d)(2)(ii) added	4966
558.366	(c) table amended	4966
558.485	(a)(17) removed	15684
558.618	(d)(3) amended	13679
558.635	(b)(2) amended	15684
564	Removed	4294
	Technical correction	6801

